



කෘෂිකර්ම දෙපාර්තමේන්තුව
விவசாயத் திணைக்களம்
Department of Agriculture



(<https://doa.gov.lk/>)

(/ (h

si tt

[<< Crop Page \(/hordi-crops/\)](#)

n ps

h ://



- ta

Brinjal

cr mi

Solanum melongena

Brinjal is considered as a highly marketable nutritious crop

br rdi

in -

Released Varieties

jal cr

/) op

-

bri

nj

al/

)

- Amanda F1
- Anjalee-F1
- HORDI Lenairi 1
- Thinnaweli purple
- Padagoda
- SM 164

(/Hordi-Variety2-Brinjal/)

Climatic requirements/ Areas suitable for cultivation



Crop can be grown up to 1300 m above sea level.

Soil

Well drained soil is suitable, pH range from 5.5 – 5.8

Seed requirement

350g/ha

Nursery Management

3m x 1m beds in better sunlight conditions are suitable. Beds should be sterilized. Add 1:1 ratio of surface soil and cow dung. Seeds should be placed in rows in 6 inches spacing.

Land preparation

Zero tillery or ploughed soil should be prepared and 30m x 30m x 30m holes are suitable

Planting

25-30 days old plants can be established in prepared holes as one plant per hole

Spacing

90cm x 60cm

Fertilizer

Time of application	Urea kg/ha	TSP kg/ha	MOP kg/ha
Basal	75	325	85
After 4 weeks	75	–	–
After 8 weeks	75	–	85



Water supply

Better to supply water for better growth

Weed Control

2,4,7,9,12 days after planting , better to remove weeds

Disease Management

- **Fungal diseases**

- ▲ **1. Damping-off**

Causal organism: *Pythium* spp., *Phytophthora* spp., *Fusarium* spp., *Rhizoctonia* spp. and *Sclerotium rolfsii* **Symptoms:**

- Sudden collapsing seedlings at the base of the stem
- Root rot and seedling death
- Brownish black discoloration at the base due to *Rhizoctonia* spp



Management

- Prepare nursery beds in well drained virgin soil
- Nursery sterilization – burning, solarization or with chemicals
- Remove affected plants with the adjacent soil
- Seed treatment (should be done just before seeding)

^

-Thiram 80% WP – 5g/ kg -Thiophanate-methyl 50% WP + Thiram 30% WP – 4g/ kg -
Captan 50% WP – 6g/ kg.

- Soil treatment – Drench with a recommended fungicide around the base of the plant as spot application

– Thiophanate-methyl 50% WP + Thiram 30% WP – 50g/50l / 10m² – Thiram 80% WP
– 70g/50l / 10m² – Captan 50% WP – 60g/50l / 10m²

▲ [2. Collar rot](#)

Causal organism: *Sclerotium rolfsii* **Symptoms:**

- Yellowing of leaves
- Decaying near collar region
- Mycelia mat with white / brown Sclerotia on collar region



Management

- Destroy affected plants with the adjacent soil
- Improve drainage
- Deeply plough the land and expose to affected soil to bright sunlight
- Avoid root damage at transplanting
- Soil treatment – Drench with a recommended fungicide around the base of the plant as spot application

– Thiophanate-methyl 50% WP + Thiram 30% WP – 50g/50l / 10m² – Thiram 80% WP
– 70g/50l / 10m² – Captan 50% WP – 60g/50l / 10m²

▲ [3. Fusarium wilt](#)

Causal organism: *Fusarium oxysporum* f.sp *lycopersici*

Symptoms:



- Yellowing of leaves and wilt
- Vascular discoloration of stem
- Root rot and dying of the plant

Management

- Destroy affected plants with the adjacent soil
 - Improve drainage
 - Deeply plough the land and expose to affected soil to bright sunlight
 - Avoid root damage at transplanting
 - Soil treatment – Drench with a recommended fungicide around the base of the plant as spot application
- Thiophanate-methyl 50% WP + Thiram 30% WP – 50g/50l / 10m²
- Thiram 80% WP – 70g/50l / 10m²
- Captan 50% WP – 60g/50l / 10m²

▲ 4. Powdery mildew

Causal organism: *Podosphaera xanthii*



Symptoms:

- Circular yellow patches or spots appear on the lower surface of the leaves.



- White powdery residue on the upper leaf surface.

Management:

- Use recommended fungicides at initial stage of disease development (Refer: Pesticide recommendation DOA, 2019)
 - Apply sulfur 80% WG 50 g/10 l of water
 - Chlorothalonil 500g/l SC- 30 ml/10 l of water
 - Carbendazim 50% WP- 7 g/10 l of water

▲ 5. Anthracnose

Causal organism: *Colletotrichum gloeosporioides*

Symptoms:

- Sunken circular lesions with tan to orange to black concentric rings in their center
- Fruiting bodies with pink masses of conidia may ooze from lesions



Management

- Use of healthy seeds
- Avoid excess overhead irrigation
- Remove and destroy infected fruits
- Seed treatment (should be done just before seeding)

– Thiram 80% WP – 5g/ kg -Thiophanate-methyl 50% WP + Thiram 30% WP – 4g/ kg - Captan 50% WP – 6g/ kg.

- Foliar application of fungicides start at flowering stage in rainy season

– Fluzinam 500g/l SC – 10 ml/10l of water – Metiram 55% + Pyraclostrobin 5% WP – 20 g/10l of water – Cabendazim 50% WG – 7 g/10l of water – Thiophanate-methyl 70% WP – 6g/10l of water – Chlorothalonil 500 SC – 30ml/10l of water

▲ 6. Phomopsis blight

Causal organism: *Phomopsis vexans* **Symptoms**

- Small canker gray spots with black margins appear on stem and petioles
- Center of the spot becomes gray and black due to pycnidia development
- Soft and watery fruit spots become black and mummified



Management

- Remove and destruction of infected fruits
- Use recommended fungicides at initial stage of disease development (Refer: Pesticide recommendation DOA, 2019)

-Chlorothalonil 500g/l SC- 30 ml/10 l of water -Carbendazim 50% WP- 7 g/10 l of water – Thiophanate-methyl 70% WP of 6g/10l of water

• **Bacterial diseases**

▲ 7. Bacterial wilt

Causal organism: *Ralstonia solanacearum* **Symptoms:**

- Slight wilting of single branch/ branches



- Sudden and permanent wilting of entire plant
- Discoloration of the vascular tissues
- Slimy viscous bacterial ooze comes out from the cut end of affected parts when immersed in clear



Management

- Remove and destroy affected plants with soil
- Destroy crop debris after harvesting
- Crop rotation with non-susceptible crop (cruciferous vegetables and okra) help in reducing the disease incidence
- Use of resistant varieties
- Disinfect all farm implements/tools with bleach after they have been
- No chemical controlling method

• Phytoplasma disease

▲ 8. Little leaf disease (witches' broom)

Causal organism: Phytoplasma The disease is transmitted by a leaf hopper(*Hishimonus phycitii*) **Symptoms**

- leaves are narrow, soft, smooth and yellow
- Internodes of the stem become shorten



- Auxiliary buds get enlarged but their petioles and leaves remain shortened and give the plant a bushy appearance.
- flowers remain green/ no flowering
- Fruiting is rare



Management • Remove and burn infected plants. • Control leaf hopper with recommended insecticides to minimize disease spread.

Harvesting

Harvesting can be started 10-12 weeks after transplanting. Fruit can be harvested 7 days intervals up to 10-12 picks.

Yield

Potential yield depends on the variety. OPV gives 18-20 t/ha and hybrids give 30- 40 t/ha



Leaflet - Diseases on Brinjal (https://drive.google.com/file/d/1BTsdmAyK05cCnGbgz7Elukml08DwuSeH/view?usp=drive_link)



Copyright © 2026 Department of Agriculture Sri Lanka (<https://doa.gov.lk>). All rights reserved.

