



කෘෂිකර්ම දෙපාර්තමේන්තුව
விவசாயத் திணைக்களம்
Department of Agriculture



(<https://doa.gov.lk/>)

(/ (h

si tt

[<< Crop Page \(/hordi-crops/\)](#)

n ps

h ://



r ta
Bean

di தமிழ்

Phaseolus vulgaris L.

Bean belongs to the family Fabaceae and Genus Phaseolus. It is widely cultivated in Badulla, Nuwara Eliya, Matale and Kandy Districts. Areas with high temperature (>32 °C) and continuous high rainfall is unsuitable for bean cultivation. Temperature should be less than 30°C at the time of flowering because high temperature adversely effect on flowering and pod setting. Although bean is mainly cultivated in Up Country Intermediate Zone, it can also be cultivated in Up Country Wet Zone.

Bean varieties can be categorized into two types based on their growth habit as vine types or “pole bean” and bush types or “Bush bean”. Vine types are the most popular among farmers and consumers as well. Pole bean varieties grow continuously and need supports (stakes) to grow upright. Pods are bigger in size and light in color. Bush beans doesn’t produce much yield and pods are dark green. Its normally cultivate as an intermediate crop between two main cropping seasons

Released Varieties



Vine types (Pole Bean)

- Balangoda Nil
 - Bandarawela Green
 - Gannoruwa BIL
 - Kekulu Bonchi
 - Gannoruwa Green
-
- Kappetipola Nil
 - HORDI Green
 - Kentucky Wonder Green- KWG
 - Lanka Butter
 - HORDI Bean-03(HORDI Pokuru)

Bush types (Bush Bean)

- Cherokee Wax
- Sanjaya
- Wade
- Top Crop

Variety Details >>
(/Hordi-Variety-Bean/)

Climatic requirements/ Areas suitable for cultivation

Up Country Intermediate Zone and Up-Country Wet Zone (Badulla, Nuwara Eliya, Matale and Kandy Districts) are the most suitable areas for cultivation

Soil

Red Yellow Podzolic soil with good drainage and loam or sandy loam texture are ideal for cultivation. Soil with lots of clay tends to be poorly drained, which can cause root rot and blossom drop. Optimum soil pH for better growth should be around 5.5-6.5.



Seed requirement

Vine type – 50kg/ha

Bush type- 75kg/ha

Nursery management/time of planting

No nursery needed, direct seeding is the commonly practiced.

Badulla District –

- Maha season – November and December
- Yala Season – March and April

Land preparation

Plough soils to make fine tilth and prepare beds which can accommodate three rows. Mark planting holes in recommended spacing.

Planting

Soak seeds for about 24 hrs. before planting. Plant about 3-4 seeds per planting hole fairly away from each other. Keep beds well-watered. Two weeks after germination, keep two healthy seedling per planting hole and remove the other seedlings.

Spacing

Vine types – 50 cm between rows and 40 cm between plants

Bush types – 50 cm between rows and 15 cm between plants

Fertilizer

Add 10 t of well decomposed organic manure five or six days before seeding and recommended quantity of synthetic fertilizers (basal dressing) one to two days before seeding.

Pole bean



Time of application	Urea (kg/Ha)	Triple supper phosphate (kg/Ha)	Muriate of potash (kg/Ha)
Basal dressing	110	270*	75*
Top dressing (three weeks after planting)	110	–	75
Bush bean			
Basal dressing	85	165*	65*
Top dressing (three weeks after planting)	85	–	65

****it is better to apply after a soil test***

Water supply

Maintaining soil moisture at the optimum level during flowering and fruiting stages will assure higher yield. Therefore, supplementary irrigation has to be provided if the rainfall is insufficient.

Weed Control

Keep the cultivation weed free to reduce pests, diseases and competition of plants and weeds for nutrient and space.

Pest Management

Several insect pests can damage plants and pods during the growth period. Use seed treatments or spray application of recommended chemicals to control those pests.

Common name	Scientific name
Bean fly	<i>Ophiomyia phaseoli</i>
Mites	<i>Tetranychus urticae</i>
Thrips	<i>Thrips</i> spp



Whiteflies	<i>Bemisia tabaci</i>
Aphids	<i>Aphis</i> spp
Pod borers	<i>Maruka vitrata</i>

Disease Management

▲ [Fungal Diseases](#)

Four fungal diseases are the main problems in bean cultivation. These diseases are very serious during rainy season (Maha) and difficult to manage without fungicides. Use fungicides recommended by the Department of Agriculture to control these diseases.

Disease	Causal agent	Affected part
Anthrachnose	<i>Coletotrichum lindemuthianum</i>	Pods, leaves and stems
Rust	<i>Uromyces appendiculatus</i>	Leaves and pods
Angular leaf spot	<i>Phaeoisariopsis griseola</i>	Leaves and pods
Root rot	<i>Fusarium oxysporum</i>	Roots and stem base
	<i>Fusarium solani</i>	

▲ [Virus Diseases](#)

- Four virus diseases are common in bean cultivation.
- Infected leaves turn into mosaic or yellow/golden color.
- Remove and destroy infected plants as soon as possible to control the disease spread.
- Use recommended insecticides against the virus vectors

Common name	Vector
Bean Common Mosaic Virus (BCMV)	Aphids
Bean Yellow Mosaic Virus (BYMV)	Aphids
Bean Golden Mosaic Virus (BGMV)	Whiteflies

Horsegram Yellow Mosaic Virus (HYMV)

Thrips

Harvesting

- Pick pods before they reach to full maturity (tender pods)
- Vine types can be harvested from 60 days after planting
- Bush types can be harvested from 45 days after planting
- Gannoruwa BIL can be harvested in every 4 days and the rest of varieties in every 7 days

Yield

- Bush types – 5-10 t/ha
- Gannoruwa green – 18-20 t/ha
- Gannoruwa BIL – 20 t/ha
- Kekulu Bonchi – 25 t/ha
- Other vine types 12-16 t/ha

Post-harvest handling

Keep harvested pods in a shade

Pack carefully in boxes or bags with adequate ventilation

Specific crop management practices

- Do staking for vine type varieties after 2-3 weeks of planting
- Use 6-7 ft high sticks for staking.
- Trellises can be used to get higher yield from variety Keppetipola Nil



Leaflet - Bean
Varieties

(https://drive.google.com/file/d/1_YyBdgXMZtMndrAWrcmqOd2ITrnwtsJj/view?usp=drive_link)





Leaflet - Disease Control on Bean (<https://drive.google.com/file/d/18UfsdqmHxFA7qXgSVuC625V0hw->

Cultivation

[Zh33H/view?usp=drive_link](https://drive.google.com/file/d/18UfsdqmHxFA7qXgSVuC625V0hw-Zh33H/view?usp=drive_link))

Copyright © 2026 Department of Agriculture Sri Lanka (<https://doa.gov.lk>). All rights reserved.

