

Homework 3

NAME/S: _____ SCORE: _____

Subject: Quantum Mechanics I

Deadline: Friday 29 July 2022 (until 7pm)

Credits: 20 points

Number of problems: 5

Type of evaluation: Formative Evaluation

- This homework consists of problems related to units 1 and 2 of the course.
- You may submit this assignment either individually or in pairs. Submitted assignments should have maximum two authors.
- Unless stated otherwise, write your answers in SI units, and consider all bolded quantities as vector quantities. Please highlight the answers.

1. (4 points) Quantum experiments: photoelectric effect

In a photoelectric effect experiment, a piece of sodium (Na) metal is illuminated by light beams with two different wavelengths $\lambda_1 = 300 \text{ nm}$ and $\lambda_2 = 600 \text{ nm}$. If a piece of sodium (Na) metal has a cutoff wavelength of $\lambda_{\text{cutoff}} = 504 \text{ nm}$, calculate:

- (a) The work function, ϕ , for the sodium metal.
- (b) The maximum kinetic energy and speed of the electrons ejected from the metal piece for each beam.
- (c) The de Broglie wavelength of the ejected electrons.
- (d) Use programming tools to make two plots: one of K_{max} versus λ , and another one of K_{max} versus ν including the data points for both light beams.

Recall that the Planck constant is $h = 6.626 \times 10^{-34} \text{ J s}$, the speed of light is $c = 3 \times 10^8 \text{ m s}^{-1}$, and the electron mass is $m_e = 9.11 \times 10^{-31} \text{ kg}$.

2. (4 points) Infinite square well potential and expectation values

In class we solved the Schrödinger equation for an infinite square well potential of width a . Such potential allows for bound solutions only as the particle cannot escape from the well. The solutions we found had the following functional form:

$$\psi_n(x) = \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi}{a}x\right) \quad (1)$$

For the n -th state given by the function above, calculate:

- (a) The expectation values associated with the position x : $\langle x \rangle$, $\langle x^2 \rangle$.
- (b) The expectation values associated with the momentum p : $\langle p \rangle$, $\langle p^2 \rangle$.
- (c) The dispersions σ_x and σ_p , and their product $\sigma_x \sigma_p$.
- (d) Use programming tools to make a plot of $(\sigma_x \sigma_p)$ vs. n . Is the uncertainty principle satisfied? Which of the $\psi_n(x)$ states comes closest to the uncertainty limit?

3. **(5 points) Free particles: Gaussian wave packets**

We studied free particles in class and showed that they are represented by wave packets. Consider the case of a free particle whose initial wave function is given by:

$$\Psi(x, 0) = \alpha e^{-\beta x^2}, \quad (2)$$

where α and β are real and positive constants.

- (a) Find α by normalising the initial wave function, $\Psi(x, 0)$.
- (b) Find $\Psi(x, t)$. Hint: compute $\phi(k)$ via Fourier analysis first, and then plug it into the wave packet function.
- (c) Find $|\Psi(x, t)|^2$. Then, plug some fiducial numbers, and sketch $|\Psi(x, t)|^2$ versus x for $t = 0$ and two later times. Qualitatively, what happens to $|\Psi(x, t)|^2$ as time progresses?
- (d) Calculate $\langle x \rangle$, $\langle x^2 \rangle$, $\langle p \rangle$, $\langle p^2 \rangle$, σ_x , and σ_p .
- (e) Does the uncertainty principle hold? At what time t does the system come closest to the uncertainty limit?

4. **(5 points) Square potential barrier: transmission and reflection coefficients**

Let us consider a time-independent square potential barrier, $V(x)$, given by the following piecewise function:

$$V(x) = \begin{cases} 0, & x < 0 \\ V_0, & 0 \leq x \leq L \\ 0, & x > L, \end{cases}$$

- (a) Sketch $V(x)$, labelling the three regions of interest as I, II, and III.
- (b) Find the stationary states that describe particles arriving from $x = -\infty$ with energy $E > V_0$, and sketch the solutions using programming tools.
- (c) Analyse the boundary conditions to compute the transmission and reflection coefficients. Sketch these coefficients versus the barrier width, L , and briefly discuss the results.
- (d) Find the stationary states that describe particles arriving from $x = -\infty$ with energy $E < V_0$, and sketch the solutions using programming tools.
- (e) Compute the transmission coefficient, and discuss how this quantum result differs with respect to classical expectations.

5. **(2 points) The Schrödinger equation and potential energy**

A particle of mass m with null energy is described by the following time-independent wave function:

$$\Psi(x) = C x e^{-\frac{x^2}{L^2}}$$

where C and L are constants.

- (a) Normalise $\Psi(x)$. Then, plug some fiducial numbers and sketch $\Psi(x)$ versus x .
- (b) Calculate the potential energy $V(x)$ of the particle, and sketch $V(x)$ versus x .