

# Homework 1

NAME/S: \_\_\_\_\_ SCORE: \_\_\_\_\_

Subject: Quantum Mechanics I

Deadline: Wednesday 1 June 2022 (until 4pm)

Credits: 20 points

Number of problems: 5

Type of evaluation: Formative Evaluation

- This homework consists of problems related to the preliminary concepts reviewed in class about quantum mechanics, plus some problems on classical mechanics.

- You may submit this assignment either individually or in pairs. Submitted assignments should have maximum two authors.

- Unless stated otherwise, write your answers in SI units, and consider all bolded quantities as vector quantities. Please highlight the answers.

## 1. (4 points) Classical harmonic oscillator

The frictionless motion of a block with a mass of 3.5 kg, attached to a spring of constant,  $k$ , is described by the following one-dimensional velocity:

$$v_x = -2 \sin\left(\frac{\pi}{6}t + \frac{5\pi}{2}\right) \text{ m s}^{-1}$$


Note that this is generally called a “horizontal spring-mass system”.

- Make a sketch of the system.
- Find the remaining equations of motion, i.e., displacement  $x$ , and acceleration  $a_x$ .
- What is the amplitude and what is the angular frequency of the oscillation?
- What is the period and what is the frequency of the oscillation?
- Use your favourite programming tool to make plots of  $x$  vs.  $t$ ,  $v_x$  vs.  $t$ , and  $a_x$  vs.  $t$ . Please do not make these plots by hand, use programming tools!
- What are the maximum values of velocity and acceleration of the system?
- What are the initial position, velocity, and acceleration?
- Calculate the spring constant,  $k$ .

NAME/S: \_\_\_\_\_

2. (4 points) Classical collision

Consider the problem of a particle A of mass  $m$ , moving at an initial velocity  $\mathbf{v}_0$ , that collides with another particle B of mass  $2m$ , initially at rest. After the collision, the particles follow the trajectories shown in the figure below. Find the value of the angle  $\theta$ .



NAME/S: \_\_\_\_\_

3. (4 points) **Group velocity of a waveguide and water waves**

Waveguides are structures that guide waves (e.g. sound or electromagnetic waves) with minimal energy loss as they restrict their transmission to a single direction. The wavelength in a waveguide is considered as a wavelength in the direction of wave propagation and is defined as follows:

$$\lambda_{\text{wg}} = \frac{\lambda_0}{\sqrt{1 - \left(\frac{\lambda_0}{\lambda_c}\right)^2}},$$

where  $\lambda_0$  is a wavelength in free space at a given frequency,  $\nu_0$ , and  $\lambda_c$  is the cutoff wavelength at frequency  $\nu_c$ . The latter is related to the dimensions of the waveguide.

- (a) Write an expression for  $\lambda_{\text{wg}}$  in terms of the frequencies  $\nu_0$  and  $\nu_c$ .
- (b) Write an expression for the group velocity,  $v_g$ , of a waveguide in terms of the light speed,  $c$ , and the phase velocity,  $v_p$ .
- (c) Find the group velocity of water waves in deep water, which have a frequency given by  $\nu = \sqrt{\frac{g}{2\pi\lambda}}$ , where  $g$  is the acceleration by gravity.
- (d) Find the group velocity of water waves in shallow water, which have a frequency given by  $\nu = \sqrt{\frac{2\pi T}{\rho\lambda^3}}$ , where  $T$  is the surface tension and  $\rho$  is the density.
- (e) Compare the results found in (c) and (d), and briefly explain why they are different.

NAME/S: \_\_\_\_\_

4. (4 points) **Compton scattering**

Suppose we have an experiment in which monochromatic light is scattered by an electron.

- (a) Find the shift in the wavelength of the light when the scattering angle is  $90^\circ$ .
- (b) Make a sketch of the experiment.
- (c) If the incident light has a  $\lambda = 500 \text{ nm}$  (i.e. photons are in the visible region), what is the fractional increase in the wavelength,  $\frac{\Delta\lambda}{\lambda}$ ?
- (d) If the incident light has a  $\lambda = 0.1 \text{ nm}$  (i.e. photons are in the X-ray region), what is the fractional increase in the wavelength,  $\frac{\Delta\lambda}{\lambda}$ ?
- (e) Why were X-rays used by Compton in his experiments?

NAME/S: \_\_\_\_\_

5. (4 points) **Compton and de Broglie wavelengths**

- (a) Calculate the Compton wavelength of an electron.
- (b) What can happen if you shine that electron with a photon with that wavelength? Why?
- (c) Calculate the de Broglie wavelength of a (macro)particle of diameter  $1\text{ }\mu\text{m}$  with a mass of  $m = 10^{-15}\text{ kg}$  that is moving at a speed of  $1\text{ mm s}^{-1}$ .
- (d) Based on the result above, are the wave properties of matter relevant in the macroscopic world? Why?