

Discrete Standing Wave Spacetime: A Model of Emergent Geometry and Black Surfaces

Wayne Brassem

August 26, 2025

Abstract

In this speculative yet mathematically grounded framework, we propose a radical reconceptualization of gravity, time, and mass through the lens of a standing wave structure of spacetime. Here, spacetime is not merely a passive backdrop but an active, oscillatory medium: a standing wave whose nodes and antinodes define the physical behavior of energy and matter.

Gravity emerges not as an attractive force but as the universal tendency for matter to flow toward regions of slower temporal progression—a gradient in the ticking of time itself. We revisit the Higgs mechanism, suggesting that matter’s interaction with the discrete nodal structure generates impedance absent for massless particles, naturally giving rise to inertia and mass as geometric consequences of perturbing the spacetime fabric.

This framework preserves the predictions of General Relativity and Quantum Mechanics while offering fresh explanations for black hole horizons (reframed as ‘Black Surfaces’), temporal evolution of the speed of light, early structure formation, and Hawking radiation. Temporal pressure gradients, variable $c(V)$, and the standing wave architecture together provide a coherent narrative that may unify apparent paradoxes in modern physics.

Finally, by linking spacetime’s deep oscillatory structure to observable phenomena—from CMB anisotropies to gravitational echoes—the theory invites reconsideration of cosmic chronology, information retention, and the very nature of discreteness.

If the wave structure leaves no trace, it is hidden; if it does, it is indispensable.

Contents

I Background and Motivation	4
1 Introduction	4
2 The Evolving Story of Cosmology	4
3 Event Horizons and the Limits of Knowledge	5
4 Manifold Structure and Singularities	7
II Standing Wave Geometry	8
5 Spacetime as a Discrete Standing Wave	8
5.1 From Continuum to Discreteness	8
5.2 Spacetime as a Wave Medium	8
5.3 Emergent Lorentz Invariance	9
5.4 The Speed of Light as a Structural Consequence	9
5.5 Geometry Over Mediation	10
5.6 Temporal Quantization and Uncertainty	10
6 Gravity as Temporal Gradients in a Standing Wave Spacetime	11
6.1 From Mediation to Emergence	11
6.2 Temporal Gradients and Free Fall	11
6.3 Mass, Impedance, and the Higgs Connection	11
6.4 Summary: Gravity as an Emergent Push	12
7 Black Surfaces and the Critical Warp Limit	12
7.1 Physical Intuition and Mode Filtering	12
7.2 Observational Anchors	13
7.3 Implications for Observables	14
III Mathematical Consequences	15
8 Cosmic Timeline	15
8.1 Looking Backward: Early–Universe Behavior	15
8.2 Modifications to the FRW Metric	16
8.3 Einstein Equations with $c(V)$	16
8.4 Reinterpreting the Cosmological Constant	17
8.5 Reassessing the Horizon Problem	17
8.6 Baryon Acoustic Oscillations and the Poisson Imprint	18
8.7 Summary	19

9 Energy as the Universal Currency	20
9.1 Lorentz Invariance and the Emergence of c	20
9.2 Einstein Equations with a Variable Coupling	21
9.3 Alternative Derivation via the Action Principle	21
9.4 Rest Energy and the Energy–Volume Relation	22
9.5 Recasting the Friedmann Energy Budget	23
9.6 Implications for Equality Epochs	23
9.7 Summary and Connections to Spacetime Impedance	24
10 High-Energy Consequences: Black Surfaces	24
10.1 Static Thin-Shell Model and Israel Junction Conditions	24
10.2 Rotating Black Surfaces: Oblate Shells and Kerr Exterior	25
10.3 Effective Speed of Light Inside Black Surfaces	25
10.4 Frequency Filtering and the Stretch Limit	26
IV Implications and Extensions	28
11 Observational and Experimental Consequences	28
11.1 Gravitational Wave Ringdowns and Echoes	28
11.2 Accretion and Electromagnetic Signatures	28
11.3 Surface Oscillations and Quasinormal Mode Spectra	28
11.4 Information Recovery and Interior Access	29
11.5 Open Problems and Research Directions	29
12 Open Questions and Theoretical Tensions	29
12.1 Does a Discrete Spacetime Demand a Reformulation of Quantum Mechanics?	30
12.2 How Does Gravity Arise from Discrete Geometry?	30
12.3 What is the Role of the Higgs Field in a Resonant Spacetime?	30
12.4 Can Vacuum Energy Be Regularized by Discreteness?	30
12.5 Is Cosmological Redshift Partially Misattributed?	30
12.6 How Does Information Escape a Black Surface?	30
12.7 Does Discreteness Imply a Preferred Scale or Frame?	31
12.8 Is There a Boundary to the Universe or Its Wave Domain?	31
12.9 Is This Model Testable?	31

Part I

Background and Motivation

1 Introduction

Modern physics rests on two towering pillars: General Relativity (GR), which governs the large-scale structure of spacetime and gravity, and Quantum Mechanics (QM), which accurately describes phenomena at the smallest scales. Despite their individual successes, these frameworks remain fundamentally incompatible when pushed to their extremes [1]. In particular, attempts to describe gravity at quantum scales or reconcile spacetime singularities with quantum information have proven deeply problematic [2, 3].

Efforts to unify these two domains have produced many theoretical frameworks—string theory, loop quantum gravity, and others—yet none have yielded definitive, testable predictions. Observational clues, such as the flatness of galactic rotation curves [4], anomalous galaxy cluster velocities [5], and the accelerated expansion of the universe [6, 7, 8], have motivated the introduction of dark matter and dark energy as explanatory constructs. However, these additions may represent symptoms of a deeper misunderstanding about the nature of spacetime itself.

In parallel, the holographic principle has emerged from black hole thermodynamics and string theory, suggesting that the information content of a volume of space may be encoded on its boundary [9]. While this principle remains speculative in many contexts, it offers tantalizing hints that spacetime may be fundamentally discrete or quantized [9, 10].

This work explores an alternative hypothesis: that spacetime is a discrete, physically real medium capable of supporting standing waves. In this model, gravity is not a fundamental force but an emergent effect caused by temporal gradients—regions where time progresses at different rates. The model replaces singularities with surface phenomena, characterized by a critical energy density per unit area, and proposes that the speed of light c may vary with the volume of the universe, denoted $c(V)$, potentially explaining cosmological observation—and modifying the case for dark matter and energy.

The following sections examine the implications of such a model, its potential observational signatures, and its ability to unify or reinterpret existing phenomena in general relativity, quantum mechanics, and cosmology.

2 The Evolving Story of Cosmology

When Edwin Hubble observed that galaxies were receding—and that the farther away they were, the faster they moved—he shattered the long-held belief in a static universe. The cosmos, it seemed, was expanding [11]. Rewinding this motion backward led to a profound and singular conclusion: the universe must have once been compressed into an incredibly dense state. Thus was born the Big Bang theory—and with it, modern cosmology.

Yet the early universe revealed further mysteries. The Cosmic Microwave Background (CMB), the relic radiation from the Big Bang, was found to be astonishingly smooth—with only minute temperature fluctuations across the sky, as confirmed by missions like COBE [12]. This unexpected uniformity gave rise to the theory of **inflation**—a brief period of ultra-rapid expansion just after the Big Bang [13]. Inflation solved the smoothness problem, but introduced new puzzles: what triggered it, and what caused it to stop?

At the same time, astronomers observed something equally puzzling: distant, highly redshifted galaxies appeared to be surprisingly well-formed. Stars and galactic structures seemed to have emerged far earlier than expected. This discrepancy hinted at the presence of unseen mass—Cold Dark Matter (CDM)[14]—which could explain the rapid formation of structure through additional gravitational pull.

For decades, cosmologists debated whether the universe was open or closed. An open universe would expand forever; a closed one would eventually collapse. To settle this, scientists turned to **Type Ia supernovae**—stellar explosions with nearly uniform intrinsic brightness. By comparing their observed brightness to their redshift, researchers could map the universe’s expansion history.

But what they found was astonishing. Instead of slowing down, the expansion of the universe was accelerating [6, 7]. This defied all expectations—and implied the existence of a previously unknown repulsive force counteracting gravity. Thus, the concept of Dark Energy was born. Today, this mysterious energy is thought to comprise nearly 70% of the universe, yet its nature remains one of the deepest puzzles in physics [15].

In an accelerating universe, we inhabit a strange and precarious observational moment. It is still early enough that we can peer deep into the cosmos—detecting ancient galaxies, measuring the cosmic microwave background, and constructing a picture of the early universe. But as expansion continues to accelerate, the observable universe shrinks. Galaxies will gradually redshift beyond visibility, and one day only our local group will remain observable. Had cosmic observations begun then, we might never have known the universe held anything beyond our own galaxy. Push far enough into this future, and even our own galactic structures will dissolve. In the ultimate extrapolation—the so-called Big Rip [16]—the fabric of space itself is torn apart, atom by atom, as even molecular bonds are overcome by expansion.

As it stands, modern cosmology describes three main eras. First, a brief and poorly understood **inflationary epoch**, during which the universe expanded faster than the speed of light [13]. Then, a **matter-dominated era**, governed by the gravitational pull of baryonic and cold dark matter, leading to galaxy formation. Finally, our current era—the era of **accelerating expansion**, driven by Dark Energy. Each era is invoked to reconcile observation with theory, yet each also introduces profound and unresolved mysteries.

3 Event Horizons and the Limits of Knowledge

In the realm of quantum mechanics, one of its foundational postulates is that the wavefunction of a free particle is non-zero across all space [17]. The probability distribution may diminish with distance, but it never fully vanishes—it pervades the entire universe.

This assertion underlies the idea that quantum particles are fundamentally delocalized until measured or interacted with.

But this universal presence of the wavefunction seems to stand in direct contradiction with one of the most striking predictions of general relativity: the event horizon [18, 19]. By definition, an event horizon is a surface beyond which no information, no signal—not even light—can escape to the outside universe. It marks the boundary of a black hole, and according to GR, anything that crosses this threshold is lost to the observable universe.

This presents a philosophical and physical dilemma: *What happens to the wavefunction inside the event horizon?* Does it persist as quantum mechanics insists, or is it obliterated, as the singularities of general relativity seem to demand?

To accept the GR view fully is to accept that spacetime becomes infinitely curved at the center of a black hole—a singularity where density becomes infinite and time stops. But in quantum theory, time is not optional—it is the very stage upon which the wavefunction evolves. If time truly stops at a singularity, then the Schrödinger equation—and all of quantum mechanics—loses its footing. A wavefunction with no time cannot evolve. It cannot be normalized. It ceases to be physical. This exposes a deep incompatibility: in general relativity, spacetime may end; in quantum mechanics, it must persist [3].

This tension is precisely where a discrete standing wave model offers a resolution. In this framework, time cannot stop. It may slow to near-standstill due to extreme temporal dilation, but the underlying oscillatory structure ensures that every region of spacetime retains some finite temporal frequency. Even beyond an event horizon, no region drops below the minimal “tick” of the temporal waveform. As a result, the wavefunction remains well-defined. Evolution continues, however slowly. Unitarity is preserved. The singularity, in this view, is not a physical endpoint but a breakdown of the continuum approximation.

Such questions point to a deeper discontinuity between QM and GR. Quantum mechanics cannot tolerate a system where the wavefunction abruptly ends or becomes undefined. The singularity, in this sense, may be less a “thing” than a symptom—a clue that we are forcing two incompatible frameworks to describe the same extreme reality.

Moreover, black holes are not as featureless as they were once imagined. The no-hair theorem [20] states that a black hole, from the outside, can be described by only three observable parameters:

- Mass
- Spin (angular momentum)
- Charge

This raises another striking question: *If the internal structure of a black hole is a singularity—an infinitesimal point—how can it exhibit macroscopic properties like spin or charge?* How does a zero-volume point rotate? How does it possess a field?

These paradoxes hint that the singularity is not the end of the story, but rather the beginning of a deeper one. The properties we attribute to black holes may not emerge from

an actual point of infinite density but from the structure of the spacetime wave itself—a standing wave compressed to its extreme, where geometry becomes a function of probability amplitude.

4 Manifold Structure and Singularities

In General Relativity, spacetime is modeled as a smooth, four-dimensional manifold, with a one-to-one, onto, and differentiable mapping to physical events [1]. This bijective structure ensures that the geometry of the manifold faithfully represents the behavior of spacetime itself. However, the appearance of a curvature singularity in physical space signals a deep problem: it corresponds to a “pole” in the manifold [19], a concept that is at odds with the manifold’s requirement of infinite differentiability. In other words, true poles cannot exist on a smooth manifold, highlighting an intrinsic tension in the classical framework.

Singularities are thus more than mere coordinate artifacts—they mark a breakdown in both geometry and predictability. The smooth manifold cannot accommodate these divergent points, which undermines the assumption of completeness and suggests that GR, despite its elegance, may be an effective theory rather than a final one.

The standing wave spacetime model addresses this issue by replacing singularities with structured nodal regions of near-zero wave amplitude. These regions preserve differentiability and maintain the bijection between manifold points and physical events, avoiding infinities and causal discontinuities altogether.

Classical singularity theorems, such as those formulated by Hawking and Penrose, formalize the unavoidable appearance of geodesic incompleteness under certain assumptions: energy conditions, trapped surfaces, and a smooth, continuous spacetime manifold [21]. In conventional GR, these theorems imply that singularities are physically inevitable consequences of gravitational collapse or cosmic expansion.

By contrast, the standing wave spacetime model departs fundamentally from these premises. It replaces the continuous manifold with a discrete, oscillatory substrate, introduces a critical surface energy density in place of divergent curvature, and reframes gravitational effects as temporal gradients rather than stress-energy curvature. In doing so, it avoids the conditions required for the classical theorems—not by contradiction, but by rejecting the axioms from which those conclusions arise. Singularities, then, may be understood not as physical inevitabilities, but as artifacts of an incomplete theoretical framework.

Part II

Standing Wave Geometry

5 Spacetime as a Discrete Standing Wave

The classical view of spacetime, as a smooth, continuous manifold, has been enormously successful. General Relativity models it as a differentiable geometry whose curvature encodes gravitation [22]. Quantum field theory treats it as a fixed backdrop upon which particles and fields interact. Yet both perspectives encounter limits: General Relativity breaks down at singularities, while quantum theory struggles with Planck-scale discreteness [23]. To move forward, it may be necessary to shift perspective—not to discard either theory, but to see spacetime itself as a resonant structure.

This chapter advances the hypothesis that spacetime is a *discrete standing wave*: a self-organizing, quantized geometry whose oscillatory nature explains both its smooth large-scale appearance and its paradox-resolving behavior at extreme scales.

5.1 From Continuum to Discreteness

Wave systems in nature rarely admit arbitrary configurations. A vibrating string supports only harmonics consistent with its length and boundary conditions; a resonant cavity permits only quantized modes. If spacetime is fundamentally wave-like, its structure may be similarly constrained.

In this view, spacetime is not infinitely divisible but possesses an intrinsic granularity. Each “node” or unit cell embodies a minimal spacetime volume, its properties fixed by the global geometry of the universe. Quantization then arises not from imposed rules but from the natural discreteness of the medium.

This perspective renders a graviton unnecessary: gravitation is not the exchange of quanta, but the geometric response of the wave structure itself. Curvature and phase gradients are sufficient to account for the apparent force of gravity, aligning with the spirit of General Relativity but extending it into the quantum domain.

5.2 Spacetime as a Wave Medium

If spacetime sustains standing waves, it must act as a medium with elastic and resonant qualities. While not material in the classical sense, its properties can be abstracted in analogy with known wave-bearing systems:

- **Discrete at fundamental scales**, ensuring quantization of modes;
- **Elastic**, so that curvature responds to stress-energy;
- **Finite**, so that only a restricted set of modes is globally consistent.

Such a medium resolves longstanding paradoxes:

- It avoids the infinities of singularities by enforcing minimum structure;
- It imposes a resolution scale consistent with quantum uncertainty;
- It explains gravitation as geometry, rather than as particle exchange.

Thus, rather than “fields on a manifold” physics becomes the resonant behavior of the manifold itself.

5.3 Emergent Lorentz Invariance

Special Relativity is built upon Lorentz invariance, traditionally introduced as a postulate about the form of physical laws. In a standing-wave spacetime, Lorentz symmetry arises naturally from the coherence of the oscillatory medium.

- Each spacetime point oscillates as a synchronized temporal resonator;
- Relative motion manifests as a phase shift between oscillators;
- Preservation of phase coherence requires time dilation and length contraction.

Thus Lorentz invariance is not imposed externally but is an emergent necessity of the geometry. Motion itself becomes a relational phase difference within the oscillating lattice, and the familiar machinery of Special Relativity follows as a consequence rather than an axiom [24].

5.4 The Speed of Light as a Structural Consequence

In conventional physics, the speed of light c is fundamental and universal. Within the standing-wave model, it is instead a structural property of the spacetime medium. As wave speed on a string depends on tension and mass density, the effective value of c depends on the impedance of the spacetime lattice.

If the universe expands, boundary conditions shift. A larger volume alters permissible resonant modes, suggesting a dynamic relationship:

$$c = c(V)$$

where V denotes the volumetric scale of the universe. This possibility carries deep implications:

- Early-universe structure formation may proceed easier if c were lower at small V ;
- Inflationary behavior could be mimicked without requiring superluminal expansion;
- Observations of Type Ia supernovae redshifts [6, 7] might be reinterpreted without invoking dark energy.

Thus c is not an immutable constant but an emergent property of spacetime’s evolving resonant structure.

5.5 Geometry Over Mediation

Conventional quantum field theory relies upon exchange particles—gravitons, photons, gluons—to mediate forces. The standing-wave model reframes this picture: interactions arise directly from geometry. Mass-energy deforms the resonant structure; other entities respond not to “forces” but to the local geometry of phase and curvature.

This return to geometry resonates with Wheeler’s dictum, “spacetime tells matter how to move, matter tells spacetime how to curve” [22]. The novelty here is discreteness: curvature itself is quantized, embedded within the oscillatory framework. This offers a possible route to unification, since both relativity and quantum theory become facets of the same resonant substrate.

5.6 Temporal Quantization and Uncertainty

In this model, time is not a smooth external parameter but an emergent property of oscillation. Each unit of spacetime undergoes periodic temporal fluctuation, so that the “flow” of time is the coordination of phase relationships across the lattice.

Just as a standing wave admits only discrete spatial modes, it admits only certain temporal frequencies. Below the fundamental oscillation period, temporal ordering loses meaning—not because of measurement imprecision, but because the underlying structure forbids it. This yields a natural temporal uncertainty principle, shielding physics from pathological infinities and aligning with quantum uncertainty.

Time thus joins position and momentum as contextual and quantized. It is not absolute, but emergent: the heartbeat of spacetime itself.




Figure 1: A schematic representation of spacetime as a standing wave. Each spatial coordinate (x, y) exhibits a temporal oscillation, represented on the vertical axis. The oscillatory structure encodes temporal granularity, setting a natural limit to resolution and uncertainty.

6 Gravity as Temporal Gradients in a Standing Wave Spacetime

The prevailing view in modern physics treats gravity as a fundamental interaction, possibly mediated by a massless spin-2 boson, the graviton [25, 26, 27]. While mathematically consistent, this picture raises persistent challenges: how could such a mediator act universally on all matter, propagate effects that seem instantaneous across vast distances, and remain both persistent and irreversible if governed by particle exchange? These difficulties motivate a shift in perspective: gravity need not be mediated at all, but may instead emerge from the geometry of time itself.

6.1 From Mediation to Emergence

Consider two neighboring regions of spacetime where clocks tick at different rates. A physical object entering such a region will drift toward slower time, not because it is pulled by a force, but because the structure of spacetime itself has a gradient. In this view, gravity is a push into regions of greater temporal dilation, much like fluid flows along a pressure gradient. General Relativity already encodes this behavior mathematically as geodesic motion in curved spacetime [28, 29]; the present model extends that intuition by proposing that curvature is not a passive geometry but an active *temporal gradient*.

6.2 Temporal Gradients and Free Fall

Let $\tau(x^\mu)$ represent the proper time measured at spacetime point x^μ . Near mass-energy concentrations, the local rate of time flow decreases, and we may define a temporal gradient as

$$\nabla_t \equiv \nabla \left(\frac{d\tau}{dt} \right).$$

This gradient points toward regions where time runs more slowly. Matter responds by accelerating in that direction:

$$\vec{a} \propto -\nabla_t.$$

Free fall is therefore not motion under a force, but the natural descent along a temporal slope. The equivalence principle [30, 31] follows directly: all matter responds identically to ∇_t , because it is not being acted upon by an external interaction but simply guided by the structure of proper time.

6.3 Mass, Impedance, and the Higgs Connection

The Higgs mechanism explains why particles acquire mass through interaction with a pervasive field [32, 33], yet it offers little intuition for why such a field exists. In the standing wave model of spacetime, a new picture emerges: spacetime itself is a discrete resonant structure, with nodes and antinodes that define fundamental intervals Δx and Δt . Massless particles such as photons traverse this structure without impedance, while massive particles disturb it

locally. This disturbance produces resistance to motion — an *impedance* — which manifests as inertia.

In this framework, the Higgs field need not be a separate entity. Rather, it is an emergent feature of the nodal structure of spacetime. Mass is the local impedance of matter against this discrete wave medium, and gravity arises when temporal gradients create regions of least resistance. Matter is effectively pushed toward slower time because that is where motion through the spacetime lattice encounters minimal impedance. Gravity thus becomes a macroscopic consequence of microscopic impedance.

6.4 Summary: Gravity as an Emergent Push

In this reframing, gravity is neither a mysterious long-range force nor a quantum exchange. It is the natural flow of matter down temporal gradients in a discrete standing wave spacetime. The Higgs mechanism and gravitational behavior are united under a single structural principle: mass, inertia, and acceleration all originate from the way matter interacts with the temporal fabric of the universe. Gravity, then, is best understood not as attraction, but as an emergent push into slower time.

7 Black Surfaces and the Critical Warp Limit

In the standing wave spacetime framework, the extreme warping traditionally attributed to black hole singularities is replaced by a finite, physical surface limit. Rather than diverging to infinite curvature, spacetime exhibits a maximum surface energy density beyond which further stretching is not possible [34, 21].

We define the *critical surface energy density* as:

$$\sigma_{\text{crit}} = \frac{E}{A} \quad \text{where } E = \text{local energy content}, \quad A = \text{surface area enclosing the region.}$$

At σ_{crit} , the local standing wave structure reaches its maximum deformation. This implies that a black hole does not harbor a mathematical singularity but terminates at a *Black Surface* — a phase transition boundary beyond which spacetime can no longer sustain additional curvature.

7.1 Physical Intuition and Mode Filtering

Spacetime in this picture behaves analogously to an elastic drumhead composed of quantized standing wave nodes. As mass-energy accumulates, the nodes are displaced, producing a local temporal gradient — the origin of gravitational effects. Approaching σ_{crit} , the system undergoes a transition similar to a drum tightening: lower-frequency modes can no longer be supported and are expelled, while only higher-frequency oscillations persist.

This expulsion of long-wavelength modes acts as a *high-pass filter* on matter-energy degrees of freedom, resonant with ideas in quasinormal mode analyses [35]. Low kinetic energy

matter is squeezed outward, forming a dense shell at the periphery. The Black Surface therefore emerges not as a mathematical artifact but as a physically stable configuration: a shell-like boundary where the discrete standing wave lattice can still sustain oscillation, while the interior flattens into a nearly featureless region.

In this sense, the Black Surface preserves the observable phenomena of horizons and shadows while avoiding infinities. The singularity is replaced by a phase transition in the standing wave structure itself.

7.2 Observational Anchors

Remarkably, existing astrophysical measurements hint at this finite limit. Supermassive black holes exhibit average volumetric densities comparable to water [36, 37] — orders of magnitude lower than nuclear densities. Such values are consistent with the idea that the interior is not a runaway collapse but instead a regulated region whose warp has saturated at σ_{crit} , pushing additional matter outward into the surrounding shell.




Figure 2: Stages of Black Surface formation in the discrete standing wave model: a smooth progression from flat spacetime through warp limit to stable shell hidden behind the emergent event horizon.

7.3 Implications for Observables

By reframing black holes as Black Surfaces, this model preserves the large-scale predictions of general relativity while offering new testable features:

- The size and morphology of photon spheres and gravitational shadows [38], which may encode the presence of a shell-like surface rather than a pointlike singularity.
- Angular momentum constraints (e.g., the Kerr bound) as manifestations of how the standing wave structure resists further stretching.
- Potential imprints of surface-mode filtering in gravitational wave spectra or accretion dynamics, where transitions across σ_{crit} could leave observable signatures [39, 40].

Part III

Mathematical Consequences

8 Cosmic Timeline

8.1 Looking Backward: Early–Universe Behavior

In this model, the speed of light is not taken as a fundamental constant but arises from the ratio of spatial to temporal intervals,

$$c \equiv \frac{\Delta x}{\Delta t}. \quad (1)$$

While the fundamental tick of time, Δt , is assumed invariant, the spatial interval Δx scales with the expansion factor $a(t)$. As the universe contracts toward the big bang, $\Delta x \rightarrow 0$ while Δt remains finite, forcing c itself to evolve with cosmic volume.

Thus, in the limit $a(t) \rightarrow 0$, we obtain

$$\lim_{a \rightarrow 0} c(V) = 0, \quad (2)$$

so that the *effective* speed of light in the early universe tends toward zero — in stark contrast to the usual intuition of diverging speeds near the big bang.

This inversion has profound consequences:

- Causal signals in the early universe propagate extremely slowly.
- The comoving particle horizon evolves in a radically different manner, directly resolving the horizon problem and removing the need for an *ad hoc* inflationary phase to explain the observed uniformity of the Cosmic Microwave Background.
- Interpreting high-redshift events under the constant- c assumption introduces systematic errors that grow exponentially with redshift.

For example, if $c(V) \propto a^{3\alpha}$, then the standard lookback time formula,

$$t_{\text{inferred}}(z) = \int_0^z \frac{dz'}{(1+z') H(z')}, \quad (3)$$

must be replaced by

$$t_{\text{true}}(z) = \int_0^z \frac{[1+z']^{-3\alpha}}{(1+z') H(z')} dz'. \quad (4)$$

The ratio $t_{\text{true}}/t_{\text{inferred}}$ falls sharply toward zero as z increases. Thus, distant supernovae appear anomalously faint when analyzed under constant- c assumptions, creating the illusion of accelerated expansion even in the absence of dark energy. In this view, what standard cosmology interprets as dark energy may be an artifact of treating c as immutable.

To examine these consequences rigorously, we now reformulate the FRW metric and Einstein equations under the condition $c = c(V)$.

8.2 Modifications to the FRW Metric

In the standard Friedmann–Robertson–Walker (FRW) cosmology, the line element is written as

$$ds^2 = -c^2 dt^2 + a(t)^2 \left(\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right), \quad (5)$$

with constant light speed c . In the standing wave spacetime model, however, the speed of light is not an immutable constant but depends on the volumetric scale of the universe,

$$c = c(V), \quad (6)$$

where V is the effective spatial volume determined by the scale factor $a(t)$. Thus the line element becomes

$$ds^2 = -c(V)^2 dt^2 + a(t)^2 \left(\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right). \quad (7)$$

This change has immediate consequences. Proper time and conformal time must be carefully distinguished when c itself evolves. The conformal time is defined by

$$\eta = \int \frac{c(V)}{a(t)} dt \quad (8)$$

which alters the causal structure of the FRW spacetime. Horizons, particle communication limits, and light cone structures all evolve differently than in the constant- c case.

8.3 Einstein Equations with $c(V)$

The Einstein field equations,

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}, \quad (9)$$

inherit the variability of c . If $c = c(V)$, the coupling between geometry and stress–energy becomes time dependent. This does not break covariance: c remains a scalar under local Lorentz transformations. Rather, it introduces an evolving *impedance* in the relationship between curvature and matter, a **phase–metric coupling** whereby the standing–wave structure of spacetime modulates the strength of interaction between geometry and energy density.

Deriving the modified Friedmann equations, we obtain

$$\left(\frac{\dot{a}}{a} \right)^2 + \frac{k c(V)^2}{a^2} = \frac{8\pi G}{3} \rho, \quad (10)$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} \left(\rho + \frac{3p}{c(V)^2} \right). \quad (11)$$

In the constant- c limit these formulae reduce to the standard Friedmann–Robertson–Walker form [41], but with $c(V)$ explicit the balance between density, pressure, and acceleration is altered.

Previous varying-speed-of-light (VSL) models (e.g. [42, 43, 44]) explored whether a time-dependent $c(t)$ could resolve puzzles such as the horizon and flatness problems. In those approaches, c was introduced as an external prescription, a function appended to the equations.

By contrast, in the present framework the variability of $c(V)$ is inevitable: it follows directly from the geometric structure of spacetime itself. With Δt invariant and Δx scaling with cosmic volume, the effective propagation speed cannot remain constant. The result is a dynamical reinterpretation of cosmological evolution in which phenomena usually ascribed to a cosmological constant or inflation may instead reflect the natural evolution of $c(V)$.

8.4 Reinterpreting the Cosmological Constant

When distant Type Ia supernovae were first observed to be dimmer than expected, the conclusion was that the universe’s expansion was accelerating [6, 7]. This led directly to the resurrection of Einstein’s cosmological constant Λ , rebranded as Dark Energy, comprising nearly 70% of the cosmic energy budget [15].

But in the discrete standing wave spacetime model, this interpretation may not be required. If the speed of light grows with the volume of the universe, i.e.

$$c = c(V), \quad \frac{dc}{dV} > 0, \quad \frac{d^2c}{dV^2} < 0, \quad (12)$$

then signals from distant supernovae will appear redshifted and dimmer not because of acceleration, but because our interpretation assumes constant light speed. The observed dimming may instead be a signature of a **time-varying propagation speed in a stretching medium**. Correcting the luminosity-distance relation to account for $c(V)$ may eliminate or greatly reduce the apparent acceleration. The “Dark Energy” problem becomes a bookkeeping error of interpretation.

Einstein’s greatest mistake may have been doubting his deepest intuition.

This reinterpretation also directly addresses the infamous *Vacuum Energy Catastrophe*. In standard treatments, quantum field theory predicts vacuum energy densities 10^{120} times larger than observed. But if the cosmological constant is not a true energy density, but an artifact of misinterpreting signals under constant c , then the contradiction vanishes [45].

8.5 Reassessing the Horizon Problem

In standard cosmology, the so-called *horizon problem* arises because the Cosmic Microwave Background (CMB) is observed to be remarkably uniform in temperature across regions that, under the constant- c assumption, appear causally disconnected at recombination [12]. The conventional reasoning is that there was insufficient time for thermalization to occur, motivating an inflationary phase to explain this uniformity [46].

This argument relies on three key assumptions:

- The speed of light c is constant throughout cosmic history.

- All energy resides in matter or radiation; spacetime itself is passive.
- Thermal equilibrium requires direct causal contact between regions.

While these considerations make the uniformity of the CMB puzzling under standard assumptions, the standing-wave spacetime model offers a different perspective.

In this framework, spacetime carries energy, which is transferred uniformly into matter as the universe expands, according to the evolving mass-energy relation

$$E(V) = m c(V)^2. \quad (13)$$

Because this energy distribution is homogeneous, matter inherits thermal uniformity naturally, independent of causal contact or the size of the particle horizon.

Minor quantum fluctuations in matter are superimposed on this uniform background, providing seeds for the anisotropies that later grow into galaxies and clusters [47]. These fluctuations coexist with an overall thermal uniformity and do not contradict the mechanism: the uniform energy transfer sets the baseline, while quantum perturbations generate structure.

Thus, under this framework, the horizon problem is no longer a puzzle. Uniformity of the CMB is a direct consequence of the underlying geometry and energy distribution of spacetime. What standard cosmology treats as a “problem” becomes instead a prediction: thermal homogeneity is expected across the observable universe without requiring inflation or artificially adjusting the propagation speed of light.

8.6 Baryon Acoustic Oscillations and the Poisson Imprint

Baryon Acoustic Oscillations (BAO) provide a snapshot of the primordial density field at the epoch of recombination. In standard cosmology, the sound horizon at that time, r_s , sets the scale of the oscillations, which then seed the later formation of galaxies and clusters [48, 49].

In the standing wave spacetime model, the effective speed of light in the early universe is reduced since it is determined by the underlying spatial geometry,

$$c(V) \rightarrow 0 \quad \text{as} \quad a(t) \rightarrow 0, \quad (14)$$

resulting in a correspondingly smaller sound horizon. The BAO imprint in the density field is therefore compressed relative to what would be inferred under a constant- c assumption.

The Poisson equation,

$$\nabla^2 \Phi = 4\pi G a^2 \rho \delta, \quad (15)$$

evolves the initial density perturbations into the large-scale structure observed today [50]. Because $\delta(\mathbf{x})$ carries the compressed BAO signature, structure formation proceeds on a timeline that appears accelerated if interpreted with constant c .




Figure 3: Causal chain connecting early-universe behavior of $c(V)$ to observed large-scale structure. The compressed BAO imprint seeds density perturbations that evolve under the Poisson equation, making early galaxies appear “ahead of schedule” if interpreted with constant- c .

Observationally, this provides a natural explanation for why galaxies and clusters in the early universe seem “ahead of schedule” [51, 52] — they are not forming unusually fast; rather, the effective early-universe clock runs slower in this model.

Thus, what appears observationally as a tension with Λ CDM and a driver for dark matter or dark energy can instead be reframed as a natural outcome of a variable- $c(V)$ early universe.

8.7 Summary

The FRW metric, once modified to include a variable speed of light, provides a natural mechanism for reproducing the apparent acceleration of the universe without requiring Dark Energy. The reinterpretation is consistent with covariance, alters the Einstein equations in a physically transparent way, and suggests fresh perspectives on BAO, CMB horizons, and the Poisson-governed growth of density perturbations. This framework thus offers a mathematically tractable and observationally testable alternative to the cosmological constant paradigm.




Figure 4: Horizon problem “resolved”: thermal uniformity of matter and the CMB emerges naturally from uniform energy distribution in standing-wave spacetime.

9 Energy as the Universal Currency

9.1 Lorentz Invariance and the Emergence of c

A natural objection to any theory with a variable speed of light is the possible violation of Lorentz invariance. The standing-wave spacetime model, however, preserves *local* Lorentz symmetry while providing a physical basis for the emergence of c .

The model introduces a natural node structure defined by characteristic intervals of space Δx and time Δt . Their ratio directly yields the local speed of light,

$$c = \frac{\Delta x}{\Delta t}. \quad (16)$$

As the universe expands, Δt remains fixed as a fundamental measure of change, while Δx grows with the volume of space. This leads to a slow global evolution of c , but one that leaves local physics unaffected.

Locally, in a tangent Minkowski frame, the metric retains the standard form,

$$ds^2 = -c_{\text{local}}^2 d\tau^2 + dx^2 + dy^2 + dz^2, \quad (17)$$

with c_{local} invariant for all observers. All experimentally tested relativistic phenomena—time dilation, length contraction, mass–energy equivalence—remain intact. The variation arises only in comparing widely separated epochs of cosmic history:

$$c = c(V), \quad \frac{dc}{dV} > 0, \quad \frac{d^2c}{dV^2} < 0, \quad (18)$$

In this sense, Lorentz invariance is not broken but reinterpreted. The constancy of Δt and the evolution of Δx follow necessarily from the standing-wave structure of spacetime, grounding c in the geometry of the universe itself rather than treating it as a fundamental axiom.

9.2 Einstein Equations with a Variable Coupling

In general relativity, the Einstein field equations [22]

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}, \quad (19)$$

show that the speed of light does more than set units: it enters as part of the coupling constant that links matter to geometry [42]. The standing-wave spacetime model modifies this structure by allowing

$$c = c(V), \quad (20)$$

so that the effective coupling evolves with the cosmic volume.

This has two key consequences:

1. **Time-dependent impedance of spacetime.** Spacetime geometry does not respond to stress-energy with fixed strength. Instead, the ratio of curvature to energy density changes as the universe expands, reflecting a dynamical impedance of spacetime itself.
2. **Redefinition of energy density.** Stress-energy conservation,

$$\nabla^\mu T_{\mu\nu} = 0, \quad (21)$$

still holds by construction. But the energy associated with matter now includes a factor of $c(V)^2$, so that the very definition of energy density evolves with cosmic volume, consistent with

$$E(V) = m c(V)^2. \quad (22)$$

It is useful to write the volume-dependent coupling explicitly as

$$\kappa(V) = \frac{8\pi G}{c(V)^4}, \quad (23)$$

so that

$$G_{\mu\nu} = \kappa(V) T_{\mu\nu}. \quad (24)$$

As $c(V)$ grows with the expansion, $\kappa(V)$ decreases, making the effective strength of gravity diminish over cosmic time. This provides a natural dynamical explanation for the relative weakness of gravity compared to the other forces, without requiring fine-tuning of constants.

9.3 Alternative Derivation via the Action Principle

The same conclusion can be obtained from the Einstein–Hilbert action [1, 53]:

$$S = \frac{c^3}{16\pi G} \int R \sqrt{-g} d^4x + S_{\text{matter}}. \quad (25)$$

If $c = c(V)$, both the prefactor c^3 and the scaling of the matter sector acquire explicit volume dependence. Variation of this action yields Einstein equations with an evolving coupling $\kappa(V)$, consistent with the analysis above [42].

Unlike the standard case, additional boundary contributions appear when varying a time-dependent c . These terms may be linked to horizon evolution and vacuum structure, suggesting a deeper connection between cosmic expansion, particle creation, and the effective energy content of spacetime.

9.4 Rest Energy and the Energy–Volume Relation

The volume dependence of the speed of light modifies the equivalence between mass and energy. For a particle of invariant rest mass m , the rest energy becomes

$$E(V) = m c^2(V). \quad (26)$$

Mass itself is not changing; rather, the energy equivalence scale evolves with cosmic volume. This shift may be interpreted as a form of *temporal pressure*: the gradient in the rate of time progression across spacetime acts as an effective pressure, driving matter toward regions of slower time (the phenomenon we perceive as gravity) while simultaneously rescaling its energy equivalence.

If rest energy grows with $c(V)$, energy conservation requires that this increase draw from a physical reservoir. We posit that the standing-wave fabric of spacetime itself stores energy, from which matter continuously extracts rest-energy. In the continuity equation,

$$\frac{d\rho_m}{dt} + 3H\rho_m = Q(a), \quad (27)$$

a source term $Q(a)$ naturally arises, with

$$Q(a) \sim \frac{d}{dt} [n_m mc^2(a)], \quad (28)$$

where n_m is the comoving number density. This mechanism uniformly deposits energy throughout space, consistent with the observed isotropy of the cosmic microwave background.

For a power-law ansatz $c(a) = c_0 a^{3\alpha}$, the matter density evolves as

$$\rho_m(a) \propto a^{-3} c^2(a) = a^{-3+6\alpha}, \quad (29)$$

recovering the same scaling, but now endowed with a direct energetic interpretation.

Finally, the functional form of $c(V)$ must satisfy

$$c = c(V), \quad \frac{dc}{dV} > 0, \quad \frac{d^2c}{dV^2} < 0, \quad (30)$$

ensuring that c always increases with cosmic volume, but at a decelerating rate. Today the change is imperceptible locally, though its signature remains in the light from distant epochs, where the slower early- c regime played a dominant role in the universe’s energetic evolution.

9.5 Recasting the Friedmann Energy Budget

The Friedmann equation in standard cosmology reads

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3} (\rho_m + \rho_\gamma + \rho_\Lambda) - \frac{k}{a^2}. \quad (31)$$

with densities scaling as

$$\rho_m(a) \propto a^{-3}, \quad \rho_\gamma(a) \propto a^{-4}, \quad \rho_\Lambda \approx \text{const.} \quad (32)$$

If instead $c = c(V)$, these relations no longer hold. Matter density acquires a time-dependent factor,

$$\rho_m(a, V) = \frac{mc^2(V)}{a^3}, \quad (33)$$

while radiation scales as

$$\rho_\gamma(a, V) \sim \frac{c(V)}{a^4}. \quad (34)$$

Even the vacuum term may be interpreted as an impedance property of spacetime,

$$\rho_\Lambda(V) \sim f(c(V)), \quad (35)$$

with form set by the phase-metric coupling of the standing wave background. The traditional separation into radiation-, matter-, and dark-energy-dominated eras is thus replaced by a coupled evolution controlled by $c(V)$.

9.6 Implications for Equality Epochs

Matter–radiation equality is normally set by

$$\rho_m(a) = \rho_\gamma(a). \quad (36)$$

With $c(V)$ this becomes

$$\frac{mc^2(V)}{a^3} = \frac{c(V)}{a^4}, \quad (37)$$

leading to

$$a_{\text{eq}} \sim \frac{1}{mc(V)}. \quad (38)$$

The epoch of equality is no longer fixed by microphysical constants but drifts with the cosmic evolution of $c(V)$. A rapidly increasing $c(V)$ pushes equality to smaller a , delaying matter domination and the onset of large-scale structure. A more gradual evolution accelerates it. The observed “early galaxy problem” thus becomes diagnostic of the slope of $c(V)$ rather than an anomaly.

9.7 Summary and Connections to Spacetime Impedance

In this section we have seen how the evolution of $c(V)$ provides a unified description of energy flow, rest-mass scaling, and the apparent anomalies in cosmic expansion. Matter gains energy from the underlying standing-wave structure of spacetime, producing the observed uniformity and shaping the growth of structure.

The qualitative mechanism linking mass, inertia, and temporal gradients was discussed in Part II (Section 6.3), where we introduced the notion of spacetime impedance as the origin of the Higgs effect. Here, that framework is extended quantitatively: $c(V)$ sets the energy scale, $\kappa(V)$ modulates gravitational coupling, and $Q(a)$ formalizes the extraction of energy from spacetime.

Together, these ideas provide a consistent energetic picture that naturally transitions to the discussion of Black Surfaces and the resolution of GR, QM, and information tensions.

10 High-Energy Consequences: Black Surfaces

The evolution of rest energy in a standing-wave spacetime destabilizes the classical notion of collapse. In conventional general relativity, increasing density drives curvature to infinity at a singularity. In our framework, increasing $c(V)$ drives the system toward a critical surface energy density,

$$\sigma_{\text{crit}} = \frac{E}{A} \sim \frac{mc^2(V)}{A}. \quad (39)$$

Once reached, further compression triggers a phase transition: the formation of a *Black Surface*. This replaces the singularity with a stable impedance boundary, tying the endpoint of collapse directly to the structure of spacetime and its standing-wave properties.

10.1 Static Thin-Shell Model and Israel Junction Conditions

To provide a mathematical foundation, consider a static, spherically symmetric thin shell in vacuum. The spacetime is split into interior and exterior regions:

- **Interior:** Flat Minkowski space,

$$ds_{\text{in}}^2 = -dt^2 + dr^2 + r^2 d\Omega^2.$$

- **Exterior:** Schwarzschild spacetime,

$$ds_{\text{out}}^2 = -\left(1 - \frac{2GM}{r}\right)dt^2 + \left(1 - \frac{2GM}{r}\right)^{-1}dr^2 + r^2 d\Omega^2.$$

Matching at the shell radius $r = R$ requires continuity of the induced metric, while the jump in extrinsic curvature gives the surface stress-energy tensor S_{ab} via the Israel junction conditions:

$$[K_{ab}] - h_{ab}[K] = -8\pi GS_{ab}.$$

For a static shell:

$$\sigma = -\frac{1}{4\pi R} \left(\sqrt{1 - \frac{2GM}{R}} - 1 \right),$$

$$P = \frac{1}{8\pi R} \left(\frac{1 - GM/R}{\sqrt{1 - 2GM/R}} - 1 \right),$$

where σ is surface energy density and P the transverse pressure. As $R \rightarrow 2GM$, the shell mimics the Schwarzschild horizon, with σ and P diverging, reproducing the classical horizon location, but the interior remains flat—no singularity forms.

10.2 Rotating Black Surfaces: Oblate Shells and Kerr Exterior

For rotation, the exterior Schwarzschild metric is replaced with the Kerr metric in Boyer–Lindquist coordinates:

$$ds^2 = - \left(1 - \frac{2Mr}{\Sigma} \right) dt^2 - \frac{4Mar \sin^2 \theta}{\Sigma} dt d\phi + \frac{\Sigma}{\Delta} dr^2 + \Sigma d\theta^2$$

$$+ \left(r^2 + a^2 + \frac{2Ma^2r \sin^2 \theta}{\Sigma} \right) \sin^2 \theta d\phi^2,$$

$$\Sigma = r^2 + a^2 \cos^2 \theta, \quad \Delta = r^2 - 2Mr + a^2, \quad a = \frac{J}{M}.$$

Oblate spheroidal coordinates are chosen to ensure that the interior matches the Kerr exterior on the shell while maintaining flatness at the center. The oblate shell’s interior is expressed as follows:

$$ds^2 = -dt^2 + \frac{a^2(\xi^2 + \eta^2)}{\xi^2 + 1} d\xi^2 + \frac{a^2(\xi^2 + \eta^2)}{1 - \eta^2} d\eta^2 + a^2(\xi^2 + 1)(1 - \eta^2) d\phi^2,$$

with shell at $\xi = \xi_0$. Transformations map (ξ, η) to standard spherical coordinates as $a \rightarrow 0$, confirming smooth recovery of spherical Minkowski spacetime.

Frame dragging naturally emerges from the rotational junction: the $g_{t\phi}$ term encodes the “twist” of inertial frames induced by the shell’s angular momentum. This arises purely from geometric continuity and rotational symmetry, with no exotic matter or singularity needed.

10.3 Effective Speed of Light Inside Black Surfaces

Within the shell, nodal separation Δx grows while Δt remains fixed, leading to an extremely large local effective speed of light,

$$c_{\text{eff}} = \frac{\Delta x}{\Delta t}.$$

Near the shell boundary, spacetime compression reduces Δx , creating a low- c barrier. Light from the interior encounters:

1. Reduced local c at the boundary (trapping),
2. A “refraction” effect due to the c -gradient.

This provides a physical underpinning for horizon opacity beyond classical GR.

10.4 Frequency Filtering and the Stretch Limit

As spacetime approaches the critical stretch σ_{crit} , it behaves as a high-pass filter. Low-frequency modes (rest-mass matter) are progressively excluded from the interior, leaving only high-energy kinetic and massless modes.




Figure 5: Effect of increasing spacetime tension on frequency support. Each curve represents a distinct tension regime, illustrating how lower-frequency modes are progressively suppressed as the stretch approaches and exceeds the critical limit.

Figure 5 illustrates how the frequency support of spacetime evolves under increasing tension. Each curve corresponds to a distinct physical regime:

- **Blue — Unstressed (Flat):** All frequency modes are equally supported. No stretch or curvature is present. This is idealized Minkowski space.
- **Green — Subcritical Stretch:** Geometry begins favoring higher-frequency modes. Lower frequencies are disfavored but still supported — no evictions yet.
- **Orange — Critical Stretch:** The curve touches the origin, indicating the first suppression of zero-frequency modes. This marks the onset of rest-mass mode eviction at this spacetime stretch — shell formation begins.
- **Red — Supercritical Stretch:** Lower-frequency modes are actively excluded. Only high-energy or massless excitations remain sustainable in the interior. The shell is now fully formed. The energy cutoff is shown here as a dashed vertical line.

This dynamic naturally explains surface-area scaling and shell-like mass distributions: low-frequency matter accumulates near the outer shell, while the interior is emptied of rest-mass content. This mechanism is also consistent with entropy scaling: since low-frequency modes are excluded, the number of accessible interior microstates scales with surface area rather than volume.

Part IV

Implications and Extensions

11 Observational and Experimental Consequences

If the discrete standing wave model is to be more than a mathematical curiosity, it must ultimately bear observable fruit. The replacement of black hole singularities with extended shell-like *Black Surfaces* introduces sharp differences in both geometry and dynamics. These differences may leave detectable signatures across multiple observational channels.

11.1 Gravitational Wave Ringdowns and Echoes

Mergers of compact objects provide the most direct window into the strong-gravity regime. In classical general relativity, black holes exhibit damped quasinormal mode (QNM) ringdowns that depend only on mass and spin. In contrast, a shell-based compact object could support:

- **Modified QNM spectra**, with additional or shifted resonant frequencies.
- **Echoes** in the post-merger signal, arising from reflections at the hard inner boundary.
- **Extended ringdown tails**, modulated by the mechanical response of the shell.

Future detectors such as LISA may have sufficient sensitivity to identify such deviations, providing a sharp observational discriminator.

11.2 Accretion and Electromagnetic Signatures

Infalling matter interacts differently with a hard surface than with an event horizon. This may produce:

- **Hard surface emissions**, where accreted material deposits energy onto the shell, generating thermal radiation or high-energy bursts.
- **Quasi-steady boundary layers**, potentially visible in X-ray spectra.
- **Time-variable hotspots**, tied to surface inhomogeneities or oscillations.

Comparisons of accretion efficiencies between candidate black holes and neutron stars may already hint at such possibilities. High-resolution imaging (e.g., from the Event Horizon Telescope) could provide direct constraints.

11.3 Surface Oscillations and Quasinormal Mode Spectra

The shell surface itself may host interface waves or pressure oscillations absent in classical horizons. These could couple to both gravitational and electromagnetic channels, produc-

ing:

- Distinctive QNM ringdown structures.
- Modulations in post-merger signals.

Numerical relativity simulations incorporating matter shells would be required to quantify these spectra, but even qualitative deviations could be observationally testable.

11.4 Information Recovery and Interior Access

The absence of a singularity reframes the black hole information paradox. A Black Surface introduces new possibilities:

- The shell may act as a reflective boundary preserving quantum correlations.
- The interior could be Minkowski vacuum, or some exotic phase, allowing partial recovery of information.
- Unlike “firewall” proposals, the surface offers a geometric rather than ad hoc resolution.

This connects to holographic dualities but also extends beyond them, offering an alternative view of complementarity.

11.5 Open Problems and Research Directions

The above should not be read as firm predictions, but as fertile research directions. A non-exhaustive list includes:

1. **QNM and echo analyses**, searching gravitational wave data (LIGO/Virgo, and especially LISA) for non-Kerr structures.
2. **High-precision accretion modeling**, incorporating hard boundaries into X-ray and EHT studies.
3. **Numerical modeling of matter shells**, within both classical and modified GR frameworks.
4. **Exploration of shell mechanics and energy conditions**, particularly in dynamical collapse scenarios.

Together, these constitute perhaps the most testable implications of the discrete standing wave model.

12 Open Questions and Theoretical Tensions

If Part III set out the architecture of a discrete standing wave spacetime, and the previous section traced its observational fingerprints, the present section steps back. It asks not what the model *predicts*, but what it *requires*. The questions below are not failures but frontiers:

invitations to mathematical refinement, physical insight, and perhaps even experimental discovery.

12.1 Does a Discrete Spacetime Demand a Reformulation of Quantum Mechanics?

Quantum theory presumes a continuous background for wavefunction evolution, with time treated as a smooth parameter. If time instead emerges from an oscillatory, discrete substrate, can the Schrödinger equation still apply? Or must a new operator formalism be constructed, one adapted to a fundamentally discrete temporal flow?

12.2 How Does Gravity Arise from Discrete Geometry?

Here, gravity emerges not as a force but as a pressure gradient driven by temporal differentials. Yet a first-principles derivation remains absent. Is there a discrete analog to curvature, or to the stress-energy tensor, from which field equations can be derived? This is essential for a full reconciliation with general relativity.

12.3 What is the Role of the Higgs Field in a Resonant Spacetime?

If inertial mass results from impedance against spacetime oscillation, the Higgs mechanism may itself be secondary. Is the Higgs field simply another standing wave mode? If so, this could recast spontaneous symmetry breaking not as a brute fact, but as a resonance phenomenon within the fabric of spacetime.

12.4 Can Vacuum Energy Be Regularized by Discreteness?

The vacuum energy catastrophe reflects a failure of continuum theory. In a discrete standing wave model, both a minimum frequency and a maximum warp depth exist. These natural cutoffs may eliminate infinitesimal mode contributions, suggesting a route to resolving the paradox.

12.5 Is Cosmological Redshift Partially Misattributed?

If $c = c(V)$, then cosmic redshift may not arise solely from metric expansion. Some fraction could reflect historical variation in c , lessening or removing the need for dark energy. The convexity of supernova Hubble plots may hint at precisely this effect.

12.6 How Does Information Escape a Black Surface?

Unlike an event horizon, a Black Surface is a finite shell. If the interior is not a true void, tunneling or resonant leakage may allow information escape, preserving unitarity without paradox. This raises the prospect of reconciling quantum mechanics with collapse without firewalls or singularities.

12.7 Does Discreteness Imply a Preferred Scale or Frame?

A fundamental oscillation frequency might define a cosmological frame, apparently at odds with relativity. Yet local Lorentz invariance may survive as an emergent property of the wave structure, just as fluid dynamics inherits Galilean symmetry despite molecular discreteness.

12.8 Is There a Boundary to the Universe or Its Wave Domain?

A standing wave requires boundary conditions. What sets these in an expanding universe? Possibilities include holographic boundaries, topological closure, or embedding in a higher-dimensional domain. The observable universe may itself be only a subharmonic of a larger wave domain.

12.9 Is This Model Testable?

Ultimately, a theory stands or falls by experiment. Observable signatures may include:

- Gravitational lensing near Black Surfaces deviating from Kerr predictions.
- Primordial light or gravitational waves exhibiting coherence limits inconsistent with a continuous metric.
- QNM ringdowns with echo structures absent from general relativity.

Discreteness is either invisible—or it is everything.

References

- [1] Robert M. Wald. *General Relativity*. University of Chicago Press, 1984. ISBN 9780226870335.
- [2] Albert Einstein. Die feldgleichungen der gravitation. *Sitzungsberichte der Preussischen Akademie der Wissenschaften zu Berlin*, pages 844–847, 1915.
- [3] Stephen W. Hawking. Breakdown of predictability in gravitational collapse. *Physical Review D*, 14(10):2460–2473, 1976. doi: 10.1103/PhysRevD.14.2460.
- [4] Vera C. Rubin and W. Kent Ford Jr. Rotation of the andromeda nebula from a spectroscopic survey of emission regions. *The Astrophysical Journal*, 159:379–403, 1970. doi: 10.1086/150317.
- [5] Fritz Zwicky. Die rotverschiebung von extragalaktischen nebeln. *Helvetica Physica Acta*, 6:110–127, 1933.
- [6] Saul Perlmutter et al. Measurements of ω and λ from 42 high-redshift supernovae. *The Astrophysical Journal*, 517(2):565–586, 1999. doi: 10.1086/307221.
- [7] Adam G. Riess et al. Observational evidence from supernovae for an accelerating universe and a cosmological constant. *The Astronomical Journal*, 116(3):1009–1038, 1998. doi: 10.1086/300499.
- [8] Brian P. Schmidt and et al. The high-z supernova search: Measuring cosmic deceleration and global curvature of the universe using type ia supernovae. *The Astrophysical Journal*, 507(1):46–63, 1998. doi: 10.1086/306308.
- [9] Gerard 't Hooft. Dimensional reduction in quantum gravity. *arXiv preprint gr-qc/9310026*, 1993. Presented at the Salamfest, Trieste.
- [10] Juan M. Maldacena. The large n limit of superconformal field theories and supergravity. *International Journal of Theoretical Physics*, 38:1113–1133, 1999. doi: 10.1023/A:1026654312961.
- [11] Edwin Hubble. A relation between distance and radial velocity among extra-galactic nebulae. *Proceedings of the National Academy of Sciences*, 15(3):168–173, 1929. doi: 10.1073/pnas.15.3.168.
- [12] G. F. Smoot et al. Structure in the cobe differential microwave radiometer first-year maps. *The Astrophysical Journal*, 396:L1–L5, 1992. doi: 10.1086/186504.
- [13] Alan H. Guth. Inflationary universe: A possible solution to the horizon and flatness problems. *Physical Review D*, 23(2):347–356, 1981. doi: 10.1103/PhysRevD.23.347.
- [14] George R. Blumenthal, Sandra M. Faber, Joel R. Primack, and Martin J. Rees. Formation of galaxies and large-scale structure with cold dark matter. *Nature*, 311:517–525, 1984. doi: 10.1038/311517a0.
- [15] Planck Collaboration. Planck 2018 results. vi. cosmological parameters. *Astronomy & Astrophysics*, 641:A6, 2020. doi: 10.1051/0004-6361/201833910.

- [16] Robert R. Caldwell, Marc Kamionkowski, and Nevin N. Weinberg. Phantom energy and cosmic doomsday. *Physical Review Letters*, 91(7):071301, 2003. doi: 10.1103/PhysRevLett.91.071301.
- [17] LibreTexts. 2.2: The postulates of quantum mechanics, 2023. URL [https://chem.libretexts.org/Bookshelves/Physical_and_Theoretical_Chemistry_Textbook_Maps/Quantum_Chemistry_with_Applications_in_Spectroscopy_\(Fleming\)/02:_Particle_in_a_Box/2.02:_The_Postulates_of_Quantum_Mechanics](https://chem.libretexts.org/Bookshelves/Physical_and_Theoretical_Chemistry_Textbook_Maps/Quantum_Chemistry_with_Applications_in_Spectroscopy_(Fleming)/02:_Particle_in_a_Box/2.02:_The_Postulates_of_Quantum_Mechanics). Accessed: 2025-04-21.
- [18] Encyclopædia Britannica. Event horizon — definition, explanation, & facts, 2025. URL <https://www.britannica.com/topic/event-horizon-black-hole>. Accessed: 2025-04-21.
- [19] Karl Schwarzschild. Über das gravitationsfeld eines massenpunktes nach der einsteinschen theorie. *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften*, pages 189–196, 1916.
- [20] Wikipedia contributors. No-hair theorem, 2025. URL https://en.wikipedia.org/wiki/No-hair_theorem. Accessed: 2025-04-21.
- [21] Stephen W. Hawking and Roger Penrose. The singularities of gravitational collapse and cosmology. *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences*, 314(1519):529–548, 1970. doi: 10.1098/rspa.1970.0021.
- [22] Charles W. Misner, Kip S. Thorne, and John Archibald Wheeler. *Gravitation*. W. H. Freeman, 1973. ISBN 978-0716703440.
- [23] Giovanni Amelino-Camelia. Relativity in space-times with short-distance structure governed by an observer-independent (planckian) length scale. *International Journal of Modern Physics D*, 11(1):35–60, 2002. doi: 10.1142/S0218271802001330.
- [24] Albert Einstein. Zur elektrodynamik bewegter körper. *Annalen der Physik*, 322(10): 891–921, 1905. doi: 10.1002/andp.19053221004. English translation: On the Electrodynamics of Moving Bodies.
- [25] R. P. Feynman, F. B. Morinigo, and W. G. Wagner. *Feynman Lectures on Gravitation*. Addison-Wesley, 1963.
- [26] Steven Weinberg. Photons and gravitons in perturbation theory: Derivation of maxwell's and einstein's equations. *Physical Review*, 138(4B):B988–B1002, 1965.
- [27] Wikipedia contributors. Graviton. <https://en.wikipedia.org/wiki/Graviton>. Accessed 2025-08-20.
- [28] Albert Einstein. The foundation of the general theory of relativity. *Annalen der Physik*, 49:769–822, 1916.
- [29] Charles W. Misner, Kip S. Thorne, and John Archibald Wheeler. *Gravitation*. W. H. Freeman, 1973.

- [30] Albert Einstein. On the principle of relativity and its consequences in modern physics. *Jahrbuch der Radioaktivität und Elektronik*, 4:411–462, 1907.
- [31] Clifford M. Will. *The Confrontation between General Relativity and Experiment*, volume 17. Living Reviews in Relativity, 2014.
- [32] Peter W. Higgs. Broken symmetries and the masses of gauge bosons. *Physical Review Letters*, 13:508–509, 1964.
- [33] François Englert and Robert Brout. Broken symmetry and the mass of gauge vector mesons. *Physical Review Letters*, 13:321–323, 1964.
- [34] Roger Penrose. Gravitational collapse and space-time singularities. *Phys. Rev. Lett.*, 14:57–59, 1965.
- [35] K. D. Kokkotas and B. G. Schmidt. Quasi-normal modes of stars and black holes. *Living Rev. Relativity*, 2:2, 1999.
- [36] Andrew King. Black holes, galaxy formation, and the mbh–sigma relation. *Astrophysical Journal Letters*, 635:L121–L123, 2006.
- [37] Feryal Özel, Dimitrios Psaltis, and Ramesh Narayan. Black hole masses and the fundamental plane of black hole accretion. *Astrophysical Journal*, 725(2):1918–1927, 2010.
- [38] Event Horizon Telescope Collaboration. First m87 event horizon telescope results. i. the shadow of the supermassive black hole. *Astrophysical Journal Letters*, 875(1):L1, 2019.
- [39] LIGO Scientific Collaboration and Virgo Collaboration. Observation of gravitational waves from a binary black hole merger. *Phys. Rev. Lett.*, 116:061102, 2016.
- [40] Vitor Cardoso, Edgardo Franzin, and Paolo Pani. Is the gravitational-wave ringdown a probe of the event horizon? *Phys. Rev. Lett.*, 116:171101, 2016.
- [41] Alexander Friedmann, Howard P. Robertson, and Arthur G. Walker. Foundational papers on the expanding universe and cosmological metrics. *Collected references*, 1922–1937. See Friedmann (1922, 1924); Robertson (1935); Walker (1937).
- [42] Andreas Albrecht and Joao Magueijo. A time varying speed of light as a solution to cosmological puzzles. *Physical Review D*, 59(4):043516, 1999. doi: 10.1103/PhysRevD.59.043516.
- [43] John D. Barrow. Cosmologies with varying light speed. *Physical Review D*, 59(4):043515, 1999. doi: 10.1103/PhysRevD.59.043515.
- [44] Joao Magueijo. *Faster Than the Speed of Light: The Story of a Scientific Speculation*. Perseus Publishing, 2003. ISBN 9780738205250.
- [45] Steven Weinberg. The cosmological constant problem. *Reviews of Modern Physics*, 61(1):1–23, 1989. doi: 10.1103/RevModPhys.61.1.
- [46] Andrew R. Liddle and David H. Lyth. *Cosmological Inflation and Large-Scale Structure*. Cambridge University Press, Cambridge, UK, 2000.

- [47] Viatcheslav F. Mukhanov, H. A. Feldman, and R. H. Brandenberger. Theory of cosmological perturbations. *Physics Reports*, 215(5-6):203–333, 1992. doi: 10.1016/0370-1573(92)90044-Z.
- [48] Daniel J. Eisenstein et al. Detection of the baryon acoustic peak in the large-scale correlation function of sdss luminous red galaxies. *Astrophysical Journal*, 633:560–574, 2005. doi: 10.1086/466512.
- [49] Martin White. Baryon acoustic oscillations: A standard ruler method for determining the expansion rate of the universe. *New Astronomy Reviews*, 49:360–366, 2005. doi: 10.1016/j.newar.2005.05.005.
- [50] P. J. E. Peebles. *The Large-Scale Structure of the Universe*. Princeton University Press, 1980.
- [51] J. Adams et al. Early massive galaxy formation revealed by jwst. *Nature Astronomy*, 2023. doi: 10.1038/s41550-023-01958-0.
- [52] R. Naidu et al. Massive galaxies at cosmic dawn: Evidence for rapid early formation. *Astrophysical Journal Letters*, 939:L15, 2022. doi: 10.3847/2041-8213/ac9e44.
- [53] Sean M. Carroll. *Spacetime and Geometry: An Introduction to General Relativity*. Addison-Wesley, San Francisco, CA, 2004. ISBN 978-0805387329.
- [54] R. K. Sachs and A. M. Wolfe. Perturbations of a cosmological model and angular variations of the microwave background. *Astrophysical Journal*, 147:73–90, 1967. doi: 10.1086/148982.
- [55] Scott Dodelson. *Modern Cosmology*. Academic Press, San Diego, CA, 2003. ISBN 978-0122191411.