MATH 180 Strategies of Problem Solving

W.R. Casper

Department of Mathematics
California State University Fullerton

October 28, 2026

Outline

1. SoPS Lecture 19
   - Functions
   - Cardinality
   - Wrapping up

Review

Last time

Review

Last time

- Sets, Subsets, Intersection, Union, Complements

Review

Last time

- Sets, Subsets, Intersection, Union, Complements
- Cardinality of finite sets, power set

Review

Last time

- Sets, Subsets, Intersection, Union, Complements
- Cardinality of finite sets, power set
- Universal quantifiers, Russell's paradox

Outline

1. SoPS Lecture 19
   - Functions
   - Cardinality
   - Wrapping up

Intuition



Intuition


THE SANDWICH FUNCTION
SANDWICH MACHINE
SANDWICH MACHINE
SANDWICH MACHINE
tuna
cheese
tomato
tuna sandwich
cheese sandwich
tomato sandwich
Ingredient in. Matching sandwich out.

Intuition


A BRICK IS OUTSIDE THE DOMAIN
Allowed inputs (ingredients):
tuna
cheese
tomato
...
Ingredients only!
SANDWICH FUNCTION
No output is defined.
A brick is not an allowed input.

Relations

Relations

Definition

A **relation** $\mathcal{R}$ from A to B is a subset of the Cartesian product $A \times B$.

Relations

Definition

A **relation** $\mathcal{R}$ from A to B is a subset of the Cartesian product $A \times B$.

The **domain** and **range** of $\mathcal{R}$ are

$$\text{Dom}(\mathcal{R}) = \{a \in A : \exists b \in B,\ (a, b) \in \mathcal{R}\}$$

$$\text{Rng}(\mathcal{R}) = \{b \in B : \exists a \in A,\ (a, b) \in \mathcal{R}\}$$

Relations

Relations

Example

Relations

Example

$$\mathcal{R} = \{(2, \heartsuit), (5, \diamondsuit), (5, \spadesuit), (7, \heartsuit)\}$$

is a relation from A to B for

Relations

Example

$$\mathcal{R} = \{(2, \heartsuit), (5, \diamondsuit), (5, \spadesuit), (7, \heartsuit)\}$$

is a relation from A to B for

$$A = \{2, 3, 4, 5, 6, 7, 8, 9, 10, J, Q, K, A\}$$

Relations

Example

$$\mathcal{R} = \{(2, \heartsuit), (5, \diamondsuit), (5, \spadesuit), (7, \heartsuit)\}$$

is a relation from A to B for

$$A = \{2, 3, 4, 5, 6, 7, 8, 9, 10, J, Q, K, A\}$$

$$B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$$

Relations

Example

$$\mathcal{R} = \{(2, \heartsuit), (5, \diamondsuit), (5, \spadesuit), (7, \heartsuit)\}$$

is a relation from A to B for

$$A = \{2, 3, 4, 5, 6, 7, 8, 9, 10, J, Q, K, A\}$$

$$B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$$

Domain and range:

Relations

Example

$$\mathcal{R} = \{(2, \heartsuit), (5, \diamondsuit), (5, \spadesuit), (7, \heartsuit)\}$$

is a relation from A to B for

$$A = \{2, 3, 4, 5, 6, 7, 8, 9, 10, J, Q, K, A\}$$

$$B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$$

Domain and range:

$$\text{Dom}(\mathcal{R}) = \{2, 5, 7\}$$

Relations

Example

$$\mathcal{R} = \{(2, \heartsuit), (5, \diamondsuit), (5, \spadesuit), (7, \heartsuit)\}$$

is a relation from A to B for

$$A = \{2, 3, 4, 5, 6, 7, 8, 9, 10, J, Q, K, A\}$$

$$B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$$

Domain and range:

$$\text{Dom}(\mathcal{R}) = \{2, 5, 7\}$$

$$\text{Rng}(\mathcal{R}) = \{\heartsuit, \diamondsuit, \spadesuit\}$$

Relations

Relations

Example

Relations

Example

$$\mathcal{S} = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2/4 = 1\}$$

is a relation from $\mathbb{R}$ to $\mathbb{R}$.

Relations

Example

$$\mathcal{S} = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2/4 = 1\}$$

is a relation from $\mathbb{R}$ to $\mathbb{R}$.

It describes an ellipse!

Relations

Example

$$\mathcal{S} = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2/4 = 1\}$$

is a relation from $\mathbb{R}$ to $\mathbb{R}$.

It describes an ellipse!
Domain and range:

Relations

Example

$$\mathcal{S} = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2/4 = 1\}$$

is a relation from $\mathbb{R}$ to $\mathbb{R}$.

It describes an ellipse!
Domain and range:

$$\text{Dom}(\mathcal{S}) = [-1, 1]$$

Relations

Example

$$\mathcal{S} = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2/4 = 1\}$$

is a relation from $\mathbb{R}$ to $\mathbb{R}$.

It describes an ellipse!
Domain and range:

$$\text{Dom}(\mathcal{S}) = [-1, 1]$$

$$\text{Rng}(\mathcal{S}) = [-2, 2]$$

Relations

Relations

Example

Relations

Example

$$\mathcal{T} = \{(A, B) : \text{A,B are sets of integers and } A \subseteq B\}$$

is a relation from $\mathcal{P}(\mathbb{Z})$ to $\mathcal{P}(\mathbb{Z})$.

Relations

Example

$$\mathcal{T} = \{(A, B) : \text{A,B are sets of integers and } A \subseteq B\}$$

is a relation from $\mathcal{P}(\mathbb{Z})$ to $\mathcal{P}(\mathbb{Z})$.

$$(\{1\}, \{1, 2\}) \in \mathcal{T}, \quad (\varnothing, \{2, 3, 5\}) \in \mathcal{T}, \quad (\{2\}, \{1\}) \notin \mathcal{T}$$

Relations

Example

$$\mathcal{T} = \{(A, B) : \text{A,B are sets of integers and } A \subseteq B\}$$

is a relation from $\mathcal{P}(\mathbb{Z})$ to $\mathcal{P}(\mathbb{Z})$.

$$(\{1\}, \{1, 2\}) \in \mathcal{T}, \quad (\varnothing, \{2, 3, 5\}) \in \mathcal{T}, \quad (\{2\}, \{1\}) \notin \mathcal{T}$$

Domain and range:

Relations

Example

$$\mathcal{T} = \{(A, B) : \text{A,B are sets of integers and } A \subseteq B\}$$

is a relation from $\mathcal{P}(\mathbb{Z})$ to $\mathcal{P}(\mathbb{Z})$.

$$(\{1\}, \{1, 2\}) \in \mathcal{T}, \quad (\varnothing, \{2, 3, 5\}) \in \mathcal{T}, \quad (\{2\}, \{1\}) \notin \mathcal{T}$$

Domain and range:

$$\text{Dom}(\mathcal{T}) = \mathcal{P}(\mathbb{Z})$$

Relations

Example

$$\mathcal{T} = \{(A, B) : \text{A,B are sets of integers and } A \subseteq B\}$$

is a relation from $\mathcal{P}(\mathbb{Z})$ to $\mathcal{P}(\mathbb{Z})$.

$$(\{1\}, \{1, 2\}) \in \mathcal{T}, \quad (\varnothing, \{2, 3, 5\}) \in \mathcal{T}, \quad (\{2\}, \{1\}) \notin \mathcal{T}$$

Domain and range:

$$\text{Dom}(\mathcal{T}) = \mathcal{P}(\mathbb{Z})$$

$$\text{Rng}(\mathcal{T}) = \mathcal{P}(\mathbb{Z})$$

Relation notation

Notation

We write $a\mathcal{R}b$ as another way of saying $(a, b) \in \mathcal{R}$.

Relation notation

Notation

We write $a\mathcal{R}b$ as another way of saying $(a, b) \in \mathcal{R}$.

Example: if $\mathcal{R} = \{(1, 3), (2, 1), (0, 0)\}$ then

Relation notation

Notation

We write $a\mathcal{R}b$ as another way of saying $(a, b) \in \mathcal{R}$.

Example: if $\mathcal{R} = \{(1, 3), (2, 1), (0, 0)\}$ then

- $1\mathcal{R}3$ is **true**

Relation notation

Notation

We write $a\mathcal{R}b$ as another way of saying $(a, b) \in \mathcal{R}$.

Example: if $\mathcal{R} = \{(1, 3), (2, 1), (0, 0)\}$ then

- $1\mathcal{R}3$ is **true**
- $1\mathcal{R}2$ is **false**

Relation notation

Notation

We write $a\mathcal{R}b$ as another way of saying $(a, b) \in \mathcal{R}$.

Example: if $\mathcal{R} = \{(1, 3), (2, 1), (0, 0)\}$ then

- $1\mathcal{R}3$ is **true**
- $1\mathcal{R}2$ is **false**
- $0\mathcal{R}0$ is **true**

Relation notation

Notation

We write $a\mathcal{R}b$ as another way of saying $(a, b) \in \mathcal{R}$.

Example: if $\mathcal{R} = \{(1, 3), (2, 1), (0, 0)\}$ then

- $1\mathcal{R}3$ is **true**
- $1\mathcal{R}2$ is **false**
- $0\mathcal{R}0$ is **true**
- $1\mathcal{R}1$ is **false**

Functions

Functions

Definition

Functions

Definition

A function f from A to B is a relation from A to B satisfying:

Functions

Definition

A function f from A to B is a relation from A to B satisfying:

- for each $a \in A$ there exists $b \in B$ with $(a, b) \in f$

Functions

Definition

A function f from A to B is a relation from A to B satisfying:

- for each $a \in A$ there exists $b \in B$ with $(a, b) \in f$
- for each $a \in A$, if both $(a, b) \in f$ and $(a, c) \in f$ then $b = c$

Functions

Definition

A function f from A to B is a relation from A to B satisfying:

- for each $a \in A$ there exists $b \in B$ with $(a, b) \in f$
- for each $a \in A$, if both $(a, b) \in f$ and $(a, c) \in f$ then $b = c$

Notation:

Functions

Definition

A function f from A to B is a relation from A to B satisfying:

- for each $a \in A$ there exists $b \in B$ with $(a, b) \in f$
- for each $a \in A$, if both $(a, b) \in f$ and $(a, c) \in f$ then $b = c$

Notation:

- we write $f : A \to B$

Functions

Definition

A function f from A to B is a relation from A to B satisfying:

- for each $a \in A$ there exists $b \in B$ with $(a, b) \in f$
- for each $a \in A$, if both $(a, b) \in f$ and $(a, c) \in f$ then $b = c$

Notation:

- we write $f : A \to B$
- redundant notation!

Functions

Definition

A function f from A to B is a relation from A to B satisfying:

- for each $a \in A$ there exists $b \in B$ with $(a, b) \in f$
- for each $a \in A$, if both $(a, b) \in f$ and $(a, c) \in f$ then $b = c$

Notation:

- we write $f : A \to B$
- redundant notation!

$$f(a) = b \quad afb \quad \text{and} \quad (a, b) \in f$$

all mean the same thing!

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit), (3, \diamondsuit)\}$$

is a function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit), (3, \diamondsuit)\}$$

is a function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

- everything in A is in the domain

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit), (3, \diamondsuit)\}$$

is a function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

- everything in A is in the domain
- everything gets sent to only one place

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit), (3, \diamondsuit)\}$$

is a function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

- everything in A is in the domain
- everything gets sent to only one place
- we can write $f : A \to B$

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit), (3, \diamondsuit)\}$$

is a function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

- everything in A is in the domain
- everything gets sent to only one place
- we can write $f : A \to B$
- we can write $f(1) = \heartsuit$ or $1 f \heartsuit$ or $(1, \heartsuit) \in f$

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit), (1, \diamondsuit), (3, \clubsuit)\}$$

is NOT function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit), (1, \diamondsuit), (3, \clubsuit)\}$$

is NOT function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

- everything in A is in the domain

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit), (1, \diamondsuit), (3, \clubsuit)\}$$

is NOT function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

- everything in A is in the domain
- however! f tries to send 1 to both $\heartsuit$ and to $\diamondsuit$

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit), (1, \diamondsuit), (3, \clubsuit)\}$$

is NOT function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

- everything in A is in the domain
- however! f tries to send 1 to both $\heartsuit$ and to $\diamondsuit$
- $1 f \heartsuit$ and $1 f \diamondsuit$

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit), (1, \diamondsuit), (3, \clubsuit)\}$$

is NOT function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

- everything in A is in the domain
- however! f tries to send 1 to both $\heartsuit$ and to $\diamondsuit$
- $1f\heartsuit$ and $1f\diamondsuit$
- $f(1) = \heartsuit$ and $f(1) = \diamondsuit$

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit), (1, \diamondsuit), (3, \clubsuit)\}$$

is NOT function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

- everything in A is in the domain
- however! f tries to send 1 to both $\heartsuit$ and to $\diamondsuit$
- $1 f \heartsuit$ and $1 f \diamondsuit$
- $f(1) = \heartsuit$ and $f(1) = \diamondsuit$

NOT a function!

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit)\}$$

is NOT function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit)\}$$

is NOT function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

- everything gets sent to only one place

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit)\}$$

is NOT function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

- everything gets sent to only one place
- NOT everything in A is in the domain

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit)\}$$

is NOT function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

- everything gets sent to only one place
- NOT everything in A is in the domain
- $f(3)$ isn't defined!

Functions

Example

The relation

$$f = \{(1, \heartsuit), (2, \spadesuit)\}$$

is NOT function from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

- everything gets sent to only one place
- NOT everything in A is in the domain
- $f(3)$ isn't defined!

NOT a function!

Functions

Challenge

Find how many functions there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Functions

Challenge

Find how many functions there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Solution

Functions

Challenge

Find how many functions there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Solution

Just have to decide the value of $f(1)$, $f(2)$, and $f(3)$.

Functions

Challenge

Find how many functions there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Solution

Just have to decide the value of $f(1)$, $f(2)$, and $f(3)$.

$$4 \times 4 \times 4 = 64 \text{ possibilities.}$$

Functions

Challenge

Find how many relations there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Functions

Challenge

Find how many relations there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Solution

Functions

Challenge

Find how many relations there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Solution

How many subsets of $A \times B$?

Functions

Challenge

Find how many relations there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Solution

How many subsets of $A \times B$?
How many elements in $\mathcal{P}(A \times B)$?

Functions

Challenge

Find how many relations there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Solution

How many subsets of $A \times B$?
How many elements in $\mathcal{P}(A \times B)$?

$$|\mathcal{P}(A \times B)| = 2^{|A \times B|} = 2^{3 \times 4} = 4096$$

Rules

Defining functions by rules

Often, we define a function by specifying a **rule** for what $f(x)$ sends a value x to.

Rules

Defining functions by rules

Often, we define a function by specifying a **rule** for what $f(x)$ sends a value x to.

Examples:

Rules

Defining functions by rules

Often, we define a function by specifying a **rule** for what $f(x)$ sends a value x to.

Examples:

- $f : \mathbb{R} \to \mathbb{R}$ defined by $f(x) = x^2$

Rules

Defining functions by rules

Often, we define a function by specifying a **rule** for what $f(x)$ sends a value x to.

Examples:

- $f : \mathbb{R} \to \mathbb{R}$ defined by $f(x) = x^2$
- $f : \mathbb{Z} \to \mathbb{Q}$ defined by

$$f(x) = \begin{cases} \frac{2x+1}{x-2}, & x \neq 2 \\ 3, & x = 2 \end{cases}$$

Outline

1. SoPS Lecture 19
   - Functions
   - Cardinality
   - Wrapping up

Properties of functions

Properties of functions

Definition

A function $f : A \to B$ is called

Properties of functions

Definition

A function $f : A \to B$ is called

- **injective** or **one-to-one** if for all $a_1, a_2 \in A$ if $f(a_1) = f(a_2)$ then $a_1 = a_2$

Properties of functions

Definition

A function $f : A \to B$ is called

- **injective** or **one-to-one** if for all $a_1, a_2 \in A$ if $f(a_1) = f(a_2)$ then $a_1 = a_2$
- **surjective** or **onto** if for all $b \in B$ there exists $a \in A$ with $f(a) = b$

Properties of functions

Definition

A function $f : A \to B$ is called

- **injective** or **one-to-one** if for all $a_1, a_2 \in A$ if $f(a_1) = f(a_2)$ then $a_1 = a_2$
- **surjective** or **onto** if for all $b \in B$ there exists $a \in A$ with $f(a) = b$
- **bijective** if its both injective and surjective

Properties of functions

These are statements about the range!

Properties of functions

These are statements about the range!

Properties of functions

These are statements about the range!

- **injective** means each $b \in B$ is hit **at most one time**

Properties of functions

These are statements about the range!

- **injective** means each $b \in B$ is hit **at most one time**
- **surjective** means each $b \in B$ is hit **at least one time**

Properties of functions

These are statements about the range!

- **injective** means each $b \in B$ is hit **at most one time**
- **surjective** means each $b \in B$ is hit **at least one time**
- **bijective** means each $b \in B$ is hit **exactly one time**

Example

Problem

What properties does the following function have?

$$f : \mathbb{R} \to \mathbb{R}, \quad f(x) = x^2$$

Example

Problem

What properties does the following function have?

$$f : \mathbb{R} \to \mathbb{R}, \quad f(x) = x^2$$

Solution

Neither injective nor surjective

Example

Problem

What properties does the following function have?

$$f : \mathbb{R} \to [0, \infty), \quad f(x) = x^2$$

Example

Problem

What properties does the following function have?

$$f : \mathbb{R} \to [0, \infty), \quad f(x) = x^2$$

Solution

Surjective but not injective

Example

Problem

What properties does the following function have?

$$f : (-\infty, 0] \to [0, \infty), \quad f(x) = x^2$$

Example

Problem

What properties does the following function have?

$$f : (-\infty, 0] \to [0, \infty), \quad f(x) = x^2$$

Solution

Both injective and surjective. Bijective!

Functions

Challenge

Find how many injective functions there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Functions

Challenge

Find how many injective functions there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Solution

Functions

Challenge

Find how many injective functions there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Solution

Choose where 1 goes. Then choose somewhere different for 2. Then yet another different place for 3.
Possibilities are $4 \times 3 \times 2 = 24$.

Functions

Challenge

Find how many surjective functions there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Functions

Challenge

Find how many surjective functions there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Solution

Functions

Challenge

Find how many surjective functions there are from $A = \{1, 2, 3\}$ to $B = \{\heartsuit, \clubsuit, \diamondsuit, \spadesuit\}$.

Solution

None!

Cardinality

Definition

We say two sets have the same **cardinality** if there exists a bijection $f : A \to B$

- for finite sets, bijection iff $|A| = |B|$
- for infinite sets, things get interesting

Hilbert's Infinite Hotel

Problem

David Hilbert owns a rather particular hotel with infinitely many rooms labeled $1, 2, 3, \ldots$.
One night, the hotel is completely booked when a guest shows up and asks for a room. Hilbert says "No problem!" and makes room for the new guest. How does he do it?

Hilbert's Infinite Hotel

Hilbert's Infinite Hotel

Hilbert's Infinite Hotel

Hilbert's Infinite Hotel

Mathematically, we've built a bijection!

$$f : \mathbb{N} \cup \{0\} \to \mathbb{N}$$

$$f(x) = x + 1$$

The new guest is 0. The old guests are in rooms 2 and up. The new guest is in room 1.

Hilbert's Infinite Hotel

Problem

Later the same night, a bus full of infinitely more guests show up. Hilbert makes room again! How does he do it?

Hilbert's Infinite Hotel

Hilbert's Infinite Hotel

Hilbert's Infinite Hotel

Hilbert's Infinite Hotel

Mathematically, we've built another bijection!

$$f : \mathbb{Z} \to \mathbb{N}$$

$$f(x) = \begin{cases} 2x, & x > 0 \\ 2x + 1, & x \leq 0 \end{cases}$$

The old guests are in even rooms. The new guests are in odd rooms.

Infinite rooms ... infinite complaints

Outline

1. SoPS Lecture 19
   - Functions
   - Cardinality
   - Wrapping up

Final thoughts

Reflection

Final thoughts

Reflection

- Describe the difference between a relation and a function
- Give an example of a relation which is not a function. Explain why it isn't.
- Give an example of a function which is surjective but not injective.

Submit your responses by **Tonight!**