MATH 180 Strategies of Problem Solving

W.R. Casper

Department of Mathematics
California State University Fullerton

November 2, 2026

Outline

1. SoPS Lecture 20
   - Finite Sets
   - Pidgeonhole Principle
   - Wrapping up

Review

Last time

Review

Last time

- Relations, Functions

Review

Last time

- Relations, Functions
- Injective, Surjective, Bijective

Review

Last time

- Relations, Functions
- Injective, Surjective, Bijective
- Cardinality, Hilbert's Hotel

Outline

1. SoPS Lecture 20
   - Finite Sets
   - Pidgeonhole Principle
   - Wrapping up

Finite means what?

Question

How should we define a set to be **finite**?

Finite sets

We have some sets we know are finite.

Finite sets

We have some sets we know are finite.
Define

$$\mathbb{N}_1 = \{1\}, \quad \mathbb{N}_2 = \{1, 2\}, \quad \mathbb{N}_3 = \{1, 2, 3\}$$

Finite sets

We have some sets we know are finite.
Define

$$\mathbb{N}_1 = \{1\}, \quad \mathbb{N}_2 = \{1, 2\}, \quad \mathbb{N}_3 = \{1, 2, 3\}$$

In general, for any $k \in \mathbb{N}$ define

$$\mathbb{N}_k = \{1, 2, 3, \ldots, k\}.$$

Finite sets

We have some sets we know are finite.
Define

$$\mathbb{N}_1 = \{1\}, \quad \mathbb{N}_2 = \{1, 2\}, \quad \mathbb{N}_3 = \{1, 2, 3\}$$

In general, for any $k \in \mathbb{N}$ define

$$\mathbb{N}_k = \{1, 2, 3, \ldots, k\}.$$

Definition

A set A is called **finite** if it is empty or there exists a $k \in \mathbb{N}$ for which there is a bijection from $\mathbb{N}_k$ to A.

Example

Problem

Show that $\mathbb{N}_j$ is finite for all j.

Example

Problem

Show that $\mathbb{N}_j$ is finite for all j.

Must show that there's a $k \in \mathbb{N}$ with a bijection from $\mathbb{N}_k$ to $\mathbb{N}_j$.

Example

Problem

Show that $\mathbb{N}_j$ is finite for all j.

Must show that there's a $k \in \mathbb{N}$ with a bijection from $\mathbb{N}_k$ to $\mathbb{N}_j$.

Question

What value of k should we choose?

Example

Problem

Show that $\mathbb{N}_j$ is finite for all j.

Must show that there's a $k \in \mathbb{N}$ with a bijection from $\mathbb{N}_k$ to $\mathbb{N}_j$.

Question

What value of k should we choose?

Answer

We should choose $k = j$. That's the only value we should expect to have a bijection!

Example

Problem

Show that $\mathbb{N}_j$ is finite for all j.

Must show that there's a $k \in \mathbb{N}$ with a bijection from $\mathbb{N}_k$ to $\mathbb{N}_j$.

Question

What bijection is there from $\mathbb{N}_k$ to $\mathbb{N}_k$?

Example

Problem

Show that $\mathbb{N}_j$ is finite for all j.

Must show that there's a $k \in \mathbb{N}$ with a bijection from $\mathbb{N}_k$ to $\mathbb{N}_j$.

Question

What bijection is there from $\mathbb{N}_k$ to $\mathbb{N}_k$?

Answer

There's lots! We just need to demonstrate that there's at least one.

Example

Problem

Show that $\mathbb{N}_j$ is finite for all j.

Must show that there's a $k \in \mathbb{N}$ with a bijection from $\mathbb{N}_k$ to $\mathbb{N}_j$.

Question

What bijection is there from $\mathbb{N}_k$ to $\mathbb{N}_k$?

Answer

There's lots! We just need to demonstrate that there's at least one.
The **identity function**

$$f : \mathbb{N}_k \to \mathbb{N}_k, \quad f(x) = x$$

in particular is a bijection

Example

Problem

Show that $A = \{\heartsuit, \spadesuit, \diamondsuit, \clubsuit\}$ is finite.

Must show that there's a $k \in \mathbb{N}$ with a bijection from $\mathbb{N}_k$ to A.

Question

What k should we choose?

Example

Problem

Show that $A = \{\heartsuit, \spadesuit, \diamondsuit, \clubsuit\}$ is finite.

Must show that there's a $k \in \mathbb{N}$ with a bijection from $\mathbb{N}_k$ to A.

Question

What k should we choose?

Answer

We should choose $k = 4$.

Example

Problem

Show that $A = \{\heartsuit, \spadesuit, \diamondsuit, \clubsuit\}$ is finite.

Must show that there's a $k \in \mathbb{N}$ with a bijection from $\mathbb{N}_k$ to A.

Question

What bijection from $\mathbb{N}_4$ to A is there?

Example

Problem

Show that $A = \{\heartsuit, \spadesuit, \diamondsuit, \clubsuit\}$ is finite.

Must show that there's a $k \in \mathbb{N}$ with a bijection from $\mathbb{N}_k$ to A.

Question

What bijection from $\mathbb{N}_4$ to A is there?

Answer

There's lots! We just need to show that there's at least one.

Example

Problem

Show that $A = \{\heartsuit, \spadesuit, \diamondsuit, \clubsuit\}$ is finite.

Must show that there's a $k \in \mathbb{N}$ with a bijection from $\mathbb{N}_k$ to A.

Question

What bijection from $\mathbb{N}_4$ to A is there?

Answer

There's lots! We just need to show that there's at least one. For example, $f : \mathbb{N}_4 \to A$ defined by

$$f(1) = \heartsuit, \quad f(2) = \spadesuit, \quad f(3) = \diamondsuit, \quad f(4) = \clubsuit$$

Outline

1. SoPS Lecture 20
   - Finite Sets
   - Pidgeonhole Principle
   - Wrapping up

Starting Example

Starting Example

Question

Why is there no bijection from $\mathbb{N}_2$ to $\mathbb{N}_1$?

Starting Example

Question

Why is there no bijection from $\mathbb{N}_2$ to $\mathbb{N}_1$?

Solution

Starting Example

Question

Why is there no bijection from $\mathbb{N}_2$ to $\mathbb{N}_1$?

Solution

Take a function $f : \mathbb{N}_2 \to \mathbb{N}_1$.

Starting Example

Question

Why is there no bijection from $\mathbb{N}_2$ to $\mathbb{N}_1$?

Solution

Take a function $f : \mathbb{N}_2 \to \mathbb{N}_1$.

$$f : \{1, 2\} \to \{1\}$$

Starting Example

Question

Why is there no bijection from $\mathbb{N}_2$ to $\mathbb{N}_1$?

Solution

Take a function $f : \mathbb{N}_2 \to \mathbb{N}_1$.

$$f : \{1, 2\} \to \{1\}$$

$$f(1) = 1 \quad \text{and} \quad f(2) = 1$$

Starting Example

Question

Why is there no bijection from $\mathbb{N}_2$ to $\mathbb{N}_1$?

Solution

Take a function $f : \mathbb{N}_2 \to \mathbb{N}_1$.

$$f : \{1, 2\} \to \{1\}$$

$$f(1) = 1 \quad \text{and} \quad f(2) = 1$$

Therefore f fails to be injective.

Second Example

Question

Why is there no bijection from $\mathbb{N}_3$ to $\mathbb{N}_2$?

Second Example

Question

Why is there no bijection from $\mathbb{N}_3$ to $\mathbb{N}_2$?

Solution

Second Example

Question

Why is there no bijection from $\mathbb{N}_3$ to $\mathbb{N}_2$?

Solution

Take a function $f : \mathbb{N}_3 \to \mathbb{N}_2$.

Second Example

Question

Why is there no bijection from $\mathbb{N}_3$ to $\mathbb{N}_2$?

Solution

Take a function $f : \mathbb{N}_3 \to \mathbb{N}_2$.

$$f : \{1, 2, 3\} \to \{1, 2\}$$

Second Example

Question

Why is there no bijection from $\mathbb{N}_3$ to $\mathbb{N}_2$?

Solution

Take a function $f : \mathbb{N}_3 \to \mathbb{N}_2$.

$$f : \{1, 2, 3\} \to \{1, 2\}$$

If $f(1) = f(2)$, then not injective.

Second Example

Question

Why is there no bijection from $\mathbb{N}_3$ to $\mathbb{N}_2$?

Solution

Take a function $f : \mathbb{N}_3 \to \mathbb{N}_2$.

$$f : \{1, 2, 3\} \to \{1, 2\}$$

If $f(1) = f(2)$, then not injective.
Otherwise either $f(1) = 1, f(2) = 2$ or $f(1) = 2, f(2) = 1$.

Second Example

Question

Why is there no bijection from $\mathbb{N}_3$ to $\mathbb{N}_2$?

Solution

Take a function $f : \mathbb{N}_3 \to \mathbb{N}_2$.

$$f : \{1, 2, 3\} \to \{1, 2\}$$

If $f(1) = f(2)$, then not injective.
Otherwise either $f(1) = 1, f(2) = 2$ or $f(1) = 2, f(2) = 1$.
In either case, $f(3)$ must repeat one of the values!

Second Example

Question

Why is there no bijection from $\mathbb{N}_3$ to $\mathbb{N}_2$?

Solution

Take a function $f : \mathbb{N}_3 \to \mathbb{N}_2$.

$$f : \{1, 2, 3\} \to \{1, 2\}$$

If $f(1) = f(2)$, then not injective.
Otherwise either $f(1) = 1, f(2) = 2$ or $f(1) = 2, f(2) = 1$.
In either case, $f(3)$ must repeat one of the values!
Therefore f fails to be injective.

Pidgeonhole Principle

Theorem (Pidgeonhole Principle)

Let j and k be integers with $k > j$. If $f : \mathbb{N}_k \to \mathbb{N}_j$, then f is not injective.

Application

Problem

Show that at any party, two people in the party will always end up having the same number of acquaintances attending the party.

Application

Problem

Show that at any party, two people in the party will always end up having the same number of acquaintances attending the party.

Fine print:

Application

Problem

Show that at any party, two people in the party will always end up having the same number of acquaintances attending the party.

Fine print:

- we always consider someone to be their own acquaintance

Application

Solution

Application

Solution

Label the guests at the party as $1, 2, \ldots, n$.

Application

Solution

Label the guests at the party as $1, 2, \ldots, n$.
Define a function

$$f(k) = \text{number of people } k \text{ knows at the party.}$$

Application

Solution

Label the guests at the party as $1, 2, \ldots, n$.
Define a function

$$f(k) = \text{number of people } k \text{ knows at the party.}$$

Consider two cases:

Application

Solution

Label the guests at the party as $1, 2, \ldots, n$.
Define a function

$$f(k) = \text{number of people } k \text{ knows at the party.}$$

Consider two cases:

- nobody knows everyone at the party

Application

Solution

Label the guests at the party as $1, 2, \ldots, n$.
Define a function

$$f(k) = \text{number of people } k \text{ knows at the party.}$$

Consider two cases:

- nobody knows everyone at the party
- somebody knows everyone at the party

Application

Application

Solution

Application

Solution

If nobody knows everyone at the party, then the values of f are $1, 2, \ldots, n-1$.

Application

Solution

If nobody knows everyone at the party, then the values of f are $1, 2, \ldots, n-1$.
Therefore f is a function

$$f : \mathbb{N}_n \to \mathbb{N}_{n-1}.$$

Application

Solution

If nobody knows everyone at the party, then the values of f are $1, 2, \ldots, n-1$.
Therefore f is a function

$$f : \mathbb{N}_n \to \mathbb{N}_{n-1}.$$

The Pidgeonhole Principle says that f is not injective!

Application

Solution

If nobody knows everyone at the party, then the values of f are $1, 2, \ldots, n-1$.
Therefore f is a function

$$f : \mathbb{N}_n \to \mathbb{N}_{n-1}.$$

The Pidgeonhole Principle says that f is not injective!
Therefore there are two different values a and b with $f(a) = f(b)$.

Application

Solution

If nobody knows everyone at the party, then the values of f are $1, 2, \ldots, n-1$.
Therefore f is a function

$$f : \mathbb{N}_n \to \mathbb{N}_{n-1}.$$

The Pidgeonhole Principle says that f is not injective!
Therefore there are two different values a and b with $f(a) = f(b)$.
Guests a and b have the same number of acquaintances at the party!

Application

Solution

Application

Solution

If somebody knows everyone at the party, then the values of f are $2, 3, \ldots, n$.

Application

Solution

If somebody knows everyone at the party, then the values of f are $2, 3, \ldots, n$.
Define $g(x) = f(x) - 1$, which takes values $1, 2, \ldots, n-1$.

Application

Solution

If somebody knows everyone at the party, then the values of f are $2, 3, \ldots, n$.
Define $g(x) = f(x) - 1$, which takes values $1, 2, \ldots, n-1$.

$$g : \mathbb{N}_n \to \mathbb{N}_{n-1}.$$

Application

Solution

If somebody knows everyone at the party, then the values of f are $2, 3, \ldots, n$.
Define $g(x) = f(x) - 1$, which takes values $1, 2, \ldots, n-1$.

$$g : \mathbb{N}_n \to \mathbb{N}_{n-1}.$$

The Pidgeonhole Principle says that g is not injective!

Application

Solution

If somebody knows everyone at the party, then the values of f are $2, 3, \ldots, n$.
Define $g(x) = f(x) - 1$, which takes values $1, 2, \ldots, n-1$.

$$g : \mathbb{N}_n \to \mathbb{N}_{n-1}.$$

The Pidgeonhole Principle says that g is not injective!
Therefore there are two different values a and b with $g(a) = g(b)$.

Application

Solution

If somebody knows everyone at the party, then the values of f are $2, 3, \ldots, n$.
Define $g(x) = f(x) - 1$, which takes values $1, 2, \ldots, n-1$.

$$g : \mathbb{N}_n \to \mathbb{N}_{n-1}.$$

The Pidgeonhole Principle says that g is not injective!
Therefore there are two different values a and b with $g(a) = g(b)$.
This means $f(a) - 1 = f(b) - 1$, so $f(a) = f(b)$

Application

Solution

If somebody knows everyone at the party, then the values of f are $2, 3, \ldots, n$.
Define $g(x) = f(x) - 1$, which takes values $1, 2, \ldots, n-1$.

$$g : \mathbb{N}_n \to \mathbb{N}_{n-1}.$$

The Pidgeonhole Principle says that g is not injective!
Therefore there are two different values a and b with $g(a) = g(b)$.
This means $f(a) - 1 = f(b) - 1$, so $f(a) = f(b)$ Guests a and b have the same number of acquaintances at the party!

Application

Problem

Let $p = 8675309$. Prove that there is a number n of the form

$$n = 111111\ldots1$$

which is divisible by p.

Application

Problem

Let $p = 8675309$. Prove that there is a number n of the form

$$n = 111111\ldots1$$

which is divisible by p.

Observation:

Application

Problem

Let $p = 8675309$. Prove that there is a number n of the form

$$n = 111111\ldots1$$

which is divisible by p.

Observation:

- the number p is prime, which might explain why Jenny didn't want to change it.

Application

Application

Solution

Define $n_1 = 1$, $n_2 = 11$, $n_3 = 111$, and so on.
For each k, define r_k to be the remainder of n_k divided by p.

Application

Solution

Define $n_1 = 1$, $n_2 = 11$, $n_3 = 111$, and so on.
For each k, define r_k to be the remainder of n_k divided by p.

$$B = \{r_1, r_2, \ldots, r_{p+1}\} \quad \text{has at most } p \text{ elements}$$

Application

Solution

Define $n_1 = 1$, $n_2 = 11$, $n_3 = 111$, and so on.
For each k, define r_k to be the remainder of n_k divided by p.

$$B = \{r_1, r_2, \ldots, r_{p+1}\} \quad \text{has at most } p \text{ elements}$$

Pidgeonhole principle: there's $j < k$ with $r_j = r_k$.

Application

Solution

Define $n_1 = 1$, $n_2 = 11$, $n_3 = 111$, and so on.
For each k, define r_k to be the remainder of n_k divided by p.

$$B = \{r_1, r_2, \ldots, r_{p+1}\} \quad \text{has at most } p \text{ elements}$$

Pidgeonhole principle: there's $j < k$ with $r_j = r_k$.

$$p \text{ divides } n_k - n_j = \overbrace{111\ldots1}^{(k-j) \text{ many}}\ \overbrace{00\ldots0}^{j \text{ many}} = n_{k-j} \cdot 10^j$$

Application

Solution

Define $n_1 = 1$, $n_2 = 11$, $n_3 = 111$, and so on.
For each k, define r_k to be the remainder of n_k divided by p.

$$B = \{r_1, r_2, \ldots, r_{p+1}\} \quad \text{has at most } p \text{ elements}$$

Pidgeonhole principle: there's $j < k$ with $r_j = r_k$.

$$p \text{ divides } n_k - n_j = \overbrace{111\ldots1}^{(k-j) \text{ many}}\ \overbrace{00\ldots0}^{j \text{ many}} = n_{k-j} \cdot 10^j$$

Therefore p divides $n_{k-j} = \overbrace{111\ldots1}^{(k-j) \text{ many}}$

Natural numbers are infinite

Natural numbers are infinite

Theorem

The set $\mathbb{N}$ of natural numbers are infinite.

Natural numbers are infinite

Theorem

The set $\mathbb{N}$ of natural numbers are infinite.

Proof

To see this, suppose that $\mathbb{N}$ was finite.

Natural numbers are infinite

Theorem

The set $\mathbb{N}$ of natural numbers are infinite.

Proof

To see this, suppose that $\mathbb{N}$ was finite.
Then there would be an integer $k \in \mathbb{N}$ and a bijection between $\mathbb{N}$ and $\mathbb{N}_k$ for some $k \in \mathbb{N}$.

Natural numbers are infinite

Theorem

The set $\mathbb{N}$ of natural numbers are infinite.

Proof

To see this, suppose that $\mathbb{N}$ was finite.
Then there would be an integer $k \in \mathbb{N}$ and a bijection between $\mathbb{N}$ and $\mathbb{N}_k$ for some $k \in \mathbb{N}$.

$$f : \mathbb{N} \to \mathbb{N}_k \quad \text{bijection}$$

Natural numbers are infinite

Theorem

The set $\mathbb{N}$ of natural numbers are infinite.

Proof

To see this, suppose that $\mathbb{N}$ was finite.
Then there would be an integer $k \in \mathbb{N}$ and a bijection between $\mathbb{N}$ and $\mathbb{N}_k$ for some $k \in \mathbb{N}$.

$$f : \mathbb{N} \to \mathbb{N}_k \quad \text{bijection}$$

Use f to define a new function

$$g : \mathbb{N}_{k+1} \to \mathbb{N}_k, \quad g(k) = f(k).$$

Natural numbers are infinite

Natural numbers are infinite

Theorem

The set $\mathbb{N}$ of natural numbers are infinite.

Natural numbers are infinite

Theorem

The set $\mathbb{N}$ of natural numbers are infinite.

Proof

Natural numbers are infinite

Theorem

The set $\mathbb{N}$ of natural numbers are infinite.

Proof

If $g(x) = g(y)$, then $f(x) = f(y)$.

Natural numbers are infinite

Theorem

The set $\mathbb{N}$ of natural numbers are infinite.

Proof

If $g(x) = g(y)$, then $f(x) = f(y)$.
Since f is bijective, it's injective, so $x = y$.

Natural numbers are infinite

Theorem

The set $\mathbb{N}$ of natural numbers are infinite.

Proof

If $g(x) = g(y)$, then $f(x) = f(y)$.
Since f is bijective, it's injective, so $x = y$.
This shows g is injective.

Natural numbers are infinite

Theorem

The set $\mathbb{N}$ of natural numbers are infinite.

Proof

If $g(x) = g(y)$, then $f(x) = f(y)$.
Since f is bijective, it's injective, so $x = y$.
This shows g is injective.
Waaaaait a minute! This violates the Pidgeonhole Principle!

Natural numbers are infinite

Theorem

The set $\mathbb{N}$ of natural numbers are infinite.

Proof

If $g(x) = g(y)$, then $f(x) = f(y)$.
Since f is bijective, it's injective, so $x = y$.
This shows g is injective.
Waaaaait a minute! This violates the Pidgeonhole Principle!
Our original assumption that $\mathbb{N}$ was finite must be false.

Outline

1. SoPS Lecture 20
   - Finite Sets
   - Pidgeonhole Principle
   - Wrapping up

Final thoughts

Reflection

Final thoughts

Reflection

- Give the formal definition of a finite set.
- Explain why there's a bijection between $\mathbb{N}_j$ and $\mathbb{N}_k$ if and only if $j = k$.

Submit your responses by **Tonight!**