

# MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics  
California State University Fullerton

September 30, 2024

# Outline

## 1 Real Analysis Lecture 9

- More on Open Sets
- Closed Sets
- Compactness

# Outline

## 1 Real Analysis Lecture 9

- More on Open Sets
- Closed Sets
- Compactness

# Finite intersections of open sets are open

## Theorem (Open Intersection Theorem)

*Suppose that  $U_1, U_2, \dots, U_n$  are open sets. Then  $\bigcap_{i=1}^n U_i$  is open.*

# Finite intersections of open sets are open

## Theorem (Open Intersection Theorem)

*Suppose that  $U_1, U_2, \dots, U_n$  are open sets. Then  $\bigcap_{i=1}^n U_i$  is open.*

Proof.

# Finite intersections of open sets are open

## Theorem (Open Intersection Theorem)

*Suppose that  $U_1, U_2, \dots, U_n$  are open sets. Then  $\bigcap_{i=1}^n U_i$  is open.*

## Proof.

Suppose that  $\vec{x} \in \bigcap_{i=1}^n U_i$ .

# Finite intersections of open sets are open

## Theorem (Open Intersection Theorem)

*Suppose that  $U_1, U_2, \dots, U_n$  are open sets. Then  $\bigcap_{i=1}^n U_i$  is open.*

## Proof.

Suppose that  $\vec{x} \in \bigcap_{i=1}^n U_i$ .  
Then for all  $i$ ,  $\vec{x} \in U_i$ .

# Finite intersections of open sets are open

## Theorem (Open Intersection Theorem)

*Suppose that  $U_1, U_2, \dots, U_n$  are open sets. Then  $\bigcap_{i=1}^n U_i$  is open.*

## Proof.

Suppose that  $\vec{x} \in \bigcap_{i=1}^n U_i$ .

Then for all  $i$ ,  $\vec{x} \in U_i$ .

Since  $U_i$  is open, this means  $\vec{x}$  is an interior point of  $U_i$ .



# Finite intersections of open sets are open

## Theorem (Open Intersection Theorem)

*Suppose that  $U_1, U_2, \dots, U_n$  are open sets. Then  $\bigcap_{i=1}^n U_i$  is open.*

## Proof.

Suppose that  $\vec{x} \in \bigcap_{i=1}^n U_i$ .

Then for all  $i$ ,  $\vec{x} \in U_i$ .

Since  $U_i$  is open, this means  $\vec{x}$  is an interior point of  $U_i$ .

Therefore there exists  $r_i > 0$  such that  $B(\vec{x}; r_i) \subseteq U_i$ .

# Finite intersections of open sets are open

## Theorem (Open Intersection Theorem)

*Suppose that  $U_1, U_2, \dots, U_n$  are open sets. Then  $\bigcap_{i=1}^n U_i$  is open.*

## Proof.

Suppose that  $\vec{x} \in \bigcap_{i=1}^n U_i$ .

Then for all  $i$ ,  $\vec{x} \in U_i$ .

Since  $U_i$  is open, this means  $\vec{x}$  is an interior point of  $U_i$ .

Therefore there exists  $r_i > 0$  such that  $B(\vec{x}; r_i) \subseteq U_i$ .

Take  $r = \min\{r_i : 1 \leq i \leq n\}$ .

# Finite intersections of open sets are open

## Theorem (Open Intersection Theorem)

*Suppose that  $U_1, U_2, \dots, U_n$  are open sets. Then  $\bigcap_{i=1}^n U_i$  is open.*

## Proof.

Suppose that  $\vec{x} \in \bigcap_{i=1}^n U_i$ .

Then for all  $i$ ,  $\vec{x} \in U_i$ .

Since  $U_i$  is open, this means  $\vec{x}$  is an interior point of  $U_i$ .

Therefore there exists  $r_i > 0$  such that  $B(\vec{x}; r_i) \subseteq U_i$ .

Take  $r = \min\{r_i : 1 \leq i \leq n\}$ .

This means  $B(\vec{x}; r) \subseteq B(\vec{x}; r_i) \subseteq U_i$  for all  $i$ .

# Finite intersections of open sets are open

## Theorem (Open Intersection Theorem)

*Suppose that  $U_1, U_2, \dots, U_n$  are open sets. Then  $\bigcap_{i=1}^n U_i$  is open.*

## Proof.

Suppose that  $\vec{x} \in \bigcap_{i=1}^n U_i$ .

Then for all  $i$ ,  $\vec{x} \in U_i$ .

Since  $U_i$  is open, this means  $\vec{x}$  is an interior point of  $U_i$ .

Therefore there exists  $r_i > 0$  such that  $B(\vec{x}; r_i) \subseteq U_i$ .

Take  $r = \min\{r_i : 1 \leq i \leq n\}$ .

This means  $B(\vec{x}; r) \subseteq B(\vec{x}; r_i) \subseteq U_i$  for all  $i$ .

Therefore  $B(\vec{x}; r) \subseteq \bigcap_{i=1}^n U_i$ .

# Finite intersections of open sets are open

## Theorem (Open Intersection Theorem)

*Suppose that  $U_1, U_2, \dots, U_n$  are open sets. Then  $\bigcap_{i=1}^n U_i$  is open.*

## Proof.

Suppose that  $\vec{x} \in \bigcap_{i=1}^n U_i$ .

Then for all  $i$ ,  $\vec{x} \in U_i$ .

Since  $U_i$  is open, this means  $\vec{x}$  is an interior point of  $U_i$ .

Therefore there exists  $r_i > 0$  such that  $B(\vec{x}; r_i) \subseteq U_i$ .

Take  $r = \min\{r_i : 1 \leq i \leq n\}$ .

This means  $B(\vec{x}; r) \subseteq B(\vec{x}; r_i) \subseteq U_i$  for all  $i$ .

Therefore  $B(\vec{x}; r) \subseteq \bigcap_{i=1}^n U_i$ .

Thus  $\vec{x}$  is an interior point of  $\bigcap_{i=1}^n U_i$ .

# Finite intersections of open sets are open

## Theorem (Open Intersection Theorem)

*Suppose that  $U_1, U_2, \dots, U_n$  are open sets. Then  $\bigcap_{i=1}^n U_i$  is open.*

## Proof.

Suppose that  $\vec{x} \in \bigcap_{i=1}^n U_i$ .

Then for all  $i$ ,  $\vec{x} \in U_i$ .

Since  $U_i$  is open, this means  $\vec{x}$  is an interior point of  $U_i$ .

Therefore there exists  $r_i > 0$  such that  $B(\vec{x}; r_i) \subseteq U_i$ .

Take  $r = \min\{r_i : 1 \leq i \leq n\}$ .

This means  $B(\vec{x}; r) \subseteq B(\vec{x}; r_i) \subseteq U_i$  for all  $i$ .

Therefore  $B(\vec{x}; r) \subseteq \bigcap_{i=1}^n U_i$ .

Thus  $\vec{x}$  is an interior point of  $\bigcap_{i=1}^n U_i$ .

Since  $\vec{x}$  is arbitrary, this proves that  $\bigcap_{i=1}^n U_i$  is open. □

# Challenge

## Problem

Show that the infinite intersection

$$\bigcap_{n=1}^{\infty} \left(-\frac{1}{n}, \frac{1}{n}\right)$$

is not open.

# Component intervals

Let  $U \subseteq \mathbb{R}$  be open.



# Component intervals

Let  $U \subseteq \mathbb{R}$  be open.

## Definition

A **component interval** of  $U$  is an interval  $I$  with  $I \subseteq U$  and with the property that if  $J$  is an interval and  $I \subsetneq J$ , then  $J \not\subseteq U$ .

# Component intervals

Let  $U \subseteq \mathbb{R}$  be open.

## Definition

A **component interval** of  $U$  is an interval  $I$  with  $I \subseteq U$  and with the property that if  $J$  is an interval and  $I \subsetneq J$ , then  $J \not\subseteq U$ .

- $(0, 1)$  is a component interval of  $\mathbb{R} \setminus \{0, 1, 2, 3\}$

# Component intervals

Let  $U \subseteq \mathbb{R}$  be open.

## Definition

A **component interval** of  $U$  is an interval  $I$  with  $I \subseteq U$  and with the property that if  $J$  is an interval and  $I \subsetneq J$ , then  $J \not\subseteq U$ .

- $(0, 1)$  is a component interval of  $\mathbb{R} \setminus \{0, 1, 2, 3\}$
- $(-\infty, 0)$  is also a component interval of  $\mathbb{R} \setminus \{0, 1, 2, 3\}$

# Component intervals

Let  $U \subseteq \mathbb{R}$  be open.

## Definition

A **component interval** of  $U$  is an interval  $I$  with  $I \subseteq U$  and with the property that if  $J$  is an interval and  $I \subsetneq J$ , then  $J \not\subseteq U$ .

- $(0, 1)$  is a component interval of  $\mathbb{R} \setminus \{0, 1, 2, 3\}$
- $(-\infty, 0)$  is also a component interval of  $\mathbb{R} \setminus \{0, 1, 2, 3\}$
- we will show all open sets of  $\mathbb{R}$  are made of component intervals!

# Component intervals

## Lemma

*If  $I_1$  and  $I_2$  are two component intervals of an open subset  $U \subseteq \mathbb{R}$ , then  $I_1 = I_2$  or  $I_1 \cap I_2 = \emptyset$ .*

# Component intervals

## Lemma

*If  $I_1$  and  $I_2$  are two component intervals of an open subset  $U \subseteq \mathbb{R}$ , then  $I_1 = I_2$  or  $I_1 \cap I_2 = \emptyset$ .*

## Proof.

# Component intervals

## Lemma

*If  $I_1$  and  $I_2$  are two component intervals of an open subset  $U \subseteq \mathbb{R}$ , then  $I_1 = I_2$  or  $I_1 \cap I_2 = \emptyset$ .*

## Proof.

Suppose that  $I_1 \cap I_2 \neq \emptyset$ .

# Component intervals

## Lemma

*If  $I_1$  and  $I_2$  are two component intervals of an open subset  $U \subseteq \mathbb{R}$ , then  $I_1 = I_2$  or  $I_1 \cap I_2 = \emptyset$ .*

## Proof.

Suppose that  $I_1 \cap I_2 \neq \emptyset$ .

Then  $J := I_1 \cup I_2$  is an interval.



# Component intervals

## Lemma

*If  $I_1$  and  $I_2$  are two component intervals of an open subset  $U \subseteq \mathbb{R}$ , then  $I_1 = I_2$  or  $I_1 \cap I_2 = \emptyset$ .*

## Proof.

Suppose that  $I_1 \cap I_2 \neq \emptyset$ .

Then  $J := I_1 \cup I_2$  is an interval.

Also  $I_1 \subseteq U$  and  $I_2 \subseteq U$ , so  $J \subseteq U$ .

# Component intervals

## Lemma

*If  $I_1$  and  $I_2$  are two component intervals of an open subset  $U \subseteq \mathbb{R}$ , then  $I_1 = I_2$  or  $I_1 \cap I_2 = \emptyset$ .*

## Proof.

Suppose that  $I_1 \cap I_2 \neq \emptyset$ .

Then  $J := I_1 \cup I_2$  is an interval.

Also  $I_1 \subseteq U$  and  $I_2 \subseteq U$ , so  $J \subseteq U$ .

Since  $I_i$  is a component interval and  $I_i \subseteq J \subseteq U$ , we have  $I_i = J$ .

# Component intervals

## Lemma

*If  $I_1$  and  $I_2$  are two component intervals of an open subset  $U \subseteq \mathbb{R}$ , then  $I_1 = I_2$  or  $I_1 \cap I_2 = \emptyset$ .*

## Proof.

Suppose that  $I_1 \cap I_2 \neq \emptyset$ .

Then  $J := I_1 \cup I_2$  is an interval.

Also  $I_1 \subseteq U$  and  $I_2 \subseteq U$ , so  $J \subseteq U$ .

Since  $I_j$  is a component interval and  $I_j \subseteq J \subseteq U$ , we have  $I_j = J$ .

In particular  $I_1 = J = I_2$ . □

# Component intervals

## Theorem (Apostol 3.10)

*If  $U \subseteq \mathbb{R}$  is open and  $x \in U$ , then there is a unique component interval of  $U$  containing  $x$ .*

# Component intervals

## Theorem (Apostol 3.10)

*If  $U \subseteq \mathbb{R}$  is open and  $x \in U$ , then there is a unique component interval of  $U$  containing  $x$ .*

## Proof.

Uniqueness follows from previous Lemma, so we only need existence.

# Component intervals

## Theorem (Apostol 3.10)

*If  $U \subseteq \mathbb{R}$  is open and  $x \in U$ , then there is a unique component interval of  $U$  containing  $x$ .*

## Proof.

Uniqueness follows from previous Lemma, so we only need existence.

Suppose that  $x \in U$  and consider

$$A = \{a : (a, x) \subseteq U\}, \quad \text{and} \quad B = \{b : (x, b) \subseteq U\}$$

# Component intervals

## Theorem (Apostol 3.10)

*If  $U \subseteq \mathbb{R}$  is open and  $x \in U$ , then there is a unique component interval of  $U$  containing  $x$ .*

## Proof.

Uniqueness follows from previous Lemma, so we only need existence.

Suppose that  $x \in U$  and consider

$$A = \{a : (a, x) \subseteq U\}, \quad \text{and} \quad B = \{b : (x, b) \subseteq U\}$$

If  $A$  is not bounded below, let  $a = -\infty$ . Otherwise, let  $a = \inf(A)$ .

# Component intervals

## Theorem (Apostol 3.10)

*If  $U \subseteq \mathbb{R}$  is open and  $x \in U$ , then there is a unique component interval of  $U$  containing  $x$ .*

## Proof.

Uniqueness follows from previous Lemma, so we only need existence.

Suppose that  $x \in U$  and consider

$$A = \{a : (a, x) \subseteq U\}, \quad \text{and} \quad B = \{b : (x, b) \subseteq U\}$$

If  $A$  is not bounded below, let  $a = -\infty$ . Otherwise, let  $a = \inf(A)$ .

If  $B$  is not bounded above, let  $b = \infty$ . Otherwise, let  $b = \sup(B)$ .



# Component intervals

## Theorem (Apostol 3.10)

*If  $U \subseteq \mathbb{R}$  is open and  $x \in U$ , then there is a unique component interval of  $U$  containing  $x$ .*

## Proof.

Uniqueness follows from previous Lemma, so we only need existence.

Suppose that  $x \in U$  and consider

$$A = \{a : (a, x) \subseteq U\}, \quad \text{and} \quad B = \{b : (x, b) \subseteq U\}$$

If  $A$  is not bounded below, let  $a = -\infty$ . Otherwise, let  $a = \inf(A)$ .

If  $B$  is not bounded above, let  $b = \infty$ . Otherwise, let  $b = \sup(B)$ .

Claim:  $(a, b)$  is a component interval of  $U$  containing  $x$ . □

# Challenge

## Problem

Prove that  $(a, b)$  is a component interval of  $U$ .

# Open subsets of $\mathbb{R}$

## Theorem (Representation Theorem for Open Sets in $\mathbb{R}$ )

*If  $U \subseteq \mathbb{R}$  is open, then  $U$  is the union of a countable family of disjoint open intervals.*

# Open subsets of $\mathbb{R}$

## Theorem (Representation Theorem for Open Sets in $\mathbb{R}$ )

*If  $U \subseteq \mathbb{R}$  is open, then  $U$  is the union of a countable family of disjoint open intervals.*

### Proof.

From the previous theorem, each  $x \in U$  is contained in a unique component interval  $I_x$  of  $U$ .

# Open subsets of $\mathbb{R}$

## Theorem (Representation Theorem for Open Sets in $\mathbb{R}$ )

*If  $U \subseteq \mathbb{R}$  is open, then  $U$  is the union of a countable family of disjoint open intervals.*

### Proof.

From the previous theorem, each  $x \in U$  is contained in a unique component interval  $I_x$  of  $U$ .

Therefore

$$U = \bigcup_{x \in U} I_x$$

is a union of open intervals!

# Open subsets of $\mathbb{R}$

## Theorem (Representation Theorem for Open Sets in $\mathbb{R}$ )

*If  $U \subseteq \mathbb{R}$  is open, then  $U$  is the union of a countable family of disjoint open intervals.*

### Proof.

From the previous theorem, each  $x \in U$  is contained in a unique component interval  $I_x$  of  $U$ .

Therefore

$$U = \bigcup_{x \in U} I_x$$

is a union of open intervals!

Uh oh ... this isn't a **countable** union.



# Open subsets of $\mathbb{R}$

## Theorem (Representation Theorem for Open Sets in $\mathbb{R}$ )

*If  $U \subseteq \mathbb{R}$  is open, then  $U$  is the union of a countable family of disjoint open intervals.*

### Proof.

Instead, we consider rational points  $U \cap \mathbb{Q}$ .

# Open subsets of $\mathbb{R}$

## Theorem (Representation Theorem for Open Sets in $\mathbb{R}$ )

*If  $U \subseteq \mathbb{R}$  is open, then  $U$  is the union of a countable family of disjoint open intervals.*

### Proof.

Instead, we consider rational points  $U \cap \mathbb{Q}$ .

If  $x \in U$ , then  $x \in I_x \subseteq U$ .



# Open subsets of $\mathbb{R}$

## Theorem (Representation Theorem for Open Sets in $\mathbb{R}$ )

*If  $U \subseteq \mathbb{R}$  is open, then  $U$  is the union of a countable family of disjoint open intervals.*

### Proof.

Instead, we consider rational points  $U \cap \mathbb{Q}$ .

If  $x \in U$ , then  $x \in I_x \subseteq U$ .

Any interval contains a rational number  $r \in I_x$ .

# Open subsets of $\mathbb{R}$

## Theorem (Representation Theorem for Open Sets in $\mathbb{R}$ )

*If  $U \subseteq \mathbb{R}$  is open, then  $U$  is the union of a countable family of disjoint open intervals.*

### Proof.

Instead, we consider rational points  $U \cap \mathbb{Q}$ .

If  $x \in U$ , then  $x \in I_x \subseteq U$ .

Any interval contains a rational number  $r \in I_x$ .

Since different component intervals must be disjoint,  $I_x = I_r$ .

# Open subsets of $\mathbb{R}$

## Theorem (Representation Theorem for Open Sets in $\mathbb{R}$ )

*If  $U \subseteq \mathbb{R}$  is open, then  $U$  is the union of a countable family of disjoint open intervals.*

### Proof.

Instead, we consider rational points  $U \cap \mathbb{Q}$ .

If  $x \in U$ , then  $x \in I_x \subseteq U$ .

Any interval contains a rational number  $r \in I_x$ .

Since different component intervals must be disjoint,  $I_x = I_r$ .

Thus  $x \in \bigcup_{r \in U \cap \mathbb{Q}} I_r \subseteq U$ .

# Open subsets of $\mathbb{R}$

## Theorem (Representation Theorem for Open Sets in $\mathbb{R}$ )

*If  $U \subseteq \mathbb{R}$  is open, then  $U$  is the union of a countable family of disjoint open intervals.*

### Proof.

Instead, we consider rational points  $U \cap \mathbb{Q}$ .

If  $x \in U$ , then  $x \in I_x \subseteq U$ .

Any interval contains a rational number  $r \in I_x$ .

Since different component intervals must be disjoint,  $I_x = I_r$ .

Thus  $x \in \bigcup_{r \in U \cap \mathbb{Q}} I_r \subseteq U$ .

Since  $x$  was arbitrary,

$$U = \bigcup_{r \in U \cap \mathbb{Q}} I_r$$

# Outline

## 1 Real Analysis Lecture 9

- More on Open Sets
- **Closed Sets**
- Compactness

# Closed sets

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **closed** if it is the complement of an open set, or equivalently if its complement  $\mathbb{R}^n \setminus A$  is open.

# Closed sets

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **closed** if it is the complement of an open set, or equivalently if its complement  $\mathbb{R}^n \setminus A$  is open.

## Examples:

# Closed sets

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **closed** if it is the complement of an open set, or equivalently if its complement  $\mathbb{R}^n \setminus A$  is open.

## Examples:

- singleton sets!



# Closed sets

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **closed** if it is the complement of an open set, or equivalently if its complement  $\mathbb{R}^n \setminus A$  is open.

## Examples:

- singleton sets!
- products of closed intervals

# Challenge!

## Problem

Prove that a singleton set

$$A = \{\vec{a}\}$$

in  $\mathbb{R}^n$  is closed.

# Challenge!

## Problem

Prove that a singleton set

$$A = \{\vec{a}\}$$

in  $\mathbb{R}^n$  is closed.

## Solution

We must show  $U = \mathbb{R}^n \setminus \{\vec{a}\}$  is open.

# Challenge!

## Problem

Prove that a singleton set

$$A = \{\vec{a}\}$$

in  $\mathbb{R}^n$  is closed.

## Solution

We must show  $U = \mathbb{R}^n \setminus \{\vec{a}\}$  is open.

Suppose  $\vec{x} \in U$ .

# Challenge!

## Problem

Prove that a singleton set

$$A = \{\vec{a}\}$$

in  $\mathbb{R}^n$  is closed.

## Solution

We must show  $U = \mathbb{R}^n \setminus \{\vec{a}\}$  is open.

Suppose  $\vec{x} \in U$ . We must show  $\vec{x}$  is an interior point of  $U$ .

# Challenge!

## Problem

Prove that a singleton set

$$A = \{\vec{a}\}$$

in  $\mathbb{R}^n$  is closed.

## Solution

We must show  $U = \mathbb{R}^n \setminus \{\vec{a}\}$  is open.

Suppose  $\vec{x} \in U$ . We must show  $\vec{x}$  is an interior point of  $U$ .

Take  $r = |\vec{x} - \vec{a}|$ .

# Challenge!

## Problem

Prove that a singleton set

$$A = \{\vec{a}\}$$

in  $\mathbb{R}^n$  is closed.

## Solution

We must show  $U = \mathbb{R}^n \setminus \{\vec{a}\}$  is open.

Suppose  $\vec{x} \in U$ . We must show  $\vec{x}$  is an interior point of  $U$ .

Take  $r = |\vec{x} - \vec{a}|$ . Then  $B(\vec{x}; r)$  does not contain  $\vec{a}$ .

# Challenge!

## Problem

Prove that a singleton set

$$A = \{\vec{a}\}$$

in  $\mathbb{R}^n$  is closed.

## Solution

We must show  $U = \mathbb{R}^n \setminus \{\vec{a}\}$  is open.

Suppose  $\vec{x} \in U$ . We must show  $\vec{x}$  is an interior point of  $U$ .

Take  $r = |\vec{x} - \vec{a}|$ . Then  $B(\vec{x}; r)$  does not contain  $\vec{a}$ .

Therefore  $B(\vec{x}; r) \subseteq U$ , so that  $\vec{x}$  is an interior point.



# Challenge!

## Problem

Prove that a singleton set

$$A = \{\vec{a}\}$$

in  $\mathbb{R}^n$  is closed.

## Solution

We must show  $U = \mathbb{R}^n \setminus \{\vec{a}\}$  is open.

Suppose  $\vec{x} \in U$ . We must show  $\vec{x}$  is an interior point of  $U$ .

Take  $r = |\vec{x} - \vec{a}|$ . Then  $B(\vec{x}; r)$  does not contain  $\vec{a}$ .

Therefore  $B(\vec{x}; r) \subseteq U$ , so that  $\vec{x}$  is an interior point.

Since  $\vec{x}$  was arbitrary, this proves  $U$  is open.

# Challenge!

## Problem

Prove that the **closed square**

$$[a, b] \times [c, d] = \{(x, y) \in \mathbb{R}^2 : a \leq x \leq b, c \leq y \leq d\}$$

is a closed set.

# Challenge!

## Problem

Prove that the **closed square**

$$[a, b] \times [c, d] = \{(x, y) \in \mathbb{R}^2 : a \leq x \leq b, c \leq y \leq d\}$$

is a closed set.

## Solution

$\mathbb{R}^2 \setminus ([a, b] \times [c, d])$  is the same as

$$((-\infty, a) \times \mathbb{R}) \cup ((b, \infty) \times \mathbb{R}) \cup (\mathbb{R} \times (-\infty, c)) \cup (\mathbb{R} \times (d, \infty))$$

# Challenge!

## Problem

Prove that the **closed square**

$$[a, b] \times [c, d] = \{(x, y) \in \mathbb{R}^2 : a \leq x \leq b, c \leq y \leq d\}$$

is a closed set.

## Solution

$\mathbb{R}^2 \setminus ([a, b] \times [c, d])$  is the same as

$$((-\infty, a) \times \mathbb{R}) \cup ((b, \infty) \times \mathbb{R}) \cup (\mathbb{R} \times (-\infty, c)) \cup (\mathbb{R} \times (d, \infty))$$

Union of open sets is open!

# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in A$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in A$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

It is called an **accumulation point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$  **different from  $\vec{x}$** .

# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in A$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

It is called an **accumulation point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$  **different from  $\vec{x}$** .

## Examples:

# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in A$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

It is called an **accumulation point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$  **different from  $\vec{x}$** .

## Examples:

- every point in  $A$  is adherent



# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in A$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

It is called an **accumulation point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$  **different from  $\vec{x}$** .

## Examples:

- every point in  $A$  is adherent
- accumulation points are adherent points

# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in A$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

It is called an **accumulation point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$  **different from  $\vec{x}$** .

## Examples:

- every point in  $A$  is adherent
- accumulation points are adherent points
- $-1$  is an accumulation point of  $(-1, 1)$ .

# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in A$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

It is called an **accumulation point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$  **different from  $\vec{x}$** .

## Examples:

- every point in  $A$  is adherent
- accumulation points are adherent points
- $-1$  is an accumulation point of  $(-1, 1)$ .
- suprema and infima are accumulation points!

# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in A$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

It is called an **accumulation point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$  **different from  $\vec{x}$** .

## Examples:

- every point in  $A$  is adherent
- accumulation points are adherent points
- $-1$  is an accumulation point of  $(-1, 1)$ .
- suprema and infima are accumulation points!
- $0$  is an accumulation point of  $\{1/1, 1/2, 1/3, \dots\}$

# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

# Closure of a set

## Definition

The **closure** of a set  $A$  is the set  $\bar{A}$  of all adherent points of  $A$

# Closure of a set

## Definition

The **closure** of a set  $A$  is the set  $\overline{A}$  of all adherent points of  $A$

## Theorem (Apostol Theorem 3.18, 3.20, 3.22)

*A set is closed if and only if it contains all of its adherent points (ie.  $A = \overline{A}$ ), or equivalently if and only if  $A$  contains all of its accumulation points.*

# Outline

## 1 Real Analysis Lecture 9

- More on Open Sets
- Closed Sets
- Compactness



# Intuition of compactness

We are moving toward a very important notion called **compactness**.

# Intuition of compactness

We are moving toward a very important notion called **compactness**.

This important concept is captured in many different ways:

# Intuition of compactness

We are moving toward a very important notion called **compactness**.

This important concept is captured in many different ways:

- no “missing points” in the set

# Intuition of compactness

We are moving toward a very important notion called **compactness**.

This important concept is captured in many different ways:

- no “missing points” in the set
- sequences have convergent subsequences

# Intuition of compactness

We are moving toward a very important notion called **compactness**.

This important concept is captured in many different ways:

- no “missing points” in the set
- sequences have convergent subsequences
- Cauchy sequences converge

# Intuition of compactness

We are moving toward a very important notion called **compactness**.

This important concept is captured in many different ways:

- no “missing points” in the set
- sequences have convergent subsequences
- Cauchy sequences converge
- open covers have finite subcovers

# Intuition of compactness

We are moving toward a very important notion called **compactness**.

This important concept is captured in many different ways:

- no “missing points” in the set
- sequences have convergent subsequences
- Cauchy sequences converge
- open covers have finite subcovers
- in  $\mathbb{R}^n$ : compact = closed and bounded

# Open covers

## Definition

A **open cover** of a set  $A \subseteq \mathbb{R}^n$  is a family  $\{U_i : i \in I\}$  of open sets with  $A \subseteq \bigcup_{i \in I} U_i$ .



# Open covers

## Definition

A **open cover** of a set  $A \subseteq \mathbb{R}^n$  is a family  $\{U_i : i \in I\}$  of open sets with  $A \subseteq \bigcup_{i \in I} U_i$ .

A **subcover** is a family  $\{U_j : j \in J\}$  with  $J \subseteq I$  which still covers  $A$ , ie.  $A \subseteq \bigcup_{j \in J} U_j$ .

# Open covers

## Definition

A **open cover** of a set  $A \subseteq \mathbb{R}^n$  is a family  $\{U_i : i \in I\}$  of open sets with  $A \subseteq \bigcup_{i \in I} U_i$ .

A **subcover** is a family  $\{U_j : j \in J\}$  with  $J \subseteq I$  which still covers  $A$ , ie.  $A \subseteq \bigcup_{j \in J} U_j$ .

## Examples:

# Open covers

## Definition

A **open cover** of a set  $A \subseteq \mathbb{R}^n$  is a family  $\{U_i : i \in I\}$  of open sets with  $A \subseteq \bigcup_{i \in I} U_i$ .

A **subcover** is a family  $\{U_j : j \in J\}$  with  $J \subseteq I$  which still covers  $A$ , ie.  $A \subseteq \bigcup_{j \in J} U_j$ .

## Examples:

- $\{\mathbb{R}\}$  is an open cover of the interval  $(0, 1]$

# Open covers

## Definition

A **open cover** of a set  $A \subseteq \mathbb{R}^n$  is a family  $\{U_i : i \in I\}$  of open sets with  $A \subseteq \bigcup_{i \in I} U_i$ .

A **subcover** is a family  $\{U_j : j \in J\}$  with  $J \subseteq I$  which still covers  $A$ , ie.  $A \subseteq \bigcup_{j \in J} U_j$ .

## Examples:

- $\{\mathbb{R}\}$  is an open cover of the interval  $(0, 1]$
- $\{(\frac{1}{n}, \frac{n+1}{n}) : n \in \mathbb{Z}_+\}$  is an open cover of  $(0, 1]$

# Lindelöf Covering Theorem

## Theorem

*Let  $A \subseteq \mathbb{R}^n$  be a set and suppose  $\{U_i : i \in I\}$  is an open covering of  $A$ . Then there exists a countable subcover  $\{U_j : j \in J\}$ .*

# Lindelöf Covering Theorem

## Theorem

*Let  $A \subseteq \mathbb{R}^n$  be a set and suppose  $\{U_i : i \in I\}$  is an open covering of  $A$ . Then there exists a countable subcover  $\{U_j : j \in J\}$ .*

## Question

Can we do better than this?

# Lindelöf Covering Theorem

## Theorem

*Let  $A \subseteq \mathbb{R}^n$  be a set and suppose  $\{U_i : i \in I\}$  is an open covering of  $A$ . Then there exists a countable subcover  $\{U_j : j \in J\}$ .*

## Question

Can we do better than this?

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **compact** if every open cover of  $A$  has a *finite* subcover.

# Roadmap

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **bounded** if there exists  $\vec{a} \in \mathbb{R}^n$  and  $r > 0$  with  $A \subseteq B(\vec{a}; r)$ .



# Roadmap

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **bounded** if there exists  $\vec{a} \in \mathbb{R}^n$  and  $r > 0$  with  $A \subseteq B(\vec{a}; r)$ .

We wish to prove the **Heine-Borel Theorem** that closed and bounded sets are compact.

# Roadmap

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **bounded** if there exists  $\vec{a} \in \mathbb{R}^n$  and  $r > 0$  with  $A \subseteq B(\vec{a}; r)$ .

We wish to prove the **Heine-Borel Theorem** that closed and bounded sets are compact.

- 1 We prove the Bolzano-Weierstrass Theorem

# Roadmap

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **bounded** if there exists  $\vec{a} \in \mathbb{R}^n$  and  $r > 0$  with  $A \subseteq B(\vec{a}; r)$ .

We wish to prove the **Heine-Borel Theorem** that closed and bounded sets are compact.

- 1 We prove the Bolzano-Weierstrass Theorem
- 2 Then use Bolzano-Weierstrass to prove the Cantor Intersection Theorem

# Roadmap

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **bounded** if there exists  $\vec{a} \in \mathbb{R}^n$  and  $r > 0$  with  $A \subseteq B(\vec{a}; r)$ .

We wish to prove the **Heine-Borel Theorem** that closed and bounded sets are compact.

- 1 We prove the Bolzano-Weierstrass Theorem
- 2 Then use Bolzano-Weierstrass to prove the Cantor Intersection Theorem
- 3 We use the Cantor Intersection Theorem to prove the Heine-Borel Theorem

# Bolzano-Weierstrass Theorem

## Theorem (Bolzano-Weierstrass Theorem)

*A bounded set  $A \subseteq \mathbb{R}^n$  with infinitely many points will contain an accumulation point.*