

# MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics  
California State University Fullerton

September 18, 2024

# Outline

- 1 Real Analysis Lecture 7
  - More with cardinality
  - Set algebra
  - Open Balls and Open Sets

# Outline

- 1 Real Analysis Lecture 7
  - More with cardinality
  - Set algebra
  - Open Balls and Open Sets

# Ultimate Cantor diagonalization

The **power set**  $\mathcal{P}(A)$  of a set  $A$  is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

# Ultimate Cantor diagonalization

The **power set**  $\mathcal{P}(A)$  of a set  $A$  is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

## Theorem

*The cardinality of  $\mathcal{P}(A)$  is strictly larger than  $A$ .*

# Ultimate Cantor diagonalization

The **power set**  $\mathcal{P}(A)$  of a set  $A$  is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

## Theorem

*The cardinality of  $\mathcal{P}(A)$  is strictly larger than  $A$ .*

## Proof.

# Ultimate Cantor diagonalization

The **power set**  $\mathcal{P}(A)$  of a set  $A$  is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

## Theorem

*The cardinality of  $\mathcal{P}(A)$  is strictly larger than  $A$ .*

## Proof.

Suppose  $f : A \rightarrow \mathcal{P}(A)$  is surjective.

# Ultimate Cantor diagonalization

The **power set**  $\mathcal{P}(A)$  of a set  $A$  is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

## Theorem

*The cardinality of  $\mathcal{P}(A)$  is strictly larger than  $A$ .*

## Proof.

Suppose  $f : A \rightarrow \mathcal{P}(A)$  is surjective.

Consider the set

$$S = \{a \in A : a \notin f(a)\}.$$





# Challenge

# Challenge

Since  $f$  is surjective,  $S = f(x)$  for some  $x \in A$ .

# Challenge

Since  $f$  is surjective,  $S = f(x)$  for some  $x \in A$ .

## Problem

Consider the statement  $x \in S$ . What can you conclude?

# Challenge

## Problem

Is there a set with cardinality larger than  $\mathbb{R}$ ?

# Challenge

## Problem

Show that the cardinality of the line segment  $(0, 1)$  and the square  $(0, 1) \times (0, 1)$  is the same.

# Outline

- 1 Real Analysis Lecture 7
  - More with cardinality
  - Set algebra
  - Open Balls and Open Sets

# Unions of sets

Unions:

# Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$



# Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

# Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

$$A_1 \cup A_2 \cup \cdots \cup A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

# Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

$$A_1 \cup A_2 \cup \cdots \cup A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

NOTATION:

$$\bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \cdots \cup A_n.$$

# Intersections of sets

Intersections:

# Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

# Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

# Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

$$A_1 \cap A_2 \cap \cdots \cap A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

# Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

$$A_1 \cap A_2 \cap \cdots \cap A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

NOTATION:

$$\bigcap_{i=1}^n A_i = A_1 \cap A_2 \cap \cdots \cap A_n.$$



# Complements of sets

# Complements of sets

The **complement** of  $A$  relative to  $B$  is

$$B - A = \{b \in B : b \notin A\}.$$

# Complements of sets

The **complement** of  $A$  relative to  $B$  is

$$B - A = \{b \in B : b \notin A\}.$$

## Theorem

### *De Morgan's Laws*

# Complements of sets

The **complement** of  $A$  relative to  $B$  is

$$B - A = \{b \in B : b \notin A\}.$$

## Theorem

*De Morgan's Laws*

$$B - \bigcup_{i=1}^n A_i = \bigcap_{i=1}^n (B - A_i)$$

# Complements of sets

The **complement** of  $A$  relative to  $B$  is

$$B - A = \{b \in B : b \notin A\}.$$

## Theorem

### *De Morgan's Laws*

$$B - \bigcup_{i=1}^n A_i = \bigcap_{i=1}^n (B - A_i)$$

$$B - \bigcap_{i=1}^n A_i = \bigcup_{i=1}^n (B - A_i)$$

# Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

# Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- $I$  any set, called the **index set**

# Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- $I$  any set, called the **index set**
- $A_i$  is a set for all  $i$



# Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- $I$  any set, called the **index set**
- $A_i$  is a set for all  $i$

$$\bigcup_{i \in I} A_i = \{x : x \in A_i, \text{ for some } i \in I\}$$

# Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- $I$  any set, called the **index set**
- $A_i$  is a set for all  $i$

$$\bigcup_{i \in I} A_i = \{x : x \in A_i, \text{ for some } i \in I\}$$

$$\bigcap_{i \in I} A_i = \{x : x \in A_i, \text{ for all } i \in I\}$$

# Challenge

- index set  $I = (1, 2)$
- family of sets  $\{A_i : i \in I\}$
- $A_i = [0, i]$

## Problem

Determine  $\bigcup_{i \in I} A_i$ .

# Challenge

- index set  $I = \mathbb{Z}_+$
- family of sets  $\{A_i : i \in I\}$
- $A_i = [0, i)$

## Problem

Determine  $\bigcap_{i \in I} A_i$ .

# Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

# Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

## Theorem

*The union of a countable family of countable sets is countable.*

# Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

## Theorem

*The union of a countable family of countable sets is countable.*

## Proof.

Let  $\{A_i : i \in I\}$  be a countable family of countable sets.

# Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

## Theorem

*The union of a countable family of countable sets is countable.*

## Proof.

Let  $\{A_i : i \in I\}$  be a countable family of countable sets.  
 $I$  is countable.



# Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

## Theorem

*The union of a countable family of countable sets is countable.*

## Proof.

Let  $\{A_i : i \in I\}$  be a countable family of countable sets.  $I$  is countable. There is a surjection  $\mathbb{Z}_+ \rightarrow I$

# Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

## Theorem

*The union of a countable family of countable sets is countable.*

## Proof.

Let  $\{A_i : i \in I\}$  be a countable family of countable sets.  
 $I$  is countable. There is a surjection  $\mathbb{Z}_+ \rightarrow I$   
In other words, we can enumerate  $I = \{i_1, i_2, i_3, \dots\}$ .

# Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

## Theorem

*The union of a countable family of countable sets is countable.*

## Proof.

Let  $\{A_i : i \in I\}$  be a countable family of countable sets.  $I$  is countable. There is a surjection  $\mathbb{Z}_+ \rightarrow I$ . In other words, we can enumerate  $I = \{i_1, i_2, i_3, \dots\}$ .  $A_i$  is countable for all  $i \in I$ .

# Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

## Theorem

*The union of a countable family of countable sets is countable.*

## Proof.

Let  $\{A_i : i \in I\}$  be a countable family of countable sets.

$I$  is countable. There is a surjection  $\mathbb{Z}_+ \rightarrow I$

In other words, we can enumerate  $I = \{i_1, i_2, i_3, \dots\}$ .

$A_i$  is countable for all  $i \in I$ .

So again  $A_i = \{a_{i1}, a_{i2}, a_{i3}, \dots\}$ .

# Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

## Theorem

*The union of a countable family of countable sets is countable.*

## Proof.

Let  $\{A_i : i \in I\}$  be a countable family of countable sets.

$I$  is countable. There is a surjection  $\mathbb{Z}_+ \rightarrow I$

In other words, we can enumerate  $I = \{i_1, i_2, i_3, \dots\}$ .

$A_i$  is countable for all  $i \in I$ .

So again  $A_i = \{a_{i1}, a_{i2}, a_{i3}, \dots\}$ .

Define  $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \bigcup_{i \in I} A_i$

# Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

## Theorem

*The union of a countable family of countable sets is countable.*

## Proof.

Let  $\{A_i : i \in I\}$  be a countable family of countable sets.

$I$  is countable. There is a surjection  $\mathbb{Z}_+ \rightarrow I$

In other words, we can enumerate  $I = \{i_1, i_2, i_3, \dots\}$ .

$A_i$  is countable for all  $i \in I$ .

So again  $A_i = \{a_{i1}, a_{i2}, a_{i3}, \dots\}$ .

Define  $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \bigcup_{i \in I} A_i$

$f(j, k) = a_{i_j, k}$ .

# Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

## Theorem

*The union of a countable family of countable sets is countable.*

## Proof.

Let  $\{A_i : i \in I\}$  be a countable family of countable sets.

$I$  is countable. There is a surjection  $\mathbb{Z}_+ \rightarrow I$

In other words, we can enumerate  $I = \{i_1, i_2, i_3, \dots\}$ .

$A_i$  is countable for all  $i \in I$ .

So again  $A_i = \{a_{i1}, a_{i2}, a_{i3}, \dots\}$ .

Define  $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \bigcup_{i \in I} A_i$

$f(j, k) = a_{i_j, k}$ . Surjection!



# Challenge!

A real number is called **algebraic** if it is a root of a polynomial with integer coefficients.



# Challenge!

A real number is called **algebraic** if it is a root of a polynomial with integer coefficients.

## Problem

Prove that the set of all algebraic numbers is countable.

# Challenge!

A real number is called **algebraic** if it is a root of a polynomial with integer coefficients.

## Problem

Prove that the set of all algebraic numbers is countable.

Hint: use the previous theorem!

# Outline

- 1 Real Analysis Lecture 7
  - More with cardinality
  - Set algebra
  - Open Balls and Open Sets

# Euclidean space

# Euclidean space

$n$ -dimensional **euclidean space** is

# Euclidean space

$n$ -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

# Euclidean space

$n$ -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

# Euclidean space

$n$ -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given  $\vec{x} = (x_1, \dots, x_n)$  and  $\vec{y} = (y_1, \dots, y_n)$



# Euclidean space

$n$ -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given  $\vec{x} = (x_1, \dots, x_n)$  and  $\vec{y} = (y_1, \dots, y_n)$

Ⓐ  $\vec{x} + \vec{y} = (x_1 + y_1, \dots, x_n + y_n)$

# Euclidean space

$n$ -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given  $\vec{x} = (x_1, \dots, x_n)$  and  $\vec{y} = (y_1, \dots, y_n)$

(a)  $\vec{x} + \vec{y} = (x_1 + y_1, \dots, x_n + y_n)$

(b)  $a\vec{x} = (ax_1, \dots, ax_n)$

# Euclidean space

$n$ -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given  $\vec{x} = (x_1, \dots, x_n)$  and  $\vec{y} = (y_1, \dots, y_n)$

- (a)  $\vec{x} + \vec{y} = (x_1 + y_1, \dots, x_n + y_n)$
- (b)  $a\vec{x} = (ax_1, \dots, ax_n)$
- (c)  $\vec{x} - \vec{y} = \vec{x} + (-1)\vec{y} = (x_1 - y_1, \dots, x_n - y_n)$

# Euclidean space

$n$ -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given  $\vec{x} = (x_1, \dots, x_n)$  and  $\vec{y} = (y_1, \dots, y_n)$

- (a)  $\vec{x} + \vec{y} = (x_1 + y_1, \dots, x_n + y_n)$
- (b)  $a\vec{x} = (ax_1, \dots, ax_n)$
- (c)  $\vec{x} - \vec{y} = \vec{x} + (-1)\vec{y} = (x_1 - y_1, \dots, x_n - y_n)$
- (d)  $\vec{0} = (0, \dots, 0)$

# Euclidean space

$n$ -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given  $\vec{x} = (x_1, \dots, x_n)$  and  $\vec{y} = (y_1, \dots, y_n)$

- (a)  $\vec{x} + \vec{y} = (x_1 + y_1, \dots, x_n + y_n)$
- (b)  $a\vec{x} = (ax_1, \dots, ax_n)$
- (c)  $\vec{x} - \vec{y} = \vec{x} + (-1)\vec{y} = (x_1 - y_1, \dots, x_n - y_n)$
- (d)  $\vec{0} = (0, \dots, 0)$
- (e)  $\vec{x} \cdot \vec{y} = \sum_{i=1}^n x_i y_i$

# Euclidean space

$n$ -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given  $\vec{x} = (x_1, \dots, x_n)$  and  $\vec{y} = (y_1, \dots, y_n)$

- (a)  $\vec{x} + \vec{y} = (x_1 + y_1, \dots, x_n + y_n)$
- (b)  $a\vec{x} = (ax_1, \dots, ax_n)$
- (c)  $\vec{x} - \vec{y} = \vec{x} + (-1)\vec{y} = (x_1 - y_1, \dots, x_n - y_n)$
- (d)  $\vec{0} = (0, \dots, 0)$
- (e)  $\vec{x} \cdot \vec{y} = \sum_{i=1}^n x_i y_i$
- (f)  $|\vec{x}| = (\vec{x} \cdot \vec{x})^{1/2} = (\sum_{i=1}^n x_i^2)^{1/2}$

# Metric properties

# Metric properties

The norm  $|\vec{x}|$  is an example of a **metric**.



# Metric properties

The norm  $|\vec{x}|$  is an example of a **metric**.  
It satisfies several important properties:

# Metric properties

The norm  $|\vec{x}|$  is an example of a **metric**.  
It satisfies several important properties:

## Theorem

# Metric properties

The norm  $|\vec{x}|$  is an example of a **metric**.  
It satisfies several important properties:

## Theorem

(a) *(positivity)  $|\vec{x}| \geq 0$  with equality iff  $\vec{x} = \vec{0}$*

# Metric properties

The norm  $|\vec{x}|$  is an example of a **metric**.  
It satisfies several important properties:

## Theorem

- (a) (positivity)  $|\vec{x}| \geq 0$  with equality iff  $\vec{x} = \vec{0}$
- (b) (symmetry)  $|\vec{x} + \vec{y}| = |\vec{y} + \vec{x}|$

# Metric properties

The norm  $|\vec{x}|$  is an example of a **metric**.  
It satisfies several important properties:

## Theorem

- (a) *(positivity)*  $|\vec{x}| \geq 0$  with equality iff  $\vec{x} = \vec{0}$
- (b) *(symmetry)*  $|\vec{x} + \vec{y}| = |\vec{y} + \vec{x}|$
- (c) *(triangle inequality)*  $|\vec{x} + \vec{y}| \leq |\vec{x}| + |\vec{y}|$

# Metric properties

The norm  $|\vec{x}|$  is an example of a **metric**.  
It satisfies several important properties:

## Theorem

- (a) *(positivity)*  $|\vec{x}| \geq 0$  with equality iff  $\vec{x} = \vec{0}$
- (b) *(symmetry)*  $|\vec{x} + \vec{y}| = |\vec{y} + \vec{x}|$
- (c) *(triangle inequality)*  $|\vec{x} + \vec{y}| \leq |\vec{x}| + |\vec{y}|$

It also satisfies

# Metric properties

The norm  $|\vec{x}|$  is an example of a **metric**.  
It satisfies several important properties:

## Theorem

- (a) *(positivity)*  $|\vec{x}| \geq 0$  with equality iff  $\vec{x} = \vec{0}$
- (b) *(symmetry)*  $|\vec{x} + \vec{y}| = |\vec{y} + \vec{x}|$
- (c) *(triangle inequality)*  $|\vec{x} + \vec{y}| \leq |\vec{x}| + |\vec{y}|$

It also satisfies

# Metric properties

The norm  $|\vec{x}|$  is an example of a **metric**.  
It satisfies several important properties:

## Theorem

- (a) *(positivity)*  $|\vec{x}| \geq 0$  with equality iff  $\vec{x} = \vec{0}$
- (b) *(symmetry)*  $|\vec{x} + \vec{y}| = |\vec{y} + \vec{x}|$
- (c) *(triangle inequality)*  $|\vec{x} + \vec{y}| \leq |\vec{x}| + |\vec{y}|$

It also satisfies

- (a) *(scaling)*  $|c\vec{x}| = |c| |\vec{x}|$



# Metric properties

The norm  $|\vec{x}|$  is an example of a **metric**.  
It satisfies several important properties:

## Theorem

- (a) *(positivity)*  $|\vec{x}| \geq 0$  with equality iff  $\vec{x} = \vec{0}$
- (b) *(symmetry)*  $|\vec{x} + \vec{y}| = |\vec{y} + \vec{x}|$
- (c) *(triangle inequality)*  $|\vec{x} + \vec{y}| \leq |\vec{x}| + |\vec{y}|$

It also satisfies

- (a) *(scaling)*  $|c\vec{x}| = |c| |\vec{x}|$
- (b) *(Cauchy-Schwartz)*  $|\vec{x} \cdot \vec{y}| \leq |\vec{x}| |\vec{y}|$