MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics California State University Fullerton

September 8, 2024

Outline

- Real Analysis Lecture 3
 - Decimal expansions
 - The Triangle Inequality
 - Sets, Relations, Functions

Outline

- Real Analysis Lecture 3
 - Decimal expansions
 - The Triangle Inequality
 - Sets, Relations, Functions

A finite decimal expansion is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \dots + \frac{a_n}{10^n},$$

where $a_0 \in \mathbb{Z}_+$ and $0 \le a_k \le 9$ for $1 \le k \le n$.

A finite decimal expansion is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \dots + \frac{a_n}{10^n},$$

where $a_0 \in \mathbb{Z}_+$ and $0 \le a_k \le 9$ for $1 \le k \le n$.

Notation:

$$a_0.a_1a_2a_3...a_n$$
.

A finite decimal expansion is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \dots + \frac{a_n}{10^n},$$

where $a_0 \in \mathbb{Z}_+$ and $0 \le a_k \le 9$ for $1 \le k \le n$.

Notation:

$$a_0.a_1a_2a_3...a_n$$
.

Any positive real number x > 0 can be approximated by a finite decimal expansion.

Theorem (Apostol Theorem 1.20)

For any real x > 0 and $n \in \mathbb{Z}_+$, there exists a finite decimal expansion $r_n = a_0.a_1a_2...a_n$ with

$$r_n \leq x < r_n + \frac{1}{10^n}.$$

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \le x\}.$$



Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \le x\}.$$

The set A is a set of integers which is bounded above by x, so it has a maximum a_0 .

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \le x\}.$$

The set A is a set of integers which is bounded above by x, so it has a maximum a_0 .

Then clearly $x_1 = a - a_0 \in [0, 1)$.



Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set A is a set of integers which is bounded above by x, so it has a maximum a_0 .

Then clearly $x_1 = a - a_0 \in [0, 1)$.

Define a_1, a_2, a_3, \ldots and x_1, x_2, x_3, \ldots recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and
$$x_{k+1} = 10x_k - a_k$$
.



Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set A is a set of integers which is bounded above by x, so it has a maximum a_0 .

Then clearly $x_1 = a - a_0 \in [0, 1)$.

Define a_1, a_2, a_3, \ldots and x_1, x_2, x_3, \ldots recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and $x_{k+1} = 10x_k - a_k$. Then $0 \le a_k \le 9$ for all $k \ge 1$



Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set A is a set of integers which is bounded above by x, so it has a maximum a_0 .

Then clearly $x_1 = a - a_0 \in [0, 1)$.

Define a_1, a_2, a_3, \ldots and x_1, x_2, x_3, \ldots recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and $x_{k+1} = 10x_k - a_k$. Then $0 \le a_k \le 9$ for all $k \ge 1$ and

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \le x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n+1}{10^n}$$



We say that x > 0 has the decimal expansion $a_0.a_1a_2a_3...$ and write

$$x=a_0.a_1a_2a_3a_4\ldots$$

We say that x > 0 has the decimal expansion $a_0.a_1a_2a_3...$ and write

$$x=a_0.a_1a_2a_3a_4\ldots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n}{10^n} \le x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n+1}{10^n}.$$

We say that x > 0 has the decimal expansion $a_0.a_1a_2a_3...$ and write

$$x=a_0.a_1a_2a_3a_4\ldots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n}{10^n} \le x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n+1}{10^n}.$$

We say that x > 0 has the decimal expansion $a_0.a_1a_2a_3...$ and write

$$x=a_0.a_1a_2a_3a_4\ldots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n}{10^n} \le x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n+1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

we haven't defined limits

We say that x > 0 has the decimal expansion $a_0.a_1a_2a_3...$ and write

$$x=a_0.a_1a_2a_3a_4\ldots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n}{10^n} \le x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n+1}{10^n}.$$

- we haven't defined limits
- with this definition, numbers have unique decimal expansions!

We say that x > 0 has the decimal expansion $a_0.a_1a_2a_3...$ and write

$$x=a_0.a_1a_2a_3a_4\ldots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n}{10^n} \le x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n+1}{10^n}.$$

- we haven't defined limits
- with this definition, numbers have unique decimal expansions!

$$1 \neq 0.999999999...$$



We say that x > 0 has the decimal expansion $a_0.a_1a_2a_3...$ and write

$$x=a_0.a_1a_2a_3a_4\ldots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n}{10^n} \le x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n+1}{10^n}.$$

- we haven't defined limits
- with this definition, numbers have unique decimal expansions!

$$1 \neq 0.99999999...$$

$$1 \nless 0 + \frac{9}{10} + \frac{9}{100} + \dots + \frac{9+1}{100} = 1$$

Outline

- Real Analysis Lecture 3
 - Decimal expansions
 - The Triangle Inequality
 - Sets, Relations, Functions

Absolute value

The **absolute value** of $x \in \mathbb{R}$ is

$$|x| = \begin{cases} x, & x \ge 0 \\ -x, & x < 0 \end{cases}$$

In particular,

$$0 \le |x|$$

and also

$$-|x| \le x \le |x|.$$

Problem

Prove Apostol Theorem 1.21 that if $a \ge 0$, then $|x| \le a$ if and only if $-a \le x \le a$.

Problem

Prove Apostol Theorem 1.21 that if $a \ge 0$, then $|x| \le a$ if and only if $-a \le x \le a$.

Solution

If
$$|x| \le a$$
 then $-a \le -|x|$

Problem

Prove Apostol Theorem 1.21 that if $a \ge 0$, then $|x| \le a$ if and only if $-a \le x \le a$.

Solution

If $|x| \le a$ then $-a \le -|x|$ and therefore

$$-a \le -|x| \le x \le |x| \le a$$

Problem

Prove Apostol Theorem 1.21 that if $a \ge 0$, then $|x| \le a$ if and only if $-a \le x \le a$.

Solution

If $|x| \le a$ then $-a \le -|x|$ and therefore

$$-a \le -|x| \le x \le |x| \le a$$

Conversely, if $-a \le x \le a$ then



Problem

Prove Apostol Theorem 1.21 that if $a \ge 0$, then $|x| \le a$ if and only if $-a \le x \le a$.

Solution

If $|x| \le a$ then $-a \le -|x|$ and therefore

$$-a \le -|x| \le x \le |x| \le a$$

Conversely, if $-a \le x \le a$ then

$$x \ge 0 \Rightarrow |x| = x \le a$$

Problem

Prove Apostol Theorem 1.21 that if $a \ge 0$, then $|x| \le a$ if and only if $-a \le x \le a$.

Solution

If $|x| \le a$ then $-a \le -|x|$ and therefore

$$-a \le -|x| \le x \le |x| \le a$$

Conversely, if $-a \le x \le a$ then

$$x \ge 0 \Rightarrow |x| = x \le a$$

$$x \le 0 \Rightarrow |x| = -x \le -(-a) = a$$

Theorem (Triangle inequality)

For any real numbers $x, y \in \mathbb{R}$ we have

$$|x+y|\leq |x|+|y|.$$

Theorem (Triangle inequality)

For any real numbers $x, y \in \mathbb{R}$ we have

$$|x+y|\leq |x|+|y|.$$

Proof.

$$-|x| \le x \le |x|$$
 and $-|y| \le y \le |y|$



Theorem (Triangle inequality)

For any real numbers $x, y \in \mathbb{R}$ we have

$$|x+y|\leq |x|+|y|.$$

Proof.

$$-|x| \le x \le |x|$$
 and $-|y| \le y \le |y|$

Adding these together, we get

$$-(|x|+|y|) \le x+y \le |x|+|y|$$



Theorem (Triangle inequality)

For any real numbers $x, y \in \mathbb{R}$ we have

$$|x+y| \le |x| + |y|.$$

Proof.

$$-|x| \le x \le |x|$$
 and $-|y| \le y \le |y|$

Adding these together, we get

$$-(|x|+|y|) \le x+y \le |x|+|y|$$

It follows from the previous theorem that

$$|x+y| \le |x| + |y|$$



Advanced triangle inequality

Theorem (Triangle inequality)

For any real numbers $x_1, x_2, \ldots, x_n \in \mathbb{R}$ we have

$$|x_1 + x_2 + \cdots + x_n| \le |x_1| + |x_2| + \cdots + |x_n|.$$

Advanced triangle inequality

Theorem (Triangle inequality)

For any real numbers $x_1, x_2, \ldots, x_n \in \mathbb{R}$ we have

$$|x_1 + x_2 + \cdots + x_n| \le |x_1| + |x_2| + \cdots + |x_n|.$$

Proof.

Induction.



Higher-dimensional triangle inequality

Theorem (Cauchy-Schwartz Inequality (Apostol Theorem 1.23))

If x_1, \ldots, x_n and y_1, \ldots, y_n are real numbers, then

$$\left(\sum_{k=1}^n x_k y_k\right)^2 \le \left(\sum_{k=1}^n x_k^2\right) \left(\sum_{k=1}^n y_k^2\right)$$

If y_k isn't always zero, then equality holds if and only if there exists $t \in \mathbb{R}$ with $x_k = ty_k$ for all k.

Higher-dimensional triangle inequality

Theorem (Cauchy-Schwartz Inequality (Apostol Theorem 1.23))

If x_1, \ldots, x_n and y_1, \ldots, y_n are real numbers, then

$$\left(\sum_{k=1}^n x_k y_k\right)^2 \le \left(\sum_{k=1}^n x_k^2\right) \left(\sum_{k=1}^n y_k^2\right)$$

If y_k isn't always zero, then equality holds if and only if there exists $t \in \mathbb{R}$ with $x_k = ty_k$ for all k.

Vector version:

$$(\vec{x}\cdot\vec{y})^2 \leq |\vec{x}|^2 |\vec{y}|^2.$$



$$\sum_{k=1}^{n} (x_k + ty_k)^2 \ge 0, \text{ with equality iff all terms zero}$$

$$\sum_{k=1}^{n} (x_k + ty_k)^2 \ge 0, \quad \text{with equality iff all terms zero}$$

$$\sum_{k=1}^{n} x_k^2 + 2t \sum_{k=1}^{n} x_k y_k + t^2 \sum_{k=1}^{n} y_k^2 \ge 0$$

$$\sum_{k=1}^{n} (x_k + ty_k)^2 \ge 0, \text{ with equality iff all terms zero}$$

$$\sum_{k=1}^{n} x_k^2 + 2t \sum_{k=1}^{n} x_k y_k + t^2 \sum_{k=1}^{n} y_k^2 \ge 0$$

Take
$$t = -(\sum_{k=1}^{n} x_{k} y_{k}) / (\sum_{k=1}^{n} y_{k}^{2})$$
:



$$\sum_{k=1}^{n} (x_k + ty_k)^2 \ge 0, \text{ with equality iff all terms zero}$$

$$\sum_{k=1}^{n} x_k^2 + 2t \sum_{k=1}^{n} x_k y_k + t^2 \sum_{k=1}^{n} y_k^2 \ge 0$$

Take
$$t = -\left(\sum_{k=1}^{n} x_{k} y_{k}\right) / \left(\sum_{k=1}^{n} y_{k}^{2}\right)$$
:

$$\sum_{k=1}^{n} x_k^2 - \frac{\left(\sum_{k=1}^{n} x_k y_k\right)^2}{\left(\sum_{k=1}^{n} y_k^2\right)} \ge 0.$$



Theorem (Minkowski inequality)

For any real numbers $x_1, x_2, \dots, x_n \in \mathbb{R}$ and $y_1, y_2, \dots, y_n \in \mathbb{R}$ we have

$$\left(\sum_{k=1}^{n}(x_k+y_k)^2\right)^{1/2} \leq \left(\sum_{k=1}^{n}x_k^2\right)^{1/2} + \left(\sum_{k=1}^{n}y_k^2\right)^{1/2}$$

Theorem (Minkowski inequality)

For any real numbers $x_1, x_2, \ldots, x_n \in \mathbb{R}$ and $y_1, y_2, \ldots, y_n \in \mathbb{R}$ we have

$$\left(\sum_{k=1}^{n}(x_k+y_k)^2\right)^{1/2} \leq \left(\sum_{k=1}^{n}x_k^2\right)^{1/2} + \left(\sum_{k=1}^{n}y_k^2\right)^{1/2}$$

Vector version:

$$|\vec{x} + \vec{y}| \le |\vec{x}| + |\vec{y}|.$$

Theorem (Minkowski inequality)

For any real numbers $x_1, x_2, \ldots, x_n \in \mathbb{R}$ and $y_1, y_2, \ldots, y_n \in \mathbb{R}$ we have

$$\left(\sum_{k=1}^{n}(x_k+y_k)^2\right)^{1/2} \leq \left(\sum_{k=1}^{n}x_k^2\right)^{1/2} + \left(\sum_{k=1}^{n}y_k^2\right)^{1/2}$$

Vector version:

$$|\vec{x} + \vec{y}| \le |\vec{x}| + |\vec{y}|.$$

A higher dimensional triangle inequality!

Proof.

By the triangle inequality:



Proof.

By the triangle inequality:

$$\begin{split} \sum_{k=1}^{n} (x_k + y_k)^2 &\leq \sum_{k=1}^{n} (|x_k| + |y_k|)|x_k + y_k| \\ &= \sum_{k=1}^{n} |x_k| \cdot |x_k + y_k| + \sum_{k=1}^{n} |y_k| \cdot |x_k + y_k| \end{split}$$



Proof.

By the triangle inequality:

$$\begin{split} \sum_{k=1}^{n} (x_k + y_k)^2 &\leq \sum_{k=1}^{n} (|x_k| + |y_k|)|x_k + y_k| \\ &= \sum_{k=1}^{n} |x_k| \cdot |x_k + y_k| + \sum_{k=1}^{n} |y_k| \cdot |x_k + y_k| \end{split}$$

Now applying the Cauchy-Schwartz inequality:



Proof.

By the triangle inequality:

$$\begin{split} \sum_{k=1}^{n} (x_k + y_k)^2 &\leq \sum_{k=1}^{n} (|x_k| + |y_k|)|x_k + y_k| \\ &= \sum_{k=1}^{n} |x_k| \cdot |x_k + y_k| + \sum_{k=1}^{n} |y_k| \cdot |x_k + y_k| \end{split}$$

Now applying the Cauchy-Schwartz inequality:

$$\leq \left(\sum_{k=1}^{n} |x_{k}|^{2}\right)^{1/2} \left(\sum_{k=1}^{n} |x_{k} + y_{k}|^{2}\right)^{1/2} + \left(\sum_{k=1}^{n} |y_{k}|^{2}\right)^{1/2} \left(\sum_{k=1}^{n} |x_{k} + y_{k}|^{2}\right)^{1/2}$$

$$= \left[\left(\sum_{k=1}^{n} |x_{k}|^{2}\right)^{1/2} + \left(\sum_{k=1}^{n} |y_{k}|^{2}\right)^{1/2}\right] \left(\sum_{k=1}^{n} |x_{k} + y_{k}|^{2}\right)^{1/2}$$

Proof.

By the triangle inequality:

$$\begin{split} \sum_{k=1}^{n} (x_k + y_k)^2 &\leq \sum_{k=1}^{n} (|x_k| + |y_k|)|x_k + y_k| \\ &= \sum_{k=1}^{n} |x_k| \cdot |x_k + y_k| + \sum_{k=1}^{n} |y_k| \cdot |x_k + y_k| \end{split}$$

Now applying the Cauchy-Schwartz inequality:

$$\leq \left(\sum_{k=1}^{n} |x_k|^2 \right)^{1/2} \left(\sum_{k=1}^{n} |x_k + y_k|^2 \right)^{1/2} + \left(\sum_{k=1}^{n} |y_k|^2 \right)^{1/2} \left(\sum_{k=1}^{n} |x_k + y_k|^2 \right)^{1/2}$$

$$= \left[\left(\sum_{k=1}^{n} |x_k|^2 \right)^{1/2} + \left(\sum_{k=1}^{n} |y_k|^2 \right)^{1/2} \right] \left(\sum_{k=1}^{n} |x_k + y_k|^2 \right)^{1/2}$$



Outline

- Real Analysis Lecture 3
 - Decimal expansions
 - The Triangle Inequality
 - Sets, Relations, Functions

Intuitively, a **set** *A* is a "collection of things" which we call the **elements** of *A*.

Intuitively, a **set** *A* is a "collection of things" which we call the **elements** of *A*.

in practice, this is a bad definition (Russell's Paradox)

Intuitively, a **set** *A* is a "collection of things" which we call the **elements** of *A*.

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Intuitively, a **set** *A* is a "collection of things" which we call the **elements** of *A*.

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Intuitively, a **set** *A* is a "collection of things" which we call the **elements** of *A*.

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

 \bullet \mathbb{R} , \mathbb{Z}_+ , \mathbb{Z} , \mathbb{Q}

Intuitively, a **set** *A* is a "collection of things" which we call the **elements** of *A*.

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

- \bullet \mathbb{R} , \mathbb{Z}_+ , \mathbb{Z} , \mathbb{Q}
- $(1,5], (0,\infty)$

Intuitively, a **set** *A* is a "collection of things" which we call the **elements** of *A*.

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

- \bullet \mathbb{R} , \mathbb{Z}_+ , \mathbb{Z} , \mathbb{Q}
- $(1,5], (0,\infty)$
- empty set Ø

Intuitively, a **set** *A* is a "collection of things" which we call the **elements** of *A*.

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

- \bullet \mathbb{R} , \mathbb{Z}_+ , \mathbb{Z} , \mathbb{Q}
- $(1,5], (0,\infty)$
- empty set Ø
- $\{\heartsuit, \mathsf{Fall}, \{\varnothing\}\}$

Intuitively, a **set** *A* is a "collection of things" which we call the **elements** of *A*.

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

- \bullet \mathbb{R} , \mathbb{Z}_+ , \mathbb{Z} , \mathbb{Q}
- $(1,5], (0,\infty)$
- empty set Ø
- {♡, Fall, {∅}}
- $\{n \in \mathbb{Z} : n \text{ is prime}\}$

In the true minimalistic philosophy of mathematics, we want everything to be a set.

In the true minimalistic philosophy of mathematics, we want everything to be a set.

integers

In the true minimalistic philosophy of mathematics, we want everything to be a set.

integers

$$0=\varnothing, 1=\{\varnothing\}, 2=\{\varnothing, \{\varnothing\}\}, \dots$$

In the true minimalistic philosophy of mathematics, we want everything to be a set.

integers

$$0=\varnothing, 1=\{\varnothing\}, 2=\{\varnothing, \{\varnothing\}\}, \dots$$

ordered pairs

In the true minimalistic philosophy of mathematics, we want everything to be a set.

integers

$$0=\varnothing, 1=\{\varnothing\}, 2=\{\varnothing, \{\varnothing\}\}, \dots$$

ordered pairs

$$(a,b) = \{a, \{a,b\}\}$$

In the true minimalistic philosophy of mathematics, we want everything to be a set.

integers

$$0=\varnothing, 1=\{\varnothing\}, 2=\{\varnothing, \{\varnothing\}\}, \dots$$

ordered pairs

$$(a,b) = \{a, \{a,b\}\}$$

even relations and functions are sets!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

$$(a,b) = (c,d)$$
 if and only if $\{a,\{a,b\}\} = \{c,\{c,d\}\}$

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

$$(a,b) = (c,d)$$
 if and only if $\{a, \{a,b\}\} = \{c, \{c,d\}\}$
Clearly, if $a = c$ and $b = d$, then $\{a, \{a,b\}\} = \{c, \{c,d\}\}$

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

$$(a,b) = (c,d)$$
 if and only if $\{a,\{a,b\}\} = \{c,\{c,d\}\}$
Clearly, if $a = c$ and $b = d$, then $\{a,\{a,b\}\} = \{c,\{c,d\}\}$
The tough part is the opposite direction!



Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

$$(a,b)=(c,d)$$
 if and only if $\{a,\{a,b\}\}=\{c,\{c,d\}\}$
Clearly, if $a=c$ and $b=d$, then $\{a,\{a,b\}\}=\{c,\{c,d\}\}$
The tough part is the opposite direction!
Suppose $\{a,\{a,b\}\}=\{c,\{c,d\}\}$.



Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

$$(a,b)=(c,d)$$
 if and only if $\{a,\{a,b\}\}=\{c,\{c,d\}\}$
Clearly, if $a=c$ and $b=d$, then $\{a,\{a,b\}\}=\{c,\{c,d\}\}$
The tough part is the opposite direction!
Suppose $\{a,\{a,b\}\}=\{c,\{c,d\}\}$.
Two possible cases:



Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

$$(a,b)=(c,d)$$
 if and only if $\{a,\{a,b\}\}=\{c,\{c,d\}\}$
Clearly, if $a=c$ and $b=d$, then $\{a,\{a,b\}\}=\{c,\{c,d\}\}$
The tough part is the opposite direction!
Suppose $\{a,\{a,b\}\}=\{c,\{c,d\}\}$.
Two possible cases:

Case I:
$$a = c$$
 and $\{a, b\} = \{c, d\}$
Case II: $a = \{c, d\}$ and $\{a, b\} = c$

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

Solution

(a,b)=(c,d) if and only if $\{a,\{a,b\}\}=\{c,\{c,d\}\}$ Clearly, if a=c and b=d, then $\{a,\{a,b\}\}=\{c,\{c,d\}\}$ The tough part is the opposite direction! Suppose $\{a,\{a,b\}\}=\{c,\{c,d\}\}$. Two possible cases:

Case I:
$$a = c$$
 and $\{a, b\} = \{c, d\}$

Case II:
$$a = \{c, d\}$$
 and $\{a, b\} = c$

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

Case I:
$$a = c$$
 and $\{a, b\} = \{c, d\}$



Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

Case I:
$$a = c$$
 and $\{a, b\} = \{c, d\}$

Since
$$\{a, b\} = \{c, d\}$$
, we know $b \in \{c, d\}$.



Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

Case I:
$$a = c$$
 and $\{a, b\} = \{c, d\}$

Since
$$\{a, b\} = \{c, d\}$$
, we know $b \in \{c, d\}$.

Therefore
$$b = c$$
 or $b = d$.

If
$$b = d$$
, we're done!



Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

Case I:
$$a = c$$
 and $\{a, b\} = \{c, d\}$

Since
$$\{a, b\} = \{c, d\}$$
, we know $b \in \{c, d\}$.

Therefore
$$b = c$$
 or $b = d$.

If
$$b = d$$
, we're done! ... so assume instead that $b = c$



Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

Solution

Case I:
$$a = c$$
 and $\{a, b\} = \{c, d\}$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Therefore b = c or b = d.

If b = d, we're done! ... so assume instead that b = c

Then a = c implies a = b.



Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

Solution

Case I:
$$a = c$$
 and $\{a, b\} = \{c, d\}$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Therefore b = c or b = d.

If b = d, we're done! ... so assume instead that b = c

Then a = c implies a = b.

Therefore $\{c, d\} = \{a, b\} = \{a, a\} = \{a\}.$



Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

Solution

Case I:
$$a = c$$
 and $\{a, b\} = \{c, d\}$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Therefore b = c or b = d.

If b = d, we're done! ... so assume instead that b = c

Then a = c implies a = b.

Therefore $\{c, d\} = \{a, b\} = \{a, a\} = \{a\}.$

It follows that d = c = b = a.



Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

Case II:
$$a = \{c, d\}$$
 and $\{a, b\} = c$

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

Solution

Case II:
$$a = \{c, d\}$$
 and $\{a, b\} = c$

This would imply that $c \in a$ and $a \in c$.

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

Solution

Case II:
$$a = \{c, d\}$$
 and $\{a, b\} = c$

This would imply that $c \in a$ and $a \in c$.

This can be shown to contradict the ZF Axioms of Set Theory.

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a,b)=(c,d)$$
 if and only if $a=c$ and $b=d$

Solution

Case II:
$$a = \{c, d\}$$
 and $\{a, b\} = c$

This would imply that $c \in a$ and $a \in c$.

This can be shown to contradict the ZF Axioms of Set Theory. Specifically the regularity axiom for the set $\{a, c\}...$

The **Cartesian product** of *A* and *B* is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

The **Cartesian product** of *A* and *B* is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** \mathcal{R} from A to B is a subset of $A \times B$.

The **Cartesian product** of *A* and *B* is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** \mathcal{R} from A to B is a subset of $A \times B$.

NOTATION: aRb means $(a,b) \in R$.

The **Cartesian product** of *A* and *B* is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** \mathcal{R} from A to B is a subset of $A \times B$.

NOTATION: aRb means $(a,b) \in R$.

Domain and codomain:

The **Cartesian product** of *A* and *B* is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** \mathcal{R} from A to B is a subset of $A \times B$.

NOTATION: aRb means $(a,b) \in R$.

Domain and codomain:

$$dom(\mathcal{R}) = \{a \in A : \exists b \in B, \ a\mathcal{R}b\}$$
$$codom(\mathcal{R}) = \{b \in B : \exists a \in A, \ a\mathcal{R}b\}$$

The **Cartesian product** of *A* and *B* is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** \mathcal{R} from A to B is a subset of $A \times B$.

NOTATION: aRb means $(a,b) \in R$.

Domain and codomain:

$$dom(\mathcal{R}) = \{ a \in A : \exists b \in B, \ a\mathcal{R}b \}$$
$$codom(\mathcal{R}) = \{ b \in B : \exists a \in A, \ a\mathcal{R}b \}$$

A relation \mathcal{R} from A to A is called a **relation on** A. A relation on A is

A relation \mathcal{R} from A to A is called a **relation on** A. A relation on A is

• **reflexive** if aRa for all $a \in A$

A relation \mathcal{R} from A to A is called a **relation on** A. A relation on A is

- **reflexive** if aRa for all $a \in A$
- **symmetric** if aRb implies bRa for all $a, b \in A$

A relation \mathcal{R} from A to A is called a **relation on** A. A relation on A is

- **reflexive** if aRa for all $a \in A$
- **symmetric** if aRb implies bRa for all $a, b \in A$
- transitive if aRb and bRc implies aRc for all $a, b, c \in A$

A relation \mathcal{R} from A to A is called a **relation on** A. A relation on A is

- **reflexive** if aRa for all $a \in A$
- **symmetric** if aRb implies bRa for all $a, b \in A$
- **transitive** if aRb and bRc implies aRc for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

A relation \mathcal{R} from A to A is called a **relation on** A. A relation on A is

- **reflexive** if aRa for all $a \in A$
- **symmetric** if aRb implies bRa for all $a, b \in A$
- **transitive** if aRb and bRc implies aRc for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties. Examples:

A relation \mathcal{R} from A to A is called a **relation on** A. A relation on A is

- **reflexive** if aRa for all $a \in A$
- **symmetric** if aRb implies bRa for all $a, b \in A$
- **transitive** if aRb and bRc implies aRc for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties. Examples:

A relation \mathcal{R} from A to A is called a **relation on** A. A relation on A is

- **reflexive** if aRa for all $a \in A$
- **symmetric** if aRb implies bRa for all $a, b \in A$
- transitive if aRb and bRc implies aRc for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Examples:

• < = $\{(x, y) : y - x \in (0, \infty)\}$ is transitive but not reflexive or symmetric on \mathbb{R}

A relation \mathcal{R} from A to A is called a **relation on** A. A relation on A is

- **reflexive** if aRa for all $a \in A$
- **symmetric** if aRb implies bRa for all $a, b \in A$
- transitive if aRb and bRc implies aRc for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Examples:

- < = $\{(x, y) : y x \in (0, \infty)\}$ is transitive but not reflexive or symmetric on $\mathbb R$
- \leq = { $(x, y) : y x \in [0, \infty)$ } is reflexive and transitive but not symmetric on \mathbb{R}

Problem

Give an example of a relation on \mathbb{R} which is symmetric and transitive but not reflexive.

Problem

Give an example of a relation on \mathbb{R} which is reflexive and symmetric but not transitive.

A **function** from A to B is a relation \mathcal{R} from A to B with the property

A **function** from A to B is a relation \mathcal{R} from A to B with the property

aRb and aRc implies b = c.

A **function** from A to B is a relation \mathcal{R} from A to B with the property

$$aRb$$
 and aRc implies $b = c$.

If a relation R is a function, we usually use a symbol like f.

NOTATION: $f: A \rightarrow B$ means f is a function from A to B

NOTATION: f(a) = b means $(a, b) \in f$.

The set

$$img(f) = \{f(a) : a \in A\}$$

is called the range or image of f



A **function** from A to B is a relation \mathcal{R} from A to B with the property

$$aRb$$
 and aRc implies $b = c$.

If a relation R is a function, we usually use a symbol like f.

NOTATION:
$$f(a) = b$$
 means $(a, b) \in f$.

The set

$$img(f) = \{f(a) : a \in A\}$$

is called the range or image of f



A **function** from A to B is a relation \mathcal{R} from A to B with the property

aRb and aRc implies b = c.

If a relation R is a function, we usually use a symbol like f.

NOTATION: $f: A \rightarrow B$ means f is a function from A to B

NOTATION: f(a) = b means $(a, b) \in f$.

A **function** from A to B is a relation \mathcal{R} from A to B with the property

$$aRb$$
 and aRc implies $b = c$.

If a relation \mathcal{R} is a function, we usually use a symbol like f.

NOTATION: $f: A \rightarrow B$ means f is a function from A to B

NOTATION: f(a) = b means $(a, b) \in f$.

The set

$$img(f) = \{f(a) : a \in A\}$$

is called the range or image of f



Problem

Determine all the equivalence relations on $\mathbb R$ which are also functions.

Challenge!

Problem

Determine all the equivalence relations on $\mathbb R$ which are also functions.

Solution

 $f = \mathcal{R}$ must be reflexive, so $x\mathcal{R}x$ for all x

Challenge!

Problem

Determine all the equivalence relations on \mathbb{R} which are also functions.

Solution

 $f = \mathcal{R}$ must be reflexive, so $x\mathcal{R}x$ for all x. This means f(x) = x for all x.

Challenge!

Problem

Determine all the equivalence relations on \mathbb{R} which are also functions.

Solution

 $f = \mathcal{R}$ must be reflexive, so $x\mathcal{R}x$ for all x

This means f(x) = x for all x.

Thus the only function which is an equivalence relation is the identity function

$$f(x) = x$$
.



A function f from A to B is called **one-to-one** or **injective** if

A function f from A to B is called **one-to-one** or **injective** if

$$f(x) = f(y)$$
 implies $x = y$ for all $x, y \in A$.

A function f from A to B is called **one-to-one** or **injective** if

$$f(x) = f(y)$$
 implies $x = y$ for all $x, y \in A$.

It is called **onto** or **surjective** if img(f) = B, or equivalently

for all
$$b \in B$$
 there exists $a \in A$ with $f(a) = b$.

A function f from A to B is called **one-to-one** or **injective** if

$$f(x) = f(y)$$
 implies $x = y$ for all $x, y \in A$.

It is called **onto** or **surjective** if img(f) = B, or equivalently

for all
$$b \in B$$
 there exists $a \in A$ with $f(a) = b$.

If it satisfies both properties, it is called **bijective**.

The **converse** of a relation \mathcal{R} from A to B is the relation

The **converse** of a relation \mathcal{R} from A to B is the relation

$$\check{\mathcal{R}} = \{(b, a) \in \mathcal{B} \times \mathcal{A} : a\mathcal{R}b\}.$$

The **converse** of a relation \mathcal{R} from A to B is the relation

$$\check{\mathcal{R}} = \{(b, a) \in B \times A : a\mathcal{R}b\}.$$

If f is function, then \check{f} may or not be a function.

The **converse** of a relation \mathcal{R} from A to B is the relation

$$\check{\mathcal{R}} = \{(b, a) \in \mathcal{B} \times \mathcal{A} : a\mathcal{R}b\}.$$

If f is function, then \check{f} may or not be a function. If it is, we call it the **inverse** of f.

The composition of $f: A \rightarrow B$ and $g: B \rightarrow C$ is the function

The composition of $f: A \rightarrow B$ and $g: B \rightarrow C$ is the function

$$g \circ f : A \rightarrow C$$

The composition of $f: A \rightarrow B$ and $g: B \rightarrow C$ is the function

$$g \circ f : A \rightarrow C$$

with domain A and codomain C defined by

The composition of $f : A \rightarrow B$ and $g : B \rightarrow C$ is the function

$$g \circ f : A \rightarrow C$$

with domain A and codomain C defined by

$$(g\circ f)(x)=g(f(x)).$$