

# MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics  
California State University Fullerton

October 2, 2024

# Outline

- 1 Real Analysis Lecture 9
  - More on Closed Sets
  - Compactness

# Outline

- 1 Real Analysis Lecture 9
  - More on Closed Sets
  - Compactness

# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in \mathbb{R}^n$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in \mathbb{R}^n$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

It is called an **accumulation point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$  **different from  $\vec{x}$** .

# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in \mathbb{R}^n$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

It is called an **accumulation point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$  **different from  $\vec{x}$** .

## Examples:

# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in \mathbb{R}^n$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

It is called an **accumulation point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$  **different from  $\vec{x}$** .

## Examples:

- every point in  $A$  is adherent

# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in \mathbb{R}^n$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

It is called an **accumulation point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$  **different from  $\vec{x}$** .

## Examples:

- every point in  $A$  is adherent
- accumulation points are adherent points



# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in \mathbb{R}^n$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

It is called an **accumulation point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$  **different from  $\vec{x}$** .

## Examples:

- every point in  $A$  is adherent
- accumulation points are adherent points
- $-1$  is an accumulation point of  $(-1, 1)$ .

# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in \mathbb{R}^n$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

It is called an **accumulation point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$  **different from  $\vec{x}$** .

## Examples:

- every point in  $A$  is adherent
- accumulation points are adherent points
- $-1$  is an accumulation point of  $(-1, 1)$ .
- suprema and infima are accumulation points!

# Adherent and Accumulation Points

## Definition

A point  $\vec{x} \in \mathbb{R}^n$  is called an **adherent point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$ .

It is called an **accumulation point** if for all  $r > 0$  the ball  $B(\vec{x}; r)$  contains at least one element of  $A$  **different from  $\vec{x}$** .

## Examples:

- every point in  $A$  is adherent
- accumulation points are adherent points
- $-1$  is an accumulation point of  $(-1, 1)$ .
- suprema and infima are accumulation points!
- $0$  is an accumulation point of  $\{1/1, 1/2, 1/3, \dots\}$

# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if and only if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if and only if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

Proof.

# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if and only if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

## Proof.

Obviously, if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ , then it contains at least one point of  $A$  different from  $\vec{x}$ , so it's an accumulation point.

# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if and only if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

## Proof.

Obviously, if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ , then it contains at least one point of  $A$  different from  $\vec{x}$ , so it's an accumulation point.

Suppose instead that  $\vec{x}$  is an accumulation point and let  $r > 0$ .

# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if and only if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

## Proof.

Obviously, if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ , then it contains at least one point of  $A$  different from  $\vec{x}$ , so it's an accumulation point.

Suppose instead that  $\vec{x}$  is an accumulation point and let  $r > 0$ . It suffices to show that the set  $C = (B(\vec{x}; r) \setminus \{\vec{x}\}) \cap A$  is infinite.





# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

Proof.

# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

## Proof.

Suppose that  $C$  is finite.

# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

## Proof.

Suppose that  $C$  is finite.

Then the set  $\{|\vec{y} - \vec{x}| : \vec{y} \in C\}$  is also finite.

# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

## Proof.

Suppose that  $C$  is finite.

Then the set  $\{|\vec{y} - \vec{x}| : \vec{y} \in C\}$  is also finite.

It is also nonempty and has only positive values.

# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

## Proof.

Suppose that  $C$  is finite.

Then the set  $\{|\vec{y} - \vec{x}| : \vec{y} \in C\}$  is also finite.

It is also nonempty and has only positive values. This means that it has a minimum value  $s > 0$ .

# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

## Proof.

Suppose that  $C$  is finite.

Then the set  $\{|\vec{y} - \vec{x}| : \vec{y} \in C\}$  is also finite.

It is also nonempty and has only positive values. This means that it has a minimum value  $s > 0$ .

Since  $\vec{x}$  is an accumulation point,  $B(\vec{x}; s) \cap A$  has an element  $\vec{y}$  different from  $\vec{x}$ .

# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

## Proof.

Suppose that  $C$  is finite.

Then the set  $\{|\vec{y} - \vec{x}| : \vec{y} \in C\}$  is also finite.

It is also nonempty and has only positive values. This means that it has a minimum value  $s > 0$ .

Since  $\vec{x}$  is an accumulation point,  $B(\vec{x}; s) \cap A$  has an element  $\vec{y}$  different from  $\vec{x}$ .

However,  $\vec{y} \in C$  and  $|\vec{y} - \vec{x}| < s$ , contradicting the minimality of  $s$ .



# Characterizing accumulation points

## Theorem (Apostol Theorem 3.17)

*A point  $\vec{x}$  is an accumulation point of  $A$  if for all  $r > 0$ , the ball  $B(\vec{x}; r)$  contains infinitely many points of  $A$ .*

## Proof.

Suppose that  $C$  is finite.

Then the set  $\{|\vec{y} - \vec{x}| : \vec{y} \in C\}$  is also finite.

It is also nonempty and has only positive values. This means that it has a minimum value  $s > 0$ .

Since  $\vec{x}$  is an accumulation point,  $B(\vec{x}; s) \cap A$  has an element  $\vec{y}$  different from  $\vec{x}$ .

However,  $\vec{y} \in C$  and  $|\vec{y} - \vec{x}| < s$ , contradicting the minimality of  $s$ .

We conclude that  $C$  is infinite. □

# Closure of a set

## Definition

The **closure** of a set  $A$  is the set  $\overline{A}$  of all adherent points of  $A$

# Closure of a set

## Definition

The **closure** of a set  $A$  is the set  $\overline{A}$  of all adherent points of  $A$

## Theorem (Apostol Theorem 3.18, 3.20, 3.22)

*A set is closed if and only if it contains all of its adherent points (ie.  $A = \overline{A}$ ), or equivalently if and only if  $A$  contains all of its accumulation points.*

# Closure of a set

## Proof.

Suppose that  $A$  is closed.

Then  $\mathbb{R}^n \setminus A$  is open.

If  $\vec{x} \in \mathbb{R}^n \setminus A$ , then there exist  $r > 0$  such that  $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$ .

This means that  $B(\vec{x}; r) \cap A = \emptyset$ .

Thus  $\vec{x}$  is not an adherent point of  $A$ .

Thus every adherent point of  $A$  is an element of  $A$ . □

# Closures are closed

## Theorem

*If  $A \subseteq \mathbb{R}^n$  is any set, then  $\overline{A}$  is closed.*

## Proof.

Exercise! □

# Outline

- 1 Real Analysis Lecture 9
  - More on Closed Sets
  - Compactness

# Intuition of compactness

We are moving toward a very important notion called **compactness**.

# Intuition of compactness

We are moving toward a very important notion called **compactness**.

This important concept is captured in many different ways:



# Intuition of compactness

We are moving toward a very important notion called **compactness**.

This important concept is captured in many different ways:

- no “missing points” in the set

# Intuition of compactness

We are moving toward a very important notion called **compactness**.

This important concept is captured in many different ways:

- no “missing points” in the set
- sequences have convergent subsequences

# Intuition of compactness

We are moving toward a very important notion called **compactness**.

This important concept is captured in many different ways:

- no “missing points” in the set
- sequences have convergent subsequences
- Cauchy sequences converge

# Intuition of compactness

We are moving toward a very important notion called **compactness**.

This important concept is captured in many different ways:

- no “missing points” in the set
- sequences have convergent subsequences
- Cauchy sequences converge
- open covers have finite subcovers

# Intuition of compactness

We are moving toward a very important notion called **compactness**.

This important concept is captured in many different ways:

- no “missing points” in the set
- sequences have convergent subsequences
- Cauchy sequences converge
- open covers have finite subcovers
- in  $\mathbb{R}^n$ : compact = closed and bounded

# Open covers

## Definition

A **open cover** of a set  $A \subseteq \mathbb{R}^n$  is a family  $\{U_i : i \in I\}$  of open sets with  $A \subseteq \bigcup_{i \in I} U_i$ .

# Open covers

## Definition

A **open cover** of a set  $A \subseteq \mathbb{R}^n$  is a family  $\{U_i : i \in I\}$  of open sets with  $A \subseteq \bigcup_{i \in I} U_i$ .

A **subcover** is a family  $\{U_j : j \in J\}$  with  $J \subseteq I$  which still covers  $A$ , ie.  $A \subseteq \bigcup_{j \in J} U_j$ .

# Open covers

## Definition

A **open cover** of a set  $A \subseteq \mathbb{R}^n$  is a family  $\{U_i : i \in I\}$  of open sets with  $A \subseteq \bigcup_{i \in I} U_i$ .

A **subcover** is a family  $\{U_j : j \in J\}$  with  $J \subseteq I$  which still covers  $A$ , ie.  $A \subseteq \bigcup_{j \in J} U_j$ .

## Examples:



# Open covers

## Definition

A **open cover** of a set  $A \subseteq \mathbb{R}^n$  is a family  $\{U_i : i \in I\}$  of open sets with  $A \subseteq \bigcup_{i \in I} U_i$ .

A **subcover** is a family  $\{U_j : j \in J\}$  with  $J \subseteq I$  which still covers  $A$ , ie.  $A \subseteq \bigcup_{j \in J} U_j$ .

## Examples:

- $\{\mathbb{R}\}$  is an open cover of the interval  $(0, 1]$

# Open covers

## Definition

A **open cover** of a set  $A \subseteq \mathbb{R}^n$  is a family  $\{U_i : i \in I\}$  of open sets with  $A \subseteq \bigcup_{i \in I} U_i$ .

A **subcover** is a family  $\{U_j : j \in J\}$  with  $J \subseteq I$  which still covers  $A$ , ie.  $A \subseteq \bigcup_{j \in J} U_j$ .

## Examples:

- $\{\mathbb{R}\}$  is an open cover of the interval  $(0, 1]$
- $\{(\frac{1}{n}, \frac{n+1}{n}) : n \in \mathbb{Z}_+\}$  is an open cover of  $(0, 1]$

# Lindelöf Covering Theorem

## Theorem

*Let  $A \subseteq \mathbb{R}^n$  be a set and suppose  $\{U_i : i \in I\}$  is an open covering of  $A$ . Then there exists a countable subcover  $\{U_j : j \in J\}$ .*

# Lindelöf Covering Theorem

## Theorem

*Let  $A \subseteq \mathbb{R}^n$  be a set and suppose  $\{U_i : i \in I\}$  is an open covering of  $A$ . Then there exists a countable subcover  $\{U_j : j \in J\}$ .*

## Question

Can we do better than this?

# Lindelöf Covering Theorem

## Theorem

*Let  $A \subseteq \mathbb{R}^n$  be a set and suppose  $\{U_i : i \in I\}$  is an open covering of  $A$ . Then there exists a countable subcover  $\{U_j : j \in J\}$ .*

## Question

Can we do better than this?

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **compact** if every open cover of  $A$  has a *finite* subcover.

# Roadmap

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **bounded** if there exists  $\vec{a} \in \mathbb{R}^n$  and  $r > 0$  with  $A \subseteq B(\vec{a}; r)$ .

# Roadmap

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **bounded** if there exists  $\vec{a} \in \mathbb{R}^n$  and  $r > 0$  with  $A \subseteq B(\vec{a}; r)$ .

We wish to prove the **Heine-Borel Theorem** that closed and bounded sets are compact.

# Roadmap

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **bounded** if there exists  $\vec{a} \in \mathbb{R}^n$  and  $r > 0$  with  $A \subseteq B(\vec{a}; r)$ .

We wish to prove the **Heine-Borel Theorem** that closed and bounded sets are compact.

- 1 We prove the Bolzano-Weierstrass Theorem



# Roadmap

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **bounded** if there exists  $\vec{a} \in \mathbb{R}^n$  and  $r > 0$  with  $A \subseteq B(\vec{a}; r)$ .

We wish to prove the **Heine-Borel Theorem** that closed and bounded sets are compact.

- 1 We prove the Bolzano-Weierstrass Theorem
- 2 Then use Bolzano-Weierstrass to prove the Cantor Intersection Theorem

# Roadmap

## Definition

A set  $A \subseteq \mathbb{R}^n$  is called **bounded** if there exists  $\vec{a} \in \mathbb{R}^n$  and  $r > 0$  with  $A \subseteq B(\vec{a}; r)$ .

We wish to prove the **Heine-Borel Theorem** that closed and bounded sets are compact.

- 1 We prove the Bolzano-Weierstrass Theorem
- 2 Then use Bolzano-Weierstrass to prove the Cantor Intersection Theorem
- 3 We use the Cantor Intersection Theorem to prove the Heine-Borel Theorem

# Bolzano-Weierstrass Theorem

## Theorem (Bolzano-Weierstrass Theorem)

*A bounded set  $A \subseteq \mathbb{R}^n$  with infinitely many points will contain an accumulation point.*