

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

September 23, 2024

Outline

- 1 Real Analysis Lecture 7
 - Open Balls and Open Sets

Outline

- 1 Real Analysis Lecture 7
 - Open Balls and Open Sets

Euclidean space

Euclidean space

n -dimensional **euclidean space** is

Euclidean space

n -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Euclidean space

n -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Euclidean space

n -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given $\vec{x} = (x_1, \dots, x_n)$ and $\vec{y} = (y_1, \dots, y_n)$

Euclidean space

n -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given $\vec{x} = (x_1, \dots, x_n)$ and $\vec{y} = (y_1, \dots, y_n)$

(a) $\vec{x} + \vec{y} = (x_1 + y_1, \dots, x_n + y_n)$

Euclidean space

n -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given $\vec{x} = (x_1, \dots, x_n)$ and $\vec{y} = (y_1, \dots, y_n)$

- (a) $\vec{x} + \vec{y} = (x_1 + y_1, \dots, x_n + y_n)$
- (b) $a\vec{x} = (ax_1, \dots, ax_n)$

Euclidean space

n -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given $\vec{x} = (x_1, \dots, x_n)$ and $\vec{y} = (y_1, \dots, y_n)$

- (a) $\vec{x} + \vec{y} = (x_1 + y_1, \dots, x_n + y_n)$
- (b) $a\vec{x} = (ax_1, \dots, ax_n)$
- (c) $\vec{x} - \vec{y} = \vec{x} + (-1)\vec{y} = (x_1 - y_1, \dots, x_n - y_n)$

Euclidean space

n -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given $\vec{x} = (x_1, \dots, x_n)$ and $\vec{y} = (y_1, \dots, y_n)$

- (a) $\vec{x} + \vec{y} = (x_1 + y_1, \dots, x_n + y_n)$
- (b) $a\vec{x} = (ax_1, \dots, ax_n)$
- (c) $\vec{x} - \vec{y} = \vec{x} + (-1)\vec{y} = (x_1 - y_1, \dots, x_n - y_n)$
- (d) $\vec{0} = (0, \dots, 0)$

Euclidean space

n -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given $\vec{x} = (x_1, \dots, x_n)$ and $\vec{y} = (y_1, \dots, y_n)$

- (a) $\vec{x} + \vec{y} = (x_1 + y_1, \dots, x_n + y_n)$
- (b) $a\vec{x} = (ax_1, \dots, ax_n)$
- (c) $\vec{x} - \vec{y} = \vec{x} + (-1)\vec{y} = (x_1 - y_1, \dots, x_n - y_n)$
- (d) $\vec{0} = (0, \dots, 0)$
- (e) $\vec{x} \cdot \vec{y} = \sum_{i=1}^n x_i y_i$

Euclidean space

n -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given $\vec{x} = (x_1, \dots, x_n)$ and $\vec{y} = (y_1, \dots, y_n)$

- (a) $\vec{x} + \vec{y} = (x_1 + y_1, \dots, x_n + y_n)$
- (b) $a\vec{x} = (ax_1, \dots, ax_n)$
- (c) $\vec{x} - \vec{y} = \vec{x} + (-1)\vec{y} = (x_1 - y_1, \dots, x_n - y_n)$
- (d) $\vec{0} = (0, \dots, 0)$
- (e) $\vec{x} \cdot \vec{y} = \sum_{i=1}^n x_i y_i$
- (f) $|\vec{x}| = (\vec{x} \cdot \vec{x})^{1/2} = (\sum_{i=1}^n x_i^2)^{1/2}$

Metric properties

Metric properties

The norm $|\vec{x}|$ is an example of a **metric**.

Metric properties

The norm $|\vec{x}|$ is an example of a **metric**.
It satisfies several important properties:

Metric properties

The norm $|\vec{x}|$ is an example of a **metric**.
It satisfies several important properties:

Theorem

Metric properties

The norm $|\vec{x}|$ is an example of a **metric**.
It satisfies several important properties:

Theorem

(a) *(positivity) $|\vec{x}| \geq 0$ with equality iff $\vec{x} = \vec{0}$*

Metric properties

The norm $|\vec{x}|$ is an example of a **metric**.
It satisfies several important properties:

Theorem

- (a) *(positivity)* $|\vec{x}| \geq 0$ with equality iff $\vec{x} = \vec{0}$
- (b) *(symmetry)* $|\vec{x} + \vec{y}| = |\vec{y} + \vec{x}|$

Metric properties

The norm $|\vec{x}|$ is an example of a **metric**.
It satisfies several important properties:

Theorem

- (a) *(positivity)* $|\vec{x}| \geq 0$ with equality iff $\vec{x} = \vec{0}$
- (b) *(symmetry)* $|\vec{x} + \vec{y}| = |\vec{y} + \vec{x}|$
- (c) *(triangle inequality)* $|\vec{x} + \vec{y}| \leq |\vec{x}| + |\vec{y}|$

Metric properties

The norm $|\vec{x}|$ is an example of a **metric**.
It satisfies several important properties:

Theorem

- (a) *(positivity)* $|\vec{x}| \geq 0$ with equality iff $\vec{x} = \vec{0}$
- (b) *(symmetry)* $|\vec{x} + \vec{y}| = |\vec{y} + \vec{x}|$
- (c) *(triangle inequality)* $|\vec{x} + \vec{y}| \leq |\vec{x}| + |\vec{y}|$

It also satisfies

Metric properties

The norm $|\vec{x}|$ is an example of a **metric**.
It satisfies several important properties:

Theorem

- (a) *(positivity)* $|\vec{x}| \geq 0$ with equality iff $\vec{x} = \vec{0}$
- (b) *(symmetry)* $|\vec{x} + \vec{y}| = |\vec{y} + \vec{x}|$
- (c) *(triangle inequality)* $|\vec{x} + \vec{y}| \leq |\vec{x}| + |\vec{y}|$

It also satisfies

Metric properties

The norm $|\vec{x}|$ is an example of a **metric**.
It satisfies several important properties:

Theorem

- (a) *(positivity)* $|\vec{x}| \geq 0$ with equality iff $\vec{x} = \vec{0}$
- (b) *(symmetry)* $|\vec{x} + \vec{y}| = |\vec{y} + \vec{x}|$
- (c) *(triangle inequality)* $|\vec{x} + \vec{y}| \leq |\vec{x}| + |\vec{y}|$

It also satisfies

- (a) *(scaling)* $|c\vec{x}| = |c| |\vec{x}|$

Metric properties

The norm $|\vec{x}|$ is an example of a **metric**.
It satisfies several important properties:

Theorem

- (a) *(positivity)* $|\vec{x}| \geq 0$ with equality iff $\vec{x} = \vec{0}$
- (b) *(symmetry)* $|\vec{x} + \vec{y}| = |\vec{y} + \vec{x}|$
- (c) *(triangle inequality)* $|\vec{x} + \vec{y}| \leq |\vec{x}| + |\vec{y}|$

It also satisfies

- (a) *(scaling)* $|c\vec{x}| = |c| |\vec{x}|$
- (b) *(Cauchy-Schwartz)* $|\vec{x} \cdot \vec{y}| \leq |\vec{x}| |\vec{y}|$

Open sets

An **open ball** of radius r centered at \vec{a} is

Open sets

An **open ball** of radius r centered at \vec{a} is

$$B(\vec{a}; r) = \{\vec{x} : |\vec{x} - \vec{a}| < r\}.$$

Open sets

An **open ball** of radius r centered at \vec{a} is

$$B(\vec{a}; r) = \{\vec{x} : |\vec{x} - \vec{a}| < r\}.$$

Definition

Open sets

An **open ball** of radius r centered at \vec{a} is

$$B(\vec{a}; r) = \{\vec{x} : |\vec{x} - \vec{a}| < r\}.$$

Definition

A point \vec{a} in a subset $A \subseteq \mathbb{R}^n$ is called an **interior point** of A if there exists $r > 0$ with $B(\vec{a}; r) \subseteq A$.

Open sets

An **open ball** of radius r centered at \vec{a} is

$$B(\vec{a}; r) = \{\vec{x} : |\vec{x} - \vec{a}| < r\}.$$

Definition

A point \vec{a} in a subset $A \subseteq \mathbb{R}^n$ is called an **interior point** of A if there exists $r > 0$ with $B(\vec{a}; r) \subseteq A$. If every point of A is an interior point, then A is called an **open set**.

Challenge!

Problem

Prove that the empty set \emptyset and the whole space \mathbb{R}^n are open.

Challenge!

Problem

Let $\vec{a} \in \mathbb{R}^n$. Show that the singleton set

$$A = \{\vec{a}\}$$

is not open.

Challenge!

Problem

Let $a, b, c, d \in \mathbb{R}$ with $a < b$ and $c < d$. Prove that the **open square**

$$(a, b) \times (c, d) = \{(x, y) : a < x < b, \ c < y < d\}$$

is an open set

Challenge!

Problem

Prove that an open ball is an open set.

Unions of open sets are open

Theorem (Open Union Theorem)

*Suppose that $\{U_i : i \in I\}$ is an arbitrary family of open sets.
Then $\bigcup_{i \in I} U_i$ is open.*

Unions of open sets are open

Theorem (Open Union Theorem)

Suppose that $\{U_i : i \in I\}$ is an arbitrary family of open sets. Then $\bigcup_{i \in I} U_i$ is open.

Proof.

Unions of open sets are open

Theorem (Open Union Theorem)

*Suppose that $\{U_i : i \in I\}$ is an arbitrary family of open sets.
Then $\bigcup_{i \in I} U_i$ is open.*

Proof.

Suppose that $\vec{x} \in \bigcup_{i \in I} U_i$.

Unions of open sets are open

Theorem (Open Union Theorem)

*Suppose that $\{U_i : i \in I\}$ is an arbitrary family of open sets.
Then $\bigcup_{i \in I} U_i$ is open.*

Proof.

Suppose that $\vec{x} \in \bigcup_{i \in I} U_i$.
Then there exists $j \in I$ with $\vec{x} \in U_j$.

Unions of open sets are open

Theorem (Open Union Theorem)

*Suppose that $\{U_i : i \in I\}$ is an arbitrary family of open sets.
Then $\bigcup_{i \in I} U_i$ is open.*

Proof.

Suppose that $\vec{x} \in \bigcup_{i \in I} U_i$.

Then there exists $j \in I$ with $\vec{x} \in U_j$.

Since U_j is open, this means \vec{x} is an interior point of U_j .

Unions of open sets are open

Theorem (Open Union Theorem)

*Suppose that $\{U_i : i \in I\}$ is an arbitrary family of open sets.
Then $\bigcup_{i \in I} U_i$ is open.*

Proof.

Suppose that $\vec{x} \in \bigcup_{i \in I} U_i$.

Then there exists $j \in I$ with $\vec{x} \in U_j$.

Since U_j is open, this means \vec{x} is an interior point of U_j .

Therefore there exists $r > 0$ such that $B(\vec{x}; r) \subseteq U_j$.

Unions of open sets are open

Theorem (Open Union Theorem)

*Suppose that $\{U_i : i \in I\}$ is an arbitrary family of open sets.
Then $\bigcup_{i \in I} U_i$ is open.*

Proof.

Suppose that $\vec{x} \in \bigcup_{i \in I} U_i$.

Then there exists $j \in I$ with $\vec{x} \in U_j$.

Since U_j is open, this means \vec{x} is an interior point of U_j .

Therefore there exists $r > 0$ such that $B(\vec{x}; r) \subseteq U_j$.

This means $B(\vec{x}; r) \subseteq \bigcup_{i \in I} U_i$.

Unions of open sets are open

Theorem (Open Union Theorem)

*Suppose that $\{U_i : i \in I\}$ is an arbitrary family of open sets.
Then $\bigcup_{i \in I} U_i$ is open.*

Proof.

Suppose that $\vec{x} \in \bigcup_{i \in I} U_i$.

Then there exists $j \in I$ with $\vec{x} \in U_j$.

Since U_j is open, this means \vec{x} is an interior point of U_j .

Therefore there exists $r > 0$ such that $B(\vec{x}; r) \subseteq U_j$.

This means $B(\vec{x}; r) \subseteq \bigcup_{i \in I} U_i$.

Thus \vec{x} is an interior point of $\bigcup_{i \in I} U_i$.

Unions of open sets are open

Theorem (Open Union Theorem)

*Suppose that $\{U_i : i \in I\}$ is an arbitrary family of open sets.
Then $\bigcup_{i \in I} U_i$ is open.*

Proof.

Suppose that $\vec{x} \in \bigcup_{i \in I} U_i$.

Then there exists $j \in I$ with $\vec{x} \in U_j$.

Since U_j is open, this means \vec{x} is an interior point of U_j .

Therefore there exists $r > 0$ such that $B(\vec{x}; r) \subseteq U_j$.

This means $B(\vec{x}; r) \subseteq \bigcup_{i \in I} U_i$.

Thus \vec{x} is an interior point of $\bigcup_{i \in I} U_i$.

Since \vec{x} is arbitrary, this proves that $\bigcup_{i \in I} U_i$ is open. □

Finite intersections of open sets are open

Theorem (Open Intersection Theorem)

Suppose that U_1, U_2, \dots, U_n are open sets. Then $\bigcap_{i=1}^n U_i$ is open.

Finite intersections of open sets are open

Theorem (Open Intersection Theorem)

Suppose that U_1, U_2, \dots, U_n are open sets. Then $\bigcap_{i=1}^n U_i$ is open.

Proof.

Finite intersections of open sets are open

Theorem (Open Intersection Theorem)

Suppose that U_1, U_2, \dots, U_n are open sets. Then $\bigcap_{i=1}^n U_i$ is open.

Proof.

Suppose that $\vec{x} \in \bigcap_{i=1}^n U_i$.

Finite intersections of open sets are open

Theorem (Open Intersection Theorem)

Suppose that U_1, U_2, \dots, U_n are open sets. Then $\bigcap_{i=1}^n U_i$ is open.

Proof.

Suppose that $\vec{x} \in \bigcap_{i=1}^n U_i$.
Then for all i , $\vec{x} \in U_i$.

Finite intersections of open sets are open

Theorem (Open Intersection Theorem)

Suppose that U_1, U_2, \dots, U_n are open sets. Then $\bigcap_{i=1}^n U_i$ is open.

Proof.

Suppose that $\vec{x} \in \bigcap_{i=1}^n U_i$.

Then for all i , $\vec{x} \in U_i$.

Since U_i is open, this means \vec{x} is an interior point of U_i .

Finite intersections of open sets are open

Theorem (Open Intersection Theorem)

Suppose that U_1, U_2, \dots, U_n are open sets. Then $\bigcap_{i=1}^n U_i$ is open.

Proof.

Suppose that $\vec{x} \in \bigcap_{i=1}^n U_i$.

Then for all i , $\vec{x} \in U_i$.

Since U_i is open, this means \vec{x} is an interior point of U_i .

Therefore there exists $r_i > 0$ such that $B(\vec{x}; r_i) \subseteq U_i$.

Finite intersections of open sets are open

Theorem (Open Intersection Theorem)

Suppose that U_1, U_2, \dots, U_n are open sets. Then $\bigcap_{i=1}^n U_i$ is open.

Proof.

Suppose that $\vec{x} \in \bigcap_{i=1}^n U_i$.

Then for all i , $\vec{x} \in U_i$.

Since U_i is open, this means \vec{x} is an interior point of U_i .

Therefore there exists $r_i > 0$ such that $B(\vec{x}; r_i) \subseteq U_i$.

Take $r = \min\{r_i : 1 \leq i \leq n\}$.

Finite intersections of open sets are open

Theorem (Open Intersection Theorem)

Suppose that U_1, U_2, \dots, U_n are open sets. Then $\bigcap_{i=1}^n U_i$ is open.

Proof.

Suppose that $\vec{x} \in \bigcap_{i=1}^n U_i$.

Then for all i , $\vec{x} \in U_i$.

Since U_i is open, this means \vec{x} is an interior point of U_i .

Therefore there exists $r_i > 0$ such that $B(\vec{x}; r_i) \subseteq U_i$.

Take $r = \min\{r_i : 1 \leq i \leq n\}$.

This means $B(\vec{x}; r) \subseteq B(\vec{x}; r_i) \subseteq U_i$ for all i .

Finite intersections of open sets are open

Theorem (Open Intersection Theorem)

Suppose that U_1, U_2, \dots, U_n are open sets. Then $\bigcap_{i=1}^n U_i$ is open.

Proof.

Suppose that $\vec{x} \in \bigcap_{i=1}^n U_i$.

Then for all i , $\vec{x} \in U_i$.

Since U_i is open, this means \vec{x} is an interior point of U_i .

Therefore there exists $r_i > 0$ such that $B(\vec{x}; r_i) \subseteq U_i$.

Take $r = \min\{r_i : 1 \leq i \leq n\}$.

This means $B(\vec{x}; r) \subseteq B(\vec{x}; r_i) \subseteq U_i$ for all i .

Therefore $B(\vec{x}; r) \subseteq \bigcap_{i=1}^n U_i$.

Finite intersections of open sets are open

Theorem (Open Intersection Theorem)

Suppose that U_1, U_2, \dots, U_n are open sets. Then $\bigcap_{i=1}^n U_i$ is open.

Proof.

Suppose that $\vec{x} \in \bigcap_{i=1}^n U_i$.

Then for all i , $\vec{x} \in U_i$.

Since U_i is open, this means \vec{x} is an interior point of U_i .

Therefore there exists $r_i > 0$ such that $B(\vec{x}; r_i) \subseteq U_i$.

Take $r = \min\{r_i : 1 \leq i \leq n\}$.

This means $B(\vec{x}; r) \subseteq B(\vec{x}; r_i) \subseteq U_i$ for all i .

Therefore $B(\vec{x}; r) \subseteq \bigcap_{i=1}^n U_i$.

Thus \vec{x} is an interior point of $\bigcap_{i=1}^n U_i$.

Finite intersections of open sets are open

Theorem (Open Intersection Theorem)

Suppose that U_1, U_2, \dots, U_n are open sets. Then $\bigcap_{i=1}^n U_i$ is open.

Proof.

Suppose that $\vec{x} \in \bigcap_{i=1}^n U_i$.

Then for all i , $\vec{x} \in U_i$.

Since U_i is open, this means \vec{x} is an interior point of U_i .

Therefore there exists $r_i > 0$ such that $B(\vec{x}; r_i) \subseteq U_i$.

Take $r = \min\{r_i : 1 \leq i \leq n\}$.

This means $B(\vec{x}; r) \subseteq B(\vec{x}; r_i) \subseteq U_i$ for all i .

Therefore $B(\vec{x}; r) \subseteq \bigcap_{i=1}^n U_i$.

Thus \vec{x} is an interior point of $\bigcap_{i=1}^n U_i$.

Since \vec{x} is arbitrary, this proves that $\bigcap_{i=1}^n U_i$ is open. □

Challenge

Problem

Show that the infinite intersection

$$\bigcap_{n=1}^{\infty} \left(-\frac{1}{n}, \frac{1}{n}\right)$$

is not open.

Component intervals

Let $U \subseteq \mathbb{R}$ be open.

Component intervals

Let $U \subseteq \mathbb{R}$ be open.

Definition

A **component interval** of U is an interval I with $I \subseteq U$ and with the property that if J is an interval and $I \subseteq J$, then $J \not\subseteq U$.

Component intervals

Let $U \subseteq \mathbb{R}$ be open.

Definition

A **component interval** of U is an interval I with $I \subseteq U$ and with the property that if J is an interval and $I \subseteq J$, then $J \not\subseteq U$.

- $(0, 1)$ is a component interval of $\mathbb{R} \setminus \{0, 1, 2, 3\}$

Component intervals

Let $U \subseteq \mathbb{R}$ be open.

Definition

A **component interval** of U is an interval I with $I \subseteq U$ and with the property that if J is an interval and $I \subseteq J$, then $J \not\subseteq U$.

- $(0, 1)$ is a component interval of $\mathbb{R} \setminus \{0, 1, 2, 3\}$
- $(-\infty, 0)$ is also a component interval of $\mathbb{R} \setminus \{0, 1, 2, 3\}$

Component intervals

Let $U \subseteq \mathbb{R}$ be open.

Definition

A **component interval** of U is an interval I with $I \subseteq U$ and with the property that if J is an interval and $I \subseteq J$, then $J \not\subseteq U$.

- $(0, 1)$ is a component interval of $\mathbb{R} \setminus \{0, 1, 2, 3\}$
- $(-\infty, 0)$ is also a component interval of $\mathbb{R} \setminus \{0, 1, 2, 3\}$
- we will show all open sets of \mathbb{R} are made of component intervals!

Component intervals

Lemma

If I_1 and I_2 are two component intervals of an open subset $U \subseteq \mathbb{R}$, then $I_1 = I_2$ or $I_1 \cap I_2 = \emptyset$.

Component intervals

Lemma

If I_1 and I_2 are two component intervals of an open subset $U \subseteq \mathbb{R}$, then $I_1 = I_2$ or $I_1 \cap I_2 = \emptyset$.

Proof.

Component intervals

Lemma

If I_1 and I_2 are two component intervals of an open subset $U \subseteq \mathbb{R}$, then $I_1 = I_2$ or $I_1 \cap I_2 = \emptyset$.

Proof.

Suppose that $I_1 \cap I_2 \neq \emptyset$.

Component intervals

Lemma

If I_1 and I_2 are two component intervals of an open subset $U \subseteq \mathbb{R}$, then $I_1 = I_2$ or $I_1 \cap I_2 = \emptyset$.

Proof.

Suppose that $I_1 \cap I_2 \neq \emptyset$.
Then $J := I_1 \cup I_2$ is an interval.

Component intervals

Lemma

If I_1 and I_2 are two component intervals of an open subset $U \subseteq \mathbb{R}$, then $I_1 = I_2$ or $I_1 \cap I_2 = \emptyset$.

Proof.

Suppose that $I_1 \cap I_2 \neq \emptyset$.

Then $J := I_1 \cup I_2$ is an interval.

Also $I_1 \subseteq U$ and $I_2 \subseteq U$, so $J \subseteq U$.

Component intervals

Lemma

If I_1 and I_2 are two component intervals of an open subset $U \subseteq \mathbb{R}$, then $I_1 = I_2$ or $I_1 \cap I_2 = \emptyset$.

Proof.

Suppose that $I_1 \cap I_2 \neq \emptyset$.

Then $J := I_1 \cup I_2$ is an interval.

Also $I_1 \subseteq U$ and $I_2 \subseteq U$, so $J \subseteq U$.

Since I_1 is a component interval and $I_1 \subseteq J \subseteq U$, we have $I_1 = J$.

Component intervals

Lemma

If I_1 and I_2 are two component intervals of an open subset $U \subseteq \mathbb{R}$, then $I_1 = I_2$ or $I_1 \cap I_2 = \emptyset$.

Proof.

Suppose that $I_1 \cap I_2 \neq \emptyset$.

Then $J := I_1 \cup I_2$ is an interval.

Also $I_1 \subseteq U$ and $I_2 \subseteq U$, so $J \subseteq U$.

Since I_i is a component interval and $I_i \subseteq J \subseteq U$, we have $I_i = J$.
In particular $I_1 = J = I_2$. □

Component intervals

Theorem (Apostol 3.10)

If $U \subseteq \mathbb{R}$ is open and $x \in U$, then there is a unique component interval of U containing x .

Component intervals

Theorem (Apostol 3.10)

If $U \subseteq \mathbb{R}$ is open and $x \in U$, then there is a unique component interval of U containing x .

Proof.

Uniqueness follows from previous Lemma, so we only need existence.

Component intervals

Theorem (Apostol 3.10)

If $U \subseteq \mathbb{R}$ is open and $x \in U$, then there is a unique component interval of U containing x .

Proof.

Uniqueness follows from previous Lemma, so we only need existence.

Suppose that $x \in U$ and consider

$$A = \{a : (a, x) \subseteq S\}, \quad \text{and} \quad B = \{b : (x, b) \subseteq S\}$$

Component intervals

Theorem (Apostol 3.10)

If $U \subseteq \mathbb{R}$ is open and $x \in U$, then there is a unique component interval of U containing x .

Proof.

Uniqueness follows from previous Lemma, so we only need existence.

Suppose that $x \in U$ and consider

$$A = \{a : (a, x) \subseteq S\}, \quad \text{and} \quad B = \{b : (x, b) \subseteq S\}$$

If A is not bounded below, let $a = -\infty$. Otherwise, let $a = \inf(A)$.

Component intervals

Theorem (Apostol 3.10)

If $U \subseteq \mathbb{R}$ is open and $x \in U$, then there is a unique component interval of U containing x .

Proof.

Uniqueness follows from previous Lemma, so we only need existence.

Suppose that $x \in U$ and consider

$$A = \{a : (a, x) \subseteq S\}, \quad \text{and} \quad B = \{b : (x, b) \subseteq S\}$$

If A is not bounded below, let $a = -\infty$. Otherwise, let $a = \inf(A)$.

If B is not bounded above, let $b = \infty$. Otherwise, let $b = \sup(B)$.

Component intervals

Theorem (Apostol 3.10)

If $U \subseteq \mathbb{R}$ is open and $x \in U$, then there is a unique component interval of U containing x .

Proof.

Uniqueness follows from previous Lemma, so we only need existence.

Suppose that $x \in U$ and consider

$$A = \{a : (a, x) \subseteq S\}, \quad \text{and} \quad B = \{b : (x, b) \subseteq S\}$$

If A is not bounded below, let $a = -\infty$. Otherwise, let $a = \inf(A)$.

If B is not bounded above, let $b = \infty$. Otherwise, let $b = \sup(B)$.

Claim: (a, b) is a component interval of U containing x . □

Challenge

Problem

Prove that (a, b) is a component interval of U .

Open subsets of \mathbb{R}

Theorem (Representation Theorem for Open Intervals in \mathbb{R})

If $U \subseteq \mathbb{R}$ is open, then U is the union of a countable family of disjoint open intervals.