

# MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics  
California State University Fullerton

October 21, 2024

# Outline

- 1 Real Analysis Lecture 14
  - More on limits
  - Cauchy sequences

# Outline

- 1 Real Analysis Lecture 14
  - More on limits
  - Cauchy sequences

# Existence of limits

## Theorem

*If a sequence  $\{x_n\}$  in a metric space  $(M, d)$  converges to a value  $L \in M$ , then the range*

$$X = \{x_1, x_2, \dots\} \subseteq M$$

*is a bounded set and  $L$  is an adherent point of  $X$ .*

# Existence of limits

## Theorem

*If a sequence  $\{x_n\}$  in a metric space  $(M, d)$  converges to a value  $L \in M$ , then the range*

$$X = \{x_1, x_2, \dots\} \subseteq M$$

*is a bounded set and  $L$  is an adherent point of  $X$ .*

When the range of a sequence is bounded, we call the sequence  $\{x_n\}$  bounded.

# Existence of limits

Proof.

Suppose that  $\lim_{n \rightarrow \infty} x_n = L$ .

# Existence of limits

Proof.

Suppose that  $\lim_{n \rightarrow \infty} x_n = L$ .

Let  $\epsilon = 2024$ .

# Existence of limits

Proof.

Suppose that  $\lim_{n \rightarrow \infty} x_n = L$ .

Let  $\epsilon = 2024$ .

Then there exists  $N \in \mathbb{Z}_+$  with  $d(x_n, L) \leq 2024$  for all  $n \geq N$ .



# Existence of limits

Proof.

Suppose that  $\lim_{n \rightarrow \infty} x_n = L$ .

Let  $\epsilon = 2024$ .

Then there exists  $N \in \mathbb{Z}_+$  with  $d(x_n, L) \leq 2024$  for all  $n \geq N$ .

If we take

$$R = \max\{d(x_1, L), d(x_2, L), \dots, d(x_{N-1}, L), 2024\},$$

# Existence of limits

Proof.

Suppose that  $\lim_{n \rightarrow \infty} x_n = L$ .

Let  $\epsilon = 2024$ .

Then there exists  $N \in \mathbb{Z}_+$  with  $d(x_n, L) \leq 2024$  for all  $n \geq N$ .

If we take

$$R = \max\{d(x_1, L), d(x_2, L), \dots, d(x_{N-1}, L), 2024\},$$

then  $d(x_n, L) \leq R$  for all  $N$ .

# Existence of limits

Proof.

Suppose that  $\lim_{n \rightarrow \infty} x_n = L$ .

Let  $\epsilon = 2024$ .

Then there exists  $N \in \mathbb{Z}_+$  with  $d(x_n, L) \leq 2024$  for all  $n \geq N$ .

If we take

$$R = \max\{d(x_1, L), d(x_2, L), \dots, d(x_{N-1}, L), 2024\},$$

then  $d(x_n, L) \leq R$  for all  $N$ .

Therefore  $X \subseteq B_M(L, R)$  and in particular  $X$  is bounded. □

# Existence of limits

Sort of a converse of the previous theorem.

# Existence of limits

Sort of a converse of the previous theorem.

## Theorem

*If  $S$  is a subset of a metric space  $(M, d)$  and  $L \in M$  is an adherent point of  $S$ , then there exists a sequence  $\{x_n\}$  of elements of  $S$  which converges to  $L$ .*

# Existence of limits

Proof.

Suppose  $L$  is an adherent point of  $S$ .

# Existence of limits

## Proof.

Suppose  $L$  is an adherent point of  $S$ .

Then for all  $n \in \mathbb{Z}_+$ , the ball  $B_M(x, \frac{1}{n})$  contains at least one element of  $S$ .

# Existence of limits

## Proof.

Suppose  $L$  is an adherent point of  $S$ .

Then for all  $n \in \mathbb{Z}_+$ , the ball  $B_M(x, \frac{1}{n})$  contains at least one element of  $S$ .

For each  $n \in \mathbb{Z}_+$ , choose  $x_n \in S \cap B_M(x, \frac{1}{n})$ .



# Existence of limits

## Proof.

Suppose  $L$  is an adherent point of  $S$ .

Then for all  $n \in \mathbb{Z}_+$ , the ball  $B_M(x, \frac{1}{n})$  contains at least one element of  $S$ .

For each  $n \in \mathbb{Z}_+$ , choose  $x_n \in S \cap B_M(x, \frac{1}{n})$ .

We claim  $\lim x_n = L$ .

# Existence of limits

## Proof.

Suppose  $L$  is an adherent point of  $S$ .

Then for all  $n \in \mathbb{Z}_+$ , the ball  $B_M(x, \frac{1}{n})$  contains at least one element of  $S$ .

For each  $n \in \mathbb{Z}_+$ , choose  $x_n \in S \cap B_M(x, \frac{1}{n})$ .

We claim  $\lim x_n = L$ .

To see this, let  $\epsilon > 0$ .

# Existence of limits

## Proof.

Suppose  $L$  is an adherent point of  $S$ .

Then for all  $n \in \mathbb{Z}_+$ , the ball  $B_M(x, \frac{1}{n})$  contains at least one element of  $S$ .

For each  $n \in \mathbb{Z}_+$ , choose  $x_n \in S \cap B_M(x, \frac{1}{n})$ .

We claim  $\lim x_n = L$ .

To see this, let  $\epsilon > 0$ .

Then choose  $N \in \mathbb{Z}_+$  with  $N > 1/\epsilon$ .

# Existence of limits

## Proof.

Suppose  $L$  is an adherent point of  $S$ .

Then for all  $n \in \mathbb{Z}_+$ , the ball  $B_M(x, \frac{1}{n})$  contains at least one element of  $S$ .

For each  $n \in \mathbb{Z}_+$ , choose  $x_n \in S \cap B_M(x, \frac{1}{n})$ .

We claim  $\lim x_n = L$ .

To see this, let  $\epsilon > 0$ .

Then choose  $N \in \mathbb{Z}_+$  with  $N > 1/\epsilon$ .

It follows that for all  $n \geq N$

$$d(x_n, L) < 1/n < 1/N < \epsilon.$$

# Existence of limits

## Proof.

Suppose  $L$  is an adherent point of  $S$ .

Then for all  $n \in \mathbb{Z}_+$ , the ball  $B_M(x, \frac{1}{n})$  contains at least one element of  $S$ .

For each  $n \in \mathbb{Z}_+$ , choose  $x_n \in S \cap B_M(x, \frac{1}{n})$ .

We claim  $\lim x_n = L$ .

To see this, let  $\epsilon > 0$ .

Then choose  $N \in \mathbb{Z}_+$  with  $N > 1/\epsilon$ .

It follows that for all  $n \geq N$

$$d(x_n, L) < 1/n < 1/N < \epsilon.$$

Since  $\epsilon > 0$  was arbitrary, this proves convergence. □

# Existence of limits

Recall a **subsequence** of a sequence  $\{x_n\}$  is a sequence of the form  $\{x_{k(n)}\}$ , for  $k(1) < k(2) < \dots$  positive integers.

# Existence of limits

Recall a **subsequence** of a sequence  $\{x_n\}$  is a sequence of the form  $\{x_{k(n)}\}$ , for  $k(1) < k(2) < \dots$  positive integers.

## Theorem

*A sequence  $\{x_n\}$  in a metric space  $(M, d)$  converges to a value  $L \in M$  if and only if every subsequence  $\{x_{k(n)}\}$  also converges to  $L$ .*

# Existence of limits

Proof.

Suppose that every subsequence  $\{x_{k(n)}\}$  of  $\{x_n\}$  converges to  $L$ .



# Existence of limits

## Proof.

Suppose that every subsequence  $\{x_{k(n)}\}$  of  $\{x_n\}$  converges to  $L$ .

A sequence is a subsequence of itself! Thus  $\{x_n\}$  will converge to  $L$ .

# Existence of limits

## Proof.

Suppose that every subsequence  $\{x_{k(n)}\}$  of  $\{x_n\}$  converges to  $L$ .

A sequence is a subsequence of itself! Thus  $\{x_n\}$  will converge to  $L$ .

Conversely, suppose that  $\{x_n\}$  converges to  $L$ .

# Existence of limits

## Proof.

Suppose that every subsequence  $\{x_{k(n)}\}$  of  $\{x_n\}$  converges to  $L$ .

A sequence is a subsequence of itself! Thus  $\{x_n\}$  will converge to  $L$ .

Conversely, suppose that  $\{x_n\}$  converges to  $L$ .

Let  $\{x_{k(n)}\}$  be a subsequence of  $\{x_n\}$ .

# Existence of limits

## Proof.

Suppose that every subsequence  $\{x_{k(n)}\}$  of  $\{x_n\}$  converges to  $L$ .

A sequence is a subsequence of itself! Thus  $\{x_n\}$  will converge to  $L$ .

Conversely, suppose that  $\{x_n\}$  converges to  $L$ .

Let  $\{x_{k(n)}\}$  be a subsequence of  $\{x_n\}$ .

To show it limits to  $L$ , let  $\epsilon > 0$ .

# Existence of limits

## Proof.

Suppose that every subsequence  $\{x_{k(n)}\}$  of  $\{x_n\}$  converges to  $L$ .

A sequence is a subsequence of itself! Thus  $\{x_n\}$  will converge to  $L$ .

Conversely, suppose that  $\{x_n\}$  converges to  $L$ .

Let  $\{x_{k(n)}\}$  be a subsequence of  $\{x_n\}$ .

To show it limits to  $L$ , let  $\epsilon > 0$ .

Then there exists  $N \in \mathbb{Z}_+$  such that  $n \geq N$  implies  $d(x_n, L) < \epsilon$ .

# Existence of limits

## Proof.

Suppose that every subsequence  $\{x_{k(n)}\}$  of  $\{x_n\}$  converges to  $L$ .

A sequence is a subsequence of itself! Thus  $\{x_n\}$  will converge to  $L$ .

Conversely, suppose that  $\{x_n\}$  converges to  $L$ .

Let  $\{x_{k(n)}\}$  be a subsequence of  $\{x_n\}$ .

To show it limits to  $L$ , let  $\epsilon > 0$ .

Then there exists  $N \in \mathbb{Z}_+$  such that  $n \geq N$  implies  $d(x_n, L) < \epsilon$ .

Moreover,  $k(n) \geq n$  so  $k(n) \geq N$  and therefore  $d(x_{k(n)}, L) < \epsilon$ .

# Existence of limits

## Proof.

Suppose that every subsequence  $\{x_{k(n)}\}$  of  $\{x_n\}$  converges to  $L$ .

A sequence is a subsequence of itself! Thus  $\{x_n\}$  will converge to  $L$ .

Conversely, suppose that  $\{x_n\}$  converges to  $L$ .

Let  $\{x_{k(n)}\}$  be a subsequence of  $\{x_n\}$ .

To show it limits to  $L$ , let  $\epsilon > 0$ .

Then there exists  $N \in \mathbb{Z}_+$  such that  $n \geq N$  implies  $d(x_n, L) < \epsilon$ .

Moreover,  $k(n) \geq n$  so  $k(n) \geq N$  and therefore  $d(x_{k(n)}, L) < \epsilon$ .

Since  $\epsilon > 0$  was arbitrary, this proves convergence.  $\square$

# Monotone sequences

## Definition



# Monotone sequences

## Definition

A sequence  $\{x_n\}$  of real numbers is called **increasing** or **monotone increasing** if  $x_n \leq x_{n+1}$  for all  $n$ .

# Monotone sequences

## Definition

A sequence  $\{x_n\}$  of real numbers is called **increasing** or **monotone increasing** if  $x_n \leq x_{n+1}$  for all  $n$ .

It is called **strictly increasing** if  $x_n < x_{n+1}$  for all  $n$ .

# Monotone sequences

## Definition

A sequence  $\{x_n\}$  of real numbers is called **increasing** or **monotone increasing** if  $x_n \leq x_{n+1}$  for all  $n$ .

It is called **strictly increasing** if  $x_n < x_{n+1}$  for all  $n$ .

A sequence  $\{x_n\}$  of real numbers is called **decreasing** or **monotone decreasing** if  $x_n \geq x_{n+1}$  for all  $n$ .

# Monotone sequences

## Definition

A sequence  $\{x_n\}$  of real numbers is called **increasing** or **monotone increasing** if  $x_n \leq x_{n+1}$  for all  $n$ .

It is called **strictly increasing** if  $x_n < x_{n+1}$  for all  $n$ .

A sequence  $\{x_n\}$  of real numbers is called **decreasing** or **monotone decreasing** if  $x_n \geq x_{n+1}$  for all  $n$ .

It is called **strictly decreasing** if  $x_n > x_{n+1}$  for all  $n$ .

# Monotone sequences

## Definition

A sequence  $\{x_n\}$  of real numbers is called **increasing** or **monotone increasing** if  $x_n \leq x_{n+1}$  for all  $n$ .

It is called **strictly increasing** if  $x_n < x_{n+1}$  for all  $n$ .

A sequence  $\{x_n\}$  of real numbers is called **decreasing** or **monotone decreasing** if  $x_n \geq x_{n+1}$  for all  $n$ .

It is called **strictly decreasing** if  $x_n > x_{n+1}$  for all  $n$ .

In any of these cases, we call the sequence **monotone**.

# Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

# Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

Theorem (Monotone convergence theorem)

# Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

## Theorem (Monotone convergence theorem)

*Consider the metric space formed by  $\mathbb{R}$  with Euclidean distance.*



# Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

## Theorem (Monotone convergence theorem)

*Consider the metric space formed by  $\mathbb{R}$  with Euclidean distance.*

*A monotone increasing sequence  $\{x_n\}$  of real numbers which is bounded above converges to  $L = \sup\{x_1, x_2, \dots\}$ .*

# Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

## Theorem (Monotone convergence theorem)

*Consider the metric space formed by  $\mathbb{R}$  with Euclidean distance.*

*A monotone increasing sequence  $\{x_n\}$  of real numbers which is bounded above converges to  $L = \sup\{x_1, x_2, \dots\}$ .*

*A monotone decreasing sequence  $\{x_n\}$  which is bounded below converges to  $L = \inf\{x_1, x_2, \dots\}$ .*

# Monotone sequences

Proof.

# Monotone sequences

Proof.

Suppose that  $\{x_n\}$  is a monotone increasing sequence which is bounded above and let  $X = \{x_1, x_2, \dots\}$ .

# Monotone sequences

## Proof.

Suppose that  $\{x_n\}$  is a monotone increasing sequence which is bounded above and let  $X = \{x_1, x_2, \dots\}$ . Then by the Completeness Axiom  $L = \sup X$  exists.

# Monotone sequences

## Proof.

Suppose that  $\{x_n\}$  is a monotone increasing sequence which is bounded above and let  $X = \{x_1, x_2, \dots\}$ . Then by the Completeness Axiom  $L = \sup X$  exists.

Let  $\epsilon > 0$ .

# Monotone sequences

## Proof.

Suppose that  $\{x_n\}$  is a monotone increasing sequence which is bounded above and let  $X = \{x_1, x_2, \dots\}$ . Then by the Completeness Axiom  $L = \sup X$  exists.

Let  $\epsilon > 0$ .

Then  $L - \epsilon$  is not a supremum of  $X$ , so there exists  $x_N \in X$  such that  $x_N > L - \epsilon$ .

# Monotone sequences

## Proof.

Suppose that  $\{x_n\}$  is a monotone increasing sequence which is bounded above and let  $X = \{x_1, x_2, \dots\}$ . Then by the Completeness Axiom  $L = \sup X$  exists.

Let  $\epsilon > 0$ .

Then  $L - \epsilon$  is not a supremum of  $X$ , so there exists  $x_N \in X$  such that  $x_N > L - \epsilon$ .

Since the sequence is increasing, for any  $n > N$ , we have that  $L - \epsilon < x_N \leq x_n < L$ .



# Monotone sequences

## Proof.

Suppose that  $\{x_n\}$  is a monotone increasing sequence which is bounded above and let  $X = \{x_1, x_2, \dots\}$ . Then by the Completeness Axiom  $L = \sup X$  exists.

Let  $\epsilon > 0$ .

Then  $L - \epsilon$  is not a supremum of  $X$ , so there exists  $x_N \in X$  such that  $x_N > L - \epsilon$ .

Since the sequence is increasing, for any  $n > N$ , we have that  $L - \epsilon < x_N \leq x_n < L$ .

It follows that

$$d(x_n, L) = |x_n - L| = L - x_n < \epsilon.$$

# Monotone sequences

## Proof.

Suppose that  $\{x_n\}$  is a monotone increasing sequence which is bounded above and let  $X = \{x_1, x_2, \dots\}$ . Then by the Completeness Axiom  $L = \sup X$  exists.

Let  $\epsilon > 0$ .

Then  $L - \epsilon$  is not a supremum of  $X$ , so there exists  $x_N \in X$  such that  $x_N > L - \epsilon$ .

Since the sequence is increasing, for any  $n > N$ , we have that  $L - \epsilon < x_N \leq x_n < L$ .

It follows that

$$d(x_n, L) = |x_n - L| = L - x_n < \epsilon.$$

Since  $\epsilon > 0$  was arbitrary, this proves  $x_n$  converges to  $L$ . □

# Outline

- 1 Real Analysis Lecture 14
  - More on limits
  - Cauchy sequences

# Cauchy sequence

A sequence  $\{x_n\}$  in a metric space  $(M, d)$  is called Cauchy if the terms get arbitrarily close together as  $n$  gets arbitrarily large.

# Cauchy sequence

A sequence  $\{x_n\}$  in a metric space  $(M, d)$  is called Cauchy if the terms get arbitrarily close together as  $n$  gets arbitrarily large.

More formally:

# Cauchy sequence

A sequence  $\{x_n\}$  in a metric space  $(M, d)$  is called Cauchy if the terms get arbitrarily close together as  $n$  gets arbitrarily large.

More formally:

## Definition

A sequence  $\{x_n\}$  is called a **Cauchy sequence** if

$$\forall \epsilon > 0, \text{ there exists } N \in \mathbb{Z}_+ \text{ with } m, n \geq N \Rightarrow d(x_m, x_n) < \epsilon.$$

# Cauchy sequence

A sequence  $\{x_n\}$  in a metric space  $(M, d)$  is called Cauchy if the terms get arbitrarily close together as  $n$  gets arbitrarily large.

More formally:

## Definition

A sequence  $\{x_n\}$  is called a **Cauchy sequence** if

$$\forall \epsilon > 0, \text{ there exists } N \in \mathbb{Z}_+ \text{ with } m, n \geq N \Rightarrow d(x_m, x_n) < \epsilon.$$

- every convergent sequence is Cauchy

# Cauchy sequence

A sequence  $\{x_n\}$  in a metric space  $(M, d)$  is called Cauchy if the terms get arbitrarily close together as  $n$  gets arbitrarily large.

More formally:

## Definition

A sequence  $\{x_n\}$  is called a **Cauchy sequence** if

$$\forall \epsilon > 0, \text{ there exists } N \in \mathbb{Z}_+ \text{ with } m, n \geq N \Rightarrow d(x_m, x_n) < \epsilon.$$

- every convergent sequence is Cauchy
- is every Cauchy sequence convergent?



# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

Suppose  $\{x_n\}$  is Cauchy.

# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

Suppose  $\{x_n\}$  is Cauchy.  
Let  $\epsilon = 2024$ .

# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

Suppose  $\{x_n\}$  is Cauchy.

Let  $\epsilon = 2024$ .

Then  $\exists N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < 2024$  for all  $m, n \geq N$ .

# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

Suppose  $\{x_n\}$  is Cauchy.

Let  $\epsilon = 2024$ .

Then  $\exists N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < 2024$  for all  $m, n \geq N$ .

Therefore for all  $n \geq N$ , we have  $d(x_n, x_N) \leq 2024$ .

# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

Suppose  $\{x_n\}$  is Cauchy.

Let  $\epsilon = 2024$ .

Then  $\exists N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < 2024$  for all  $m, n \geq N$ .

Therefore for all  $n \geq N$ , we have  $d(x_n, x_N) \leq 2024$ .

So if we define

$$R = \max\{d(x_1, x_N), d(x_2, x_N), \dots, d(x_{n-1}, x_N), 2024\},$$

# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

Suppose  $\{x_n\}$  is Cauchy.

Let  $\epsilon = 2024$ .

Then  $\exists N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < 2024$  for all  $m, n \geq N$ .

Therefore for all  $n \geq N$ , we have  $d(x_n, x_N) \leq 2024$ .

So if we define

$$R = \max\{d(x_1, x_N), d(x_2, x_N), \dots, d(x_{n-1}, x_N), 2024\},$$

then  $d(x_n, x_N) \leq 2024$  for all  $n \geq 1$ .



# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

Suppose  $\{x_n\}$  is Cauchy.

Let  $\epsilon = 2024$ .

Then  $\exists N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < 2024$  for all  $m, n \geq N$ .

Therefore for all  $n \geq N$ , we have  $d(x_n, x_N) \leq 2024$ .

So if we define

$$R = \max\{d(x_1, x_N), d(x_2, x_N), \dots, d(x_{n-1}, x_N), 2024\},$$

then  $d(x_n, x_N) \leq 2024$  for all  $n \geq 1$ .

Hence  $X = \{x_1, x_2, \dots\} \subseteq B_M(x_N, R)$ , and is bounded.