

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

September 15, 2024

Outline

- 1 Real Analysis Lecture 5
 - Sets, Relations, Functions

Outline

- 1 Real Analysis Lecture 5
 - Sets, Relations, Functions

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$
- empty set \emptyset

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$
- empty set \emptyset
- $\{\heartsuit, \text{Fall}, \{\emptyset\}\}$

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$
- empty set \emptyset
- $\{\heartsuit, \text{Fall}, \{\emptyset\}\}$
- $\{n \in \mathbb{Z} : n \text{ is prime}\}$

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

- ordered pairs

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

- ordered pairs

$$(a, b) = \{a, \{a, b\}\}$$

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

- ordered pairs

$$(a, b) = \{a, \{a, b\}\}$$

- even relations and functions are sets!

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$$(a, b) = (c, d) \text{ if and only if } \{a, \{a, b\}\} = \{c, \{c, d\}\}$$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$(a, b) = (c, d)$ if and only if $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if $a = c$ and $b = d$, then $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$(a, b) = (c, d)$ if and only if $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if $a = c$ and $b = d$, then $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$(a, b) = (c, d)$ if and only if $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if $a = c$ and $b = d$, then $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

Suppose $\{a, \{a, b\}\} = \{c, \{c, d\}\}$.

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$(a, b) = (c, d)$ if and only if $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if $a = c$ and $b = d$, then $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

Suppose $\{a, \{a, b\}\} = \{c, \{c, d\}\}$.

Two possible cases:

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$(a, b) = (c, d)$ if and only if $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if $a = c$ and $b = d$, then $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

Suppose $\{a, \{a, b\}\} = \{c, \{c, d\}\}$.

Two possible cases:

Case I: $a = c$ and $\{a, b\} = \{c, d\}$

Case II: $a = \{c, d\}$ and $\{a, b\} = c$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$(a, b) = (c, d)$ if and only if $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if $a = c$ and $b = d$, then $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

Suppose $\{a, \{a, b\}\} = \{c, \{c, d\}\}$.

Two possible cases:

Case I: $a = c$ and $\{a, b\} = \{c, d\}$

Case II: $a = \{c, d\}$ and $\{a, b\} = c$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$$\text{Case I: } a = c \text{ and } \{a, b\} = \{c, d\}$$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

Case I: $a = c$ and $\{a, b\} = \{c, d\}$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

Case I: $a = c$ and $\{a, b\} = \{c, d\}$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Therefore $b = c$ or $b = d$.

If $b = d$, we're done!

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$$\text{Case I: } a = c \text{ and } \{a, b\} = \{c, d\}$$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Therefore $b = c$ or $b = d$.

If $b = d$, we're done! ... so assume instead that $b = c$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$$\text{Case I: } a = c \text{ and } \{a, b\} = \{c, d\}$$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Therefore $b = c$ or $b = d$.

If $b = d$, we're done! ... so assume instead that $b = c$

Then $a = c$ implies $a = b$.

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$$\text{Case I: } a = c \text{ and } \{a, b\} = \{c, d\}$$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Therefore $b = c$ or $b = d$.

If $b = d$, we're done! ... so assume instead that $b = c$

Then $a = c$ implies $a = b$.

Therefore $\{c, d\} = \{a, b\} = \{a, a\} = \{a\}$.

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

Case I: $a = c$ and $\{a, b\} = \{c, d\}$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Therefore $b = c$ or $b = d$.

If $b = d$, we're done! ... so assume instead that $b = c$

Then $a = c$ implies $a = b$.

Therefore $\{c, d\} = \{a, b\} = \{a, a\} = \{a\}$.

It follows that $d = c = b = a$.

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$$\text{Case II: } a = \{c, d\} \text{ and } \{a, b\} = c$$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$$\text{Case II: } a = \{c, d\} \text{ and } \{a, b\} = c$$

This would imply that $c \in a$ and $a \in c$.

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$$\text{Case II: } a = \{c, d\} \text{ and } \{a, b\} = c$$

This would imply that $c \in a$ and $a \in c$.

This can be shown to contradict the ZF Axioms of Set Theory.

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$$\text{Case II: } a = \{c, d\} \text{ and } \{a, b\} = c$$

This would imply that $c \in a$ and $a \in c$.

This can be shown to contradict the ZF Axioms of Set Theory.
Specifically the regularity axiom for the set $\{a, c\}$...

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** \mathcal{R} from A to B is a subset of $A \times B$.

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** \mathcal{R} from A to B is a subset of $A \times B$.

NOTATION: $a\mathcal{R}b$ means $(a, b) \in \mathcal{R}$.

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** \mathcal{R} from A to B is a subset of $A \times B$.

NOTATION: $a\mathcal{R}b$ means $(a, b) \in \mathcal{R}$.

Domain and codomain:

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** \mathcal{R} from A to B is a subset of $A \times B$.

NOTATION: $a\mathcal{R}b$ means $(a, b) \in \mathcal{R}$.

Domain and codomain:

$$\text{dom}(\mathcal{R}) = \{a \in A : \exists b \in B, a\mathcal{R}b\}$$

$$\text{codom}(\mathcal{R}) = \{b \in B : \exists a \in A, a\mathcal{R}b\}$$

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** \mathcal{R} from A to B is a subset of $A \times B$.

NOTATION: $a\mathcal{R}b$ means $(a, b) \in \mathcal{R}$.

Domain and codomain:

$$\begin{aligned}\text{dom}(\mathcal{R}) &= \{a \in A : \exists b \in B, a\mathcal{R}b\} \\ \text{codom}(\mathcal{R}) &= \{b \in B : \exists a \in A, a\mathcal{R}b\}\end{aligned}$$

Relations

A relation \mathcal{R} from A to A is called a **relation on A** . A relation on A is

Relations

A relation \mathcal{R} from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$

Relations

A relation \mathcal{R} from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$
- **symmetric** if $a\mathcal{R}b$ implies $b\mathcal{R}a$ for all $a, b \in A$

Relations

A relation \mathcal{R} from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$
- **symmetric** if $a\mathcal{R}b$ implies $b\mathcal{R}a$ for all $a, b \in A$
- **transitive** if $a\mathcal{R}b$ and $b\mathcal{R}c$ implies $a\mathcal{R}c$ for all $a, b, c \in A$

Relations

A relation \mathcal{R} from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$
- **symmetric** if $a\mathcal{R}b$ implies $b\mathcal{R}a$ for all $a, b \in A$
- **transitive** if $a\mathcal{R}b$ and $b\mathcal{R}c$ implies $a\mathcal{R}c$ for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Relations

A relation \mathcal{R} from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$
- **symmetric** if $a\mathcal{R}b$ implies $b\mathcal{R}a$ for all $a, b \in A$
- **transitive** if $a\mathcal{R}b$ and $b\mathcal{R}c$ implies $a\mathcal{R}c$ for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Examples:

Relations

A relation \mathcal{R} from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$
- **symmetric** if $a\mathcal{R}b$ implies $b\mathcal{R}a$ for all $a, b \in A$
- **transitive** if $a\mathcal{R}b$ and $b\mathcal{R}c$ implies $a\mathcal{R}c$ for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Examples:

Relations

A relation \mathcal{R} from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$
- **symmetric** if $a\mathcal{R}b$ implies $b\mathcal{R}a$ for all $a, b \in A$
- **transitive** if $a\mathcal{R}b$ and $b\mathcal{R}c$ implies $a\mathcal{R}c$ for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Examples:

- $< = \{(x, y) : y - x \in (0, \infty)\}$ is transitive but not reflexive or symmetric on \mathbb{R}

Relations

A relation \mathcal{R} from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$
- **symmetric** if $a\mathcal{R}b$ implies $b\mathcal{R}a$ for all $a, b \in A$
- **transitive** if $a\mathcal{R}b$ and $b\mathcal{R}c$ implies $a\mathcal{R}c$ for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Examples:

- $< = \{(x, y) : y - x \in (0, \infty)\}$ is transitive but not reflexive or symmetric on \mathbb{R}
- $\leq = \{(x, y) : y - x \in [0, \infty)\}$ is reflexive and transitive but not symmetric on \mathbb{R}

Challenge!

Problem

Give an example of a relation on \mathbb{R} which is symmetric and transitive but not reflexive.

Challenge!

Problem

Give an example of a relation on \mathbb{R} which is reflexive and symmetric but not transitive.