

# MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics  
California State University Fullerton

August 25, 2024

# Outline

- 1 Real Analysis Lecture 1
  - Origin of Real Numbers
  - Properties of real numbers
  - Integers

# Outline

- 1 Real Analysis Lecture 1
  - Origin of Real Numbers
  - Properties of real numbers
  - Integers

# Prehistoric numbers



20,000 BC tallies on Ishango bone

𐎶 1	𐎶𐎶 11	𐎶𐎶𐎶 21	𐎶𐎶𐎶𐎶 31	𐎶𐎶𐎶𐎶𐎶 41	𐎶𐎶𐎶𐎶𐎶𐎶 51
𐎶𐎶 2	𐎶𐎶𐎶 12	𐎶𐎶𐎶𐎶 22	𐎶𐎶𐎶𐎶𐎶 32	𐎶𐎶𐎶𐎶𐎶𐎶 42	𐎶𐎶𐎶𐎶𐎶𐎶𐎶 52
𐎶𐎶𐎶 3	𐎶𐎶𐎶𐎶 13	𐎶𐎶𐎶𐎶𐎶 23	𐎶𐎶𐎶𐎶𐎶𐎶 33	𐎶𐎶𐎶𐎶𐎶𐎶𐎶 43	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 53
𐎶𐎶𐎶𐎶 4	𐎶𐎶𐎶𐎶𐎶 14	𐎶𐎶𐎶𐎶𐎶𐎶 24	𐎶𐎶𐎶𐎶𐎶𐎶𐎶 34	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 44	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 54
𐎶𐎶𐎶𐎶𐎶 5	𐎶𐎶𐎶𐎶𐎶𐎶 15	𐎶𐎶𐎶𐎶𐎶𐎶𐎶 25	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 35	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 45	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 55
𐎶𐎶𐎶𐎶𐎶𐎶 6	𐎶𐎶𐎶𐎶𐎶𐎶𐎶 16	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 26	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 36	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 46	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 56
𐎶𐎶𐎶𐎶𐎶𐎶𐎶 7	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 17	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 27	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 37	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 47	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 57
𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 8	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 18	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 28	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 38	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 48	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 58
𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 9	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 19	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 29	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 39	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 49	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 59
𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 10	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 20	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 30	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 40	𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 50	



$$\frac{1}{3}$$



$$\frac{1}{5}$$



$$\frac{1}{6}$$



$$\frac{1}{10}$$

3400 BC Sumerian system

1000 BC Egyptian fractions

# Invention of zero

# Invention of zero

- 1770 BC - concept of zero in heiroglyph

# Invention of zero

- 1770 BC - concept of zero in heiroglyph
- 36 BC - Mayan/Olmec heiroglyph for zero

# Invention of zero

- 1770 BC - concept of zero in heiroglyph
- 36 BC - Mayan/Olmec heiroglyph for zero
- 150 AD - Ptolemy using it in astronomy



# Invention of zero

- 1770 BC - concept of zero in heiroglyph
- 36 BC - Mayan/Olmec heiroglyph for zero
- 150 AD - Ptolemy using it in astronomy
- medieval scholars - debating the existence of 0



**Babylonian Zero**



**Mayan Zero**



**Hebrew Zero**



**Egyptian Zero**

# Invention of negatives

# Invention of negatives

- 100-50 BC - concept of negatives in China

# Invention of negatives

- 100-50 BC - concept of negatives in China
- 628 AD - used in India in quadratic formula

# Invention of negatives

- 100-50 BC - concept of negatives in China
- 628 AD - used in India in quadratic formula
- 9th century - Islamic mathematicians invent algebra, but hesitant about negatives

# Invention of negatives

- 100-50 BC - concept of negatives in China
- 628 AD - used in India in quadratic formula
- 9th century - Islamic mathematicians invent algebra, but hesitant about negatives
- 12th century - Islamic mathematicians add negative solutions of quadratics, but discard them

# Invention of negatives

- 100-50 BC - concept of negatives in China
- 628 AD - used in India in quadratic formula
- 9th century - Islamic mathematicians invent algebra, but hesitant about negatives
- 12th century - Islamic mathematicians add negative solutions of quadratics, but discard them
- 1202,1225 - Fibonacci allows negatives as solutions for financial problems

# Invention of negatives

- 100-50 BC - concept of negatives in China
- 628 AD - used in India in quadratic formula
- 9th century - Islamic mathematicians invent algebra, but hesitant about negatives
- 12th century - Islamic mathematicians add negative solutions of quadratics, but discard them
- 1202, 1225 - Fibonacci allows negatives as solutions for financial problems
- up to 18th century - rejected by western sources, referred to as "absurd numbers"



# Invention of rationals

# Invention of rationals

- 1000 BC - appear in Egyptian hieroglyphs

# Invention of rationals

- 1000 BC - appear in Egyptian hieroglyphs
- Greek and Indian mathematicians used them regularly in studying number theory

# Invention of rationals

- 1000 BC - appear in Egyptian hieroglyphs
- Greek and Indian mathematicians used them regularly in studying number theory
- 300 BC - appear in Euclid's elements



# Discovery of irrationals

# Discovery of irrationals

- 800 BC - Indian mathematicians discuss nonrationality of square roots

# Discovery of irrationals

- 800 BC - Indian mathematicians discuss nonrationality of square roots
- 500 BC - Greek mathematicians discover the existence of irrational numbers (sides of the pentagram)

# Discovery of irrationals

- 800 BC - Indian mathematicians discuss nonrationality of square roots
- 500 BC - Greek mathematicians discover the existence of irrational numbers (sides of the pentagram)
- discovered by Hippasus, who legend says was thrown into the sea for his discovery



# Discovery of irrationals

- 800 BC - Indian mathematicians discuss nonrationality of square roots
- 500 BC - Greek mathematicians discover the existence of irrational numbers (sides of the pentagram)
- discovered by Hippasus, who legend says was thrown into the sea for his discovery
- 14th-16th century - Indian mathematicians discover infinite series expressions for  $\pi$

# Discovery of irrationals

- 800 BC - Indian mathematicians discuss nonrationality of square roots
- 500 BC - Greek mathematicians discover the existence of irrational numbers (sides of the pentagram)
- discovered by Hippasus, who legend says was thrown into the sea for his discovery
- 14th-16th century - Indian mathematicians discover infinite series expressions for  $\pi$
- 17th century - European mathematicians distinguish between transcendentals and algebraic numbers

# Outline

- 1 Real Analysis Lecture 1
  - Origin of Real Numbers
  - Properties of real numbers
  - Integers

# Philosophy of our presentation

# Philosophy of our presentation

- we will not construct the reals (ref. Foundations of Analysis by Landau for a rigorous construction from the ground up using minimal axioms and Dedekind cuts)

# Philosophy of our presentation

- we will not construct the reals (ref. Foundations of Analysis by Landau for a rigorous construction from the ground up using minimal axioms and Dedekind cuts)
- instead, we will take for granted that the reals exist and describe ten fundamental rules (axioms) for how it behaves

# Philosophy of our presentation

- we will not construct the reals (ref. Foundations of Analysis by Landau for a rigorous construction from the ground up using minimal axioms and Dedekind cuts)
- instead, we will take for granted that the reals exist and describe ten fundamental rules (axioms) for how it behaves
- the integers, rationals, etc. will be defined in terms of the reals

# Field axioms

A **field** is a set  $F$  with a way to do addition  $+$  and multiplication  $\times$  with



# Field axioms

A **field** is a set  $F$  with a way to do addition  $+$  and multiplication  $\times$  with

**A1 commutativity:**  $x + y = y + x$  and  $xy = yx$

# Field axioms

A **field** is a set  $F$  with a way to do addition  $+$  and multiplication  $\times$  with

**A1 commutativity:**  $x + y = y + x$  and  $xy = yx$

**A2 associativity:**  $(x + y) + z = x + (y + z)$  and  $(xy)z = x(yz)$

# Field axioms

A **field** is a set  $F$  with a way to do addition  $+$  and multiplication  $\times$  with

- A1 **commutativity:**  $x + y = y + x$  and  $xy = yx$
- A2 **associativity:**  $(x + y) + z = x + (y + z)$  and  $(xy)z = x(yz)$
- A3 **distributivity:**  $x(y + z) = xy + xz$

# Field axioms

A **field** is a set  $F$  with a way to do addition  $+$  and multiplication  $\times$  with

- Ⓐ1 **commutativity**:  $x + y = y + x$  and  $xy = yx$
- Ⓐ2 **associativity**:  $(x + y) + z = x + (y + z)$  and  $(xy)z = x(yz)$
- Ⓐ3 **distributivity**:  $x(y + z) = xy + xz$
- Ⓐ4 **additive inverse**: given  $x, y \in F$ , there exists a unique  $z \in F$  with  $x = y + z$

Notation:  $z := x - y$

# Field axioms

A **field** is a set  $F$  with a way to do addition  $+$  and multiplication  $\times$  with

- A1 **commutativity**:  $x + y = y + x$  and  $xy = yx$
- A2 **associativity**:  $(x + y) + z = x + (y + z)$  and  $(xy)z = x(yz)$
- A3 **distributivity**:  $x(y + z) = xy + xz$
- A4 **additive inverse**: given  $x, y \in F$ , there exists a unique  $z \in F$  with  $x = y + z$

Notation:  $z := x - y$

- A5 **multiplicative inverse**: given  $x, y \in F$  with  $y \neq 0$ , there exists a unique  $z \in F$  with  $x = yz$

Notation:  $z := x/y$

# Field axioms

A **field** is a set  $F$  with a way to do addition  $+$  and multiplication  $\times$  with

- A1 **commutativity:**  $x + y = y + x$  and  $xy = yx$
- A2 **associativity:**  $(x + y) + z = x + (y + z)$  and  $(xy)z = x(yz)$
- A3 **distributivity:**  $x(y + z) = xy + xz$
- A4 **additive inverse:** given  $x, y \in F$ , there exists a unique  $z \in F$  with  $x = y + z$

Notation:  $z := x - y$

- A5 **multiplicative inverse:** given  $x, y \in F$  with  $y \neq 0$ , there exists a unique  $z \in F$  with  $x = yz$

Notation:  $z := x/y$

**Special notation:**  $0 := x - x$  and  $1 := x/x$ , and  $-x := 0 - x$

# Challenge!

## Problem

use the axioms to show that  $(x + y)z = xz + yz$ .

# Challenge!

## Problem

use the axioms to show that  $(x + y)z = xz + yz$ .

## Solution

$$(x + y)z \stackrel{A1}{=} z(x + y) \stackrel{A3}{=} zx + zy \stackrel{A1}{=} xz + yz.$$



# Challenge!

## Problem

use the axioms to show that the value of  $x - x$  doesn't depend on  $x$ .

# Challenge!

## Problem

use the axioms to show that the value of  $x - x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x - x$  and  $z_2 = y - y$ .

# Challenge!

## Problem

use the axioms to show that the value of  $x - x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x - x$  and  $z_2 = y - y$ .  
Then  $x = x + z_1$  and  $y = y + z_2$ .

# Challenge!

## Problem

use the axioms to show that the value of  $x - x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x - x$  and  $z_2 = y - y$ .

Then  $x = x + z_1$  and  $y = y + z_2$ .

Therefore we have

$$(x + y) + z_1 \stackrel{A2}{=} x + (y + z_1) \stackrel{A1}{=} x + (z_1 + y) \stackrel{A2}{=} (x + z_1) + y = x + y,$$

# Challenge!

## Problem

use the axioms to show that the value of  $x - x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x - x$  and  $z_2 = y - y$ .

Then  $x = x + z_1$  and  $y = y + z_2$ .

Therefore we have

$$(x + y) + z_1 \stackrel{A2}{=} x + (y + z_1) \stackrel{A1}{=} x + (z_1 + y) \stackrel{A2}{=} (x + z_1) + y = x + y,$$

$$(x + y) + z_2 \stackrel{A2}{=} x + (y + z_2) = x + y,$$

# Challenge!

## Problem

use the axioms to show that the value of  $x - x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x - x$  and  $z_2 = y - y$ .

Then  $x = x + z_1$  and  $y = y + z_2$ .

Therefore we have

$$(x + y) + z_1 \stackrel{A2}{=} x + (y + z_1) \stackrel{A1}{=} x + (z_1 + y) \stackrel{A2}{=} (x + z_1) + y = x + y,$$

$$(x + y) + z_2 \stackrel{A2}{=} x + (y + z_2) = x + y,$$

so that

$$z_1 = (x + y) - (x + y) = z_2.$$



# Challenge!

## Problem

use the axioms to show that the value of  $x/x$  doesn't depend on  $x$ .

# Challenge!

## Problem

use the axioms to show that the value of  $x/x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x/x$  and  $z_2 = y/y$ .



# Challenge!

## Problem

use the axioms to show that the value of  $x/x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x/x$  and  $z_2 = y/y$ .  
Then  $x = z_1 x$  and  $y = z_2 y$ .

# Challenge!

## Problem

use the axioms to show that the value of  $x/x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x/x$  and  $z_2 = y/y$ .

Then  $x = z_1 x$  and  $y = z_2 y$ .

Therefore we have

$$z_1(xy) \stackrel{A_2}{=} (z_1 x)y = xy,$$

# Challenge!

## Problem

use the axioms to show that the value of  $x/x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x/x$  and  $z_2 = y/y$ .

Then  $x = z_1 x$  and  $y = z_2 y$ .

Therefore we have

$$z_1(xy) \stackrel{A2}{=} (z_1 x)y = xy,$$

$$z_2(xy) \stackrel{A2}{=} (z_2 x)y \stackrel{A1}{=} (xz_2)y \stackrel{A2}{=} x(z_2 y) = xy,$$

# Challenge!

## Problem

use the axioms to show that the value of  $x/x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x/x$  and  $z_2 = y/y$ .

Then  $x = z_1 x$  and  $y = z_2 y$ .

Therefore we have

$$z_1(xy) \stackrel{A2}{=} (z_1 x)y = xy,$$

$$z_2(xy) \stackrel{A2}{=} (z_2 x)y \stackrel{A1}{=} (xz_2)y \stackrel{A2}{=} x(z_2 y) = xy,$$

so that

$$z_1 = (xy)/(xy) = z_2.$$

# Field examples

Lots of examples of fields!

# Field examples

Lots of examples of fields!

- **rational numbers**

$$\mathbb{Q} = \{a/b : a, b \text{ integers with } b \neq 0\}.$$

# Field examples

Lots of examples of fields!

- **rational numbers**

$$\mathbb{Q} = \{a/b : a, b \text{ integers with } b \neq 0\}.$$

- **real numbers**  $\mathbb{R}$  (what are they?)

# Field examples

Lots of examples of fields!

- **rational numbers**

$$\mathbb{Q} = \{a/b : a, b \text{ integers with } b \neq 0\}.$$

- **real numbers**  $\mathbb{R}$  (what are they?)

- **complex numbers**

$$\mathbb{C} = \{a + ib : a, b \text{ real numbers}\}.$$



# Field examples

Lots of examples of fields!

- **rational numbers**

$$\mathbb{Q} = \{a/b : a, b \text{ integers with } b \neq 0\}.$$

- **real numbers**  $\mathbb{R}$  (what are they?)

- **complex numbers**

$$\mathbb{C} = \{a + ib : a, b \text{ real numbers}\}.$$

- **field of Boolean numbers**

$$\mathbb{F}_2 = \{0, 1\}$$

$$0 + 0 = 0, \quad 0 + 1 = 1, \quad 1 + 0 = 1, \quad 1 + 1 = 0$$

$$0 \cdot 0 = 0, \quad 0 \cdot 1 = 0, \quad 1 \cdot 0 = 0, \quad 1 \cdot 1 = 1$$

# Challenge!

## Problem

explain why the set of integers isn't a field

# Challenge!

## Problem

explain why the set of integers isn't a field

## Solution

There aren't multiplicative inverses!

# Challenge!

## Problem

explain why the set of integers isn't a field

## Solution

There aren't multiplicative inverses! For example,  $1/2$  doesn't make sense because there isn't an integer  $z$  with  $1 = 2z$ .

# Order axioms

An **ordered field** is a field  $F$  with a relation  $<$

# Order axioms

An **ordered field** is a field  $F$  with a relation  $<$

Ⓐ **trichotomy**: for any  $x, y \in F$ , exactly one of the following holds:

$$x < y, \quad y < x, \quad \text{or} \quad x = y$$

# Order axioms

An **ordered field** is a field  $F$  with a relation  $<$

- Ⓐ6 **trichotomy**: for any  $x, y \in F$ , exactly one of the following holds:

$$x < y, \quad y < x, \quad \text{or} \quad x = y$$

- Ⓐ7 **compatible with  $+$** :  $x < y$  implies  $x + z < y + z$

# Order axioms

An **ordered field** is a field  $F$  with a relation  $<$

- A6 trichotomy:** for any  $x, y \in F$ , exactly one of the following holds:

$$x < y, \quad y < x, \quad \text{or} \quad x = y$$

- A7 compatible with  $+$ :**  $x < y$  implies  $x + z < y + z$
- A8 compatible with  $\times$ :**  $0 < x$  and  $0 < y$  implies  $0 < xy$



# Order axioms

An **ordered field** is a field  $F$  with a relation  $<$

- A6 trichotomy:** for any  $x, y \in F$ , exactly one of the following holds:

$$x < y, \quad y < x, \quad \text{or} \quad x = y$$

- A7 compatible with  $+$ :**  $x < y$  implies  $x + z < y + z$
- A8 compatible with  $\times$ :**  $0 < x$  and  $0 < y$  implies  $0 < xy$
- A9 transitivity:**  $x < y$  and  $y < z$  implies  $x < z$

# Order axioms

An **ordered field** is a field  $F$  with a relation  $<$

**A6 trichotomy:** for any  $x, y \in F$ , exactly one of the following holds:

$$x < y, \quad y < x, \quad \text{or} \quad x = y$$

**A7 compatible with  $+$ :**  $x < y$  implies  $x + z < y + z$

**A8 compatible with  $\times$ :**  $0 < x$  and  $0 < y$  implies  $0 < xy$

**A9 transitivity:**  $x < y$  and  $y < z$  implies  $x < z$

**Special notation:**

- $x > y$  means  $y < x$
- $x \leq y$  means  $x < y$  or  $x = y$
- $x \geq y$  means  $y \leq x$

# Challenge!

## Problem

explain why  $\mathbb{F}_2$  cannot be made into an ordered field

# Challenge!

## Problem

explain why  $\mathbb{F}_2$  cannot be made into an ordered field

## Solution

There is no way to put an order relation  $<$  on it!

# Challenge!

## Problem

explain why  $\mathbb{F}_2$  cannot be made into an ordered field

## Solution

There is no way to put an order relation  $<$  on it!

The trichotomy (Axiom 6) would say either  $0 < 1$  or  $1 < 0$ .

# Challenge!

## Problem

explain why  $\mathbb{F}_2$  cannot be made into an ordered field

## Solution

There is no way to put an order relation  $<$  on it!

The trichotomy (Axiom 6) would say either  $0 < 1$  or  $1 < 0$ .

If  $0 < 1$ , then by Axiom 7

$$1 = 0 + 1 < 1 + 1 = 0$$

but this means both  $0 < 1$  and  $1 < 0$ , which violates the trichotomy.

# Challenge!

## Problem

explain why  $\mathbb{F}_2$  cannot be made into an ordered field

## Solution

There is no way to put an order relation  $<$  on it!

The trichotomy (Axiom 6) would say either  $0 < 1$  or  $1 < 0$ .

If  $0 < 1$ , then by Axiom 7

$$1 = 0 + 1 < 1 + 1 = 0$$

but this means both  $0 < 1$  and  $1 < 0$ , which violates the trichotomy. Same outcome if  $1 < 0$ .

# Apostol Theorem 1.1

## Theorem

*If  $a < b + \epsilon$  for all  $\epsilon > 0$ , then  $a \leq b$ .*



# Apostol Theorem 1.1

## Theorem

*If  $a < b + \epsilon$  for all  $\epsilon > 0$ , then  $a \leq b$ .*

## Proof.

(Contradiction.) Assume the statement is false.

# Apostol Theorem 1.1

## Theorem

*If  $a < b + \epsilon$  for all  $\epsilon > 0$ , then  $a \leq b$ .*

## Proof.

(Contradiction.) Assume the statement is false.  
Then the trichotomy says  $a > b$ .

# Apostol Theorem 1.1

## Theorem

*If  $a < b + \epsilon$  for all  $\epsilon > 0$ , then  $a \leq b$ .*

## Proof.

(Contradiction.) Assume the statement is false.  
Then the trichotomy says  $a > b$ .  
This means  $a - b > 0$  by A7.

# Apostol Theorem 1.1

## Theorem

*If  $a < b + \epsilon$  for all  $\epsilon > 0$ , then  $a \leq b$ .*

## Proof.

(Contradiction.) Assume the statement is false.

Then the trichotomy says  $a > b$ .

This means  $a - b > 0$  by A7.

This means  $(a - b)/2 > 0$  by A8.

# Apostol Theorem 1.1

## Theorem

*If  $a < b + \epsilon$  for all  $\epsilon > 0$ , then  $a \leq b$ .*

## Proof.

(Contradiction.) Assume the statement is false.

Then the trichotomy says  $a > b$ .

This means  $a - b > 0$  by A7.

This means  $(a - b)/2 > 0$  by A8.

This means

$$a < b + \frac{a - b}{2} = \frac{a + b}{2} \stackrel{\text{A7\&A8}}{<} \frac{a + a}{2} = a.$$

# Apostol Theorem 1.1

## Theorem

*If  $a < b + \epsilon$  for all  $\epsilon > 0$ , then  $a \leq b$ .*

## Proof.

(Contradiction.) Assume the statement is false.

Then the trichotomy says  $a > b$ .

This means  $a - b > 0$  by A7.

This means  $(a - b)/2 > 0$  by A8.

This means

$$a < b + \frac{a - b}{2} = \frac{a + b}{2} \stackrel{\text{A7\&A8}}{<} \frac{a + a}{2} = a.$$

This says  $a < a$ , which contradicts the trichotomy. □

# Real numbers

The **real numbers** are the unique field  $\mathbb{R}$  satisfying Axiom 1-Axiom 9, plus an extra axiom called the completeness axiom.

$\mathbb{R}$  = the one and only complete, ordered field

A complete ordered field  $F$  satisfies

# Real numbers

The **real numbers** are the unique field  $\mathbb{R}$  satisfying Axiom 1-Axiom 9, plus an extra axiom called the completeness axiom.

$\mathbb{R}$  = the one and only complete, ordered field

A complete ordered field  $F$  satisfies

- A6 completeness axiom:** every bounded set of numbers  $S \subseteq F$  has a supremum  $\sup(S)$



# Real numbers

The **real numbers** are the unique field  $\mathbb{R}$  satisfying Axiom 1-Axiom 9, plus an extra axiom called the completeness axiom.

$\mathbb{R}$  = the one and only complete, ordered field

A complete ordered field  $F$  satisfies

- **A6 completeness axiom:** every bounded set of numbers  $S \subseteq F$  has a supremum  $\sup(S)$
- Understanding what this is challenging, but will be fundamental to the entire course!

# Real numbers

The **real numbers** are the unique field  $\mathbb{R}$  satisfying Axiom 1-Axiom 9, plus an extra axiom called the completeness axiom.

$\mathbb{R}$  = the one and only complete, ordered field

A complete ordered field  $F$  satisfies

- **A6 completeness axiom:** every bounded set of numbers  $S \subseteq F$  has a supremum  $\sup(S)$ 
  - Understanding what this is challenging, but will be fundamental to the entire course!
  - Intuitively, it represents the fact that the real line has no holes or gaps.

# Real intervals

Given an ordered field, we can **intervals**

# Real intervals

Given an ordered field, we can **intervals**

- **open and closed intervals**

$$(a, b) = \{x \in \mathbb{R} : a < x < b\} \quad \text{and} \quad [a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$$

# Real intervals

Given an ordered field, we can **intervals**

- **open and closed intervals**

$$(a, b) = \{x \in \mathbb{R} : a < x < b\} \quad \text{and} \quad [a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$$

- **half-open intervals**

$$[a, b) = \{x \in \mathbb{R} : a \leq x < b\} \quad \text{and} \quad (a, b] = \{x \in \mathbb{R} : a < x \leq b\}$$

# Real intervals

Given an ordered field, we can **intervals**

- **open and closed intervals**

$$(a, b) = \{x \in \mathbb{R} : a < x < b\} \quad \text{and} \quad [a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$$

- **half-open intervals**

$$[a, b) = \{x \in \mathbb{R} : a \leq x < b\} \quad \text{and} \quad (a, b] = \{x \in \mathbb{R} : a < x \leq b\}$$

- **semi-infinite open intervals**

$$(-\infty, a) = \{x \in \mathbb{R} : x < a\} \quad \text{and} \quad (a, \infty) = \{x \in \mathbb{R} : a < x\}$$

# Real intervals

Given an ordered field, we can **intervals**

- **open and closed intervals**

$$(a, b) = \{x \in \mathbb{R} : a < x < b\} \quad \text{and} \quad [a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$$

- **half-open intervals**

$$[a, b) = \{x \in \mathbb{R} : a \leq x < b\} \quad \text{and} \quad (a, b] = \{x \in \mathbb{R} : a < x \leq b\}$$

- **semi-infinite open intervals**

$$(-\infty, a) = \{x \in \mathbb{R} : x < a\} \quad \text{and} \quad (a, \infty) = \{x \in \mathbb{R} : a < x\}$$

- **semi-infinite closed intervals**

$$(-\infty, a] = \{x \in \mathbb{R} : x \leq a\} \quad \text{and} \quad [a, \infty) = \{x \in \mathbb{R} : a \leq x\}.$$

# Outline

- 1 Real Analysis Lecture 1
  - Origin of Real Numbers
  - Properties of real numbers
  - Integers



# Inductive sets

A subset  $S$  of  $\mathbb{R}$  is called a **inductive set** if

- $1 \in S$
- if  $x \in S$  then  $x + 1 \in S$

# Inductive sets

A subset  $S$  of  $\mathbb{R}$  is called a **inductive set** if

- $1 \in S$
- if  $x \in S$  then  $x + 1 \in S$

Examples:

$$\mathbb{R}, \quad \mathbb{Q}, \quad (0, \infty), \quad \mathbb{Z}, \quad \mathbb{N}, \dots$$

# Integers and rationals

How do we reverse-engineer the integers from the reals?

# Integers and rationals

How do we reverse-engineer the integers from the reals?

- the set of **positive integers** is

$$\mathbb{Z}_+ = \{x : x \text{ belongs to every inductive set}\}$$

# Integers and rationals

How do we reverse-engineer the integers from the reals?

- the set of **positive integers** is

$$\mathbb{Z}_+ = \{x : x \text{ belongs to every inductive set}\}$$

- integers are

$$\mathbb{Z} = \{x : x \text{ is zero or } \pm x \text{ is a positive integer}\}$$

# Integers and rationals

How do we reverse-engineer the integers from the reals?

- the set of **positive integers** is

$$\mathbb{Z}_+ = \{x : x \text{ belongs to every inductive set}\}$$

- integers are

$$\mathbb{Z} = \{x : x \text{ is zero or } \pm x \text{ is a positive integer}\}$$

- rationals are

$$\mathbb{Q} = \{a/b : a, b \in \mathbb{Z}, b \neq 0\}.$$

# Induction

## Principle of Induction:

If  $S$  is an inductive set, then  $\mathbb{Z}_+ \subseteq S$

# Induction

## Principle of Induction:

If  $S$  is an inductive set, then  $\mathbb{Z}_+ \subseteq S$

### Question

Waaaaait...How is this this induction?!



# Induction

## Principle of Induction:

If  $S$  is an inductive set, then  $\mathbb{Z}_+ \subseteq S$

### Question

Waaaaait...How is this this induction?!

**Example:** Let's prove  $n(n+1)$  is divisible by 2 for all positive integers  $n$ , using the Principle of Induction.

# Induction

## Principle of Induction:

If  $S$  is an inductive set, then  $\mathbb{Z}_+ \subseteq S$

### Question

Waaaaait...How is this this induction?!

**Example:** Let's prove  $n(n+1)$  is divisible by 2 for all positive integers  $n$ , using the Principle of Induction.

**Proof:**

# Induction

## Principle of Induction:

If  $S$  is an inductive set, then  $\mathbb{Z}_+ \subseteq S$

### Question

Waaaaait...How is this this induction?!

**Example:** Let's prove  $n(n+1)$  is divisible by 2 for all positive integers  $n$ , using the Principle of Induction.

**Proof:**

Let

$$S = \{n \in \mathbb{Z}_+ : 2|n(n+1)\}$$

.

# Induction

## Principle of Induction:

If  $S$  is an inductive set, then  $\mathbb{Z}_+ \subseteq S$

### Question

Waaaaait...How is this this induction?!

**Example:** Let's prove  $n(n+1)$  is divisible by 2 for all positive integers  $n$ , using the Principle of Induction.

### Proof:

Let

$$S = \{n \in \mathbb{Z}_+ : 2|n(n+1)\}$$

. We see that  $1 \in S$  because 2 divides  $1(1+1)$ .

# Induction

## Principle of Induction:

If  $S$  is an inductive set, then  $\mathbb{Z}_+ \subseteq S$

### Question

Waaaaait...How is this this induction?!

**Example:** Let's prove  $n(n+1)$  is divisible by 2 for all positive integers  $n$ , using the Principle of Induction.

**Proof:**

Let

$$S = \{n \in \mathbb{Z}_+ : 2|n(n+1)\}$$

. We see that  $1 \in S$  because 2 divides  $1(1+1)$ .

Now suppose  $x \in S$ . (This is our usual inductive assumption).



# Induction

**Example:** Let's prove  $n(n+1)$  is divisible by 2 for all positive integers  $n$ , using the Principle of Induction.

**Proof continued:**

# Induction

**Example:** Let's prove  $n(n + 1)$  is divisible by 2 for all positive integers  $n$ , using the Principle of Induction.

**Proof continued:**

Then 2 divides  $x(x + 1)$ , so  $x(x + 1) = 2k$  for some integer  $k$ .

# Induction

**Example:** Let's prove  $n(n + 1)$  is divisible by 2 for all positive integers  $n$ , using the Principle of Induction.

**Proof continued:**

Then 2 divides  $x(x + 1)$ , so  $x(x + 1) = 2k$  for some integer  $k$ .  
This means  $(x + 1)(x + 2) = x(x + 1) + 2(x + 1) = 2(k + x + 1)$ .



# Induction

**Example:** Let's prove  $n(n+1)$  is divisible by 2 for all positive integers  $n$ , using the Principle of Induction.

**Proof continued:**

Then 2 divides  $x(x+1)$ , so  $x(x+1) = 2k$  for some integer  $k$ .  
This means  $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$ .  
Thus 2 divides  $(x+1)(x+2)$ , showing that  $x+1 \in S$ .

# Induction

**Example:** Let's prove  $n(n+1)$  is divisible by 2 for all positive integers  $n$ , using the Principle of Induction.

**Proof continued:**

Then 2 divides  $x(x+1)$ , so  $x(x+1) = 2k$  for some integer  $k$ .  
This means  $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$ .  
Thus 2 divides  $(x+1)(x+2)$ , showing that  $x+1 \in S$ .  
Since  $x+1$  was arbitrary, this shows that  $S$  is an inductive set.

# Induction

**Example:** Let's prove  $n(n+1)$  is divisible by 2 for all positive integers  $n$ , using the Principle of Induction.

**Proof continued:**

Then 2 divides  $x(x+1)$ , so  $x(x+1) = 2k$  for some integer  $k$ .  
This means  $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$ .  
Thus 2 divides  $(x+1)(x+2)$ , showing that  $x+1 \in S$ .  
Since  $x+1$  was arbitrary, this shows that  $S$  is an inductive set.  
By the Principle of Induction, this means  $\mathbb{Z}_+ \subseteq S$ .

# Induction

**Example:** Let's prove  $n(n+1)$  is divisible by 2 for all positive integers  $n$ , using the Principle of Induction.

**Proof continued:**

Then 2 divides  $x(x+1)$ , so  $x(x+1) = 2k$  for some integer  $k$ .  
This means  $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$ .  
Thus 2 divides  $(x+1)(x+2)$ , showing that  $x+1 \in S$ .  
Since  $x+1$  was arbitrary, this shows that  $S$  is an inductive set.  
By the Principle of Induction, this means  $\mathbb{Z}_+ \subseteq S$ .  
In other words  $n \in S$  for every positive integer  $n$ .

# Induction

**Example:** Let's prove  $n(n+1)$  is divisible by 2 for all positive integers  $n$ , using the Principle of Induction.

**Proof continued:**

Then 2 divides  $x(x+1)$ , so  $x(x+1) = 2k$  for some integer  $k$ .  
This means  $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$ .  
Thus 2 divides  $(x+1)(x+2)$ , showing that  $x+1 \in S$ .  
Since  $x+1$  was arbitrary, this shows that  $S$  is an inductive set.  
By the Principle of Induction, this means  $\mathbb{Z}_+ \subseteq S$ .  
In other words  $n \in S$  for every positive integer  $n$ .  
Hence 2 divides  $n(n+1)$  for every positive integers  $n$ .

# Prime numbers

A positive integer  $p$  is **prime** if its only positive divisors are 1 and  $p$ .

Theorem (Apostol Theorem 1.5)

*Every integer is prime or a product of primes*

Theorem (Apostol Theorem 1.8)

*If  $p$  is prime and  $p$  divides  $ab$ , then  $p$  divides  $a$  or  $p$  divides  $b$ .*

Theorem (Fundamental Theorem of Arithmetic (Apostol Theorem 1.9))

*Every integer  $n > 1$  has a unique factorization as a product of primes, up to reordering.*