

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

October 14, 2024

Outline

- 1 Real Analysis Lecture 13
 - Metric subspaces
 - Limits of sequences

Outline

- 1 Real Analysis Lecture 13
 - Metric subspaces
 - Limits of sequences

Metric subspace

Definition

Let (M, d) be a metric space.

Metric subspace

Definition

Let (M, d) be a metric space.

If $S \subseteq M$ is a subset, then $d : M \times M \rightarrow \mathbb{R}$ restricts to a metric $d : S \times S \rightarrow \mathbb{R}$, called the **relative metric**.

Metric subspace

Definition

Let (M, d) be a metric space.

If $S \subseteq M$ is a subset, then $d : M \times M \rightarrow \mathbb{R}$ restricts to a metric $d : S \times S \rightarrow \mathbb{R}$, called the **relative metric**.

The space (S, d) with the relative metric is called a **metric subspace**.

Metric subspace

Definition

Let (M, d) be a metric space.

If $S \subseteq M$ is a subset, then $d : M \times M \rightarrow \mathbb{R}$ restricts to a metric $d : S \times S \rightarrow \mathbb{R}$, called the **relative metric**.

The space (S, d) with the relative metric is called a **metric subspace**.

Open balls in S :

$$B_S(x; r) = B_M(x; r) \cap S.$$

Example

Let $M = \mathbb{R}$ and d be the Euclidean metric.

Example

Let $M = \mathbb{R}$ and d be the Euclidean metric.

Consider the metric subspace (S, d) for $S = [-1, 1) \cup \{2\}$.

Example

Let $M = \mathbb{R}$ and d be the Euclidean metric.

Consider the metric subspace (S, d) for $S = [-1, 1) \cup \{2\}$.

Open balls:

Example

Let $M = \mathbb{R}$ and d be the Euclidean metric.

Consider the metric subspace (S, d) for $S = [-1, 1) \cup \{2\}$.

Open balls:

$$\{2\}$$

Example

Let $M = \mathbb{R}$ and d be the Euclidean metric.

Consider the metric subspace (S, d) for $S = [-1, 1) \cup \{2\}$.

Open balls:

$$\{2\}$$

$$[-1, s), \quad -1 < s < 1$$

Example

Let $M = \mathbb{R}$ and d be the Euclidean metric.

Consider the metric subspace (S, d) for $S = [-1, 1) \cup \{2\}$.

Open balls:

$$\{2\}$$

$$[-1, s), \quad -1 < s < 1$$

$$(r, s), \quad -1 < r < s < 1$$

Example

Let $M = \mathbb{R}$ and d be the Euclidean metric.

Consider the metric subspace (S, d) for $S = [-1, 1) \cup \{2\}$.

Open balls:

$$\{2\}$$

$$[-1, s), \quad -1 < s < 1$$

$$(r, s), \quad -1 < r < s < 1$$

$$[-1, s) \cup \{2\}, \quad -1 < s < 1$$

Example

Let $M = \mathbb{R}$ and d be the Euclidean metric.

Consider the metric subspace (S, d) for $S = [-1, 1) \cup \{2\}$.

Open balls:

$$\{2\}$$

$$[-1, s), \quad -1 < s < 1$$

$$(r, s), \quad -1 < r < s < 1$$

$$[-1, s) \cup \{2\}, \quad -1 < s < 1$$

$$(r, s) \cup \{2\}, \quad -1 < r < s < 1$$

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\implies : Assume V is an open subset of S .

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\implies : Assume V is an open subset of S .

Then for all $x \in V$, there exists $r_x > 0$ such that $B_S(x; r_x) \subseteq V$.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\implies : Assume V is an open subset of S .

Then for all $x \in V$, there exists $r_x > 0$ such that $B_S(x; r_x) \subseteq V$.

This says $B_M(x; r_x) \cap S \subseteq V$.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\implies : Assume V is an open subset of S .

Then for all $x \in V$, there exists $r_x > 0$ such that $B_S(x; r_x) \subseteq V$.

This says $B_M(x; r_x) \cap S \subseteq V$.

Define

$$U = \bigcup_{x \in V} B_M(x; r_x).$$

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\implies : Assume V is an open subset of S .

Then for all $x \in V$, there exists $r_x > 0$ such that $B_S(x; r_x) \subseteq V$.

This says $B_M(x; r_x) \cap S \subseteq V$.

Define

$$U = \bigcup_{x \in V} B_M(x; r_x).$$

This is a union of open sets, and is therefore open.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\implies : Assume V is an open subset of S .

Then for all $x \in V$, there exists $r_x > 0$ such that $B_S(x; r_x) \subseteq V$.

This says $B_M(x; r_x) \cap S \subseteq V$.

Define

$$U = \bigcup_{x \in V} B_M(x; r_x).$$

This is a union of open sets, and is therefore open.

Furthermore $U \cap S = V$.



Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\Leftarrow : Assume U is an open subset of M .

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\Leftarrow : Assume U is an open subset of M .
We need to show that $V := U \cap S$ is open.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\Leftarrow : Assume U is an open subset of M .

We need to show that $V := U \cap S$ is open.

If $x \in V$, then $x \in U$ and there exists $r > 0$ such that $B_M(x; r) \subseteq U$.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\Leftarrow : Assume U is an open subset of M .

We need to show that $V := U \cap S$ is open.

If $x \in V$, then $x \in U$ and there exists $r > 0$ such that

$$B_M(x; r) \subseteq U.$$

It follows that $B_S(x, r) = B_M(x, r) \cap S \subseteq U \cap S = V$.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\Leftarrow : Assume U is an open subset of M .

We need to show that $V := U \cap S$ is open.

If $x \in V$, then $x \in U$ and there exists $r > 0$ such that $B_M(x; r) \subseteq U$.

It follows that $B_S(x, r) = B_M(x, r) \cap S \subseteq U \cap S = V$.

This shows that x is an interior point of S in the metric subspace (S, d) .

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\Leftarrow : Assume U is an open subset of M .

We need to show that $V := U \cap S$ is open.

If $x \in V$, then $x \in U$ and there exists $r > 0$ such that $B_M(x; r) \subseteq U$.

It follows that $B_S(x, r) = B_M(x, r) \cap S \subseteq U \cap S = V$.

This shows that x is an interior point of S in the metric subspace (S, d) .

Since $x \in V$ was arbitrary, this shows that V is open in S . □

Closed sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then B is a closed subset of S if and only if there exists a closed subset A of M with $B = A \cap S$.

Closed sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then B is a closed subset of S if and only if there exists a closed subset A of M with $B = A \cap S$.

Proof.

Closed sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then B is a closed subset of S if and only if there exists a closed subset A of M with $B = A \cap S$.

Proof.

Let $B \subseteq S$.

Closed sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then B is a closed subset of S if and only if there exists a closed subset A of M with $B = A \cap S$.

Proof.

Let $B \subseteq S$.

B is closed in S if and only if $S \setminus B$ is open in B

Closed sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then B is a closed subset of S if and only if there exists a closed subset A of M with $B = A \cap S$.

Proof.

Let $B \subseteq S$.

B is closed in S if and only if $S \setminus B$ is open in B

This is true if and only if there exists an open subset U of M with $S \setminus B = U \cap S$.

Closed sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then B is a closed subset of S if and only if there exists a closed subset A of M with $B = A \cap S$.

Proof.

Let $B \subseteq S$.

B is closed in S if and only if $S \setminus B$ is open in B

This is true if and only if there exists an open subset U of M with $S \setminus B = U \cap S$.

This is true if and only if there exists an open subset U of M with $B = (M \setminus U) \cap S$.

Closed sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then B is a closed subset of S if and only if there exists a closed subset A of M with $B = A \cap S$.

Proof.

Let $B \subseteq S$.

B is closed in S if and only if $S \setminus B$ is open in B

This is true if and only if there exists an open subset U of M with $S \setminus B = U \cap S$.

This is true if and only if there exists an open subset U of M with $B = (M \setminus U) \cap S$.

this is true if and only if there exists a closed subset A of M with $B = A \cap S$.



Example

Consider the metric space (\mathbb{R}, d) with $d = d_{\text{eucl}}$ the Euclidean metric, and the metric subspace (S, d) for $S = [0, 1]$.

Example

Consider the metric space (\mathbb{R}, d) with $d = d_{\text{eucl}}$ the Euclidean metric, and the metric subspace (S, d) for $S = [0, 1]$.

- $[0, 1]$ is open in S and closed in S , but it is not open in M

Example

Consider the metric space (\mathbb{R}, d) with $d = d_{\text{eucl}}$ the Euclidean metric, and the metric subspace (S, d) for $S = [0, 1]$.

- $[0, 1]$ is open in S and closed in S , but it is not open in M
- $[0, 1/2)$ is open in S , but not in M

Example

Consider the metric space (\mathbb{R}, d) with $d = d_{\text{eucl}}$ the Euclidean metric, and the metric subspace (S, d) for $S = [0, 1]$.

- $[0, 1]$ is open in S and closed in S , but it is not open in M
- $[0, 1/2)$ is open in S , but not in M
- $(0, 1/2)$ is open in both S and in M

Example

Consider the metric space (\mathbb{R}, d) with $d = d_{\text{eucl}}$ the Euclidean metric, and the metric subspace (S, d) for $S = [0, 1]$.

- $[0, 1]$ is open in S and closed in S , but it is not open in M
- $[0, 1/2)$ is open in S , but not in M
- $(0, 1/2)$ is open in both S and in M
- $(1/2, 1]$ is open in S , but not in M

Outline

- 1 Real Analysis Lecture 13
 - Metric subspaces
 - Limits of sequences

Definition of a limit

Definition of a limit

Back in Calc. I:

Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as n gets arbitrarily large, the value of x_n gets arbitrarily close to L .

Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as n gets arbitrarily large, the value of x_n gets arbitrarily close to L .

Formally:

Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as n gets arbitrarily large, the value of x_n gets arbitrarily close to L .

Formally:

$\forall \epsilon > 0$, there exists $N \in \mathbb{Z}_+$ such that $|x_n - L| < \epsilon \forall n \geq N$.

Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as n gets arbitrarily large, the value of x_n gets arbitrarily close to L .

Formally:

$\forall \epsilon > 0$, there exists $N \in \mathbb{Z}_+$ such that $|x_n - L| < \epsilon \forall n \geq N$.

Metric space version:

Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as n gets arbitrarily large, the value of x_n gets arbitrarily close to L .

Formally:

$\forall \epsilon > 0$, there exists $N \in \mathbb{Z}_+$ such that $|x_n - L| < \epsilon \forall n \geq N$.

Metric space version:

$\forall \epsilon > 0$, there exists $N \in \mathbb{Z}_+$ such that $d(x_n, L) < \epsilon \forall n \geq N$.

Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as n gets arbitrarily large, the value of x_n gets arbitrarily close to L .

Formally:

$\forall \epsilon > 0$, there exists $N \in \mathbb{Z}_+$ such that $|x_n - L| < \epsilon \forall n \geq N$.

Metric space version:

$\forall \epsilon > 0$, there exists $N \in \mathbb{Z}_+$ such that $d(x_n, L) < \epsilon \forall n \geq N$.

We say $\{x_n\}$ **converges** to L and L is the **limit of the sequence**.

Example

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Proof.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Proof.

Let $\epsilon > 0$.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Then $n > N$ implies

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Then $n > N$ implies

$$d(x_n, 0) = |1/n - 0| = 1/n < 1/N < \epsilon.$$

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Then $n > N$ implies

$$d(x_n, 0) = |1/n - 0| = 1/n < 1/N < \epsilon.$$

Since $\epsilon > 0$ was arbitrary, this proves $\{x_n\}$ converges to 0.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Then $n > N$ implies

$$d(x_n, 0) = |1/n - 0| = 1/n < 1/N < \epsilon.$$

Since $\epsilon > 0$ was arbitrary, this proves $\{x_n\}$ converges to 0. \square

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = (n - 1)/n$ converges to 1.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = (n - 1)/n$ converges to 1.

Proof.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = (n - 1)/n$ converges to 1.

Proof.

Let $\epsilon > 0$.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = (n - 1)/n$ converges to 1.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = (n - 1)/n$ converges to 1.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Then $n > N$ implies

$$d(x_n, 1) = |(n - 1)/n - 1| = 1/n < 1/N < \epsilon.$$

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = (n - 1)/n$ converges to 1.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Then $n > N$ implies

$$d(x_n, 1) = |(n - 1)/n - 1| = 1/n < 1/N < \epsilon.$$

Since $\epsilon > 0$ was arbitrary, this proves $\{x_n\}$ converges to 1. \square

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ does not converge.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ does not converge.

Proof.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ does not converge.

Proof.

Suppose that it converges to a number L .

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ does not converge.

Proof.

Suppose that it converges to a number L .

Then for all $\epsilon > 0$, there must exist an $N \in \mathbb{Z}_+$ such that $d(x_n, L) < \epsilon$ for all $N \geq n$.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ does not converge.

Proof.

Suppose that it converges to a number L .

Then for all $\epsilon > 0$, there must exist an $N \in \mathbb{Z}_+$ such that $d(x_n, L) < \epsilon$ for all $N \geq n$.

Take $\epsilon = 1$ and choose $N \in \mathbb{Z}_+$ such that $d(x_n, L) < 1$ for all $N \geq n$.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ does not converge.

Proof.

Suppose that it converges to a number L .

Then for all $\epsilon > 0$, there must exist an $N \in \mathbb{Z}_+$ such that $d(x_n, L) < \epsilon$ for all $N \geq n$.

Take $\epsilon = 1$ and choose $N \in \mathbb{Z}_+$ such that $d(x_n, L) < 1$ for all $N \geq n$.

With the discrete metric $d(x_n, L) < 1$ implies $x_n = L$.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ does not converge.

Proof.

Suppose that it converges to a number L .

Then for all $\epsilon > 0$, there must exist an $N \in \mathbb{Z}_+$ such that $d(x_n, L) < \epsilon$ for all $N \geq n$.

Take $\epsilon = 1$ and choose $N \in \mathbb{Z}_+$ such that $d(x_n, L) < 1$ for all $N \geq n$.

With the discrete metric $d(x_n, L) < 1$ implies $x_n = L$.

This means $x_n = 1/n = L$ for all $n \geq N$, which is a contradiction! □

Uniqueness of limits

Theorem (Uniqueness of limits)

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to M , then $L = M$.

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to M , then $L = M$.

Proof.

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to M , then $L = M$.

Proof.

Suppose that $\{x_n\}$ converges to L and M .

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to M , then $L = M$.

Proof.

Suppose that $\{x_n\}$ converges to L and M .

Assume further that $L \neq M$ and let $\epsilon = d(L, M)/2$.

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to M , then $L = M$.

Proof.

Suppose that $\{x_n\}$ converges to L and M .

Assume further that $L \neq M$ and let $\epsilon = d(L, M)/2$.

Then there exists N_1 such that $d(x_n, L) < \epsilon$ for $n > N_1$.

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to M , then $L = M$.

Proof.

Suppose that $\{x_n\}$ converges to L and M .

Assume further that $L \neq M$ and let $\epsilon = d(L, M)/2$.

Then there exists N_1 such that $d(x_n, L) < \epsilon$ for $n > N_1$.

Likewise, there exists N_2 such that $d(x_n, M) < \epsilon$ for $n > N_2$.

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to M , then $L = M$.

Proof.

Suppose that $\{x_n\}$ converges to L and M .

Assume further that $L \neq M$ and let $\epsilon = d(L, M)/2$.

Then there exists N_1 such that $d(x_n, L) < \epsilon$ for $n > N_1$.

Likewise, there exists N_2 such that $d(x_n, M) < \epsilon$ for $n > N_2$.

Then for $n > \max\{N_1, N_2\}$, the triangle inequality says

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to M , then $L = M$.

Proof.

Suppose that $\{x_n\}$ converges to L and M .

Assume further that $L \neq M$ and let $\epsilon = d(L, M)/2$.

Then there exists N_1 such that $d(x_n, L) < \epsilon$ for $n > N_1$.

Likewise, there exists N_2 such that $d(x_n, M) < \epsilon$ for $n > N_2$.

Then for $n > \max\{N_1, N_2\}$, the triangle inequality says

$$d(L, M) \leq d(L, x_n) + d(x_n, M) < 2\epsilon = d(L, M).$$

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to M , then $L = M$.

Proof.

Suppose that $\{x_n\}$ converges to L and M .

Assume further that $L \neq M$ and let $\epsilon = d(L, M)/2$.

Then there exists N_1 such that $d(x_n, L) < \epsilon$ for $n > N_1$.

Likewise, there exists N_2 such that $d(x_n, M) < \epsilon$ for $n > N_2$.

Then for $n > \max\{N_1, N_2\}$, the triangle inequality says

$$d(L, M) \leq d(L, x_n) + d(x_n, M) < 2\epsilon = d(L, M).$$

This is a contradiction.



Existence of limits

Theorem

If a sequence $\{x_n\}$ in a metric space (M, d) converges to a value $L \in M$, then the range

$$X = \{x_1, x_2, \dots\} \subseteq M$$

is a bounded set and L is an adherent point of X .

Existence of limits

Theorem

If a sequence $\{x_n\}$ in a metric space (M, d) converges to a value $L \in M$, then the range

$$X = \{x_1, x_2, \dots\} \subseteq M$$

is a bounded set and L is an adherent point of X .

When the range of a sequence is bounded, we call the sequence $\{x_n\}$ bounded.

Challenge!

Try to prove the previous theorem.

Existence of limits

Sort of a converse of the previous theorem.

Existence of limits

Sort of a converse of the previous theorem.

Theorem

If S is a subset of a metric space (M, d) and $L \in M$ is an adherent point of S , then there exists a sequence $\{x_n\}$ of elements of S which converges to L .

Existence of limits

Recall a **subsequence** of a sequence $\{x_n\}$ is a sequence of the form $\{x_{k(n)}\}$, for $k(1) < k(2) < \dots$ positive integers.

Existence of limits

Recall a **subsequence** of a sequence $\{x_n\}$ is a sequence of the form $\{x_{k(n)}\}$, for $k(1) < k(2) < \dots$ positive integers.

Theorem

A sequence $\{x_n\}$ in a metric space (M, d) converges to a value $L \in M$ if and only if every subsequence $\{x_{k(n)}\}$ also converges to L .

Challenge!

Try to prove the previous theorem.

Monotone sequences

Definition

Monotone sequences

Definition

A sequence $\{x_n\}$ of real numbers is called **increasing** or **monotone increasing** if $x_n \leq x_{n+1}$ for all n .

Monotone sequences

Definition

A sequence $\{x_n\}$ of real numbers is called **increasing** or **monotone increasing** if $x_n \leq x_{n+1}$ for all n .

It is called **strictly increasing** if $x_n < x_{n+1}$ for all n .

Monotone sequences

Definition

A sequence $\{x_n\}$ of real numbers is called **increasing** or **monotone increasing** if $x_n \leq x_{n+1}$ for all n .

It is called **strictly increasing** if $x_n < x_{n+1}$ for all n .

A sequence $\{x_n\}$ of real numbers is called **decreasing** or **monotone decreasing** if $x_n \geq x_{n+1}$ for all n .

Monotone sequences

Definition

A sequence $\{x_n\}$ of real numbers is called **increasing** or **monotone increasing** if $x_n \leq x_{n+1}$ for all n .

It is called **strictly increasing** if $x_n < x_{n+1}$ for all n .

A sequence $\{x_n\}$ of real numbers is called **decreasing** or **monotone decreasing** if $x_n \geq x_{n+1}$ for all n .

It is called **strictly decreasing** if $x_n > x_{n+1}$ for all n .

Monotone sequences

Definition

A sequence $\{x_n\}$ of real numbers is called **increasing** or **monotone increasing** if $x_n \leq x_{n+1}$ for all n .

It is called **strictly increasing** if $x_n < x_{n+1}$ for all n .

A sequence $\{x_n\}$ of real numbers is called **decreasing** or **monotone decreasing** if $x_n \geq x_{n+1}$ for all n .

It is called **strictly decreasing** if $x_n > x_{n+1}$ for all n .

In any of these cases, we call the sequence **monotone**.

Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

Theorem (Monotone convergence theorem)

Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

Theorem (Monotone convergence theorem)

Consider the metric space formed by \mathbb{R} with Euclidean distance.

Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

Theorem (Monotone convergence theorem)

Consider the metric space formed by \mathbb{R} with Euclidean distance.

A monotone increasing sequence $\{x_n\}$ of real numbers which is bounded above converges to $L = \sup\{x_1, x_2, \dots\}$.

Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

Theorem (Monotone convergence theorem)

Consider the metric space formed by \mathbb{R} with Euclidean distance.

A monotone increasing sequence $\{x_n\}$ of real numbers which is bounded above converges to $L = \sup\{x_1, x_2, \dots\}$.

A monotone decreasing sequence $\{x_n\}$ which is bounded below converges to $L = \inf\{x_1, x_2, \dots\}$.

Monotone sequences

Proof.

Monotone sequences

Proof.

Suppose that $\{x_n\}$ is a monotone increasing sequence which is bounded above and let $X = \{x_1, x_2, \dots\}$.

Monotone sequences

Proof.

Suppose that $\{x_n\}$ is a monotone increasing sequence which is bounded above and let $X = \{x_1, x_2, \dots\}$. Then by the Completeness Axiom $L = \sup X$ exists.

Monotone sequences

Proof.

Suppose that $\{x_n\}$ is a monotone increasing sequence which is bounded above and let $X = \{x_1, x_2, \dots\}$. Then by the Completeness Axiom $L = \sup X$ exists.

Let $\epsilon > 0$.

Monotone sequences

Proof.

Suppose that $\{x_n\}$ is a monotone increasing sequence which is bounded above and let $X = \{x_1, x_2, \dots\}$. Then by the Completeness Axiom $L = \sup X$ exists.

Let $\epsilon > 0$.

Then $L - \epsilon$ is not a supremum of X , so there exists $x_N \in X$ such that $x_N > L - \epsilon$.

Monotone sequences

Proof.

Suppose that $\{x_n\}$ is a monotone increasing sequence which is bounded above and let $X = \{x_1, x_2, \dots\}$. Then by the Completeness Axiom $L = \sup X$ exists.

Let $\epsilon > 0$.

Then $L - \epsilon$ is not a supremum of X , so there exists $x_N \in X$ such that $x_N > L - \epsilon$.

Since the sequence is increasing, for any $n > N$, we have that $L - \epsilon < x_N \leq x_n < L$.

Monotone sequences

Proof.

Suppose that $\{x_n\}$ is a monotone increasing sequence which is bounded above and let $X = \{x_1, x_2, \dots\}$. Then by the Completeness Axiom $L = \sup X$ exists.

Let $\epsilon > 0$.

Then $L - \epsilon$ is not a supremum of X , so there exists $x_N \in X$ such that $x_N > L - \epsilon$.

Since the sequence is increasing, for any $n > N$, we have that $L - \epsilon < x_N \leq x_n < L$.

It follows that

$$d(x_n, L) = |x_n - L| = L - x_n < \epsilon.$$

Monotone sequences

Proof.

Suppose that $\{x_n\}$ is a monotone increasing sequence which is bounded above and let $X = \{x_1, x_2, \dots\}$. Then by the Completeness Axiom $L = \sup X$ exists.

Let $\epsilon > 0$.

Then $L - \epsilon$ is not a supremum of X , so there exists $x_N \in X$ such that $x_N > L - \epsilon$.

Since the sequence is increasing, for any $n > N$, we have that $L - \epsilon < x_N \leq x_n < L$.

It follows that

$$d(x_n, L) = |x_n - L| = L - x_n < \epsilon.$$

Since $\epsilon > 0$ was arbitrary, this proves x_n converges to L . □