

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

August 28, 2024

Outline

- 1 Real Analysis Lecture 2
 - Types of real numbers
 - Integers
 - Upper Bound and Supremum

Outline

- 1 Real Analysis Lecture 2
 - Types of real numbers
 - Integers
 - Upper Bound and Supremum

Real numbers

The **real numbers** are the unique field \mathbb{R} satisfying Axiom 1-Axiom 9, plus an extra axiom called the completeness axiom.

\mathbb{R} = the one and only complete, ordered field

A complete ordered field F satisfies

Real numbers

The **real numbers** are the unique field \mathbb{R} satisfying Axiom 1-Axiom 9, plus an extra axiom called the completeness axiom.

\mathbb{R} = the one and only complete, ordered field

A complete ordered field F satisfies

A10 completeness axiom: every set of numbers $S \subseteq F$ which is bounded above has a supremum $\sup(S)$

Real numbers

The **real numbers** are the unique field \mathbb{R} satisfying Axiom 1-Axiom 9, plus an extra axiom called the completeness axiom.

\mathbb{R} = the one and only complete, ordered field

A complete ordered field F satisfies

- **completeness axiom:** every set of numbers $S \subseteq F$ which is bounded above has a supremum $\sup(S)$
- Understanding what this is challenging, but will be fundamental to the entire course!

Real numbers

The **real numbers** are the unique field \mathbb{R} satisfying Axiom 1-Axiom 9, plus an extra axiom called the completeness axiom.

\mathbb{R} = the one and only complete, ordered field

A complete ordered field F satisfies

- **completeness axiom:** every set of numbers $S \subseteq F$ which is bounded above has a supremum $\sup(S)$
- Understanding what this is challenging, but will be fundamental to the entire course!
- Intuitively, it represents the fact that the real line has no holes or gaps.

Real intervals

Given an ordered field, we can **intervals**

Real intervals

Given an ordered field, we can **intervals**

- **open and closed intervals**

$$(a, b) = \{x \in \mathbb{R} : a < x < b\} \quad \text{and} \quad [a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$$

Real intervals

Given an ordered field, we can **intervals**

- **open and closed intervals**

$$(a, b) = \{x \in \mathbb{R} : a < x < b\} \quad \text{and} \quad [a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$$

- **half-open intervals**

$$[a, b) = \{x \in \mathbb{R} : a \leq x < b\} \quad \text{and} \quad (a, b] = \{x \in \mathbb{R} : a < x \leq b\}$$

Real intervals

Given an ordered field, we can **intervals**

- **open and closed intervals**

$$(a, b) = \{x \in \mathbb{R} : a < x < b\} \quad \text{and} \quad [a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$$

- **half-open intervals**

$$[a, b) = \{x \in \mathbb{R} : a \leq x < b\} \quad \text{and} \quad (a, b] = \{x \in \mathbb{R} : a < x \leq b\}$$

- **semi-infinite open intervals**

$$(-\infty, a) = \{x \in \mathbb{R} : x < a\} \quad \text{and} \quad (a, \infty) = \{x \in \mathbb{R} : a < x\}$$

Real intervals

Given an ordered field, we can **intervals**

- **open and closed intervals**

$$(a, b) = \{x \in \mathbb{R} : a < x < b\} \quad \text{and} \quad [a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$$

- **half-open intervals**

$$[a, b) = \{x \in \mathbb{R} : a \leq x < b\} \quad \text{and} \quad (a, b] = \{x \in \mathbb{R} : a < x \leq b\}$$

- **semi-infinite open intervals**

$$(-\infty, a) = \{x \in \mathbb{R} : x < a\} \quad \text{and} \quad (a, \infty) = \{x \in \mathbb{R} : a < x\}$$

- **semi-infinite closed intervals**

$$(-\infty, a] = \{x \in \mathbb{R} : x \leq a\} \quad \text{and} \quad [a, \infty) = \{x \in \mathbb{R} : a \leq x\}.$$

Outline

- 1 Real Analysis Lecture 2
 - Types of real numbers
 - **Integers**
 - Upper Bound and Supremum

Inductive sets

A subset S of \mathbb{R} is called a **inductive set** if

- $1 \in S$
- if $x \in S$ then $x + 1 \in S$

Inductive sets

A subset S of \mathbb{R} is called a **inductive set** if

- $1 \in S$
- if $x \in S$ then $x + 1 \in S$

Examples:

$$\mathbb{R}, \quad \mathbb{Q}, \quad (0, \infty), \quad \mathbb{Z}, \quad \mathbb{N}, \dots$$

Integers and rationals

How do we reverse-engineer the integers from the reals?

Integers and rationals

How do we reverse-engineer the integers from the reals?

- the set of **positive integers** is

$$\mathbb{Z}_+ = \{x : x \text{ belongs to every inductive set}\}$$

Integers and rationals

How do we reverse-engineer the integers from the reals?

- the set of **positive integers** is

$$\mathbb{Z}_+ = \{x : x \text{ belongs to every inductive set}\}$$

- integers are

$$\mathbb{Z} = \{x : x \text{ is zero or } \pm x \text{ is a positive integer}\}$$

Integers and rationals

How do we reverse-engineer the integers from the reals?

- the set of **positive integers** is

$$\mathbb{Z}_+ = \{x : x \text{ belongs to every inductive set}\}$$

- integers are

$$\mathbb{Z} = \{x : x \text{ is zero or } \pm x \text{ is a positive integer}\}$$

- rationals are

$$\mathbb{Q} = \{a/b : a, b \in \mathbb{Z}, b \neq 0\}.$$

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof:

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof:

Let

$$S = \{n \in \mathbb{Z}_+ : 2|n(n+1)\}$$

.

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof:

Let

$$S = \{n \in \mathbb{Z}_+ : 2|n(n+1)\}$$

. We see that $1 \in S$ because 2 divides $1(1+1)$.

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof:

Let

$$S = \{n \in \mathbb{Z}_+ : 2|n(n+1)\}$$

. We see that $1 \in S$ because 2 divides $1(1+1)$.

Now suppose $x \in S$. (This is our usual inductive assumption).

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Induction

Example: Let's prove $n(n + 1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x + 1)$, so $x(x + 1) = 2k$ for some integer k .

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .

This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.
Since $x+1$ was arbitrary, this shows that S is an inductive set.

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.
Since $x+1$ was arbitrary, this shows that S is an inductive set.
By the Principle of Induction, this means $\mathbb{Z}_+ \subseteq S$.

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.
Since $x+1$ was arbitrary, this shows that S is an inductive set.
By the Principle of Induction, this means $\mathbb{Z}_+ \subseteq S$.
In other words $n \in S$ for every positive integer n .

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.
Since $x+1$ was arbitrary, this shows that S is an inductive set.
By the Principle of Induction, this means $\mathbb{Z}_+ \subseteq S$.
In other words $n \in S$ for every positive integer n .
Hence 2 divides $n(n+1)$ for every positive integers n .

Prime numbers

A positive integer p is **prime** if its only positive divisors are 1 and p .

Theorem (Apostol Theorem 1.5)

Every integer is prime or a product of primes

Theorem (Apostol Theorem 1.8)


If p is prime and p divides ab , then p divides a or p divides b .

Theorem (Fundamental Theorem of Arithmetic (Apostol Theorem 1.9))

Every integer $n > 1$ has a unique factorization as a product of primes, up to reordering.

Types of numbers


Transcendental Numbers



Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Discovered by Hippasus, a pythagorean (a student of Pythagoras)



Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery

Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- dramatic reenactment, found online:

Irrational numbers


Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- dramatic reenactment, found online:

Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- dramatic reenactment, found online:



Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- anime style:

Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- anime style:



Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b .

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$. It follows that $b^2 n' = 1$, so b^2 divides 1.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$. It follows that $b^2 n' = 1$, so b^2 divides 1. This implies n' divides 1, so $n' = \pm 1$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$. It follows that $b^2 n' = 1$, so b^2 divides 1. This implies n' divides 1, so $n' = \pm 1$. Since $n > 0$, $n' > 0$ so $n' = 1$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$. It follows that $b^2 n' = 1$, so b^2 divides 1. This implies n' divides 1, so $n' = \pm 1$. Since $n > 0$, $n' > 0$ so $n' = 1$. Thus $n = a^2$, which is a contradiction.

Algebraic and Transcendental

Irrational numbers can be further divided into to categories

Algebraic and Transcendental

Irrational numbers can be further divided into to categories

- ① **algebraic numbers:** numbers which are roots of polynomials with integer coefficients, like

$$\sqrt{2}, \quad \sqrt{3}, \quad \text{and} \quad \sqrt[3]{\sqrt{5} + \sqrt{7}}.$$

Algebraic and Transcendental

Irrational numbers can be further divided into to categories

- 1 **algebraic numbers:** numbers which are roots of polynomials with integer coefficients, like

$$\sqrt{2}, \quad \sqrt{3}, \quad \text{and} \quad \sqrt[3]{\sqrt{5} + \sqrt{7}}.$$

- 2 **transcendental numbers:** numbers which are not algebraic, like

$$\pi, \quad e, \quad e^{\pi}, \quad \text{and maybe} \quad \pi + e?$$

Algebraic and Transcendental

Irrational numbers can be further divided into to categories

- 1 **algebraic numbers:** numbers which are roots of polynomials with integer coefficients, like

$$\sqrt{2}, \quad \sqrt{3}, \quad \text{and} \quad \sqrt[3]{\sqrt{5} + \sqrt{7}}.$$

- 2 **transcendental numbers:** numbers which are not algebraic, like

$$\pi, \quad e, \quad e^{\pi}, \quad \text{and maybe} \quad \pi + e?$$

- 3 transcendentals are mysterious ... but most real numbers are transcendental!

Outline

- 1 Real Analysis Lecture 2
 - Types of real numbers
 - Integers
 - Upper Bound and Supremum

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

If $b \in S$ also, then b is called a **maximal element** of S

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

If $b \in S$ also, then b is called a **maximal element** of S

Examples:

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

If $b \in S$ also, then b is called a **maximal element** of S

Examples:

- 34 is an upper bound of $[-1, 5]$, but not a maximal element

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

If $b \in S$ also, then b is called a **maximal element** of S

Examples:

- 34 is an upper bound of $[-1, 5]$, but not a maximal element
- 5 is a maximal element of $[-1, 5]$

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

If $b \in S$ also, then b is called a **maximal element** of S

Examples:

- 34 is an upper bound of $[-1, 5]$, but not a maximal element
- 5 is a maximal element of $[-1, 5]$
- 3 is an upper bound of $[0, 3]$, but not a maximal element

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

If $b \in S$ also, then b is called a **maximal element** of S

Examples:

- 34 is an upper bound of $[-1, 5]$, but not a maximal element
- 5 is a maximal element of $[-1, 5]$
- 3 is an upper bound of $[0, 3)$, but not a maximal element
- \mathbb{Z}_+ has no upper bound

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

If $b \in S$ also, then b is called a **maximal element** of S

Examples:

- 34 is an upper bound of $[-1, 5]$, but not a maximal element
- 5 is a maximal element of $[-1, 5]$
- 3 is an upper bound of $[0, 3)$, but not a maximal element
- \mathbb{Z}_+ has no upper bound
- $[3, 7)$ has an upper bound but no maximal element

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both maximal elements of S .

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both maximal elements of S .
Then $x \leq b_1$ and $x \leq b_2$ for all $x \in S$.

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both maximal elements of S .

Then $x \leq b_1$ and $x \leq b_2$ for all $x \in S$.

Moreover, $b_1 \in S$ and $b_2 \in S$.

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both maximal elements of S .

Then $x \leq b_1$ and $x \leq b_2$ for all $x \in S$.

Moreover, $b_1 \in S$ and $b_2 \in S$.

Since $b_1 \in S$ and b_2 is an upper bound of S , $b_1 \leq b_2$.

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both maximal elements of S .

Then $x \leq b_1$ and $x \leq b_2$ for all $x \in S$.

Moreover, $b_1 \in S$ and $b_2 \in S$.

Since $b_1 \in S$ and b_2 is an upper bound of S , $b_1 \leq b_2$.

Likewise, since $b_2 \in S$ and b_1 is an upper bound of S , $b_2 \leq b_1$.

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both maximal elements of S .

Then $x \leq b_1$ and $x \leq b_2$ for all $x \in S$.

Moreover, $b_1 \in S$ and $b_2 \in S$.

Since $b_1 \in S$ and b_2 is an upper bound of S , $b_1 \leq b_2$.

Likewise, since $b_2 \in S$ and b_1 is an upper bound of S , $b_2 \leq b_1$.

By the trichotomy, we find $b_1 = b_2$.

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both maximal elements of S .

Then $x \leq b_1$ and $x \leq b_2$ for all $x \in S$.

Moreover, $b_1 \in S$ and $b_2 \in S$.

Since $b_1 \in S$ and b_2 is an upper bound of S , $b_1 \leq b_2$.

Likewise, since $b_2 \in S$ and b_1 is an upper bound of S , $b_2 \leq b_1$.

By the trichotomy, we find $b_1 = b_2$.

Now we can say *the* maximum, $\max(S)$

Supremum

A **supremum** of a set S of real numbers is a real number $b \in \mathbb{R}$ such that

- b is an upper bound of S
- if $b' < b$, then b' is not an upper bound of S

Supremum

A **supremum** of a set S of real numbers is a real number $b \in \mathbb{R}$ such that

- b is an upper bound of S
- if $b' < b$, then b' is not an upper bound of S

In other words

a supremum is a **least upper bound**

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Hint: use the definition!

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both suprema of S .

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both suprema of S .
Then b_1 and b_2 are both upper bounds of S .

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both suprema of S .
Then b_1 and b_2 are both upper bounds of S .
Since b_1 is a least upper bound, $b_1 \leq b_2$.

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both suprema of S .

Then b_1 and b_2 are both upper bounds of S .

Since b_1 is a least upper bound, $b_1 \leq b_2$.

Since b_2 is also a least upper bound, $b_2 \leq b_1$.

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both suprema of S .

Then b_1 and b_2 are both upper bounds of S .

Since b_1 is a least upper bound, $b_1 \leq b_2$.

Since b_2 is also a least upper bound, $b_2 \leq b_1$.

By the trichotomy, we find $b_1 = b_2$.

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both suprema of S .

Then b_1 and b_2 are both upper bounds of S .

Since b_1 is a least upper bound, $b_1 \leq b_2$.

Since b_2 is also a least upper bound, $b_2 \leq b_1$.

By the trichotomy, we find $b_1 = b_2$.

Now we can say *the* supremum, $\sup(S)$

Challenge!

Problem

Show that if S has a maximum element. Then S has a supremum and $\max(S) = \sup(S)$

Challenge!

Problem

Show that if S has a maximum element. Then S has a supremum and $\max(S) = \sup(S)$

Hint: use the definition!

Challenge!

Problem

Show that if S has a maximum element. Then S has a supremum and $\max(S) = \sup(S)$

Hint: use the definition!

Solution

Let $b = \max(S)$.

Challenge!

Problem

Show that if S has a maximum element. Then S has a supremum and $\max(S) = \sup(S)$

Hint: use the definition!

Solution

Let $b = \max(S)$.

Then b is an upper bound of S and $b \in S$.

Challenge!

Problem

Show that if S has a maximum element. Then S has a supremum and $\max(S) = \sup(S)$

Hint: use the definition!

Solution

Let $b = \max(S)$.

Then b is an upper bound of S and $b \in S$.

If b' is another upper bound of S , then by definition $b \leq b'$.

Challenge!

Problem

Show that if S has a maximum element. Then S has a supremum and $\max(S) = \sup(S)$

Hint: use the definition!

Solution

Let $b = \max(S)$.

Then b is an upper bound of S and $b \in S$.

If b' is another upper bound of S , then by definition $b \leq b'$.

Thus b is the least upper bound of S .

Completeness Axiom


Now we have the machinery in place to state the Completeness Axiom.

Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.


Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.

 **completeness axiom:** if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Completeness Axiom


Now we have the machinery in place to state the Completeness Axiom.

 **completeness axiom:** if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.

 **completeness axiom:** if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

- 3 is the supremum of $(0, 3)$

Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.

A10 completeness axiom: if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

- 3 is the supremum of $(0, 3)$
- 1 is the supremum of

$$\left\{ \frac{n}{n+1} : n \in \mathbb{Z}_+ \right\}$$

Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.

completeness axiom: if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

- 3 is the supremum of $(0, 3)$
- 1 is the supremum of

$$\left\{ \frac{n}{n+1} : n \in \mathbb{Z}_+ \right\}$$

- π is the supremum of

$$\{3, 3.1, 3.14, 3.141, 3.1415, 3.14159, 3.141592, \dots\}.$$