

# MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics  
California State University Fullerton

August 26, 2024

# Outline

- 1 Real Analysis Lecture 1
  - Origin of Real Numbers
  - Properties of real numbers

# Outline

- 1 Real Analysis Lecture 1
  - Origin of Real Numbers
  - Properties of real numbers

# Prehistoric numbers



20,000 BC tallies on Ishango bone

1	11	21	31	41	51
2	12	22	32	42	52
3	13	23	33	43	53
4	14	24	34	44	54
5	15	25	35	45	55
6	16	26	36	46	56
7	17	27	37	47	57
8	18	28	38	48	58
9	19	29	39	49	59
10	20	30	40	50	



$$\frac{1}{3}$$



$$\frac{1}{5}$$



$$\frac{1}{6}$$



$$\frac{1}{10}$$

3400 BC Sumerian system

1000 BC Egyptian fractions

# Invention of zero

# Invention of zero

- 1770 BC - concept of zero in heiroglyph

# Invention of zero

- 1770 BC - concept of zero in heiroglyph
- 36 BC - Mayan/Olmec heiroglyph for zero

# Invention of zero

- 1770 BC - concept of zero in heiroglyph
- 36 BC - Mayan/Olmec heiroglyph for zero
- 150 AD - Ptolemy using it in astronomy



# Invention of zero

- 1770 BC - concept of zero in heiroglyph
- 36 BC - Mayan/Olmec heiroglyph for zero
- 150 AD - Ptolemy using it in astronomy
- medieval scholars - debating the existence of 0



**Babylonian Zero**



**Mayan Zero**



**Hebrew Zero**



**Egyptian Zero**

# Invention of negatives

# Invention of negatives

- 100-50 BC - concept of negatives in China

# Invention of negatives

- 100-50 BC - concept of negatives in China
- 628 AD - used in India in quadratic formula

# Invention of negatives

- 100-50 BC - concept of negatives in China
- 628 AD - used in India in quadratic formula
- 9th century - Islamic mathematicians invent algebra, but hesitant about negatives

# Invention of negatives

- 100-50 BC - concept of negatives in China
- 628 AD - used in India in quadratic formula
- 9th century - Islamic mathematicians invent algebra, but hesitant about negatives
- 12th century - Islamic mathematicians add negative solutions of quadratics, but discard them

# Invention of negatives

- 100-50 BC - concept of negatives in China
- 628 AD - used in India in quadratic formula
- 9th century - Islamic mathematicians invent algebra, but hesitant about negatives
- 12th century - Islamic mathematicians add negative solutions of quadratics, but discard them
- 1202, 1225 - Fibonacci allows negatives as solutions for financial problems

# Invention of negatives

- 100-50 BC - concept of negatives in China
- 628 AD - used in India in quadratic formula
- 9th century - Islamic mathematicians invent algebra, but hesitant about negatives
- 12th century - Islamic mathematicians add negative solutions of quadratics, but discard them
- 1202, 1225 - Fibonacci allows negatives as solutions for financial problems
- up to 18th century - rejected by western sources, referred to as "absurd numbers"



# Invention of rationals

# Invention of rationals

- 1000 BC - appear in Egyptian hieroglyphs

# Invention of rationals

- 1000 BC - appear in Egyptian hieroglyphs
- Greek and Indian mathematicians used them regularly in studying number theory

# Invention of rationals

- 1000 BC - appear in Egyptian hieroglyphs
- Greek and Indian mathematicians used them regularly in studying number theory
- 300 BC - appear in Euclid's elements



# Discovery of irrationals

# Discovery of irrationals

- 800 BC - Indian mathematicians discuss nonrationality of square roots

# Discovery of irrationals

- 800 BC - Indian mathematicians discuss nonrationality of square roots
- 500 BC - Greek mathematicians discover the existence of irrational numbers (sides of the pentagram)

# Discovery of irrationals

- 800 BC - Indian mathematicians discuss nonrationality of square roots
- 500 BC - Greek mathematicians discover the existence of irrational numbers (sides of the pentagram)
- discovered by Hippasus, who legend says was thrown into the sea for his discovery



# Discovery of irrationals

- 800 BC - Indian mathematicians discuss nonrationality of square roots
- 500 BC - Greek mathematicians discover the existence of irrational numbers (sides of the pentagram)
- discovered by Hippasus, who legend says was thrown into the sea for his discovery
- 14th-16th century - Indian mathematicians discover infinite series expressions for  $\pi$

# Discovery of irrationals

- 800 BC - Indian mathematicians discuss nonrationality of square roots
- 500 BC - Greek mathematicians discover the existence of irrational numbers (sides of the pentagram)
- discovered by Hippasus, who legend says was thrown into the sea for his discovery
- 14th-16th century - Indian mathematicians discover infinite series expressions for  $\pi$
- 17th century - European mathematicians distinguish between transcendentals and algebraic numbers

# Outline

- 1 Real Analysis Lecture 1
  - Origin of Real Numbers
  - Properties of real numbers

# Philosophy of our presentation

# Philosophy of our presentation

- we will not construct the reals (ref. Foundations of Analysis by Landau for a rigorous construction from the ground up using minimal axioms and Dedekind cuts)

# Philosophy of our presentation

- we will not construct the reals (ref. Foundations of Analysis by Landau for a rigorous construction from the ground up using minimal axioms and Dedekind cuts)
- instead, we will take for granted that the reals exist and describe ten fundamental rules (axioms) for how it behaves

# Philosophy of our presentation

- we will not construct the reals (ref. Foundations of Analysis by Landau for a rigorous construction from the ground up using minimal axioms and Dedekind cuts)
- instead, we will take for granted that the reals exist and describe ten fundamental rules (axioms) for how it behaves
- the integers, rationals, etc. will be defined in terms of the reals

# Field axioms

A **field** is a set  $F$  with a way to do addition  $+$  and multiplication  $\times$  with



# Field axioms

A **field** is a set  $F$  with a way to do addition  $+$  and multiplication  $\times$  with

**A1 commutativity:**  $x + y = y + x$  and  $xy = yx$

# Field axioms

A **field** is a set  $F$  with a way to do addition  $+$  and multiplication  $\times$  with

**A1 commutativity:**  $x + y = y + x$  and  $xy = yx$

**A2 associativity:**  $(x + y) + z = x + (y + z)$  and  $(xy)z = x(yz)$

# Field axioms

A **field** is a set  $F$  with a way to do addition  $+$  and multiplication  $\times$  with

- A1 **commutativity:**  $x + y = y + x$  and  $xy = yx$
- A2 **associativity:**  $(x + y) + z = x + (y + z)$  and  $(xy)z = x(yz)$
- A3 **distributivity:**  $x(y + z) = xy + xz$

# Field axioms

A **field** is a set  $F$  with a way to do addition  $+$  and multiplication  $\times$  with

- A1 **commutativity:**  $x + y = y + x$  and  $xy = yx$
- A2 **associativity:**  $(x + y) + z = x + (y + z)$  and  $(xy)z = x(yz)$
- A3 **distributivity:**  $x(y + z) = xy + xz$
- A4 **additive inverse:** given  $x, y \in F$ , there exists a unique  $z \in F$  with  $x = y + z$

Notation:  $z := x - y$

# Field axioms

A **field** is a set  $F$  with a way to do addition  $+$  and multiplication  $\times$  with

- A1 **commutativity:**  $x + y = y + x$  and  $xy = yx$
- A2 **associativity:**  $(x + y) + z = x + (y + z)$  and  $(xy)z = x(yz)$
- A3 **distributivity:**  $x(y + z) = xy + xz$
- A4 **additive inverse:** given  $x, y \in F$ , there exists a unique  $z \in F$  with  $x = y + z$

Notation:  $z := x - y$

- A5 **multiplicative inverse:** given  $x, y \in F$  with  $y \neq 0$ , there exists a unique  $z \in F$  with  $x = yz$

Notation:  $z := x/y$

# Field axioms

A **field** is a set  $F$  with a way to do addition  $+$  and multiplication  $\times$  with

- A1 **commutativity:**  $x + y = y + x$  and  $xy = yx$
- A2 **associativity:**  $(x + y) + z = x + (y + z)$  and  $(xy)z = x(yz)$
- A3 **distributivity:**  $x(y + z) = xy + xz$
- A4 **additive inverse:** given  $x, y \in F$ , there exists a unique  $z \in F$  with  $x = y + z$

Notation:  $z := x - y$

- A5 **multiplicative inverse:** given  $x, y \in F$  with  $y \neq 0$ , there exists a unique  $z \in F$  with  $x = yz$

Notation:  $z := x/y$

**Special notation:**  $0 := x - x$  and  $1 := x/x$ , and  $-x := 0 - x$

# Challenge!

## Problem

use the axioms to show that  $(x + y)z = xz + yz$ .

# Challenge!

## Problem

use the axioms to show that  $(x + y)z = xz + yz$ .

## Solution

$$(x + y)z \stackrel{A1}{=} z(x + y) \stackrel{A3}{=} zx + zy \stackrel{A1}{=} xz + yz.$$



# Challenge!

## Problem

use the axioms to show that the value of  $x - x$  doesn't depend on  $x$ .

# Challenge!

## Problem

use the axioms to show that the value of  $x - x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x - x$  and  $z_2 = y - y$ .

# Challenge!

## Problem

use the axioms to show that the value of  $x - x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x - x$  and  $z_2 = y - y$ .  
Then  $x = x + z_1$  and  $y = y + z_2$ .

# Challenge!

## Problem

use the axioms to show that the value of  $x - x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x - x$  and  $z_2 = y - y$ .

Then  $x = x + z_1$  and  $y = y + z_2$ .

Therefore we have

$$(x + y) + z_1 \stackrel{A2}{=} x + (y + z_1) \stackrel{A1}{=} x + (z_1 + y) \stackrel{A2}{=} (x + z_1) + y = x + y,$$

# Challenge!

## Problem

use the axioms to show that the value of  $x - x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x - x$  and  $z_2 = y - y$ .

Then  $x = x + z_1$  and  $y = y + z_2$ .

Therefore we have

$$(x + y) + z_1 \stackrel{A2}{=} x + (y + z_1) \stackrel{A1}{=} x + (z_1 + y) \stackrel{A2}{=} (x + z_1) + y = x + y,$$

$$(x + y) + z_2 \stackrel{A2}{=} x + (y + z_2) = x + y,$$

# Challenge!

## Problem

use the axioms to show that the value of  $x - x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x - x$  and  $z_2 = y - y$ .

Then  $x = x + z_1$  and  $y = y + z_2$ .

Therefore we have

$$(x + y) + z_1 \stackrel{A2}{=} x + (y + z_1) \stackrel{A1}{=} x + (z_1 + y) \stackrel{A2}{=} (x + z_1) + y = x + y,$$

$$(x + y) + z_2 \stackrel{A2}{=} x + (y + z_2) = x + y,$$

so that

$$z_1 = (x + y) - (x + y) = z_2.$$

# Challenge!

## Problem

use the axioms to show that the value of  $x/x$  doesn't depend on  $x$ .

# Challenge!

## Problem

use the axioms to show that the value of  $x/x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x/x$  and  $z_2 = y/y$ .



# Challenge!

## Problem

use the axioms to show that the value of  $x/x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x/x$  and  $z_2 = y/y$ .  
Then  $x = z_1 x$  and  $y = z_2 y$ .

# Challenge!

## Problem

use the axioms to show that the value of  $x/x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x/x$  and  $z_2 = y/y$ .

Then  $x = z_1 x$  and  $y = z_2 y$ .

Therefore we have

$$z_1(xy) \stackrel{A2}{=} (z_1 x)y = xy,$$

# Challenge!

## Problem

use the axioms to show that the value of  $x/x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x/x$  and  $z_2 = y/y$ .

Then  $x = z_1 x$  and  $y = z_2 y$ .

Therefore we have

$$z_1(xy) \stackrel{A2}{=} (z_1 x)y = xy,$$

$$z_2(xy) \stackrel{A2}{=} (z_2 x)y \stackrel{A1}{=} (xz_2)y \stackrel{A2}{=} x(z_2 y) = xy,$$

# Challenge!

## Problem

use the axioms to show that the value of  $x/x$  doesn't depend on  $x$ .

## Solution

Suppose  $z_1 = x/x$  and  $z_2 = y/y$ .

Then  $x = z_1 x$  and  $y = z_2 y$ .

Therefore we have

$$z_1(xy) \stackrel{A2}{=} (z_1 x)y = xy,$$

$$z_2(xy) \stackrel{A2}{=} (z_2 x)y \stackrel{A1}{=} (xz_2)y \stackrel{A2}{=} x(z_2 y) = xy,$$

so that

$$z_1 = (xy)/(xy) = z_2.$$

# Field examples

Lots of examples of fields!

# Field examples

Lots of examples of fields!

- **rational numbers**

$$\mathbb{Q} = \{a/b : a, b \text{ integers with } b \neq 0\}.$$

# Field examples

Lots of examples of fields!

- **rational numbers**

$$\mathbb{Q} = \{a/b : a, b \text{ integers with } b \neq 0\}.$$

- **real numbers**  $\mathbb{R}$  (what are they?)

# Field examples

Lots of examples of fields!

- **rational numbers**

$$\mathbb{Q} = \{a/b : a, b \text{ integers with } b \neq 0\}.$$

- **real numbers**  $\mathbb{R}$  (what are they?)

- **complex numbers**

$$\mathbb{C} = \{a + ib : a, b \text{ real numbers}\}.$$



# Field examples

Lots of examples of fields!

- **rational numbers**

$$\mathbb{Q} = \{a/b : a, b \text{ integers with } b \neq 0\}.$$

- **real numbers**  $\mathbb{R}$  (what are they?)

- **complex numbers**

$$\mathbb{C} = \{a + ib : a, b \text{ real numbers}\}.$$

- **field of Boolean numbers**

$$\mathbb{F}_2 = \{0, 1\}$$

$$0 + 0 = 0, \quad 0 + 1 = 1, \quad 1 + 0 = 1, \quad 1 + 1 = 0$$

$$0 \cdot 0 = 0, \quad 0 \cdot 1 = 0, \quad 1 \cdot 0 = 0, \quad 1 \cdot 1 = 1$$

# Challenge!

## Problem

explain why the set of integers isn't a field

# Challenge!

## Problem

explain why the set of integers isn't a field

## Solution

There aren't multiplicative inverses!

# Challenge!

## Problem

explain why the set of integers isn't a field

## Solution

There aren't multiplicative inverses! For example,  $1/2$  doesn't make sense because there isn't an integer  $z$  with  $1 = 2z$ .

# Order axioms

An **ordered field** is a field  $F$  with a relation  $<$

# Order axioms

An **ordered field** is a field  $F$  with a relation  $<$

**A6 trichotomy:** for any  $x, y \in F$ , exactly one of the following holds:

$$x < y, \quad y < x, \quad \text{or} \quad x = y$$

# Order axioms

An **ordered field** is a field  $F$  with a relation  $<$

- A6 trichotomy:** for any  $x, y \in F$ , exactly one of the following holds:

$$x < y, \quad y < x, \quad \text{or} \quad x = y$$

- A7 compatible with  $+$ :**  $x < y$  implies  $x + z < y + z$

# Order axioms

An **ordered field** is a field  $F$  with a relation  $<$

- Ⓐ6 **trichotomy**: for any  $x, y \in F$ , exactly one of the following holds:

$$x < y, \quad y < x, \quad \text{or} \quad x = y$$

- Ⓐ7 **compatible with  $+$** :  $x < y$  implies  $x + z < y + z$

- Ⓐ8 **compatible with  $\times$** :  $0 < x$  and  $0 < y$  implies  $0 < xy$



# Order axioms

An **ordered field** is a field  $F$  with a relation  $<$

- A6 trichotomy:** for any  $x, y \in F$ , exactly one of the following holds:

$$x < y, \quad y < x, \quad \text{or} \quad x = y$$

- A7 compatible with  $+$ :**  $x < y$  implies  $x + z < y + z$
- A8 compatible with  $\times$ :**  $0 < x$  and  $0 < y$  implies  $0 < xy$
- A9 transitivity:**  $x < y$  and  $y < z$  implies  $x < z$

# Order axioms

An **ordered field** is a field  $F$  with a relation  $<$

**A6 trichotomy:** for any  $x, y \in F$ , exactly one of the following holds:

$$x < y, \quad y < x, \quad \text{or} \quad x = y$$

**A7 compatible with  $+$ :**  $x < y$  implies  $x + z < y + z$

**A8 compatible with  $\times$ :**  $0 < x$  and  $0 < y$  implies  $0 < xy$

**A9 transitivity:**  $x < y$  and  $y < z$  implies  $x < z$

**Special notation:**

- $x > y$  means  $y < x$
- $x \leq y$  means  $x < y$  or  $x = y$
- $x \geq y$  means  $y \leq x$

# Challenge!

## Problem

explain why  $\mathbb{F}_2$  cannot be made into an ordered field

# Challenge!

## Problem

explain why  $\mathbb{F}_2$  cannot be made into an ordered field

## Solution

There is no way to put an order relation  $<$  on it!

# Challenge!

## Problem

explain why  $\mathbb{F}_2$  cannot be made into an ordered field

## Solution

There is no way to put an order relation  $<$  on it!

The trichotomy (Axiom 6) would say either  $0 < 1$  or  $1 < 0$ .

# Challenge!

## Problem

explain why  $\mathbb{F}_2$  cannot be made into an ordered field

## Solution

There is no way to put an order relation  $<$  on it!

The trichotomy (Axiom 6) would say either  $0 < 1$  or  $1 < 0$ .

If  $0 < 1$ , then by Axiom 7

$$1 = 0 + 1 < 1 + 1 = 0$$

but this means both  $0 < 1$  and  $1 < 0$ , which violates the trichotomy.

# Challenge!

## Problem

explain why  $\mathbb{F}_2$  cannot be made into an ordered field

## Solution

There is no way to put an order relation  $<$  on it!

The trichotomy (Axiom 6) would say either  $0 < 1$  or  $1 < 0$ .

If  $0 < 1$ , then by Axiom 7

$$1 = 0 + 1 < 1 + 1 = 0$$

but this means both  $0 < 1$  and  $1 < 0$ , which violates the trichotomy. Same outcome if  $1 < 0$ .

# Apostol Theorem 1.1

## Theorem

*If  $a < b + \epsilon$  for all  $\epsilon > 0$ , then  $a \leq b$ .*



# Apostol Theorem 1.1

## Theorem

*If  $a < b + \epsilon$  for all  $\epsilon > 0$ , then  $a \leq b$ .*

## Proof.

(Contradiction.) Assume the statement is false.

# Apostol Theorem 1.1

## Theorem

*If  $a < b + \epsilon$  for all  $\epsilon > 0$ , then  $a \leq b$ .*

## Proof.

(Contradiction.) Assume the statement is false.  
Then the trichotomy says  $a > b$ .

# Apostol Theorem 1.1

## Theorem

*If  $a < b + \epsilon$  for all  $\epsilon > 0$ , then  $a \leq b$ .*

## Proof.

(Contradiction.) Assume the statement is false.

Then the trichotomy says  $a > b$ .

This means  $a - b > 0$  by A7.

# Apostol Theorem 1.1

## Theorem

*If  $a < b + \epsilon$  for all  $\epsilon > 0$ , then  $a \leq b$ .*

## Proof.

(Contradiction.) Assume the statement is false.

Then the trichotomy says  $a > b$ .

This means  $a - b > 0$  by A7.

This means  $(a - b)/2 > 0$  by A8.

# Apostol Theorem 1.1

## Theorem

*If  $a < b + \epsilon$  for all  $\epsilon > 0$ , then  $a \leq b$ .*

## Proof.

(Contradiction.) Assume the statement is false.

Then the trichotomy says  $a > b$ .

This means  $a - b > 0$  by A7.

This means  $(a - b)/2 > 0$  by A8.

This means

$$a < b + \frac{a - b}{2} = \frac{a + b}{2} \stackrel{\text{A7\&A8}}{<} \frac{a + a}{2} = a.$$

# Apostol Theorem 1.1

## Theorem

*If  $a < b + \epsilon$  for all  $\epsilon > 0$ , then  $a \leq b$ .*

## Proof.

(Contradiction.) Assume the statement is false.

Then the trichotomy says  $a > b$ .

This means  $a - b > 0$  by A7.

This means  $(a - b)/2 > 0$  by A8.

This means

$$a < b + \frac{a - b}{2} = \frac{a + b}{2} \stackrel{\text{A7\&A8}}{<} \frac{a + a}{2} = a.$$

This says  $a < a$ , which contradicts the trichotomy. □