

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

September 18, 2024

Outline

- 1 Real Analysis Lecture 7
 - More with cardinality
 - Set algebra
 - Open Balls and Open Sets

Outline

- 1 Real Analysis Lecture 7
 - More with cardinality
 - Set algebra
 - Open Balls and Open Sets

Ultimate Cantor diagonalization

The **power set** $\mathcal{P}(A)$ of a set A is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

Ultimate Cantor diagonalization

The **power set** $\mathcal{P}(A)$ of a set A is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

Theorem

The cardinality of $\mathcal{P}(A)$ is strictly larger than A .

Ultimate Cantor diagonalization

The **power set** $\mathcal{P}(A)$ of a set A is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

Theorem

The cardinality of $\mathcal{P}(A)$ is strictly larger than A .

Proof.

Ultimate Cantor diagonalization

The **power set** $\mathcal{P}(A)$ of a set A is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

Theorem

The cardinality of $\mathcal{P}(A)$ is strictly larger than A .

Proof.

Suppose $f : A \rightarrow \mathcal{P}(A)$ is surjective.

Ultimate Cantor diagonalization

The **power set** $\mathcal{P}(A)$ of a set A is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

Theorem

The cardinality of $\mathcal{P}(A)$ is strictly larger than A .

Proof.

Suppose $f : A \rightarrow \mathcal{P}(A)$ is surjective.

Consider the set

$$S = \{a \in A : a \notin f(a)\}.$$



Challenge

Challenge

Since f is surjective, $S = f(x)$ for some $x \in A$.

Challenge

Since f is surjective, $S = f(x)$ for some $x \in A$.

Problem

Consider the statement $x \in S$. What can you conclude?

Challenge

Problem

Is there a set with cardinality larger than \mathbb{R} ?

Challenge

Problem

Show that the cardinality of the line segment $(0, 1)$ and the square $(0, 1) \times (0, 1)$ is the same.

Outline

- 1 Real Analysis Lecture 7
 - More with cardinality
 - Set algebra
 - Open Balls and Open Sets

Unions of sets

Unions:

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

$$A_1 \cup A_2 \cup \cdots \cup A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

$$A_1 \cup A_2 \cup \cdots \cup A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

NOTATION:

$$\bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \cdots \cup A_n.$$

Intersections of sets

Intersections:

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

$$A_1 \cap A_2 \cap \cdots \cap A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

$$A_1 \cap A_2 \cap \cdots \cap A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

NOTATION:

$$\bigcap_{i=1}^n A_i = A_1 \cap A_2 \cap \cdots \cap A_n.$$

Complements of sets

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Theorem

De Morgan's Laws

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Theorem

De Morgan's Laws

$$B - \bigcup_{i=1}^n A_i = \bigcap_{i=1}^n (B - A_i)$$

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Theorem

De Morgan's Laws

$$B - \bigcup_{i=1}^n A_i = \bigcap_{i=1}^n (B - A_i)$$

$$B - \bigcap_{i=1}^n A_i = \bigcup_{i=1}^n (B - A_i)$$

Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- I any set, called the **index set**

Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- I any set, called the **index set**
- A_i is a set for all i

Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- I any set, called the **index set**
- A_i is a set for all i

$$\bigcup_{i \in I} A_i = \{x : x \in A_i, \text{ for some } i \in I\}$$

Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- I any set, called the **index set**
- A_i is a set for all i

$$\bigcup_{i \in I} A_i = \{x : x \in A_i, \text{ for some } i \in I\}$$

$$\bigcap_{i \in I} A_i = \{x : x \in A_i, \text{ for all } i \in I\}$$

Challenge

- index set $I = (1, 2)$
- family of sets $\{A_i : i \in I\}$
- $A_i = [0, i]$

Problem

Determine $\bigcup_{i \in I} A_i$.

Challenge

- index set $I = \mathbb{Z}_+$
- family of sets $\{A_i : i \in I\}$
- $A_i = [0, i)$

Problem

Determine $\bigcap_{i \in I} A_i$.

Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

Theorem

The union of a countable family of countable sets is countable.

Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

Theorem

The union of a countable family of countable sets is countable.

Proof.

Let $\{A_i : i \in I\}$ be a countable family of countable sets.

Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

Theorem

The union of a countable family of countable sets is countable.

Proof.

Let $\{A_i : i \in I\}$ be a countable family of countable sets.
 I is countable.

Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

Theorem

The union of a countable family of countable sets is countable.

Proof.

Let $\{A_i : i \in I\}$ be a countable family of countable sets. I is countable. There is a surjection $\mathbb{Z}_+ \rightarrow I$

Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

Theorem

The union of a countable family of countable sets is countable.

Proof.

Let $\{A_i : i \in I\}$ be a countable family of countable sets. I is countable. There is a surjection $\mathbb{Z}_+ \rightarrow I$. In other words, we can enumerate $I = \{i_1, i_2, i_3, \dots\}$.

Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

Theorem

The union of a countable family of countable sets is countable.

Proof.

Let $\{A_i : i \in I\}$ be a countable family of countable sets.
 I is countable. There is a surjection $\mathbb{Z}_+ \rightarrow I$
In other words, we can enumerate $I = \{i_1, i_2, i_3, \dots\}$.
 A_i is countable for all $i \in I$.

Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

Theorem

The union of a countable family of countable sets is countable.

Proof.

Let $\{A_i : i \in I\}$ be a countable family of countable sets.

I is countable. There is a surjection $\mathbb{Z}_+ \rightarrow I$

In other words, we can enumerate $I = \{i_1, i_2, i_3, \dots\}$.

A_i is countable for all $i \in I$.

So again $A_i = \{a_{i1}, a_{i2}, a_{i3}, \dots\}$.

Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

Theorem

The union of a countable family of countable sets is countable.

Proof.

Let $\{A_i : i \in I\}$ be a countable family of countable sets.

I is countable. There is a surjection $\mathbb{Z}_+ \rightarrow I$

In other words, we can enumerate $I = \{i_1, i_2, i_3, \dots\}$.

A_i is countable for all $i \in I$.

So again $A_i = \{a_{i1}, a_{i2}, a_{i3}, \dots\}$.

Define $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \bigcup_{i \in I} A_i$

Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

Theorem

The union of a countable family of countable sets is countable.

Proof.

Let $\{A_i : i \in I\}$ be a countable family of countable sets.

I is countable. There is a surjection $\mathbb{Z}_+ \rightarrow I$

In other words, we can enumerate $I = \{i_1, i_2, i_3, \dots\}$.

A_i is countable for all $i \in I$.

So again $A_i = \{a_{i1}, a_{i2}, a_{i3}, \dots\}$.

Define $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \bigcup_{i \in I} A_i$

$f(j, k) = a_{i_j, k}$.

Countable unions

A **countable family of sets** is a family of sets where the index set is countable.

Theorem

The union of a countable family of countable sets is countable.

Proof.

Let $\{A_i : i \in I\}$ be a countable family of countable sets.

I is countable. There is a surjection $\mathbb{Z}_+ \rightarrow I$

In other words, we can enumerate $I = \{i_1, i_2, i_3, \dots\}$.

A_i is countable for all $i \in I$.

So again $A_i = \{a_{i1}, a_{i2}, a_{i3}, \dots\}$.

Define $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \bigcup_{i \in I} A_i$

$f(j, k) = a_{i_j, k}$. Surjection!



Challenge!

A real number is called **algebraic** if it is a root of a polynomial with integer coefficients.

Challenge!

A real number is called **algebraic** if it is a root of a polynomial with integer coefficients.

Problem

Prove that the set of all algebraic numbers is countable.

Challenge!

A real number is called **algebraic** if it is a root of a polynomial with integer coefficients.

Problem

Prove that the set of all algebraic numbers is countable.

Hint: use the previous theorem!

Outline

- 1 Real Analysis Lecture 7
 - More with cardinality
 - Set algebra
 - Open Balls and Open Sets

Euclidean space

n -dimensional **euclidean space** is

$$\mathbb{R}^n = \{(a_1, a_2, \dots, a_n) : a_1, \dots, a_n \in \mathbb{R}\}.$$

Definitions:

Given $\vec{x} = (x_1, \dots, x_n)$ and $\vec{y} = (y_1, \dots, y_n)$

- (a) $\vec{x} + \vec{y} = (x_1 + y_1, \dots, x_n + y_n)$
- (b) $a\vec{x} = (ax_1, \dots, ax_n)$
- (c) $\vec{x} - \vec{y} = \vec{x} + (-1)\vec{y} = (x_1 - y_1, \dots, x_n - y_n)$
- (d) $\vec{0} = (0, \dots, 0)$
- (e) $\vec{x} \cdot \vec{y} = \sum_{i=1}^n x_i y_i$
- (f) $|\vec{x}| = (\vec{x} \cdot \vec{x})^{1/2} = (\sum_{i=1}^n x_i^2)^{1/2}$