

# MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics  
California State University Fullerton

October 21, 2024

# Outline

- 1 Real Analysis Lecture 14
  - Cauchy sequences
  - Limit of a function

# Outline

- 1 Real Analysis Lecture 14
  - Cauchy sequences
  - Limit of a function

# Cauchy sequence

A sequence  $\{x_n\}$  in a metric space  $(M, d)$  is called Cauchy if the terms get arbitrarily close together as  $n$  gets arbitrarily large.

# Cauchy sequence

A sequence  $\{x_n\}$  in a metric space  $(M, d)$  is called Cauchy if the terms get arbitrarily close together as  $n$  gets arbitrarily large.

More formally:

# Cauchy sequence

A sequence  $\{x_n\}$  in a metric space  $(M, d)$  is called Cauchy if the terms get arbitrarily close together as  $n$  gets arbitrarily large.

More formally:

## Definition

A sequence  $\{x_n\}$  is called a **Cauchy sequence** if

$\forall \epsilon > 0$ , there exists  $N \in \mathbb{Z}_+$  with  $m, n \geq N \Rightarrow d(x_m, x_n) < \epsilon$ .

# Cauchy sequence

A sequence  $\{x_n\}$  in a metric space  $(M, d)$  is called Cauchy if the terms get arbitrarily close together as  $n$  gets arbitrarily large.

More formally:

## Definition

A sequence  $\{x_n\}$  is called a **Cauchy sequence** if

$$\forall \epsilon > 0, \text{ there exists } N \in \mathbb{Z}_+ \text{ with } m, n \geq N \Rightarrow d(x_m, x_n) < \epsilon.$$

- every convergent sequence is Cauchy

# Cauchy sequence

A sequence  $\{x_n\}$  in a metric space  $(M, d)$  is called Cauchy if the terms get arbitrarily close together as  $n$  gets arbitrarily large.

More formally:

## Definition

A sequence  $\{x_n\}$  is called a **Cauchy sequence** if

$$\forall \epsilon > 0, \text{ there exists } N \in \mathbb{Z}_+ \text{ with } m, n \geq N \Rightarrow d(x_m, x_n) < \epsilon.$$

- every convergent sequence is Cauchy
- is every Cauchy sequence convergent?



# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

Suppose  $\{x_n\}$  is Cauchy.

# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

Suppose  $\{x_n\}$  is Cauchy.  
Let  $\epsilon = 2024$ .

# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

Suppose  $\{x_n\}$  is Cauchy.

Let  $\epsilon = 2024$ .

Then  $\exists N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < 2024$  for all  $m, n \geq N$ .

# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

Suppose  $\{x_n\}$  is Cauchy.

Let  $\epsilon = 2024$ .

Then  $\exists N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < 2024$  for all  $m, n \geq N$ .

Therefore for all  $n \geq N$ , we have  $d(x_n, x_N) \leq 2024$ .

# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

Suppose  $\{x_n\}$  is Cauchy.

Let  $\epsilon = 2024$ .

Then  $\exists N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < 2024$  for all  $m, n \geq N$ .

Therefore for all  $n \geq N$ , we have  $d(x_n, x_N) \leq 2024$ .

So if we define

$$R = \max\{d(x_1, x_N), d(x_2, x_N), \dots, d(x_{n-1}, x_N), 2024\},$$

# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

Suppose  $\{x_n\}$  is Cauchy.

Let  $\epsilon = 2024$ .

Then  $\exists N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < 2024$  for all  $m, n \geq N$ .

Therefore for all  $n \geq N$ , we have  $d(x_n, x_N) \leq 2024$ .

So if we define

$$R = \max\{d(x_1, x_N), d(x_2, x_N), \dots, d(x_{n-1}, x_N), 2024\},$$

then  $d(x_n, x_N) \leq 2024$  for all  $n \geq 1$ .



# Challenge!

## Problem

Prove that if  $\{x_n\}$  is a Cauchy sequence in a metric space  $(M, d)$ , then it is bounded.

## Solution

Suppose  $\{x_n\}$  is Cauchy.

Let  $\epsilon = 2024$ .

Then  $\exists N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < 2024$  for all  $m, n \geq N$ .

Therefore for all  $n \geq N$ , we have  $d(x_n, x_N) \leq 2024$ .

So if we define

$$R = \max\{d(x_1, x_N), d(x_2, x_N), \dots, d(x_{n-1}, x_N), 2024\},$$

then  $d(x_n, x_N) \leq 2024$  for all  $n \geq 1$ .

Hence  $X = \{x_1, x_2, \dots\} \subseteq B_M(x_N, R)$ , and is bounded.

# Cauchy sequence

Every convergent sequence is Cauchy.

# Cauchy sequence

Every convergent sequence is Cauchy.

## Theorem

*Let  $(M, d)$  be a metric space and suppose that  $\{x_n\}$  is a sequence in  $M$  which converges to  $L \in M$ .*

*Then  $\{x_n\}$  is Cauchy.*

# Cauchy sequence

Every convergent sequence is Cauchy.

## Theorem

*Let  $(M, d)$  be a metric space and suppose that  $\{x_n\}$  is a sequence in  $M$  which converges to  $L \in M$ .*

*Then  $\{x_n\}$  is Cauchy.*

## Proof.

# Cauchy sequence

Every convergent sequence is Cauchy.

## Theorem

*Let  $(M, d)$  be a metric space and suppose that  $\{x_n\}$  is a sequence in  $M$  which converges to  $L \in M$ .*

*Then  $\{x_n\}$  is Cauchy.*

## Proof.

Let  $\epsilon > 0$ .

# Cauchy sequence

Every convergent sequence is Cauchy.

## Theorem

*Let  $(M, d)$  be a metric space and suppose that  $\{x_n\}$  is a sequence in  $M$  which converges to  $L \in M$ .*

*Then  $\{x_n\}$  is Cauchy.*

## Proof.

Let  $\epsilon > 0$ .

Then there exists  $N$  such that  $n \geq N$  implies  $d(x_n, L) < \epsilon/2$ .

# Cauchy sequence

Every convergent sequence is Cauchy.

## Theorem

*Let  $(M, d)$  be a metric space and suppose that  $\{x_n\}$  is a sequence in  $M$  which converges to  $L \in M$ .*

*Then  $\{x_n\}$  is Cauchy.*

## Proof.

Let  $\epsilon > 0$ .

Then there exists  $N$  such that  $n \geq N$  implies  $d(x_n, L) < \epsilon/2$ .

Therefore for any  $m, n \geq N$  we have

$$d(x_m, x_n) \leq d(x_m, L) + d(L, x_n) < \epsilon/2 + \epsilon/2 = \epsilon.$$

Since  $\epsilon > 0$  was arbitrary, this proves that  $\{x_n\}$  is Cauchy. □

# Cauchy sequence

Partial converse.



# Cauchy sequence

Partial converse.

## Theorem

*Let  $M = \mathbb{R}^n$  with the Euclidean metric, and suppose that  $\{x_n\}$  is a Cauchy sequence in  $M$ .*

*Then  $\{x_n\}$  converges.*

# Cauchy sequence

Partial converse.

## Theorem

*Let  $M = \mathbb{R}^n$  with the Euclidean metric, and suppose that  $\{x_n\}$  is a Cauchy sequence in  $M$ .*

*Then  $\{x_n\}$  converges.*

## Proof.

# Cauchy sequence

Partial converse.

## Theorem

*Let  $M = \mathbb{R}^n$  with the Euclidean metric, and suppose that  $\{x_n\}$  is a Cauchy sequence in  $M$ .*

*Then  $\{x_n\}$  converges.*

## Proof.

Consider the range  $X = \{x_1, x_2, \dots\}$  of  $\{x_n\}$ .

# Cauchy sequence

Partial converse.

## Theorem

*Let  $M = \mathbb{R}^n$  with the Euclidean metric, and suppose that  $\{x_n\}$  is a Cauchy sequence in  $M$ .*

*Then  $\{x_n\}$  converges.*

## Proof.

Consider the range  $X = \{x_1, x_2, \dots\}$  of  $\{x_n\}$ .

Since  $\{x_n\}$  is Cauchy,  $X$  is bounded.

# Cauchy sequence

Partial converse.

## Theorem

*Let  $M = \mathbb{R}^n$  with the Euclidean metric, and suppose that  $\{x_n\}$  is a Cauchy sequence in  $M$ .*

*Then  $\{x_n\}$  converges.*

## Proof.

Consider the range  $X = \{x_1, x_2, \dots\}$  of  $\{x_n\}$ .

Since  $\{x_n\}$  is Cauchy,  $X$  is bounded.

We consider two cases:

# Cauchy sequence

Partial converse.

## Theorem

*Let  $M = \mathbb{R}^n$  with the Euclidean metric, and suppose that  $\{x_n\}$  is a Cauchy sequence in  $M$ .*

*Then  $\{x_n\}$  converges.*

## Proof.

Consider the range  $X = \{x_1, x_2, \dots\}$  of  $\{x_n\}$ .

Since  $\{x_n\}$  is Cauchy,  $X$  is bounded.

We consider two cases:

- when  $X$  is finite

# Cauchy sequence

Partial converse.

## Theorem

*Let  $M = \mathbb{R}^n$  with the Euclidean metric, and suppose that  $\{x_n\}$  is a Cauchy sequence in  $M$ .*

*Then  $\{x_n\}$  converges.*

## Proof.

Consider the range  $X = \{x_1, x_2, \dots\}$  of  $\{x_n\}$ .

Since  $\{x_n\}$  is Cauchy,  $X$  is bounded.

We consider two cases:

- when  $X$  is finite
- when  $X$  is infinite



# Cauchy sequence

Proof.

**Case I:** Assume  $X$  is finite.



# Cauchy sequence

Proof.

**Case I:** Assume  $X$  is finite.

Then

$$\{d(x, y) : x, y \in X, x \neq y\}$$

is a finite set of positive values.

# Cauchy sequence

Proof.

**Case I:** Assume  $X$  is finite.

Then

$$\{d(x, y) : x, y \in X, x \neq y\}$$

is a finite set of positive values.

Therefore it has a minimum  $\epsilon > 0$ .

# Cauchy sequence

Proof.

**Case I:** Assume  $X$  is finite.

Then

$$\{d(x, y) : x, y \in X, x \neq y\}$$

is a finite set of positive values.

Therefore it has a minimum  $\epsilon > 0$ .

Since  $\{x_n\}$  is Cauchy, there exists  $N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < \epsilon$  for all  $m, n \geq N$ .

# Cauchy sequence

Proof.

**Case I:** Assume  $X$  is finite.

Then

$$\{d(x, y) : x, y \in X, x \neq y\}$$

is a finite set of positive values.

Therefore it has a minimum  $\epsilon > 0$ .

Since  $\{x_n\}$  is Cauchy, there exists  $N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < \epsilon$  for all  $m, n \geq N$ .

This implies  $x_n = x_N$  for all  $n \geq N$ .

# Cauchy sequence

Proof.

**Case I:** Assume  $X$  is finite.

Then

$$\{d(x, y) : x, y \in X, x \neq y\}$$

is a finite set of positive values.

Therefore it has a minimum  $\epsilon > 0$ .

Since  $\{x_n\}$  is Cauchy, there exists  $N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < \epsilon$  for all  $m, n \geq N$ .

This implies  $x_n = x_N$  for all  $n \geq N$ .

Hence  $\lim_{n \rightarrow \infty} x_n = x_N$ . □

# Cauchy sequence

Proof.

**Case II:** Assume  $X$  is infinite.

# Cauchy sequence

Proof.

**Case II:** Assume  $X$  is infinite.

Then by the Bolzano-Weierstrass Theorem,  $X$  has an accumulation point  $L \in M$ .

# Cauchy sequence

Proof.

**Case II:** Assume  $X$  is infinite.

Then by the Bolzano-Weierstrass Theorem,  $X$  has an accumulation point  $L \in M$ .

Let  $\epsilon > 0$ .



# Cauchy sequence

Proof.

**Case II:** Assume  $X$  is infinite.

Then by the Bolzano-Weierstrass Theorem,  $X$  has an accumulation point  $L \in M$ .

Let  $\epsilon > 0$ .

Since  $\{x_n\}$  is Cauchy, we can choose  $N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < \epsilon/2$  for all  $m, n \geq N$ .

# Cauchy sequence

Proof.

**Case II:** Assume  $X$  is infinite.

Then by the Bolzano-Weierstrass Theorem,  $X$  has an accumulation point  $L \in M$ .

Let  $\epsilon > 0$ .

Since  $\{x_n\}$  is Cauchy, we can choose  $N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < \epsilon/2$  for all  $m, n \geq N$ .

Moreover, the ball  $B(L, \epsilon/2)$  contains infinitely many points of  $X$ , so we can choose  $\ell \geq N$  with  $x_\ell \in B(L, \epsilon/2)$ .

# Cauchy sequence

Proof.

**Case II:** Assume  $X$  is infinite.

Then by the Bolzano-Weierstrass Theorem,  $X$  has an accumulation point  $L \in M$ .

Let  $\epsilon > 0$ .

Since  $\{x_n\}$  is Cauchy, we can choose  $N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < \epsilon/2$  for all  $m, n \geq N$ .

Moreover, the ball  $B(L, \epsilon/2)$  contains infinitely many points of  $X$ , so we can choose  $\ell \geq N$  with  $x_\ell \in B(L, \epsilon/2)$ .

It follows that for all  $n \geq N$ ,

$$d(x_n, L) \leq d(x_n, x_\ell) + d(x_\ell, L) \leq \epsilon/2 + \epsilon/2 = \epsilon.$$

# Cauchy sequence

Proof.

**Case II:** Assume  $X$  is infinite.

Then by the Bolzano-Weierstrass Theorem,  $X$  has an accumulation point  $L \in M$ .

Let  $\epsilon > 0$ .

Since  $\{x_n\}$  is Cauchy, we can choose  $N \in \mathbb{Z}_+$  with  $d(x_m, x_n) < \epsilon/2$  for all  $m, n \geq N$ .

Moreover, the ball  $B(L, \epsilon/2)$  contains infinitely many points of  $X$ , so we can choose  $\ell \geq N$  with  $x_\ell \in B(L, \epsilon/2)$ .

It follows that for all  $n \geq N$ ,

$$d(x_n, L) \leq d(x_n, x_\ell) + d(x_\ell, L) \leq \epsilon/2 + \epsilon/2 = \epsilon.$$

Since  $\epsilon > 0$  was arbitrary, this proves  $\{x_n\}$  converges to  $L$ . □

# Example

Consider the metric space  $M = \mathbb{Q}$  with the Euclidean metric  $d$ .

# Example

Consider the metric space  $M = \mathbb{Q}$  with the Euclidean metric  $d$ .  
Consider the sequence  $\{x_n\}$  defined by  $x_1 = 1$  and

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n}, \quad n \geq 1.$$

# Example

Consider the metric space  $M = \mathbb{Q}$  with the Euclidean metric  $d$ .  
Consider the sequence  $\{x_n\}$  defined by  $x_1 = 1$  and

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n}, \quad n \geq 1.$$

$$x_1 = 1, \quad x_2 = \frac{3}{2}, \quad x_3 = \frac{17}{12}, \dots$$

# Example

Consider the metric space  $M = \mathbb{Q}$  with the Euclidean metric  $d$ .  
Consider the sequence  $\{x_n\}$  defined by  $x_1 = 1$  and

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n}, \quad n \geq 1.$$

$$x_1 = 1, \quad x_2 = \frac{3}{2}, \quad x_3 = \frac{17}{12}, \dots$$

- *tries* to converge to  $\sqrt{2}$



# Example

Consider the metric space  $M = \mathbb{Q}$  with the Euclidean metric  $d$ .  
Consider the sequence  $\{x_n\}$  defined by  $x_1 = 1$  and

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n}, \quad n \geq 1.$$

$$x_1 = 1, \quad x_2 = \frac{3}{2}, \quad x_3 = \frac{17}{12}, \dots$$

- *tries* to converge to  $\sqrt{2}$
- however  $\sqrt{2} \notin M$ , so it doesn't converge!

# Example

Consider the metric space  $M = \mathbb{Q}$  with the Euclidean metric  $d$ .  
Consider the sequence  $\{x_n\}$  defined by  $x_1 = 1$  and

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n}, \quad n \geq 1.$$

$$x_1 = 1, \quad x_2 = \frac{3}{2}, \quad x_3 = \frac{17}{12}, \dots$$

- *tries* to converge to  $\sqrt{2}$
- however  $\sqrt{2} \notin M$ , so it doesn't converge!
- somehow  $M$  is "missing" some points

# Complete spaces

## Definition

A metric space  $(M, d)$  is called **complete** if every Cauchy sequence in  $M$  converges.

# Complete spaces

## Definition

A metric space  $(M, d)$  is called **complete** if every Cauchy sequence in  $M$  converges.

A subset  $S \subseteq M$  is called **complete** if the subspace  $(S, d)$  is a complete metric space.

# Complete spaces

## Definition

A metric space  $(M, d)$  is called **complete** if every Cauchy sequence in  $M$  converges.

A subset  $S \subseteq M$  is called **complete** if the subspace  $(S, d)$  is a complete metric space.

- $\mathbb{R}^n$  with the Euclidean metric is complete

# Complete spaces

## Definition

A metric space  $(M, d)$  is called **complete** if every Cauchy sequence in  $M$  converges.

A subset  $S \subseteq M$  is called **complete** if the subspace  $(S, d)$  is a complete metric space.

- $\mathbb{R}^n$  with the Euclidean metric is complete
- $[0, 1]$  with the Euclidean metric is complete

# Complete spaces

## Definition

A metric space  $(M, d)$  is called **complete** if every Cauchy sequence in  $M$  converges.

A subset  $S \subseteq M$  is called **complete** if the subspace  $(S, d)$  is a complete metric space.

- $\mathbb{R}^n$  with the Euclidean metric is complete
- $[0, 1]$  with the Euclidean metric is complete
- $(0, 1)$  with the Euclidean metric is not complete

# Challenge!

## Problem

Show that the interval  $(0, 1) \subseteq \mathbb{R}$  with the Euclidean metric is not a complete metric space.



# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space.*

# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

Let  $(M, d)$  be a metric space and  $S \subseteq M$  compact.

# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

Let  $(M, d)$  be a metric space and  $S \subseteq M$  compact.  
To show that  $(S, d)$  is complete, we must show that every Cauchy sequence in  $S$  converges to a value in  $S$ .

# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

Let  $(M, d)$  be a metric space and  $S \subseteq M$  compact.  
To show that  $(S, d)$  is complete, we must show that every Cauchy sequence in  $S$  converges to a value in  $S$ .  
Suppose that  $\{x_n\}$  is a Cauchy sequence in  $(S, d)$ .

# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

Let  $(M, d)$  be a metric space and  $S \subseteq M$  compact.  
To show that  $(S, d)$  is complete, we must show that every Cauchy sequence in  $S$  converges to a value in  $S$ .  
Suppose that  $\{x_n\}$  is a Cauchy sequence in  $(S, d)$ .  
If the set  $X$  of all values of  $\{x_n\}$  is finite, then the Cauchy condition implies  $\{x_n\}$  converges.

# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

Let  $(M, d)$  be a metric space and  $S \subseteq M$  compact.

To show that  $(S, d)$  is complete, we must show that every Cauchy sequence in  $S$  converges to a value in  $S$ .

Suppose that  $\{x_n\}$  is a Cauchy sequence in  $(S, d)$ .

If the set  $X$  of all values of  $\{x_n\}$  is finite, then the Cauchy condition implies  $\{x_n\}$  converges.

Assume instead  $X$  is infinite.



# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

Let  $(M, d)$  be a metric space and  $S \subseteq M$  compact.

To show that  $(S, d)$  is complete, we must show that every Cauchy sequence in  $S$  converges to a value in  $S$ .

Suppose that  $\{x_n\}$  is a Cauchy sequence in  $(S, d)$ .

If the set  $X$  of all values of  $\{x_n\}$  is finite, then the Cauchy condition implies  $\{x_n\}$  converges.

Assume instead  $X$  is infinite.

TIME OUT!



# Complete spaces

## Lemma

*Let  $(M, d)$  be a metric space and let  $S \subseteq M$  be compact.*

# Complete spaces

## Lemma

*Let  $(M, d)$  be a metric space and let  $S \subseteq M$  be compact. Then for any infinite subset  $X \subseteq S$ , the set  $S$  has an accumulation point of  $X$ .*

# Complete spaces

## Lemma

*Let  $(M, d)$  be a metric space and let  $S \subseteq M$  be compact. Then for any infinite subset  $X \subseteq S$ , the set  $S$  has an accumulation point of  $X$ .*

## Proof.

# Complete spaces

## Lemma

*Let  $(M, d)$  be a metric space and let  $S \subseteq M$  be compact. Then for any infinite subset  $X \subseteq S$ , the set  $S$  has an accumulation point of  $X$ .*

## Proof.

To see this, suppose otherwise.

# Complete spaces

## Lemma

*Let  $(M, d)$  be a metric space and let  $S \subseteq M$  be compact. Then for any infinite subset  $X \subseteq S$ , the set  $S$  has an accumulation point of  $X$ .*

## Proof.

To see this, suppose otherwise.

Then for all  $x \in S$ , there exists  $r_x > 0$  such that  $B_S(x, r_x) \cap X$  is empty if  $x \notin X$  or equal to  $\{x\}$  if  $x \in X$ .

# Complete spaces

## Lemma

*Let  $(M, d)$  be a metric space and let  $S \subseteq M$  be compact. Then for any infinite subset  $X \subseteq S$ , the set  $S$  has an accumulation point of  $X$ .*

## Proof.

To see this, suppose otherwise.

Then for all  $x \in S$ , there exists  $r_x > 0$  such that  $B_S(x, r_x) \cap X$  is empty if  $x \notin X$  or equal to  $\{x\}$  if  $x \in X$ .

The open cover  $\{U_i : i \in I\}$  with  $I = S$  and  $U_i = B_S(i; r_i)$  of  $S$  has no finite subcover.

# Complete spaces

## Lemma

*Let  $(M, d)$  be a metric space and let  $S \subseteq M$  be compact. Then for any infinite subset  $X \subseteq S$ , the set  $S$  has an accumulation point of  $X$ .*

## Proof.

To see this, suppose otherwise.

Then for all  $x \in S$ , there exists  $r_x > 0$  such that  $B_S(x, r_x) \cap X$  is empty if  $x \notin X$  or equal to  $\{x\}$  if  $x \in X$ .

The open cover  $\{U_i : i \in I\}$  with  $I = S$  and  $U_i = B_S(i; r_i)$  of  $S$  has no finite subcover.

This is a contradiction, so  $S$  has an accumulation point of  $X$ . □



# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

Let  $L$  be an accumulation point of  $X$  in  $S$ .

# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

Let  $L$  be an accumulation point of  $X$  in  $S$ .

Let  $\epsilon > 0$ .

# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

Let  $L$  be an accumulation point of  $X$  in  $S$ .

Let  $\epsilon > 0$ .

Then  $\exists N > 0$  such that  $m, n \geq N$  implies  $d(x_m, x_n) < \epsilon/2$ .

# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

Let  $L$  be an accumulation point of  $X$  in  $S$ .

Let  $\epsilon > 0$ .

Then  $\exists N > 0$  such that  $m, n \geq N$  implies  $d(x_m, x_n) < \epsilon/2$ .

Moreover,  $B_S(L, \epsilon/2)$  contains infinitely many points of  $X$ .

# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

Let  $L$  be an accumulation point of  $X$  in  $S$ .

Let  $\epsilon > 0$ .

Then  $\exists N > 0$  such that  $m, n \geq N$  implies  $d(x_m, x_n) < \epsilon/2$ .

Moreover,  $B_S(L, \epsilon/2)$  contains infinitely many points of  $X$ .

Hence it contains  $x_m$  for some  $m \geq N$ .

# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

Let  $L$  be an accumulation point of  $X$  in  $S$ .

Let  $\epsilon > 0$ .

Then  $\exists N > 0$  such that  $m, n \geq N$  implies  $d(x_m, x_n) < \epsilon/2$ .

Moreover,  $B_S(L, \epsilon/2)$  contains infinitely many points of  $X$ .

Hence it contains  $x_m$  for some  $m \geq N$ .

It follows that for all  $n \geq N$ ,

$$d(x_n, L) \leq d(x_n, x_m) + d(x_m, L) < \epsilon/2 + \epsilon/2 = \epsilon.$$

# Complete spaces

## Theorem

*Let  $(M, d)$  be a metric space. Then for any compact subset  $S \subseteq M$ , the subspace  $(S, d)$  is complete.*

## Proof.

Let  $L$  be an accumulation point of  $X$  in  $S$ .

Let  $\epsilon > 0$ .

Then  $\exists N > 0$  such that  $m, n \geq N$  implies  $d(x_m, x_n) < \epsilon/2$ .

Moreover,  $B_S(L, \epsilon/2)$  contains infinitely many points of  $X$ .

Hence it contains  $x_m$  for some  $m \geq N$ .

It follows that for all  $n \geq N$ ,

$$d(x_n, L) \leq d(x_n, x_m) + d(x_m, L) < \epsilon/2 + \epsilon/2 = \epsilon.$$

Since  $\epsilon > 0$  was arbitrary, this proves  $\{x_n\}$  converges to  $L$ . □



# Outline

- 1 Real Analysis Lecture 14
  - Cauchy sequences
  - Limit of a function

# Limit of a function

Let  $(S, d_S)$  and  $(T, d_T)$  be metric spaces and  $A \subseteq S$ .

# Limit of a function

Let  $(S, d_S)$  and  $(T, d_T)$  be metric spaces and  $A \subseteq S$ . Also, let  $f : A \rightarrow T$  be a function.

# Limit of a function

Let  $(S, d_S)$  and  $(T, d_T)$  be metric spaces and  $A \subseteq S$ . Also, let  $f : A \rightarrow T$  be a function.

## Definition

# Limit of a function

Let  $(S, d_S)$  and  $(T, d_T)$  be metric spaces and  $A \subseteq S$ . Also, let  $f : A \rightarrow T$  be a function.

## Definition

Then for any accumulation point  $p$  of  $A$  and element  $L \in T$ , we say that the **limit as  $x$  approaches  $p$  of  $f(x)$  is  $L$**

# Limit of a function

Let  $(S, d_S)$  and  $(T, d_T)$  be metric spaces and  $A \subseteq S$ . Also, let  $f : A \rightarrow T$  be a function.

## Definition

Then for any accumulation point  $p$  of  $A$  and element  $L \in T$ , we say that the **limit as  $x$  approaches  $p$  of  $f(x)$  is  $L$**  and write

$$\lim_{x \rightarrow p} f(x) = L$$

# Limit of a function

Let  $(S, d_S)$  and  $(T, d_T)$  be metric spaces and  $A \subseteq S$ . Also, let  $f : A \rightarrow T$  be a function.

## Definition

Then for any accumulation point  $p$  of  $A$  and element  $L \in T$ , we say that the **limit as  $x$  approaches  $p$  of  $f(x)$  is  $L$**  and write

$$\lim_{x \rightarrow p} f(x) = L$$

if for all  $\epsilon > 0$  there exists  $\delta > 0$  such that

$$0 < d_S(x, p) < \delta \Rightarrow d_T(f(x), L) < \epsilon.$$

# Limit of a function

## Theorem



# Limit of a function

## Theorem

*Let  $(S, d_S)$  and  $(T, d_T)$  be metric spaces,  $A \subseteq S$ , and  $f : A \rightarrow T$  be a function.*

# Limit of a function

## Theorem

*Let  $(S, d_S)$  and  $(T, d_T)$  be metric spaces,  $A \subseteq S$ , and  $f : A \rightarrow T$  be a function. Then*

$$\lim_{x \rightarrow a} f(x) = L$$

# Limit of a function

## Theorem

*Let  $(S, d_S)$  and  $(T, d_T)$  be metric spaces,  $A \subseteq S$ , and  $f : A \rightarrow T$  be a function. Then*

$$\lim_{x \rightarrow a} f(x) = L$$

*if and only if*

$$\lim_{n \rightarrow \infty} f(x_n) = L$$

*for all sequences  $\{x_n\}$  of values in  $A \setminus \{a\}$  which converge to  $a$ .*