

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

October 15, 2024

Outline

- 1 Real Analysis Lecture 14
 - More on limits
 - Cauchy sequences

Outline

- 1 Real Analysis Lecture 14
 - More on limits
 - Cauchy sequences

Existence of limits

Theorem

If a sequence $\{x_n\}$ in a metric space (M, d) converges to a value $L \in M$, then the range

$$X = \{x_1, x_2, \dots\} \subseteq M$$

is a bounded set and L is an adherent point of X .

Existence of limits

Theorem

If a sequence $\{x_n\}$ in a metric space (M, d) converges to a value $L \in M$, then the range

$$X = \{x_1, x_2, \dots\} \subseteq M$$

is a bounded set and L is an adherent point of X .

When the range of a sequence is bounded, we call the sequence $\{x_n\}$ bounded.

Existence of limits

Proof.

Suppose that $\lim_{n \rightarrow \infty} x_n = L$.

Existence of limits

Proof.

Suppose that $\lim_{n \rightarrow \infty} x_n = L$.

Let $\epsilon = 2024$.

Existence of limits

Proof.

Suppose that $\lim_{n \rightarrow \infty} x_n = L$.

Let $\epsilon = 2024$.

Then there exists $N \in \mathbb{Z}_+$ with $d(x_n, L) \leq 2024$ for all $n \geq N$.

Existence of limits

Proof.

Suppose that $\lim_{n \rightarrow \infty} x_n = L$.

Let $\epsilon = 2024$.

Then there exists $N \in \mathbb{Z}_+$ with $d(x_n, L) \leq 2024$ for all $n \geq N$.

If we take

$$R = \max\{d(x_1, L), d(x_2, L), \dots, d(x_{N-1}, L), 2024\},$$

Existence of limits

Proof.

Suppose that $\lim_{n \rightarrow \infty} x_n = L$.

Let $\epsilon = 2024$.

Then there exists $N \in \mathbb{Z}_+$ with $d(x_n, L) \leq 2024$ for all $n \geq N$.

If we take

$$R = \max\{d(x_1, L), d(x_2, L), \dots, d(x_{N-1}, L), 2024\},$$

then $d(x_n, L) \leq R$ for all N .

Existence of limits

Proof.

Suppose that $\lim_{n \rightarrow \infty} x_n = L$.

Let $\epsilon = 2024$.

Then there exists $N \in \mathbb{Z}_+$ with $d(x_n, L) \leq 2024$ for all $n \geq N$.

If we take

$$R = \max\{d(x_1, L), d(x_2, L), \dots, d(x_{N-1}, L), 2024\},$$

then $d(x_n, L) \leq R$ for all N .

Therefore $X \subseteq B_M(L, R)$ and in particular X is bounded. □

Existence of limits

Sort of a converse of the previous theorem.

Existence of limits

Sort of a converse of the previous theorem.

Theorem

If S is a subset of a metric space (M, d) and $L \in M$ is an adherent point of S , then there exists a sequence $\{x_n\}$ of elements of S which converges to L .

Existence of limits

Proof.

Suppose L is an adherent point of S .

Existence of limits

Proof.

Suppose L is an adherent point of S .

Then for all $n \in \mathbb{Z}_+$, the ball $B_M(x, \frac{1}{n})$ contains at least one element of S .

Existence of limits

Proof.

Suppose L is an adherent point of S .

Then for all $n \in \mathbb{Z}_+$, the ball $B_M(x, \frac{1}{n})$ contains at least one element of S .

For each $n \in \mathbb{Z}_+$, choose $x_n \in S \cap B_M(x, \frac{1}{n})$.

Existence of limits

Proof.

Suppose L is an adherent point of S .

Then for all $n \in \mathbb{Z}_+$, the ball $B_M(x, \frac{1}{n})$ contains at least one element of S .

For each $n \in \mathbb{Z}_+$, choose $x_n \in S \cap B_M(x, \frac{1}{n})$.

We claim $\lim x_n = L$.

Existence of limits

Proof.

Suppose L is an adherent point of S .

Then for all $n \in \mathbb{Z}_+$, the ball $B_M(x, \frac{1}{n})$ contains at least one element of S .

For each $n \in \mathbb{Z}_+$, choose $x_n \in S \cap B_M(x, \frac{1}{n})$.

We claim $\lim x_n = L$.

To see this, let $\epsilon > 0$.

Existence of limits

Proof.

Suppose L is an adherent point of S .

Then for all $n \in \mathbb{Z}_+$, the ball $B_M(x, \frac{1}{n})$ contains at least one element of S .

For each $n \in \mathbb{Z}_+$, choose $x_n \in S \cap B_M(x, \frac{1}{n})$.

We claim $\lim x_n = L$.

To see this, let $\epsilon > 0$.

Then choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Existence of limits

Proof.

Suppose L is an adherent point of S .

Then for all $n \in \mathbb{Z}_+$, the ball $B_M(x, \frac{1}{n})$ contains at least one element of S .

For each $n \in \mathbb{Z}_+$, choose $x_n \in S \cap B_M(x, \frac{1}{n})$.

We claim $\lim x_n = L$.

To see this, let $\epsilon > 0$.

Then choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

It follows that for all $n \geq N$

$$d(x_n, L) < 1/n < 1/N < \epsilon.$$

Existence of limits

Proof.

Suppose L is an adherent point of S .

Then for all $n \in \mathbb{Z}_+$, the ball $B_M(x, \frac{1}{n})$ contains at least one element of S .

For each $n \in \mathbb{Z}_+$, choose $x_n \in S \cap B_M(x, \frac{1}{n})$.

We claim $\lim x_n = L$.

To see this, let $\epsilon > 0$.

Then choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

It follows that for all $n \geq N$

$$d(x_n, L) < 1/n < 1/N < \epsilon.$$

Since $\epsilon > 0$ was arbitrary, this proves convergence. □

Existence of limits

Recall a **subsequence** of a sequence $\{x_n\}$ is a sequence of the form $\{x_{k(n)}\}$, for $k(1) < k(2) < \dots$ positive integers.

Existence of limits

Recall a **subsequence** of a sequence $\{x_n\}$ is a sequence of the form $\{x_{k(n)}\}$, for $k(1) < k(2) < \dots$ positive integers.

Theorem

A sequence $\{x_n\}$ in a metric space (M, d) converges to a value $L \in M$ if and only if every subsequence $\{x_{k(n)}\}$ also converges to L .

Existence of limits

Proof.

Suppose that every subsequence $\{x_{k(n)}\}$ of $\{x_n\}$ converges to L .

Existence of limits

Proof.

Suppose that every subsequence $\{x_{k(n)}\}$ of $\{x_n\}$ converges to L .

A sequence is a subsequence of itself! Thus $\{x_n\}$ will converge to L .

Existence of limits

Proof.

Suppose that every subsequence $\{x_{k(n)}\}$ of $\{x_n\}$ converges to L .

A sequence is a subsequence of itself! Thus $\{x_n\}$ will converge to L .

Conversely, suppose that $\{x_n\}$ converges to L .

Existence of limits

Proof.

Suppose that every subsequence $\{x_{k(n)}\}$ of $\{x_n\}$ converges to L .

A sequence is a subsequence of itself! Thus $\{x_n\}$ will converge to L .

Conversely, suppose that $\{x_n\}$ converges to L .

Let $\{x_{k(n)}\}$ be a subsequence of $\{x_n\}$.

Existence of limits

Proof.

Suppose that every subsequence $\{x_{k(n)}\}$ of $\{x_n\}$ converges to L .

A sequence is a subsequence of itself! Thus $\{x_n\}$ will converge to L .

Conversely, suppose that $\{x_n\}$ converges to L .

Let $\{x_{k(n)}\}$ be a subsequence of $\{x_n\}$.

To show it limits to L , let $\epsilon > 0$.

Existence of limits

Proof.

Suppose that every subsequence $\{x_{k(n)}\}$ of $\{x_n\}$ converges to L .

A sequence is a subsequence of itself! Thus $\{x_n\}$ will converge to L .

Conversely, suppose that $\{x_n\}$ converges to L .

Let $\{x_{k(n)}\}$ be a subsequence of $\{x_n\}$.

To show it limits to L , let $\epsilon > 0$.

Then there exists $N \in \mathbb{Z}_+$ such that $n \geq N$ implies $d(x_n, L) < \epsilon$.

Existence of limits

Proof.

Suppose that every subsequence $\{x_{k(n)}\}$ of $\{x_n\}$ converges to L .

A sequence is a subsequence of itself! Thus $\{x_n\}$ will converge to L .

Conversely, suppose that $\{x_n\}$ converges to L .

Let $\{x_{k(n)}\}$ be a subsequence of $\{x_n\}$.

To show it limits to L , let $\epsilon > 0$.

Then there exists $N \in \mathbb{Z}_+$ such that $n \geq N$ implies $d(x_n, L) < \epsilon$.

Moreover, $k(n) \geq n$ so $k(n) \geq N$ and therefore $d(x_{k(n)}, L) < \epsilon$.

Existence of limits

Proof.

Suppose that every subsequence $\{x_{k(n)}\}$ of $\{x_n\}$ converges to L .

A sequence is a subsequence of itself! Thus $\{x_n\}$ will converge to L .

Conversely, suppose that $\{x_n\}$ converges to L .

Let $\{x_{k(n)}\}$ be a subsequence of $\{x_n\}$.

To show it limits to L , let $\epsilon > 0$.

Then there exists $N \in \mathbb{Z}_+$ such that $n \geq N$ implies $d(x_n, L) < \epsilon$.

Moreover, $k(n) \geq n$ so $k(n) \geq N$ and therefore $d(x_{k(n)}, L) < \epsilon$.

Since $\epsilon > 0$ was arbitrary, this proves convergence. \square

Monotone sequences

Definition

Monotone sequences

Definition

A sequence $\{x_n\}$ of real numbers is called **increasing** or **monotone increasing** if $x_n \leq x_{n+1}$ for all n .

Monotone sequences

Definition

A sequence $\{x_n\}$ of real numbers is called **increasing** or **monotone increasing** if $x_n \leq x_{n+1}$ for all n .

It is called **strictly increasing** if $x_n < x_{n+1}$ for all n .

Monotone sequences

Definition

A sequence $\{x_n\}$ of real numbers is called **increasing** or **monotone increasing** if $x_n \leq x_{n+1}$ for all n .

It is called **strictly increasing** if $x_n < x_{n+1}$ for all n .

A sequence $\{x_n\}$ of real numbers is called **decreasing** or **monotone decreasing** if $x_n \geq x_{n+1}$ for all n .

Monotone sequences

Definition

A sequence $\{x_n\}$ of real numbers is called **increasing** or **monotone increasing** if $x_n \leq x_{n+1}$ for all n .

It is called **strictly increasing** if $x_n < x_{n+1}$ for all n .

A sequence $\{x_n\}$ of real numbers is called **decreasing** or **monotone decreasing** if $x_n \geq x_{n+1}$ for all n .

It is called **strictly decreasing** if $x_n > x_{n+1}$ for all n .

Monotone sequences

Definition

A sequence $\{x_n\}$ of real numbers is called **increasing** or **monotone increasing** if $x_n \leq x_{n+1}$ for all n .

It is called **strictly increasing** if $x_n < x_{n+1}$ for all n .

A sequence $\{x_n\}$ of real numbers is called **decreasing** or **monotone decreasing** if $x_n \geq x_{n+1}$ for all n .

It is called **strictly decreasing** if $x_n > x_{n+1}$ for all n .

In any of these cases, we call the sequence **monotone**.

Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

Theorem (Monotone convergence theorem)

Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

Theorem (Monotone convergence theorem)

Consider the metric space formed by \mathbb{R} with Euclidean distance.

Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

Theorem (Monotone convergence theorem)

Consider the metric space formed by \mathbb{R} with Euclidean distance.

A monotone increasing sequence $\{x_n\}$ of real numbers which is bounded above converges to $L = \sup\{x_1, x_2, \dots\}$.

Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

Theorem (Monotone convergence theorem)

Consider the metric space formed by \mathbb{R} with Euclidean distance.

A monotone increasing sequence $\{x_n\}$ of real numbers which is bounded above converges to $L = \sup\{x_1, x_2, \dots\}$.

A monotone decreasing sequence $\{x_n\}$ which is bounded below converges to $L = \inf\{x_1, x_2, \dots\}$.

Monotone sequences

Proof.

Monotone sequences

Proof.

Suppose that $\{x_n\}$ is a monotone increasing sequence which is bounded above and let $X = \{x_1, x_2, \dots\}$.

Monotone sequences

Proof.

Suppose that $\{x_n\}$ is a monotone increasing sequence which is bounded above and let $X = \{x_1, x_2, \dots\}$. Then by the Completeness Axiom $L = \sup X$ exists.

Monotone sequences

Proof.

Suppose that $\{x_n\}$ is a monotone increasing sequence which is bounded above and let $X = \{x_1, x_2, \dots\}$. Then by the Completeness Axiom $L = \sup X$ exists.

Let $\epsilon > 0$.

Monotone sequences

Proof.

Suppose that $\{x_n\}$ is a monotone increasing sequence which is bounded above and let $X = \{x_1, x_2, \dots\}$. Then by the Completeness Axiom $L = \sup X$ exists.

Let $\epsilon > 0$.

Then $L - \epsilon$ is not a supremum of X , so there exists $x_N \in X$ such that $x_N > L - \epsilon$.

Monotone sequences

Proof.

Suppose that $\{x_n\}$ is a monotone increasing sequence which is bounded above and let $X = \{x_1, x_2, \dots\}$. Then by the Completeness Axiom $L = \sup X$ exists.

Let $\epsilon > 0$.

Then $L - \epsilon$ is not a supremum of X , so there exists $x_N \in X$ such that $x_N > L - \epsilon$.

Since the sequence is increasing, for any $n > N$, we have that $L - \epsilon < x_N \leq x_n < L$.

Monotone sequences

Proof.

Suppose that $\{x_n\}$ is a monotone increasing sequence which is bounded above and let $X = \{x_1, x_2, \dots\}$. Then by the Completeness Axiom $L = \sup X$ exists.

Let $\epsilon > 0$.

Then $L - \epsilon$ is not a supremum of X , so there exists $x_N \in X$ such that $x_N > L - \epsilon$.

Since the sequence is increasing, for any $n > N$, we have that $L - \epsilon < x_N \leq x_n < L$.

It follows that

$$d(x_n, L) = |x_n - L| = L - x_n < \epsilon.$$

Monotone sequences

Proof.

Suppose that $\{x_n\}$ is a monotone increasing sequence which is bounded above and let $X = \{x_1, x_2, \dots\}$. Then by the Completeness Axiom $L = \sup X$ exists.

Let $\epsilon > 0$.

Then $L - \epsilon$ is not a supremum of X , so there exists $x_N \in X$ such that $x_N > L - \epsilon$.

Since the sequence is increasing, for any $n > N$, we have that $L - \epsilon < x_N \leq x_n < L$.

It follows that

$$d(x_n, L) = |x_n - L| = L - x_n < \epsilon.$$

Since $\epsilon > 0$ was arbitrary, this proves x_n converges to L . □

Outline

- 1 Real Analysis Lecture 14
 - More on limits
 - Cauchy sequences

Cauchy sequence

A sequence $\{x_n\}$ in a metric space (M, d) is called Cauchy if the terms get arbitrarily close together as n gets arbitrarily large.

Cauchy sequence

A sequence $\{x_n\}$ in a metric space (M, d) is called Cauchy if the terms get arbitrarily close together as n gets arbitrarily large.

More formally:

Cauchy sequence

A sequence $\{x_n\}$ in a metric space (M, d) is called Cauchy if the terms get arbitrarily close together as n gets arbitrarily large.

More formally:

Definition

A sequence $\{x_n\}$ is called a **Cauchy sequence** if

$$\forall \epsilon > 0, \text{ there exists } N \in \mathbb{Z}_+ \text{ with } m, n \geq N \Rightarrow d(x_m, x_n) < \epsilon.$$

Cauchy sequence

A sequence $\{x_n\}$ in a metric space (M, d) is called Cauchy if the terms get arbitrarily close together as n gets arbitrarily large.

More formally:

Definition

A sequence $\{x_n\}$ is called a **Cauchy sequence** if

$$\forall \epsilon > 0, \text{ there exists } N \in \mathbb{Z}_+ \text{ with } m, n \geq N \Rightarrow d(x_m, x_n) < \epsilon.$$

- every convergent sequence is Cauchy

Cauchy sequence

A sequence $\{x_n\}$ in a metric space (M, d) is called Cauchy if the terms get arbitrarily close together as n gets arbitrarily large.

More formally:

Definition

A sequence $\{x_n\}$ is called a **Cauchy sequence** if

$$\forall \epsilon > 0, \text{ there exists } N \in \mathbb{Z}_+ \text{ with } m, n \geq N \Rightarrow d(x_m, x_n) < \epsilon.$$

- every convergent sequence is Cauchy
- is every Cauchy sequence convergent?

Challenge!

Problem

Prove that if $\{x_n\}$ is a Cauchy sequence in a metric space (M, d) , then it is bounded.

Challenge!

Problem

Prove that if $\{x_n\}$ is a Cauchy sequence in a metric space (M, d) , then it is bounded.

Solution

Challenge!

Problem

Prove that if $\{x_n\}$ is a Cauchy sequence in a metric space (M, d) , then it is bounded.

Solution

Suppose $\{x_n\}$ is Cauchy.

Challenge!

Problem

Prove that if $\{x_n\}$ is a Cauchy sequence in a metric space (M, d) , then it is bounded.

Solution

Suppose $\{x_n\}$ is Cauchy.
Let $\epsilon = 2024$.

Challenge!

Problem

Prove that if $\{x_n\}$ is a Cauchy sequence in a metric space (M, d) , then it is bounded.

Solution

Suppose $\{x_n\}$ is Cauchy.

Let $\epsilon = 2024$.

Then $\exists N \in \mathbb{Z}_+$ with $d(x_m, x_n) < 2024$ for all $m, n \geq N$.

Challenge!

Problem

Prove that if $\{x_n\}$ is a Cauchy sequence in a metric space (M, d) , then it is bounded.

Solution

Suppose $\{x_n\}$ is Cauchy.

Let $\epsilon = 2024$.

Then $\exists N \in \mathbb{Z}_+$ with $d(x_m, x_n) < 2024$ for all $m, n \geq N$.

Therefore for all $n \geq N$, we have $d(x_n, x_N) \leq 2024$.

Challenge!

Problem

Prove that if $\{x_n\}$ is a Cauchy sequence in a metric space (M, d) , then it is bounded.

Solution

Suppose $\{x_n\}$ is Cauchy.

Let $\epsilon = 2024$.

Then $\exists N \in \mathbb{Z}_+$ with $d(x_m, x_n) < 2024$ for all $m, n \geq N$.

Therefore for all $n \geq N$, we have $d(x_n, x_N) \leq 2024$.

So if we define

$$R = \max\{d(x_1, x_N), d(x_2, x_N), \dots, d(x_{n-1}, x_N), 2024\},$$

Challenge!

Problem

Prove that if $\{x_n\}$ is a Cauchy sequence in a metric space (M, d) , then it is bounded.

Solution

Suppose $\{x_n\}$ is Cauchy.

Let $\epsilon = 2024$.

Then $\exists N \in \mathbb{Z}_+$ with $d(x_m, x_n) < 2024$ for all $m, n \geq N$.

Therefore for all $n \geq N$, we have $d(x_n, x_N) \leq 2024$.

So if we define

$$R = \max\{d(x_1, x_N), d(x_2, x_N), \dots, d(x_{n-1}, x_N), 2024\},$$

then $d(x_n, x_N) \leq 2024$ for all $n \geq 1$.

Challenge!

Problem

Prove that if $\{x_n\}$ is a Cauchy sequence in a metric space (M, d) , then it is bounded.

Solution

Suppose $\{x_n\}$ is Cauchy.

Let $\epsilon = 2024$.

Then $\exists N \in \mathbb{Z}_+$ with $d(x_m, x_n) < 2024$ for all $m, n \geq N$.

Therefore for all $n \geq N$, we have $d(x_n, x_N) \leq 2024$.

So if we define

$$R = \max\{d(x_1, x_N), d(x_2, x_N), \dots, d(x_{n-1}, x_N), 2024\},$$

then $d(x_n, x_N) \leq 2024$ for all $n \geq 1$.

Hence $X = \{x_1, x_2, \dots\} \subseteq B_M(x_N, R)$, and is bounded.

Cauchy sequence

Every convergent sequence is Cauchy.

Cauchy sequence

Every convergent sequence is Cauchy.

Theorem

Let (M, d) be a metric space and suppose that $\{x_n\}$ is a sequence in M which converges to $L \in M$.

Then $\{x_n\}$ is Cauchy.

Cauchy sequence

Every convergent sequence is Cauchy.

Theorem

Let (M, d) be a metric space and suppose that $\{x_n\}$ is a sequence in M which converges to $L \in M$.

Then $\{x_n\}$ is Cauchy.

Proof.

Cauchy sequence

Every convergent sequence is Cauchy.

Theorem

*Let (M, d) be a metric space and suppose that $\{x_n\}$ is a sequence in M which converges to $L \in M$.
Then $\{x_n\}$ is Cauchy.*

Proof.

Let $\epsilon > 0$.

Cauchy sequence

Every convergent sequence is Cauchy.

Theorem

Let (M, d) be a metric space and suppose that $\{x_n\}$ is a sequence in M which converges to $L \in M$.

Then $\{x_n\}$ is Cauchy.

Proof.

Let $\epsilon > 0$.

Then there exists N such that $n \geq N$ implies $d(x_n, L) < \epsilon/2$.

Cauchy sequence

Every convergent sequence is Cauchy.

Theorem

Let (M, d) be a metric space and suppose that $\{x_n\}$ is a sequence in M which converges to $L \in M$.

Then $\{x_n\}$ is Cauchy.

Proof.

Let $\epsilon > 0$.

Then there exists N such that $n \geq N$ implies $d(x_n, L) < \epsilon/2$.

Therefore for any $m, n \geq N$ we have

$$d(x_m, x_n) \leq d(x_m, L) + d(L, x_n) < \epsilon/2 + \epsilon/2 = \epsilon.$$

Since $\epsilon > 0$ was arbitrary, this proves that $\{x_n\}$ is Cauchy. □

Cauchy sequence

Partial converse.

Cauchy sequence

Partial converse.

Theorem

Let $M = \mathbb{R}^n$ with the Euclidean metric, and suppose that $\{x_n\}$ is a Cauchy sequence in M .

Then $\{x_n\}$ converges.

Cauchy sequence

Partial converse.

Theorem

Let $M = \mathbb{R}^n$ with the Euclidean metric, and suppose that $\{x_n\}$ is a Cauchy sequence in M .

Then $\{x_n\}$ converges.

Proof.

Cauchy sequence

Partial converse.

Theorem

Let $M = \mathbb{R}^n$ with the Euclidean metric, and suppose that $\{x_n\}$ is a Cauchy sequence in M .

Then $\{x_n\}$ converges.

Proof.

Consider the range $X = \{x_1, x_2, \dots\}$ of $\{x_n\}$.

Cauchy sequence

Partial converse.

Theorem

Let $M = \mathbb{R}^n$ with the Euclidean metric, and suppose that $\{x_n\}$ is a Cauchy sequence in M .

Then $\{x_n\}$ converges.

Proof.

Consider the range $X = \{x_1, x_2, \dots\}$ of $\{x_n\}$.

Since $\{x_n\}$ is Cauchy, X is bounded.

Cauchy sequence

Partial converse.

Theorem

Let $M = \mathbb{R}^n$ with the Euclidean metric, and suppose that $\{x_n\}$ is a Cauchy sequence in M .

Then $\{x_n\}$ converges.

Proof.

Consider the range $X = \{x_1, x_2, \dots\}$ of $\{x_n\}$.

Since $\{x_n\}$ is Cauchy, X is bounded.

We consider two cases:

Cauchy sequence

Partial converse.

Theorem

Let $M = \mathbb{R}^n$ with the Euclidean metric, and suppose that $\{x_n\}$ is a Cauchy sequence in M .

Then $\{x_n\}$ converges.

Proof.

Consider the range $X = \{x_1, x_2, \dots\}$ of $\{x_n\}$.

Since $\{x_n\}$ is Cauchy, X is bounded.

We consider two cases:

- when X is finite

Cauchy sequence

Partial converse.

Theorem

Let $M = \mathbb{R}^n$ with the Euclidean metric, and suppose that $\{x_n\}$ is a Cauchy sequence in M .

Then $\{x_n\}$ converges.

Proof.

Consider the range $X = \{x_1, x_2, \dots\}$ of $\{x_n\}$.

Since $\{x_n\}$ is Cauchy, X is bounded.

We consider two cases:

- when X is finite
- when X is infinite



Cauchy sequence

Proof.

Case I: Assume X is finite.

Cauchy sequence

Proof.

Case I: Assume X is finite.

Then

$$\{d(x, y) : x, y \in X, x \neq y\}$$

is a finite set of positive values.

Cauchy sequence

Proof.

Case I: Assume X is finite.

Then

$$\{d(x, y) : x, y \in X, x \neq y\}$$

is a finite set of positive values.

Therefore it has a minimum $\epsilon > 0$.

Cauchy sequence

Proof.

Case I: Assume X is finite.

Then

$$\{d(x, y) : x, y \in X, x \neq y\}$$

is a finite set of positive values.

Therefore it has a minimum $\epsilon > 0$.

Since $\{x_n\}$ is Cauchy, there exists $N \in \mathbb{Z}_+$ with $d(x_m, x_n) < \epsilon$ for all $m, n \geq N$.

Cauchy sequence

Proof.

Case I: Assume X is finite.

Then

$$\{d(x, y) : x, y \in X, x \neq y\}$$

is a finite set of positive values.

Therefore it has a minimum $\epsilon > 0$.

Since $\{x_n\}$ is Cauchy, there exists $N \in \mathbb{Z}_+$ with $d(x_m, x_n) < \epsilon$ for all $m, n \geq N$.

This implies $x_n = x_N$ for all $n \geq N$.

Cauchy sequence

Proof.

Case I: Assume X is finite.

Then

$$\{d(x, y) : x, y \in X, x \neq y\}$$

is a finite set of positive values.

Therefore it has a minimum $\epsilon > 0$.

Since $\{x_n\}$ is Cauchy, there exists $N \in \mathbb{Z}_+$ with $d(x_m, x_n) < \epsilon$ for all $m, n \geq N$.

This implies $x_n = x_N$ for all $n \geq N$.

Hence $\lim_{n \rightarrow \infty} x_n = x_N$. □

Cauchy sequence

Proof.

Case II: Assume X is infinite.

Cauchy sequence

Proof.

Case II: Assume X is infinite.

Then by the Bolzano-Weierstrass Theorem, X has an accumulation point $L \in M$.

Cauchy sequence

Proof.

Case II: Assume X is infinite.

Then by the Bolzano-Weierstrass Theorem, X has an accumulation point $L \in M$.

Let $\epsilon > 0$.

Cauchy sequence

Proof.

Case II: Assume X is infinite.

Then by the Bolzano-Weierstrass Theorem, X has an accumulation point $L \in M$.

Let $\epsilon > 0$.

Since $\{x_n\}$ is Cauchy, we can choose $N \in \mathbb{Z}_+$ with $d(x_m, x_n) < \epsilon/2$ for all $m, n \geq N$.

Cauchy sequence

Proof.

Case II: Assume X is infinite.

Then by the Bolzano-Weierstrass Theorem, X has an accumulation point $L \in M$.

Let $\epsilon > 0$.

Since $\{x_n\}$ is Cauchy, we can choose $N \in \mathbb{Z}_+$ with $d(x_m, x_n) < \epsilon/2$ for all $m, n \geq N$.

Moreover, the ball $B(L, \epsilon/2)$ contains infinitely many points of X , so we can choose $\ell \geq N$ with $x_\ell \in B(L, \epsilon/2)$.

Cauchy sequence

Proof.

Case II: Assume X is infinite.

Then by the Bolzano-Weierstrass Theorem, X has an accumulation point $L \in M$.

Let $\epsilon > 0$.

Since $\{x_n\}$ is Cauchy, we can choose $N \in \mathbb{Z}_+$ with $d(x_m, x_n) < \epsilon/2$ for all $m, n \geq N$.

Moreover, the ball $B(L, \epsilon/2)$ contains infinitely many points of X , so we can choose $\ell \geq N$ with $x_\ell \in B(L, \epsilon/2)$.

It follows that for all $n \geq N$,

$$d(x_n, L) \leq d(x_n, x_\ell) + d(x_\ell, L) \leq \epsilon/2 + \epsilon/2 = \epsilon.$$

Cauchy sequence

Proof.

Case II: Assume X is infinite.

Then by the Bolzano-Weierstrass Theorem, X has an accumulation point $L \in M$.

Let $\epsilon > 0$.

Since $\{x_n\}$ is Cauchy, we can choose $N \in \mathbb{Z}_+$ with $d(x_m, x_n) < \epsilon/2$ for all $m, n \geq N$.

Moreover, the ball $B(L, \epsilon/2)$ contains infinitely many points of X , so we can choose $\ell \geq N$ with $x_\ell \in B(L, \epsilon/2)$.

It follows that for all $n \geq N$,

$$d(x_n, L) \leq d(x_n, x_\ell) + d(x_\ell, L) \leq \epsilon/2 + \epsilon/2 = \epsilon.$$

Since $\epsilon > 0$ was arbitrary, this proves $\{x_n\}$ converges to L . □

Example

Consider the metric space $M = \mathbb{Q}$ with the Euclidean metric d .

Example

Consider the metric space $M = \mathbb{Q}$ with the Euclidean metric d .
Consider the sequence $\{x_n\}$ defined by $x_1 = 1$ and

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n}, \quad n \geq 1.$$

Example

Consider the metric space $M = \mathbb{Q}$ with the Euclidean metric d .
Consider the sequence $\{x_n\}$ defined by $x_1 = 1$ and

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n}, \quad n \geq 1.$$

$$x_1 = 1, \quad x_2 = \frac{3}{2}, \quad x_3 = \frac{17}{12}, \dots$$

Example

Consider the metric space $M = \mathbb{Q}$ with the Euclidean metric d .
Consider the sequence $\{x_n\}$ defined by $x_1 = 1$ and

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n}, \quad n \geq 1.$$

$$x_1 = 1, \quad x_2 = \frac{3}{2}, \quad x_3 = \frac{17}{12}, \dots$$

- *tries* to converge to $\sqrt{2}$

Example

Consider the metric space $M = \mathbb{Q}$ with the Euclidean metric d .
Consider the sequence $\{x_n\}$ defined by $x_1 = 1$ and

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n}, \quad n \geq 1.$$

$$x_1 = 1, \quad x_2 = \frac{3}{2}, \quad x_3 = \frac{17}{12}, \dots$$

- *tries* to converge to $\sqrt{2}$
- however $\sqrt{2} \notin M$, so it doesn't converge!

Example

Consider the metric space $M = \mathbb{Q}$ with the Euclidean metric d .
Consider the sequence $\{x_n\}$ defined by $x_1 = 1$ and

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n}, \quad n \geq 1.$$

$$x_1 = 1, \quad x_2 = \frac{3}{2}, \quad x_3 = \frac{17}{12}, \dots$$

- *tries* to converge to $\sqrt{2}$
- however $\sqrt{2} \notin M$, so it doesn't converge!
- somehow M is "missing" some points

Complete spaces

Definition

A metric space (M, d) is called **complete** if every Cauchy sequence in M converges.

Complete spaces

Definition

A metric space (M, d) is called **complete** if every Cauchy sequence in M converges.

A subset $S \subseteq M$ is called **complete** if the subspace (S, d) is a complete metric space.

Complete spaces

Definition

A metric space (M, d) is called **complete** if every Cauchy sequence in M converges.

A subset $S \subseteq M$ is called **complete** if the subspace (S, d) is a complete metric space.

- \mathbb{R}^n with the Euclidean metric is complete

Complete spaces

Definition

A metric space (M, d) is called **complete** if every Cauchy sequence in M converges.

A subset $S \subseteq M$ is called **complete** if the subspace (S, d) is a complete metric space.

- \mathbb{R}^n with the Euclidean metric is complete
- $[0, 1]$ with the Euclidean metric is complete

Complete spaces

Definition

A metric space (M, d) is called **complete** if every Cauchy sequence in M converges.

A subset $S \subseteq M$ is called **complete** if the subspace (S, d) is a complete metric space.

- \mathbb{R}^n with the Euclidean metric is complete
- $[0, 1]$ with the Euclidean metric is complete
- $(0, 1)$ with the Euclidean metric is not complete

Challenge!

Problem

Show that the interval $(0, 1) \subseteq \mathbb{R}$ with the Euclidean metric is not a complete metric space.