

# MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics  
California State University Fullerton

September 16, 2024

# Outline

- 1 Real Analysis Lecture 6
  - Functions
  - Cardinality

# Outline

- 1 Real Analysis Lecture 6
  - Functions
  - Cardinality

# Functions

A **function** from  $A$  to  $B$  is a relation  $\mathcal{R}$  from  $A$  to  $B$  with the property

# Functions

A **function** from  $A$  to  $B$  is a relation  $\mathcal{R}$  from  $A$  to  $B$  with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \quad \text{implies} \quad b = c.$$

# Functions

A **function** from  $A$  to  $B$  is a relation  $\mathcal{R}$  from  $A$  to  $B$  with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

If a relation  $\mathcal{R}$  is a function, we usually use a symbol like  $f$ .

# Functions

A **function** from  $A$  to  $B$  is a relation  $\mathcal{R}$  from  $A$  to  $B$  with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

If a relation  $\mathcal{R}$  is a function, we usually use a symbol like  $f$ .

**NOTATION:**  $f : A \rightarrow B$  means  $f$  is a function from  $A$  to  $B$

**NOTATION:**  $f(a) = b$  means  $(a, b) \in f$ .

# Functions

A **function** from  $A$  to  $B$  is a relation  $\mathcal{R}$  from  $A$  to  $B$  with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

If a relation  $\mathcal{R}$  is a function, we usually use a symbol like  $f$ .

**NOTATION:**  $f : A \rightarrow B$  means  $f$  is a function from  $A$  to  $B$

**NOTATION:**  $f(a) = b$  means  $(a, b) \in f$ .

The set

$$\text{img}(f) = \{f(a) : a \in A\}$$

is called the **range** or **image** of  $f$



# Challenge!

## Problem

Determine all the equivalence relations on  $\mathbb{R}$  which are also functions.

# Challenge!

## Problem

Determine all the equivalence relations on  $\mathbb{R}$  which are also functions.

## Solution

$f = \mathcal{R}$  must be reflexive, so  $x\mathcal{R}x$  for all  $x$

# Challenge!

## Problem

Determine all the equivalence relations on  $\mathbb{R}$  which are also functions.

## Solution

$f = \mathcal{R}$  must be reflexive, so  $x\mathcal{R}x$  for all  $x$   
This means  $f(x) = x$  for all  $x$ .

# Challenge!

## Problem

Determine all the equivalence relations on  $\mathbb{R}$  which are also functions.

## Solution

$f = \mathcal{R}$  must be reflexive, so  $x\mathcal{R}x$  for all  $x$

This means  $f(x) = x$  for all  $x$ .

Thus the only function which is an equivalence relation is the identity function

$$f(x) = x.$$

# Properties of functions

A function  $f$  from  $A$  to  $B$  is called **one-to-one** or **injective** if

# Properties of functions

A function  $f$  from  $A$  to  $B$  is called **one-to-one** or **injective** if

$$f(x) = f(y) \text{ implies } x = y \text{ for all } x, y \in A.$$

# Properties of functions

A function  $f$  from  $A$  to  $B$  is called **one-to-one** or **injective** if

$$f(x) = f(y) \text{ implies } x = y \text{ for all } x, y \in A.$$

It is called **onto** or **surjective** if  $\text{img}(f) = B$ , or equivalently

$$\text{for all } b \in B \text{ there exists } a \in A \text{ with } f(a) = b.$$

# Properties of functions

A function  $f$  from  $A$  to  $B$  is called **one-to-one** or **injective** if

$$f(x) = f(y) \text{ implies } x = y \text{ for all } x, y \in A.$$

It is called **onto** or **surjective** if  $\text{img}(f) = B$ , or equivalently

$$\text{for all } b \in B \text{ there exists } a \in A \text{ with } f(a) = b.$$

If it satisfies both properties, it is called **bijective**.



# Inverses

The **converse** of a relation  $\mathcal{R}$  from  $A$  to  $B$  is the relation

# Inverses

The **converse** of a relation  $\mathcal{R}$  from  $A$  to  $B$  is the relation

$$\check{\mathcal{R}} = \{(b, a) \in B \times A : a\mathcal{R}b\}.$$

# Inverses

The **converse** of a relation  $\mathcal{R}$  from  $A$  to  $B$  is the relation

$$\check{\mathcal{R}} = \{(b, a) \in B \times A : a\mathcal{R}b\}.$$

If  $f$  is function, then  $\check{f}$  may or not be a function.

# Inverses

The **converse** of a relation  $\mathcal{R}$  from  $A$  to  $B$  is the relation

$$\check{\mathcal{R}} = \{(b, a) \in B \times A : a\mathcal{R}b\}.$$

If  $f$  is function, then  $\check{f}$  may or not be a function.  
If it is, we call it the **inverse** of  $f$ .

# Compositions

The composition of  $f : A \rightarrow B$  and  $g : B \rightarrow C$  is the function

# Compositions

The composition of  $f : A \rightarrow B$  and  $g : B \rightarrow C$  is the function

$$g \circ f : A \rightarrow C$$

# Compositions

The composition of  $f : A \rightarrow B$  and  $g : B \rightarrow C$  is the function

$$g \circ f : A \rightarrow C$$

with domain  $A$  and codomain  $C$  defined by

# Compositions

The composition of  $f : A \rightarrow B$  and  $g : B \rightarrow C$  is the function

$$g \circ f : A \rightarrow C$$

with domain  $A$  and codomain  $C$  defined by

$$(g \circ f)(x) = g(f(x)).$$



# Sequences

A **finite sequence** is a function  $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$ .

# Sequences

A **finite sequence** is a function  $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$ .

An **infinite sequence** is a function  $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$ .

# Sequences

A **finite sequence** is a function  $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$ .

An **infinite sequence** is a function  $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$ .

**NOTATION:**  $f_n$  indicates the value  $f(n)$

# Sequences

A **finite sequence** is a function  $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$ .

An **infinite sequence** is a function  $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$ .

**NOTATION:**  $f_n$  indicates the value  $f(n)$

**NOTATION:**  $\{f_n\}$  is another way of writing the function  $f$

# Sequences

A **finite sequence** is a function  $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$ .

An **infinite sequence** is a function  $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$ .

**NOTATION:**  $f_n$  indicates the value  $f(n)$

**NOTATION:**  $\{f_n\}$  is another way of writing the function  $f$

If  $k : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$  is a function which is **strictly increasing**,  
meaning

$$m < n \quad \Rightarrow \quad k(m) < k(n),$$

# Sequences

A **finite sequence** is a function  $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$ .

An **infinite sequence** is a function  $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$ .

**NOTATION:**  $f_n$  indicates the value  $f(n)$

**NOTATION:**  $\{f_n\}$  is another way of writing the function  $f$

If  $k : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$  is a function which is **strictly increasing**,  
meaning

$$m < n \Rightarrow k(m) < k(n),$$

then the composition  $f \circ k : \mathbb{Z}_+ \rightarrow \mathbb{R}$  forms a sequence called a  
**subsequence** of  $f$ .

# Sequences

A **finite sequence** is a function  $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$ .

An **infinite sequence** is a function  $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$ .

**NOTATION:**  $f_n$  indicates the value  $f(n)$

**NOTATION:**  $\{f_n\}$  is another way of writing the function  $f$

If  $k : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$  is a function which is **strictly increasing**, meaning

$$m < n \Rightarrow k(m) < k(n),$$

then the composition  $f \circ k : \mathbb{Z}_+ \rightarrow \mathbb{R}$  forms a sequence called a **subsequence** of  $f$ .

**NOTATION:**  $\{f_{k(n)}\}$  or  $\{f_{k_n}\}$  both really mean  $f \circ k$

# Outline

- 1 Real Analysis Lecture 6
  - Functions
  - Cardinality



# Cantor's Paradise

Philosophy: sets are the same size, or **cardinality**, if there is a bijection between them.

# Cantor's Paradise

Philosophy: sets are the same size, or **cardinality**, if there is a bijection between them.

- $|A|$  denotes the cardinality of  $A$

# Cantor's Paradise

Philosophy: sets are the same size, or **cardinality**, if there is a bijection between them.

- $|A|$  denotes the cardinality of  $A$
- $|A| \leq |B|$  means there is an injection  $f : A \rightarrow B$

# Cantor's Paradise

Philosophy: sets are the same size, or **cardinality**, if there is a bijection between them.

- $|A|$  denotes the cardinality of  $A$
- $|A| \leq |B|$  means there is an injection  $f : A \rightarrow B$
- $|A| \geq |B|$  means there is a surjection  $f : A \rightarrow B$

# Cantor's Paradise

Philosophy: sets are the same size, or **cardinality**, if there is a bijection between them.

- $|A|$  denotes the cardinality of  $A$
- $|A| \leq |B|$  means there is an injection  $f : A \rightarrow B$
- $|A| \geq |B|$  means there is a surjection  $f : A \rightarrow B$
- $|A| = |B|$  means there is a bijection  $f : A \rightarrow B$

# Cantor's Paradise

Philosophy: sets are the same size, or **cardinality**, if there is a bijection between them.

- $|A|$  denotes the cardinality of  $A$
- $|A| \leq |B|$  means there is an injection  $f : A \rightarrow B$
- $|A| \geq |B|$  means there is a surjection  $f : A \rightarrow B$
- $|A| = |B|$  means there is a bijection  $f : A \rightarrow B$

# Cantor's Paradise

## Theorem (Cantor-Schroeder-Bernstein Theorem)

*The following are equivalent*

- (i)  $|A| \leq |B|$  and  $|B| \leq |A|$
- (ii)  $|A| \geq |B|$  and  $|B| \geq |A|$
- (iii)  $|A| \leq |B|$  and  $|A| \geq |A|$
- (iv)  $|A| = |B|$

# Challenge

## Problem

Show that  $\mathbb{Z}_+$  and  $\mathbb{Z}_+ \times \mathbb{Z}_+$  have the same cardinality.



# Challenge

## Problem

Show that  $\mathbb{Z}_+$  and  $\mathbb{Z}_+ \times \mathbb{Z}_+$  have the same cardinality.

Hint: consider  $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ ,  $f(m, n) = 2^m 3^n$

## Solution

# Challenge

## Problem

Show that  $\mathbb{Z}_+$  and  $\mathbb{Z}_+ \times \mathbb{Z}_+$  have the same cardinality.

Hint: consider  $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ ,  $f(m, n) = 2^m 3^n$

## Solution

$2^a 3^b = 2^c 3^d$  if and only if  $a = c$  and  $b = d$  (Fund. Thm. Arith.)

# Challenge

## Problem

Show that  $\mathbb{Z}_+$  and  $\mathbb{Z}_+ \times \mathbb{Z}_+$  have the same cardinality.

Hint: consider  $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ ,  $f(m, n) = 2^m 3^n$

## Solution

$2^a 3^b = 2^c 3^d$  if and only if  $a = c$  and  $b = d$  (Fund. Thm. Arith.)  
 $f(m, n) = 2^m 3^n$  is injective

# Challenge

## Problem

Show that  $\mathbb{Z}_+$  and  $\mathbb{Z}_+ \times \mathbb{Z}_+$  have the same cardinality.

Hint: consider  $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ ,  $f(m, n) = 2^m 3^n$

## Solution

$2^a 3^b = 2^c 3^d$  if and only if  $a = c$  and  $b = d$  (Fund. Thm. Arith.)

$f(m, n) = 2^m 3^n$  is injective

$$|\mathbb{Z}_+ \times \mathbb{Z}_+| \leq |\mathbb{Z}_+|$$

# Challenge

## Problem

Show that  $\mathbb{Z}_+$  and  $\mathbb{Z}_+ \times \mathbb{Z}_+$  have the same cardinality.

Hint: consider  $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ ,  $f(m, n) = 2^m 3^n$

## Solution

$2^a 3^b = 2^c 3^d$  if and only if  $a = c$  and  $b = d$  (Fund. Thm. Arith.)

$f(m, n) = 2^m 3^n$  is injective

$$|\mathbb{Z}_+ \times \mathbb{Z}_+| \leq |\mathbb{Z}_+|$$

$g : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ ,  $g(m, n) = m$  is surjective

# Challenge

## Problem

Show that  $\mathbb{Z}_+$  and  $\mathbb{Z}_+ \times \mathbb{Z}_+$  have the same cardinality.

Hint: consider  $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ ,  $f(m, n) = 2^m 3^n$

## Solution

$2^a 3^b = 2^c 3^d$  if and only if  $a = c$  and  $b = d$  (Fund. Thm. Arith.)

$f(m, n) = 2^m 3^n$  is injective

$$|\mathbb{Z}_+ \times \mathbb{Z}_+| \leq |\mathbb{Z}_+|$$

$g : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ ,  $g(m, n) = m$  is surjective

$$|\mathbb{Z}_+ \times \mathbb{Z}_+| \geq |\mathbb{Z}_+|$$

# Challenge

## Problem

Show that  $\mathbb{Z}_+$  and  $\mathbb{Z}$  have the same cardinality.

# Challenge

## Problem

Show that  $\mathbb{Z}_+$  and  $\mathbb{Z}$  have the same cardinality.

Hint: consider  $f : \mathbb{Z}_+ \rightarrow \mathbb{Z}$

$$f(n) = \begin{cases} n/2 & n \text{ is even} \\ -(n-1)/2 & n \text{ is odd} \end{cases}$$



# Finite and infinite

# Finite and infinite

Sets with finite cardinality:

$$\{1, 2, 3\}, \quad \{\mathbb{Z}, \mathbb{R}\}, \quad \{x : x \text{ is a student at CSUF}\}$$

# Finite and infinite

Sets with finite cardinality:

$$\{1, 2, 3\}, \quad \{\mathbb{Z}, \mathbb{R}\}, \quad \{x : x \text{ is a student at CSUF}\}$$

Sets with infinite cardinality:

$$\mathbb{Z}_+, \quad \mathbb{Q}, \quad \mathbb{R}, \quad \mathbb{C}.$$

# Finite and infinite

Sets with finite cardinality:

$$\{1, 2, 3\}, \quad \{\mathbb{Z}, \mathbb{R}\}, \quad \{x : x \text{ is a student at CSUF}\}$$

Sets with infinite cardinality:

$$\mathbb{Z}_+, \quad \mathbb{Q}, \quad \mathbb{R}, \quad \mathbb{C}.$$

Cantor's discovery: there are multiple sizes of infinity!

$$\mathbb{Z}_+, \mathbb{Z}, \mathbb{Z} \times \mathbb{Z}, \mathbb{Q}, \text{ are all the same cardinality}$$

# Finite and infinite

Sets with finite cardinality:

$$\{1, 2, 3\}, \quad \{\mathbb{Z}, \mathbb{R}\}, \quad \{x : x \text{ is a student at CSUF}\}$$

Sets with infinite cardinality:

$$\mathbb{Z}_+, \quad \mathbb{Q}, \quad \mathbb{R}, \quad \mathbb{C}.$$

Cantor's discovery: there are multiple sizes of infinity!

$$\mathbb{Z}_+, \mathbb{Z}, \mathbb{Z} \times \mathbb{Z}, \mathbb{Q}, \text{ are all the same cardinality}$$

$$\mathbb{R} \text{ has larger cardinality than } \mathbb{Z}_+.$$

# Cantor diagonalization

Suppose there were a bijection

$$f : \mathbb{Z}_+ \rightarrow (0, 1)$$

# Cantor diagonalization

Suppose there were a bijection

$$f : \mathbb{Z}_+ \rightarrow (0, 1)$$

$$f(1) = 0.a_{11}a_{12}a_{13}a_{14}a_{15}\dots$$

$$f(2) = 0.a_{21}a_{22}a_{23}a_{24}a_{25}\dots$$

$$f(3) = 0.a_{31}a_{32}a_{33}a_{34}a_{35}\dots$$

$$f(4) = 0.a_{41}a_{42}a_{43}a_{44}a_{45}\dots$$

$$f(5) = 0.a_{51}a_{52}a_{53}a_{54}a_{55}\dots$$

$$\vdots$$

# Cantor diagonalization

Suppose there were a surjection

$$f : \mathbb{Z}_+ \rightarrow (0, 1)$$



# Cantor diagonalization

Suppose there were a surjection

$$f : \mathbb{Z}_+ \rightarrow (0, 1)$$

$$f(1) = 0.\textcolor{red}{a}_{11}a_{12}a_{13}a_{14}a_{15}\dots$$

$$f(2) = 0.a_{21}\textcolor{red}{a}_{22}a_{23}a_{24}a_{25}\dots$$

$$f(3) = 0.a_{31}a_{32}\textcolor{red}{a}_{33}a_{34}a_{35}\dots$$

$$f(4) = 0.a_{41}a_{42}a_{43}\textcolor{red}{a}_{44}a_{45}\dots$$

$$f(5) = 0.a_{51}a_{52}a_{53}a_{54}\textcolor{red}{a}_{55}\dots$$

$$\vdots$$

# Cantor diagonalization

Now define  $b_1, b_2, b_3 \cdots \in \{0, 9\}$  by

$$b_j = \begin{cases} 1, & a_{jj} = 0 \\ 0, & a_{jj} \neq 0 \end{cases}$$

# Cantor diagonalization

Now define  $b_1, b_2, b_3 \dots \in \{0, 9\}$  by

$$b_j = \begin{cases} 1, & a_{jj} = 0 \\ 0, & a_{jj} \neq 0 \end{cases}$$

$$b = 0.b_1 b_2 b_3 b_4 b_5 \dots$$

# Cantor diagonalization

- $b \in (0, 1)$

# Cantor diagonalization

- $b \in (0, 1)$
- $f$  is surjective

# Cantor diagonalization

- $b \in (0, 1)$
- $f$  is surjective
- there exists  $n \in \mathbb{Z}_+$  with  $f(n) = b$

# Cantor diagonalization

- $b \in (0, 1)$
- $f$  is surjective
- there exists  $n \in \mathbb{Z}_+$  with  $f(n) = b$

$$0.b_1b_2b_3b_4b_5\cdots = 0.a_{n1}a_{n2}a_{n3}a_{n4}a_{n5}\cdots$$

# Cantor diagonalization

- $b \in (0, 1)$
- $f$  is surjective
- there exists  $n \in \mathbb{Z}_+$  with  $f(n) = b$

$$0.b_1b_2b_3b_4b_5\cdots = 0.a_{n1}a_{n2}a_{n3}a_{n4}a_{n5}\cdots$$

- $b_n = a_{nn}$



# Cantor diagonalization

- $b \in (0, 1)$
- $f$  is surjective
- there exists  $n \in \mathbb{Z}_+$  with  $f(n) = b$

$$0.b_1b_2b_3b_4b_5\cdots = 0.a_{n1}a_{n2}a_{n3}a_{n4}a_{n5}\cdots$$

- $b_n = a_{nn}$  CONTRADICTION!

# Countable and uncountable

A set is

# Countable and uncountable

A set is

- **countably infinite** if it has the same cardinality as  $\mathbb{Z}_+$

# Countable and uncountable

A set is

- **countably infinite** if it has the same cardinality as  $\mathbb{Z}_+$
- **countable** if it is finite or countably infinite

# Countable and uncountable

A set is

- **countably infinite** if it has the same cardinality as  $\mathbb{Z}_+$
- **countable** if it is finite or countably infinite
- **uncountable** otherwise

# Countable and uncountable

A set is

- **countably infinite** if it has the same cardinality as  $\mathbb{Z}_+$
- **countable** if it is finite or countably infinite
- **uncountable** otherwise

Theorem (Cantor's Theorem)

$\mathbb{R}$  is uncountable