

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

October 13, 2024

Outline

- 1 Real Analysis Lecture 12
 - Metric Spaces
 - Subspaces

Outline

- 1 Real Analysis Lecture 12
 - Metric Spaces
 - Subspaces

Warm-up Challenge

Problem

Write down each of the following:

- the definition of an open cover of a set A

Warm-up Challenge

Problem

Write down each of the following:

- the definition of an open cover of a set A
- a compact set

Warm-up Challenge

Problem

Write down each of the following:

- the definition of an open cover of a set A
- a compact set
- Lindelöf Covering Theorem

Warm-up Challenge

Problem

Write down each of the following:

- the definition of an open cover of a set A
- a compact set
- Lindelöf Covering Theorem
- Bolzano-Weierstrass Theorem

Warm-up Challenge

Problem

Write down each of the following:

- the definition of an open cover of a set A
- a compact set
- Lindelöf Covering Theorem
- Bolzano-Weierstrass Theorem
- Cantor Intersection Theorem

Warm-up Challenge

Problem

Write down each of the following:

- the definition of an open cover of a set A
- a compact set
- Lindelöf Covering Theorem
- Bolzano-Weierstrass Theorem
- Cantor Intersection Theorem
- Heine-Borel Theorem

Metric space

Metric space

Definition

A **metric space** is a pair (M, d) consisting of a nonempty set M of “points”, along with a distance function $d : M \times M \rightarrow \mathbb{R}$ with the following four properties.

Metric space

Definition

A **metric space** is a pair (M, d) consisting of a nonempty set M of “points”, along with a distance function $d : M \times M \rightarrow \mathbb{R}$ with the following four properties.

Metric space

Definition

A **metric space** is a pair (M, d) consisting of a nonempty set M of “points”, along with a distance function $d : M \times M \rightarrow \mathbb{R}$ with the following four properties.

A1 $d(x, x) = 0$ for all $x \in M$

Metric space

Definition

A **metric space** is a pair (M, d) consisting of a nonempty set M of “points”, along with a distance function $d : M \times M \rightarrow \mathbb{R}$ with the following four properties.

- A1 $d(x, x) = 0$ for all $x \in M$
- A2 **positivity:** $d(x, y) > 0$ for all $x, y \in M$ with $x \neq y$

Metric space

Definition

A **metric space** is a pair (M, d) consisting of a nonempty set M of “points”, along with a distance function $d : M \times M \rightarrow \mathbb{R}$ with the following four properties.

- A1 $d(x, x) = 0$ for all $x \in M$
- A2 **positivity:** $d(x, y) > 0$ for all $x, y \in M$ with $x \neq y$
- A3 **symmetry:** $d(x, y) = d(y, x)$ for all $x, y \in M$

Metric space

Definition

A **metric space** is a pair (M, d) consisting of a nonempty set M of “points”, along with a distance function $d : M \times M \rightarrow \mathbb{R}$ with the following four properties.

- Ⓐ₁ $d(x, x) = 0$ for all $x \in M$
- Ⓐ₂ **positivity:** $d(x, y) > 0$ for all $x, y \in M$ with $x \neq y$
- Ⓐ₃ **symmetry:** $d(x, y) = d(y, x)$ for all $x, y \in M$
- Ⓐ₄ **triangle inequality:** $d(x, y) \leq d(x, z) + d(z, y)$ for all $x, y, z \in M$

Metric space

Definition

A **metric space** is a pair (M, d) consisting of a nonempty set M of “points”, along with a distance function $d : M \times M \rightarrow \mathbb{R}$ with the following four properties.

- Ⓐ₁ $d(x, x) = 0$ for all $x \in M$
- Ⓐ₂ **positivity:** $d(x, y) > 0$ for all $x, y \in M$ with $x \neq y$
- Ⓐ₃ **symmetry:** $d(x, y) = d(y, x)$ for all $x, y \in M$
- Ⓐ₄ **triangle inequality:** $d(x, y) \leq d(x, z) + d(z, y)$ for all $x, y, z \in M$

The value $d(x, y)$ is called a **metric** and describes the “distance” between x and y .

Open balls

Open balls

Definition

Let (M, d) be a metric space. The open ball of radius $r > 0$ centered at $x \in M$ is

$$B_M(x; r) = \{y \in M : d(x, y) < r\}.$$

Examples of metrics on \mathbb{R}^2

$$d_{\text{eucl}}(\vec{x}, \vec{y}) = |\vec{x} - \vec{y}| = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2}$$

Examples of metrics on \mathbb{R}^2

$$d_{\text{eucl}}(\vec{x}, \vec{y}) = |\vec{x} - \vec{y}| = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2}$$

- Euclidean metric (2-norm)

Examples of metrics on \mathbb{R}^2


$$d_{\text{eucl}}(\vec{x}, \vec{y}) = |\vec{x} - \vec{y}| = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2}$$

- Euclidean metric (2-norm)
- open balls are circles

Examples of metrics on \mathbb{R}^2

$$d_{\text{eucl}}(\vec{x}, \vec{y}) = |\vec{x} - \vec{y}| = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2}$$

- Euclidean metric (2-norm)
- open balls are circles



$B(\vec{x}; r)$

Examples of metrics on \mathbb{R}^2

$$d_{\text{taxi}}(\vec{x}, \vec{y}) = |x_1 - y_1| + |x_2 - y_2|$$

Examples of metrics on \mathbb{R}^2

$$d_{\text{taxi}}(\vec{x}, \vec{y}) = |x_1 - y_1| + |x_2 - y_2|$$

- Taxi-cab metric (1-norm, distances on city streets)

Examples of metrics on \mathbb{R}^2

$$d_{\text{taxi}}(\vec{x}, \vec{y}) = |x_1 - y_1| + |x_2 - y_2|$$

- Taxi-cab metric (1-norm, distances on city streets)
- open balls are diamonds

Examples of metrics on \mathbb{R}^2

$$d_{\text{taxi}}(\vec{x}, \vec{y}) = |x_1 - y_1| + |x_2 - y_2|$$

- Taxi-cab metric (1-norm, distances on city streets)
- open balls are diamonds



Examples of metrics on \mathbb{R}^2

$$d_{\infty}(\vec{x}, \vec{y}) = \max\{|x_1 - y_1|, |x_2 - y_2|\}$$

Examples of metrics on \mathbb{R}^2

$$d_{\infty}(\vec{x}, \vec{y}) = \max\{|x_1 - y_1|, |x_2 - y_2|\}$$

- Chebyshev metric (infinity norm)

Examples of metrics on \mathbb{R}^2


$$d_{\infty}(\vec{x}, \vec{y}) = \max\{|x_1 - y_1|, |x_2 - y_2|\}$$

- Chebyshev metric (infinity norm)
- open balls are squares

Examples of metrics on \mathbb{R}^2

$$d_{\infty}(\vec{x}, \vec{y}) = \max\{|x_1 - y_1|, |x_2 - y_2|\}$$

- Chebyshev metric (infinity norm)
- open balls are squares

 $B(\vec{x}; r)$

Examples of metrics on \mathbb{R}^2

$$d_p(\vec{x}, \vec{y}) = \sqrt[p]{|x_1 - y_1|^p + |x_2 - y_2|^p}$$

Examples of metrics on \mathbb{R}^2

$$d_p(\vec{x}, \vec{y}) = \sqrt[p]{|x_1 - y_1|^p + |x_2 - y_2|^p}$$

- p -norm, $1 \leq p < \infty$

Examples of metrics on \mathbb{R}^2


$$d_p(\vec{x}, \vec{y}) = \sqrt[p]{|x_1 - y_1|^p + |x_2 - y_2|^p}$$

- p -norm, $1 \leq p < \infty$
- open balls are rounded squares

Examples of metrics on \mathbb{R}^2

$$d_p(\vec{x}, \vec{y}) = \sqrt[p]{|x_1 - y_1|^p + |x_2 - y_2|^p}$$

- p -norm, $1 \leq p < \infty$
- open balls are rounded squares



$B(\vec{x}; r)$

Challenge!

Let M be a nonempty set and define $d : M \times M \rightarrow \mathbb{R}$ by

$$d_{\text{disc}}(x, y) = \begin{cases} 1 & x \neq y \\ 0 & x = y \end{cases}$$

Challenge!

Let M be a nonempty set and define $d : M \times M \rightarrow \mathbb{R}$ by

$$d_{\text{disc}}(x, y) = \begin{cases} 1 & x \neq y \\ 0 & x = y \end{cases}$$

Problem

Prove that d_{disc} is a metric.

Challenge!

Let M be a nonempty set and define $d : M \times M \rightarrow \mathbb{R}$ by

$$d_{\text{disc}}(x, y) = \begin{cases} 1 & x \neq y \\ 0 & x = y \end{cases}$$

Problem

Prove that d_{disc} is a metric.

This is called the **discrete metric**.

Challenge!

Problem

What do the open balls with the discrete metric look like?

Open sets

Definition

Open sets

Definition

Let (M, d) be a metric space and $A \subseteq M$.

Open sets

Definition

Let (M, d) be a metric space and $A \subseteq M$.

A point $x \in A$ is called an **interior point** if there exists $r > 0$ such that $B_M(x; r) \subseteq A$.

Open sets

Definition

Let (M, d) be a metric space and $A \subseteq M$.

A point $x \in A$ is called an **interior point** if there exists $r > 0$ such that $B_M(x; r) \subseteq A$.

The set $\text{int}(A)$ of interior points of A is called the **interior** of A .

Open sets

Definition

Let (M, d) be a metric space and $A \subseteq M$.

A point $x \in A$ is called an **interior point** if there exists $r > 0$ such that $B_M(x; r) \subseteq A$.

The set $\text{int}(A)$ of interior points of A is called the **interior** of A .

The set A is **open** if every point in A is an interior point, or equivalently $\text{int}(A) = A$.

Challenge!

Problem

Consider the metric space $(\mathbb{R}, d_{\text{disc}})$ where d_{disc} is the discrete metric.

What sets are open sets?

Closed sets

Definition

Closed sets

Definition

Let (M, d) be a metric space and $A \subseteq M$.

Closed sets

Definition

Let (M, d) be a metric space and $A \subseteq M$.

A set $A \subseteq M$ is closed if the complement $M \setminus A$ is open.

Closed sets

Definition

Let (M, d) be a metric space and $A \subseteq M$.

A set $A \subseteq M$ is closed if the complement $M \setminus A$ is open.

A point $x \in M$ is an **adherent point** if for all $r > 0$ the ball $B_M(x; r)$ contains an element of A .

Closed sets

Definition

Let (M, d) be a metric space and $A \subseteq M$.

A set $A \subseteq M$ is closed if the complement $M \setminus A$ is open.

A point $x \in M$ is an **adherent point** if for all $r > 0$ the ball $B_M(x; r)$ contains an element of A .

A point $x \in M$ is an **accumulation point** if for all $r > 0$ the ball $B_M(x; r)$ contains an element of A different from x .

Closed sets

Definition

Let (M, d) be a metric space and $A \subseteq M$.

A set $A \subseteq M$ is closed if the complement $M \setminus A$ is open.

A point $x \in M$ is an **adherent point** if for all $r > 0$ the ball $B_M(x; r)$ contains an element of A .

A point $x \in M$ is an **accumulation point** if for all $r > 0$ the ball $B_M(x; r)$ contains an element of A different from x .

The set \bar{A} of all adherent points of A is called the **closure** of A .

Challenge!

Problem

Consider the metric space $(\mathbb{R}, d_{\text{disc}})$ where d_{disc} is the discrete metric.

Which sets are closed?

Challenge!

Problem

Prove the following (Apostol Theorem 3.36). If (M, d) is a metric space, $U \subseteq M$ is open, and $C \subseteq M$ is closed, then $U \setminus C$ is open and $C \setminus U$ is closed.

Characterizing closed sets

Theorem (Apostol Theorem 3.37)

Let (M, d) be a metric space and $A \subseteq S$. Then the following are equivalent:

Characterizing closed sets

Theorem (Apostol Theorem 3.37)

Let (M, d) be a metric space and $A \subseteq S$. Then the following are equivalent:

- (a) *A is closed*

Characterizing closed sets

Theorem (Apostol Theorem 3.37)

Let (M, d) be a metric space and $A \subseteq S$. Then the following are equivalent:

- (a) *A is closed*
- (b) *A contains all of its adherent points*

Characterizing closed sets

Theorem (Apostol Theorem 3.37)

Let (M, d) be a metric space and $A \subseteq S$. Then the following are equivalent:

- (a) A is closed*
- (b) A contains all of its adherent points*
- (c) A contains all of its accumulation points*

Characterizing closed sets

Theorem (Apostol Theorem 3.37)

Let (M, d) be a metric space and $A \subseteq S$. Then the following are equivalent:

- (a) *A is closed*
- (b) *A contains all of its adherent points*
- (c) *A contains all of its accumulation points*
- (d) *$A = \bar{A}$*

Unions and intersections

Unions and intersections

Theorem

Let (M, d) be a metric space and $\{U_i : i \in I\}$ be a family of open sets in M . The union $\bigcup_{i \in I} U_i$ is open.

Unions and intersections

Theorem

Let (M, d) be a metric space and $\{U_i : i \in I\}$ be a family of open sets in M . The union $\bigcup_{i \in I} U_i$ is open.

Proof.

Identical to the proof for \mathbb{R}^n . □

Unions and intersections

Theorem

Let (M, d) be a metric space and $\{U_i : i \in I\}$ be a family of open sets in M . The union $\bigcup_{i \in I} U_i$ is open.

Proof.

Identical to the proof for \mathbb{R}^n . □

Theorem

Let (M, d) be a metric space and U_1, U_2, \dots, U_n be open sets in M . The intersection $\bigcap_{i=1}^n U_i$ is open.

Unions and intersections

Theorem

Let (M, d) be a metric space and $\{U_i : i \in I\}$ be a family of open sets in M . The union $\bigcup_{i \in I} U_i$ is open.

Proof.

Identical to the proof for \mathbb{R}^n . □

Theorem

Let (M, d) be a metric space and U_1, U_2, \dots, U_n be open sets in M . The intersection $\bigcap_{i=1}^n U_i$ is open.

Proof.

Identical to the proof for \mathbb{R}^n . □

Unions and intersections

Unions and intersections

Theorem

Let (M, d) be a metric space and $\{C_i : i \in I\}$ be a family of closed sets in M . The intersection $\bigcap_{i \in I} C_i$ is closed.

Unions and intersections

Theorem

Let (M, d) be a metric space and $\{C_i : i \in I\}$ be a family of closed sets in M . The intersection $\bigcap_{i \in I} C_i$ is closed.

Proof.

Apply De Morgan's Laws. □

Unions and intersections

Theorem

Let (M, d) be a metric space and $\{C_i : i \in I\}$ be a family of closed sets in M . The intersection $\bigcap_{i \in I} C_i$ is closed.

Proof.

Apply De Morgan's Laws. □

Theorem

Let (M, d) be a metric space and C_1, C_2, \dots, C_n be closed sets in M . The intersection $\bigcup_{i=1}^n C_i$ is closed.

Unions and intersections

Theorem

Let (M, d) be a metric space and $\{C_i : i \in I\}$ be a family of closed sets in M . The intersection $\bigcap_{i \in I} C_i$ is closed.

Proof.

Apply De Morgan's Laws. ☐

Theorem

Let (M, d) be a metric space and C_1, C_2, \dots, C_n be closed sets in M . The intersection $\bigcup_{i=1}^n C_i$ is closed.

Proof.

Apply De Morgan's Laws. ☐

Outline

- 1 Real Analysis Lecture 12
 - Metric Spaces
 - Subspaces

Metric subspace

Definition

Let (M, d) be a metric space.

Metric subspace

Definition

Let (M, d) be a metric space.

If $S \subseteq M$ is a subset, then $d : M \times M \rightarrow \mathbb{R}$ restricts to a metric $d : S \times S \rightarrow \mathbb{R}$, called the **relative metric**.

Metric subspace

Definition

Let (M, d) be a metric space.

If $S \subseteq M$ is a subset, then $d : M \times M \rightarrow \mathbb{R}$ restricts to a metric $d : S \times S \rightarrow \mathbb{R}$, called the **relative metric**.

The space (S, d) with the relative metric is called a **metric subspace**.

Metric subspace

Definition

Let (M, d) be a metric space.

If $S \subseteq M$ is a subset, then $d : M \times M \rightarrow \mathbb{R}$ restricts to a metric $d : S \times S \rightarrow \mathbb{R}$, called the **relative metric**.

The space (S, d) with the relative metric is called a **metric subspace**.

Open balls in S :

$$B_S(x; r) = B_M(x; r) \cap S.$$

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\implies : Assume V is an open subset of S .

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\implies : Assume V is an open subset of S .

Then for all $x \in V$, there exists $r_x > 0$ such that $B_S(x; r_x) \subseteq V$.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\implies : Assume V is an open subset of S .

Then for all $x \in V$, there exists $r_x > 0$ such that $B_S(x; r_x) \subseteq V$.

This says $B_M(x; r_x) \cap S \subseteq V$.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\implies : Assume V is an open subset of S .

Then for all $x \in V$, there exists $r_x > 0$ such that $B_S(x; r_x) \subseteq V$.

This says $B_M(x; r_x) \cap S \subseteq V$.

Define

$$U = \bigcup_{x \in V} B_M(x; r_x).$$

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\implies : Assume V is an open subset of S .

Then for all $x \in V$, there exists $r_x > 0$ such that $B_S(x; r_x) \subseteq V$.

This says $B_M(x; r_x) \cap S \subseteq V$.

Define

$$U = \bigcup_{x \in V} B_M(x; r_x).$$

This is a union of open sets, and is therefore open.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\implies : Assume V is an open subset of S .

Then for all $x \in V$, there exists $r_x > 0$ such that $B_S(x; r_x) \subseteq V$.

This says $B_M(x; r_x) \cap S \subseteq V$.

Define

$$U = \bigcup_{x \in V} B_M(x; r_x).$$

This is a union of open sets, and is therefore open.

Furthermore $U \cap S = V$.



Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\Leftarrow : Assume U is an open subset of M .

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\Leftarrow : Assume U is an open subset of M .
We need to show that $V := U \cap S$ is open.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\Leftarrow : Assume U is an open subset of M .

We need to show that $V := U \cap S$ is open.

If $x \in V$, then $x \in U$ and there exists $r > 0$ such that $B_M(x; r) \subseteq U$.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\Leftarrow : Assume U is an open subset of M .

We need to show that $V := U \cap S$ is open.

If $x \in V$, then $x \in U$ and there exists $r > 0$ such that $B_M(x; r) \subseteq U$.

It follows that $B_S(x, r) = B_M(x, r) \cap S \subseteq U \cap S = V$.

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\Leftarrow : Assume U is an open subset of M .

We need to show that $V := U \cap S$ is open.

If $x \in V$, then $x \in U$ and there exists $r > 0$ such that $B_M(x; r) \subseteq U$.

It follows that $B_S(x, r) = B_M(x, r) \cap S \subseteq U \cap S = V$.

This shows that x is an interior point of S in the metric subspace (S, d) .

Open sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then V is an open subset of S if and only if there exists an open subset U of M with $V = U \cap S$.

Proof.

\Leftarrow : Assume U is an open subset of M .

We need to show that $V := U \cap S$ is open.

If $x \in V$, then $x \in U$ and there exists $r > 0$ such that $B_M(x; r) \subseteq U$.

It follows that $B_S(x, r) = B_M(x, r) \cap S \subseteq U \cap S = V$.

This shows that x is an interior point of S in the metric subspace (S, d) .

Since $x \in V$ was arbitrary, this shows that V is open in S . □

Closed sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then B is a closed subset of S if and only if there exists a closed subset A of M with $B = A \cap S$.

Closed sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then B is a closed subset of S if and only if there exists a closed subset A of M with $B = A \cap S$.

Proof.

Closed sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then B is a closed subset of S if and only if there exists a closed subset A of M with $B = A \cap S$.

Proof.

Let $B \subseteq S$.

Closed sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then B is a closed subset of S if and only if there exists a closed subset A of M with $B = A \cap S$.

Proof.

Let $B \subseteq S$.

B is closed in S if and only if $S \setminus B$ is open in B

Closed sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then B is a closed subset of S if and only if there exists a closed subset A of M with $B = A \cap S$.

Proof.

Let $B \subseteq S$.

B is closed in S if and only if $S \setminus B$ is open in B

This is true if and only if there exists an open subset U of M with $S \setminus B = U \cap S$.

Closed sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then B is a closed subset of S if and only if there exists a closed subset A of M with $B = A \cap S$.

Proof.

Let $B \subseteq S$.

B is closed in S if and only if $S \setminus B$ is open in B

This is true if and only if there exists an open subset U of M with $S \setminus B = U \cap S$.

This is true if and only if there exists an open subset U of M with $B = (M \setminus U) \cap S$.

Closed sets in subspaces

Theorem

Let (M, d) be a metric space and let (S, d) be a subspace. Then B is a closed subset of S if and only if there exists a closed subset A of M with $B = A \cap S$.

Proof.

Let $B \subseteq S$.

B is closed in S if and only if $S \setminus B$ is open in B

This is true if and only if there exists an open subset U of M with $S \setminus B = U \cap S$.

This is true if and only if there exists an open subset U of M with $B = (M \setminus U) \cap S$.

this is true if and only if there exists a closed subset A of M with $B = A \cap S$.



Example

Consider the metric space (\mathbb{R}, d) with $d = d_{\text{eucl}}$ the Euclidean metric, and the metric subspace (S, d) for $S = [0, 1]$.

Example

Consider the metric space (\mathbb{R}, d) with $d = d_{\text{eucl}}$ the Euclidean metric, and the metric subspace (S, d) for $S = [0, 1]$.

- $[0, 1]$ is open in S and closed in S , but it is not open in M

Example

Consider the metric space (\mathbb{R}, d) with $d = d_{\text{eucl}}$ the Euclidean metric, and the metric subspace (S, d) for $S = [0, 1]$.

- $[0, 1]$ is open in S and closed in S , but it is not open in M
- $[0, 1/2)$ is open in S , but not in M

Example

Consider the metric space (\mathbb{R}, d) with $d = d_{\text{eucl}}$ the Euclidean metric, and the metric subspace (S, d) for $S = [0, 1]$.

- $[0, 1]$ is open in S and closed in S , but it is not open in M
- $[0, 1/2)$ is open in S , but not in M
- $(0, 1/2)$ is open in both S and in M

Example

Consider the metric space (\mathbb{R}, d) with $d = d_{\text{eucl}}$ the Euclidean metric, and the metric subspace (S, d) for $S = [0, 1]$.

- $[0, 1]$ is open in S and closed in S , but it is not open in M
- $[0, 1/2)$ is open in S , but not in M
- $(0, 1/2)$ is open in both S and in M
- $(1/2, 1]$ is open in S , but not in M