

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

August 26, 2024

Outline

- 1 Real Analysis Lecture 2
 - Types of real numbers
 - Integers
 - Upper Bound and Supremum
 - Decimal expansions

Outline

- 1 Real Analysis Lecture 2
 - Types of real numbers
 - Integers
 - Upper Bound and Supremum
 - Decimal expansions

Real numbers

The **real numbers** are the unique field \mathbb{R} satisfying Axiom 1-Axiom 9, plus an extra axiom called the completeness axiom.

\mathbb{R} = the one and only complete, ordered field

A complete ordered field F satisfies

Real numbers

The **real numbers** are the unique field \mathbb{R} satisfying Axiom 1-Axiom 9, plus an extra axiom called the completeness axiom.

\mathbb{R} = the one and only complete, ordered field

A complete ordered field F satisfies

- Ⓐ **completeness axiom**: every bounded set of numbers $S \subseteq F$ has a supremum $\sup(S)$

Real numbers

The **real numbers** are the unique field \mathbb{R} satisfying Axiom 1-Axiom 9, plus an extra axiom called the completeness axiom.

\mathbb{R} = the one and only complete, ordered field

A complete ordered field F satisfies

- Ⓐ **completeness axiom**: every bounded set of numbers $S \subseteq F$ has a supremum $\sup(S)$
- Understanding what this is challenging, but will be fundamental to the entire course!

Real numbers

The **real numbers** are the unique field \mathbb{R} satisfying Axiom 1-Axiom 9, plus an extra axiom called the completeness axiom.

\mathbb{R} = the one and only complete, ordered field

A complete ordered field F satisfies

- Ⓐ6 **completeness axiom**: every bounded set of numbers $S \subseteq F$ has a supremum $\sup(S)$
- Understanding what this is challenging, but will be fundamental to the entire course!
- Intuitively, it represents the fact that the real line has no holes or gaps.

Real intervals

Given an ordered field, we can **intervals**

Real intervals

Given an ordered field, we can **intervals**

- **open and closed intervals**

$$(a, b) = \{x \in \mathbb{R} : a < x < b\} \quad \text{and} \quad [a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$$

Real intervals

Given an ordered field, we can **intervals**

- **open and closed intervals**

$$(a, b) = \{x \in \mathbb{R} : a < x < b\} \quad \text{and} \quad [a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$$

- **half-open intervals**

$$[a, b) = \{x \in \mathbb{R} : a \leq x < b\} \quad \text{and} \quad (a, b] = \{x \in \mathbb{R} : a < x \leq b\}$$

Real intervals

Given an ordered field, we can **intervals**

- **open and closed intervals**

$$(a, b) = \{x \in \mathbb{R} : a < x < b\} \quad \text{and} \quad [a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$$

- **half-open intervals**

$$[a, b) = \{x \in \mathbb{R} : a \leq x < b\} \quad \text{and} \quad (a, b] = \{x \in \mathbb{R} : a < x \leq b\}$$

- **semi-infinite open intervals**

$$(-\infty, a) = \{x \in \mathbb{R} : x < a\} \quad \text{and} \quad (a, \infty) = \{x \in \mathbb{R} : a < x\}$$

Real intervals

Given an ordered field, we can **intervals**

- **open and closed intervals**

$$(a, b) = \{x \in \mathbb{R} : a < x < b\} \quad \text{and} \quad [a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$$

- **half-open intervals**

$$[a, b) = \{x \in \mathbb{R} : a \leq x < b\} \quad \text{and} \quad (a, b] = \{x \in \mathbb{R} : a < x \leq b\}$$

- **semi-infinite open intervals**

$$(-\infty, a) = \{x \in \mathbb{R} : x < a\} \quad \text{and} \quad (a, \infty) = \{x \in \mathbb{R} : a < x\}$$

- **semi-infinite closed intervals**

$$(-\infty, a] = \{x \in \mathbb{R} : x \leq a\} \quad \text{and} \quad [a, \infty) = \{x \in \mathbb{R} : a \leq x\}.$$

Outline

- 1 Real Analysis Lecture 2
 - Types of real numbers
 - **Integers**
 - Upper Bound and Supremum
 - Decimal expansions

Inductive sets

A subset S of \mathbb{R} is called a **inductive set** if

- $1 \in S$
- if $x \in S$ then $x + 1 \in S$

Inductive sets

A subset S of \mathbb{R} is called a **inductive set** if

- $1 \in S$
- if $x \in S$ then $x + 1 \in S$

Examples:

$$\mathbb{R}, \quad \mathbb{Q}, \quad (0, \infty), \quad \mathbb{Z}, \quad \mathbb{N}, \dots$$

Integers and rationals

How do we reverse-engineer the integers from the reals?

Integers and rationals

How do we reverse-engineer the integers from the reals?

- the set of **positive integers** is

$$\mathbb{Z}_+ = \{x : x \text{ belongs to every inductive set}\}$$

Integers and rationals

How do we reverse-engineer the integers from the reals?

- the set of **positive integers** is

$$\mathbb{Z}_+ = \{x : x \text{ belongs to every inductive set}\}$$

- integers are

$$\mathbb{Z} = \{x : x \text{ is zero or } \pm x \text{ is a positive integer}\}$$

Integers and rationals

How do we reverse-engineer the integers from the reals?

- the set of **positive integers** is

$$\mathbb{Z}_+ = \{x : x \text{ belongs to every inductive set}\}$$

- integers are

$$\mathbb{Z} = \{x : x \text{ is zero or } \pm x \text{ is a positive integer}\}$$

- rationals are

$$\mathbb{Q} = \{a/b : a, b \in \mathbb{Z}, b \neq 0\}.$$

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof:

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof:

Let

$$S = \{n \in \mathbb{Z}_+ : 2|n(n+1)\}$$

.

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof:

Let

$$S = \{n \in \mathbb{Z}_+ : 2 \mid n(n+1)\}$$

. We see that $1 \in S$ because 2 divides $1(1+1)$.

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof:

Let

$$S = \{n \in \mathbb{Z}_+ : 2|n(n+1)\}$$

. We see that $1 \in S$ because 2 divides $1(1+1)$.

Now suppose $x \in S$. (This is our usual inductive assumption).

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Induction

Example: Let's prove $n(n + 1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x + 1)$, so $x(x + 1) = 2k$ for some integer k .

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .

This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.
Since $x+1$ was arbitrary, this shows that S is an inductive set.

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.
Since $x+1$ was arbitrary, this shows that S is an inductive set.
By the Principle of Induction, this means $\mathbb{Z}_+ \subseteq S$.

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.
Since $x+1$ was arbitrary, this shows that S is an inductive set.
By the Principle of Induction, this means $\mathbb{Z}_+ \subseteq S$.
In other words $n \in S$ for every positive integer n .

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.
Since $x+1$ was arbitrary, this shows that S is an inductive set.
By the Principle of Induction, this means $\mathbb{Z}_+ \subseteq S$.
In other words $n \in S$ for every positive integer n .
Hence 2 divides $n(n+1)$ for every positive integers n .

Prime numbers

A positive integer p is **prime** if its only positive divisors are 1 and p .

Theorem (Apostol Theorem 1.5)

Every integer is prime or a product of primes

Theorem (Apostol Theorem 1.8)


If p is prime and p divides ab , then p divides a or p divides b .

Theorem (Fundamental Theorem of Arithmetic (Apostol Theorem 1.9))

Every integer $n > 1$ has a unique factorization as a product of primes, up to reordering.

Types of numbers


Transcendental Numbers



Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Discovered by Hippasus, a pythagorean (a student of Pythagoras)



Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery

Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- dramatic reenactment, found online:

Irrational numbers


Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- dramatic reenactment, found online:

Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- dramatic reenactment, found online:



Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- anime style:

Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- anime style:



Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b .

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$. It follows that $b^2 n' = 1$, so b^2 divides 1.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$. It follows that $b^2 n' = 1$, so b^2 divides 1. This implies n' divides 1, so $n' = \pm 1$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$. It follows that $b^2 n' = 1$, so b^2 divides 1. This implies n' divides 1, so $n' = \pm 1$. Since $n > 0$, $n' > 0$ so $n' = 1$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then \sqrt{n} is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$. It follows that $b^2 n' = 1$, so b^2 divides 1. This implies n' divides 1, so $n' = \pm 1$. Since $n > 0$, $n' > 0$ so $n' = 1$. Thus $n = a^2$, which is a contradiction.

Algebraic and Transcendental

Irrational numbers can be further divided into to categories

Algebraic and Transcendental

Irrational numbers can be further divided into to categories

- ① **algebraic numbers:** numbers which are roots of polynomials with integer coefficients, like

$$\sqrt{2}, \quad \sqrt{3}, \quad \text{and} \quad \sqrt[3]{\sqrt{5} + \sqrt{7}}.$$

Algebraic and Transcendental

Irrational numbers can be further divided into to categories

- 1 **algebraic numbers:** numbers which are roots of polynomials with integer coefficients, like

$$\sqrt{2}, \quad \sqrt{3}, \quad \text{and} \quad \sqrt[3]{\sqrt{5} + \sqrt{7}}.$$

- 2 **transcendental numbers:** numbers which are not algebraic, like

$$\pi, \quad e, \quad e^{\pi}, \quad \text{and maybe} \quad \pi + e?$$

Algebraic and Transcendental

Irrational numbers can be further divided into to categories

- 1 **algebraic numbers:** numbers which are roots of polynomials with integer coefficients, like

$$\sqrt{2}, \quad \sqrt{3}, \quad \text{and} \quad \sqrt[3]{\sqrt{5} + \sqrt{7}}.$$

- 2 **transcendental numbers:** numbers which are not algebraic, like

$$\pi, \quad e, \quad e^{\pi}, \quad \text{and maybe} \quad \pi + e?$$

- 3 transcendentals are mysterious ... but most real numbers are transcendental!

Outline

- 1 Real Analysis Lecture 2
 - Types of real numbers
 - Integers
 - **Upper Bound and Supremum**
 - Decimal expansions

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

If $b \in S$ also, then b is called a **maximal element** of S

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

If $b \in S$ also, then b is called a **maximal element** of S

Examples:

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

If $b \in S$ also, then b is called a **maximal element** of S

Examples:

- 34 is an upper bound of $[-1, 5]$, but not a maximal element

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

If $b \in S$ also, then b is called a **maximal element** of S

Examples:

- 34 is an upper bound of $[-1, 5]$, but not a maximal element
- 5 is a maximal element of $[-1, 5]$

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

If $b \in S$ also, then b is called a **maximal element** of S

Examples:

- 34 is an upper bound of $[-1, 5]$, but not a maximal element
- 5 is a maximal element of $[-1, 5]$
- 3 is an upper bound of $[0, 3)$, but not a maximal element

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

If $b \in S$ also, then b is called a **maximal element** of S

Examples:

- 34 is an upper bound of $[-1, 5]$, but not a maximal element
- 5 is a maximal element of $[-1, 5]$
- 3 is an upper bound of $[0, 3)$, but not a maximal element
- \mathbb{Z}_+ has no upper bound

Upper bounds

An **upper bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say S is **bounded above** by b .

If $b \in S$ also, then b is called a **maximal element** of S

Examples:

- 34 is an upper bound of $[-1, 5]$, but not a maximal element
- 5 is a maximal element of $[-1, 5]$
- 3 is an upper bound of $[0, 3)$, but not a maximal element
- \mathbb{Z}_+ has no upper bound
- $[3, 7)$ has an upper bound but no maximal element

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both maximal elements of S .

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both maximal elements of S .
Then $x \leq b_1$ and $x \leq b_2$ for all $x \in S$.

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both maximal elements of S .

Then $x \leq b_1$ and $x \leq b_2$ for all $x \in S$.

Moreover, $b_1 \in S$ and $b_2 \in S$.

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both maximal elements of S .

Then $x \leq b_1$ and $x \leq b_2$ for all $x \in S$.

Moreover, $b_1 \in S$ and $b_2 \in S$.

Since $b_1 \in S$ and b_2 is an upper bound of S , $b_1 \leq b_2$.

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both maximal elements of S .

Then $x \leq b_1$ and $x \leq b_2$ for all $x \in S$.

Moreover, $b_1 \in S$ and $b_2 \in S$.

Since $b_1 \in S$ and b_2 is an upper bound of S , $b_1 \leq b_2$.

Likewise, since $b_2 \in S$ and b_1 is an upper bound of S , $b_2 \leq b_1$.

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both maximal elements of S .

Then $x \leq b_1$ and $x \leq b_2$ for all $x \in S$.

Moreover, $b_1 \in S$ and $b_2 \in S$.

Since $b_1 \in S$ and b_2 is an upper bound of S , $b_1 \leq b_2$.

Likewise, since $b_2 \in S$ and b_1 is an upper bound of S , $b_2 \leq b_1$.

By the trichotomy, we find $b_1 = b_2$.

Challenge!

Problem

Show that if S has a maximal element, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both maximal elements of S .

Then $x \leq b_1$ and $x \leq b_2$ for all $x \in S$.

Moreover, $b_1 \in S$ and $b_2 \in S$.

Since $b_1 \in S$ and b_2 is an upper bound of S , $b_1 \leq b_2$.

Likewise, since $b_2 \in S$ and b_1 is an upper bound of S , $b_2 \leq b_1$.

By the trichotomy, we find $b_1 = b_2$.

Now we can say *the* maximum, $\max(S)$

Supremum

A **supremum** of a set S of real numbers is a real number $b \in \mathbb{R}$ such that

- b is an upper bound of S
- if $b' < b$, then b' is not an upper bound of S

Supremum

A **supremum** of a set S of real numbers is a real number $b \in \mathbb{R}$ such that

- b is an upper bound of S
- if $b' < b$, then b' is not an upper bound of S

In other words

a supremum is a **least upper bound**

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Hint: use the definition!

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both suprema of S .

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both suprema of S .
Then b_1 and b_2 are both upper bounds of S .

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both suprema of S .
Then b_1 and b_2 are both upper bounds of S .
Since b_1 is a least upper bound, $b_1 \leq b_2$.

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both suprema of S .

Then b_1 and b_2 are both upper bounds of S .

Since b_1 is a least upper bound, $b_1 \leq b_2$.

Since b_2 is also a least upper bound, $b_2 \leq b_1$.

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both suprema of S .

Then b_1 and b_2 are both upper bounds of S .

Since b_1 is a least upper bound, $b_1 \leq b_2$.

Since b_2 is also a least upper bound, $b_2 \leq b_1$.

By the trichotomy, we find $b_1 = b_2$.

Challenge!

Problem

Show that if S has a supremum, then it is unique.

Hint: use the definition!

Solution

Suppose that b_1 and b_2 are both suprema of S .

Then b_1 and b_2 are both upper bounds of S .

Since b_1 is a least upper bound, $b_1 \leq b_2$.

Since b_2 is also a least upper bound, $b_2 \leq b_1$.

By the trichotomy, we find $b_1 = b_2$.

Now we can say *the* supremum, $\sup(S)$

Challenge!

Problem

Show that if S has a maximum element. Then S has a supremum and $\max(S) = \sup(S)$

Challenge!

Problem

Show that if S has a maximum element. Then S has a supremum and $\max(S) = \sup(S)$

Hint: use the definition!

Challenge!

Problem

Show that if S has a maximum element. Then S has a supremum and $\max(S) = \sup(S)$

Hint: use the definition!

Solution

Let $b = \max(S)$.

Challenge!

Problem

Show that if S has a maximum element. Then S has a supremum and $\max(S) = \sup(S)$

Hint: use the definition!

Solution

Let $b = \max(S)$.

Then b is an upper bound of S and $b \in S$.

Challenge!

Problem

Show that if S has a maximum element. Then S has a supremum and $\max(S) = \sup(S)$

Hint: use the definition!

Solution

Let $b = \max(S)$.

Then b is an upper bound of S and $b \in S$.

If b' is another upper bound of S , then by definition $b \leq b'$.

Challenge!

Problem

Show that if S has a maximum element. Then S has a supremum and $\max(S) = \sup(S)$

Hint: use the definition!

Solution

Let $b = \max(S)$.

Then b is an upper bound of S and $b \in S$.

If b' is another upper bound of S , then by definition $b \leq b'$.

Thus b is the least upper bound of S .

Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.

Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.


Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.

- Ⓐ10 **completeness axiom:** if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.

 **completeness axiom:** if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.

Ⓐ10 **completeness axiom:** if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

- 3 is the supremum of $(0, 3)$

Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.

A10 completeness axiom: if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

- 3 is the supremum of $(0, 3)$
- 1 is the supremum of

$$\left\{ \frac{n}{n+1} : n \in \mathbb{Z}_+ \right\}$$

Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.

completeness axiom: if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

- 3 is the supremum of $(0, 3)$
- 1 is the supremum of

$$\left\{ \frac{n}{n+1} : n \in \mathbb{Z}_+ \right\}$$

- π is the supremum of

$$\{3, 3.1, 3.14, 3.141, 3.1415, 3.14159, 3.141592, \dots\}.$$

Challenge

Problem

Prove that if A is a set of integers that is bounded above, then A has a maximum.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

If $a' \in A$, then a' is an integer and therefore $a' = a + k$ for $k \in \mathbb{Z}$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

If $a' \in A$, then a' is an integer and therefore $a' = a + k$ for $k \in \mathbb{Z}$. Since b is an upper bound, $b \geq a + k > b - 1 + k$, making $0 > k - 1$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

If $a' \in A$, then a' is an integer and therefore $a' = a + k$ for $k \in \mathbb{Z}$. Since b is an upper bound, $b \geq a + k > b - 1 + k$, making $0 > k - 1$.

Therefore $k \leq 0$ and $a' \leq a$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

If $a' \in A$, then a' is an integer and therefore $a' = a + k$ for $k \in \mathbb{Z}$. Since b is an upper bound, $b \geq a + k > b - 1 + k$, making $0 > k - 1$.

Therefore $k \leq 0$ and $a' \leq a$.

It follows that a is an upper bound of A , and since $a \in A$ it is a maximum.

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

If $b \in S$ also, then b is called a **minimal element** of S .

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

If $b \in S$ also, then b is called a **minimal element** of S . An **infimum** of a set S of real numbers is a real number $b \in \mathbb{R}$ such that

- b is a lower bound of S
- if $b < b'$, then b' is not a lower bound of S

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

If $b \in S$ also, then b is called a **minimal element** of S . An **infimum** of a set S of real numbers is a real number $b \in \mathbb{R}$ such that

- b is a lower bound of S
- if $b < b'$, then b' is not a lower bound of S

In other words

an infimum is a **greatest lower bound**

Properties of Suprema

The first property of suprema is that they must be arbitrarily close to elements of the set.

Theorem (Approximation Property)

Let $S \subseteq \mathbb{R}$ be bounded above, and let $b = \sup(S)$. Then for all $\epsilon > 0$, there exists $a \in S$ with

$$b - \epsilon < a < b.$$

Properties of Suprema

Proof.

Since $b - \epsilon < b$, the definition of a supremum implies $b - \epsilon$ cannot be an upper bound.

Properties of Suprema

Proof.

Since $b - \epsilon < b$, the definition of a supremum implies $b - \epsilon$ cannot be an upper bound.

Therefore there must exist $a \in S$ with $a > b - \epsilon$. □

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Consider the set

$$S = \{s_n : n \in \mathbb{Z}_+\}.$$

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Consider the set

$$S = \{s_n : n \in \mathbb{Z}_+\}.$$

For every n ,

$$s_n \leq 1 + 1 + \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \cdots + \frac{1}{2^n} = 3 - \frac{1}{2^n} < 3$$

so S is bounded above by 3.

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Consider the set

$$S = \{s_n : n \in \mathbb{Z}_+\}.$$

For every n ,

$$s_n \leq 1 + 1 + \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \cdots + \frac{1}{2^n} = 3 - \frac{1}{2^n} < 3$$

so S is bounded above by 3. Therefore S has a supremum, $\sup(S)$.

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N .

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

If we remember Taylor series

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

If we remember Taylor series

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots$$

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

If we remember Taylor series

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots$$

We've just shown that $e^1 = e$ exists.

Properties of Suprema

Also, suprema play nicely with addition.

Theorem (Additive Property)

Let $A, B \subseteq \mathbb{R}$ be bounded above set

$$C = \{x + y : x \in A, y \in B\}.$$

Then C is bounded above and

$$\sup(C) = \sup(A) + \sup(B).$$

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

It follows that $y \leq c - a$ for all $y \in B$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

It follows that $y \leq c - a$ for all $y \in B$.

Thus $c - a$ is an upper bound of B , and it follows $b \leq c - a$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

It follows that $y \leq c - a$ for all $y \in B$.

Thus $c - a$ is an upper bound of B , and it follows $b \leq c - a$.

Therefore $a + b \leq c$. □

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

This means there exists $n \in \mathbb{Z}_+$ with $b - 1 < n$.

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

This means there exists $n \in \mathbb{Z}_+$ with $b - 1 < n$.

It follows from Axiom 7 that $b < n + 1$.

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

This means there exists $n \in \mathbb{Z}_+$ with $b - 1 < n$.

It follows from Axiom 7 that $b < n + 1$.

However, $n + 1 \in \mathbb{Z}$, so this contradicts b being an upper bound.

Archimedian property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{Z}_+$ with $n > x$.

Archimedian property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{Z}_+$ with $n > x$.

Proof.

If not, then x is an upper bound of \mathbb{Z}_+ . □

Archimedean property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{Z}_+$ with $n > x$.

Proof.

If not, then x is an upper bound of \mathbb{Z}_+ . □

Theorem (Archimedean Property of Reals)

For every $x, y \in \mathbb{R}$ with $x > 0$, there exists $n \in \mathbb{Z}_+$ with $y < nx$.

Archimedean property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{Z}_+$ with $n > x$.

Proof.

If not, then x is an upper bound of \mathbb{Z}_+ . □

Theorem (Archimedean Property of Reals)

For every $x, y \in \mathbb{R}$ with $x > 0$, there exists $n \in \mathbb{Z}_+$ with $y < nx$.

Proof.

Replace x with y/x in the previous theorem. □

Outline

- 1 Real Analysis Lecture 2
 - Types of real numbers
 - Integers
 - Upper Bound and Supremum
 - **Decimal expansions**

Decimal approximations

A **finite decimal expansion** is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \cdots + \frac{a_n}{10^n},$$

where $a_0 \in \mathbb{Z}_+$ and $0 \leq a_k \leq 9$ for $1 \leq k \leq n$.

Decimal approximations

A **finite decimal expansion** is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \cdots + \frac{a_n}{10^n},$$

where $a_0 \in \mathbb{Z}_+$ and $0 \leq a_k \leq 9$ for $1 \leq k \leq n$.

Notation:

$$a_0.a_1a_2a_3\ldots a_n.$$

Decimal approximations

A **finite decimal expansion** is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \cdots + \frac{a_n}{10^n},$$

where $a_0 \in \mathbb{Z}_+$ and $0 \leq a_k \leq 9$ for $1 \leq k \leq n$.

Notation:

$$a_0.a_1a_2a_3 \dots a_n.$$

Any positive real number $x > 0$ can be approximated by a finite decimal expansion.

Decimal approximations

Theorem (Apostol Theorem 1.20)

For any real $x > 0$ and $n \in \mathbb{Z}_+$, there exists a finite decimal expansion $r_n = a_0.a_1a_2 \dots a_n$ with

$$r_n \leq x < r_n + \frac{1}{10^n}.$$

Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set A is a set of integers which is bounded above by x , so it has a maximum a_0 .

Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set A is a set of integers which is bounded above by x , so it has a maximum a_0 .

Then clearly $x_1 = x - a_0 \in [0, 1)$.

Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set A is a set of integers which is bounded above by x , so it has a maximum a_0 .

Then clearly $x_1 = x - a_0 \in [0, 1)$.

Define a_1, a_2, a_3, \dots and x_1, x_2, x_3, \dots recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and $x_{k+1} = 10x_k - a_k$.

Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set A is a set of integers which is bounded above by x , so it has a maximum a_0 .

Then clearly $x_1 = x - a_0 \in [0, 1)$.

Define a_1, a_2, a_3, \dots and x_1, x_2, x_3, \dots recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and $x_{k+1} = 10x_k - a_k$. Then $0 \leq a_k \leq 9$ for all $k \geq 1$

Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set A is a set of integers which is bounded above by x , so it has a maximum a_0 .

Then clearly $x_1 = x - a_0 \in [0, 1)$.

Define a_1, a_2, a_3, \dots and x_1, x_2, x_3, \dots recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and $x_{k+1} = 10x_k - a_k$. Then $0 \leq a_k \leq 9$ for all $k \geq 1$ and

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n + 1}{10^n}$$

Decimal expansions

We say that $x > 0$ has the decimal expansion $a_0.a_1a_2a_3\dots$ and write

$$x = a_0.a_1a_2a_3a_4\dots$$

Decimal expansions

We say that $x > 0$ has the decimal expansion $a_0.a_1a_2a_3\dots$ and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Decimal expansions

We say that $x > 0$ has the decimal expansion $a_0.a_1a_2a_3\dots$ and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

Decimal expansions

We say that $x > 0$ has the decimal expansion $a_0.a_1a_2a_3\dots$ and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- 1 we haven't defined limits

Decimal expansions

We say that $x > 0$ has the decimal expansion $a_0.a_1a_2a_3\dots$ and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- 1 we haven't defined limits
- 2 with this definition, numbers have unique decimal expansions!

Decimal expansions

We say that $x > 0$ has the decimal expansion $a_0.a_1a_2a_3\dots$ and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- 1 we haven't defined limits
- 2 with this definition, numbers have unique decimal expansions!

$$1 \neq 0.99999999\dots$$

Decimal expansions

We say that $x > 0$ has the decimal expansion $a_0.a_1a_2a_3\dots$ and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- ① we haven't defined limits
- ② with this definition, numbers have unique decimal expansions!

$$1 \neq 0.999999999\dots$$

$$1 \not\leq 0 + \frac{9}{10} + \frac{9}{100} + \dots + \frac{9+1}{10^n} = 1.$$