

# MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics  
California State University Fullerton

August 25, 2024

# Outline


- 1 Real Analysis Lecture 2
  - Irrational numbers
  - Upper Bound and Supremum
  - Decimal expansions

# Outline

- 1 Real Analysis Lecture 2
  - Irrational numbers
  - Upper Bound and Supremum
  - Decimal expansions

# Types of numbers


## Transcendental Numbers



# Irrational numbers

Numbers which are not of the form  $a/b$  with  $a, b \in \mathbb{Z}$  are called **irrational**.

- Discovered by Hippasus, a pythagorean (a student of Pythagoras)



# Irrational numbers

Numbers which are not of the form  $a/b$  with  $a, b \in \mathbb{Z}$  are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery

# Irrational numbers

Numbers which are not of the form  $a/b$  with  $a, b \in \mathbb{Z}$  are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- dramatic reenactment, found online:

# Irrational numbers

Numbers which are not of the form  $a/b$  with  $a, b \in \mathbb{Z}$  are called **irrational**.


- Pythagoras drowned Hippasus at sea for his discovery
- dramatic reenactment, found online:



# Irrational numbers

Numbers which are not of the form  $a/b$  with  $a, b \in \mathbb{Z}$  are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- dramatic reenactment, found online:



# Irrational numbers

Numbers which are not of the form  $a/b$  with  $a, b \in \mathbb{Z}$  are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- anime style:

# Irrational numbers

Numbers which are not of the form  $a/b$  with  $a, b \in \mathbb{Z}$  are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- anime style:



# Challenge!

## Problem

Can you prove that if  $n \in \mathbb{Z}_+$  is not a perfect square, then  $\sqrt{n}$  is irrational? (Apostol Theorem 1.10)

# Challenge!

## Problem

Can you prove that if  $n \in \mathbb{Z}_+$  is not a perfect square, then  $\sqrt{n}$  is irrational? (Apostol Theorem 1.10)

## Solution

We prove by contradiction.

# Challenge!

## Problem

Can you prove that if  $n \in \mathbb{Z}_+$  is not a perfect square, then  $\sqrt{n}$  is irrational? (Apostol Theorem 1.10)

## Solution

We prove by contradiction. Assume  $\sqrt{n} = a/b$  for integers  $a, b$ .

# Challenge!

## Problem

Can you prove that if  $n \in \mathbb{Z}_+$  is not a perfect square, then  $\sqrt{n}$  is irrational? (Apostol Theorem 1.10)

## Solution

We prove by contradiction. Assume  $\sqrt{n} = a/b$  for integers  $a, b$ . Without loss of generality, we may assume  $\gcd(a, b) = 1$ .

# Challenge!

## Problem

Can you prove that if  $n \in \mathbb{Z}_+$  is not a perfect square, then  $\sqrt{n}$  is irrational? (Apostol Theorem 1.10)

## Solution

We prove by contradiction. Assume  $\sqrt{n} = a/b$  for integers  $a, b$ . Without loss of generality, we may assume  $\gcd(a, b) = 1$ . Then  $b^2 n = a^2$ .



# Challenge!

## Problem

Can you prove that if  $n \in \mathbb{Z}_+$  is not a perfect square, then  $\sqrt{n}$  is irrational? (Apostol Theorem 1.10)

## Solution

We prove by contradiction. Assume  $\sqrt{n} = a/b$  for integers  $a, b$ . Without loss of generality, we may assume  $\gcd(a, b) = 1$ . Then  $b^2 n = a^2$ . Since  $\gcd(a, b) = 1$ ,  $a^2$  must divide  $n$ , ie.  $n = n' a^2$ .

# Challenge!

## Problem

Can you prove that if  $n \in \mathbb{Z}_+$  is not a perfect square, then  $\sqrt{n}$  is irrational? (Apostol Theorem 1.10)

## Solution

We prove by contradiction. Assume  $\sqrt{n} = a/b$  for integers  $a, b$ . Without loss of generality, we may assume  $\gcd(a, b) = 1$ . Then  $b^2 n = a^2$ . Since  $\gcd(a, b) = 1$ ,  $a^2$  must divide  $n$ , ie.  $n = n' a^2$ . It follows that  $b^2 n' = 1$ , so  $b^2$  divides 1.

# Challenge!

## Problem

Can you prove that if  $n \in \mathbb{Z}_+$  is not a perfect square, then  $\sqrt{n}$  is irrational? (Apostol Theorem 1.10)

## Solution

We prove by contradiction. Assume  $\sqrt{n} = a/b$  for integers  $a, b$ . Without loss of generality, we may assume  $\gcd(a, b) = 1$ . Then  $b^2 n = a^2$ . Since  $\gcd(a, b) = 1$ ,  $a^2$  must divide  $n$ , ie.  $n = n' a^2$ . It follows that  $b^2 n' = 1$ , so  $b^2$  divides 1. This implies  $n'$  divides 1, so  $n' = \pm 1$ .

# Challenge!

## Problem

Can you prove that if  $n \in \mathbb{Z}_+$  is not a perfect square, then  $\sqrt{n}$  is irrational? (Apostol Theorem 1.10)

## Solution

We prove by contradiction. Assume  $\sqrt{n} = a/b$  for integers  $a, b$ . Without loss of generality, we may assume  $\gcd(a, b) = 1$ . Then  $b^2 n = a^2$ . Since  $\gcd(a, b) = 1$ ,  $a^2$  must divide  $n$ , ie.  $n = n' a^2$ . It follows that  $b^2 n' = 1$ , so  $b^2$  divides 1. This implies  $n'$  divides 1, so  $n' = \pm 1$ . Since  $n > 0$ ,  $n' > 0$  so  $n' = 1$ .

# Challenge!

## Problem

Can you prove that if  $n \in \mathbb{Z}_+$  is not a perfect square, then  $\sqrt{n}$  is irrational? (Apostol Theorem 1.10)

## Solution

We prove by contradiction. Assume  $\sqrt{n} = a/b$  for integers  $a, b$ . Without loss of generality, we may assume  $\gcd(a, b) = 1$ . Then  $b^2 n = a^2$ . Since  $\gcd(a, b) = 1$ ,  $a^2$  must divide  $n$ , ie.  $n = n' a^2$ . It follows that  $b^2 n' = 1$ , so  $b^2$  divides 1. This implies  $n'$  divides 1, so  $n' = \pm 1$ . Since  $n > 0$ ,  $n' > 0$  so  $n' = 1$ . Thus  $n = a^2$ , which is a contradiction.

# Algebraic and Transcendental

Irrational numbers can be further divided into to categories

# Algebraic and Transcendental

Irrational numbers can be further divided into to categories

- 1 **algebraic numbers:** numbers which are roots of polynomials with integer coefficients, like

$$\sqrt{2}, \quad \sqrt{3}, \quad \text{and} \quad \sqrt[3]{\sqrt{5} + \sqrt{7}}.$$

# Algebraic and Transcendental

Irrational numbers can be further divided into to categories

- 1 **algebraic numbers:** numbers which are roots of polynomials with integer coefficients, like

$$\sqrt{2}, \quad \sqrt{3}, \quad \text{and} \quad \sqrt[3]{\sqrt{5} + \sqrt{7}}.$$

- 2 **transcendental numbers:** numbers which are not algebraic, like

$$\pi, \quad e, \quad e^{\pi}, \quad \text{and maybe} \quad \pi + e?$$



# Algebraic and Transcendental

Irrational numbers can be further divided into to categories

- 1 **algebraic numbers:** numbers which are roots of polynomials with integer coefficients, like

$$\sqrt{2}, \quad \sqrt{3}, \quad \text{and} \quad \sqrt[3]{\sqrt{5} + \sqrt{7}}.$$

- 2 **transcendental numbers:** numbers which are not algebraic, like

$$\pi, \quad e, \quad e^{\pi}, \quad \text{and maybe} \quad \pi + e?$$

- 3 transcendentals are mysterious ... but most real numbers are transcendental!

# Outline

- 1 Real Analysis Lecture 2
  - Irrational numbers
  - Upper Bound and Supremum
  - Decimal expansions

# Upper bounds

An **upper bound** for a set  $S \subseteq \mathbb{R}$  is a number  $b$  such that

$$x \leq b \quad \text{for all } x \in S.$$

# Upper bounds

An **upper bound** for a set  $S \subseteq \mathbb{R}$  is a number  $b$  such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say  $S$  is **bounded above** by  $b$ .

# Upper bounds

An **upper bound** for a set  $S \subseteq \mathbb{R}$  is a number  $b$  such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say  $S$  is **bounded above** by  $b$ .

If  $b \in S$  also, then  $b$  is called a **maximal element** of  $S$

# Upper bounds

An **upper bound** for a set  $S \subseteq \mathbb{R}$  is a number  $b$  such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say  $S$  is **bounded above** by  $b$ .

If  $b \in S$  also, then  $b$  is called a **maximal element** of  $S$

**Examples:**

# Upper bounds

An **upper bound** for a set  $S \subseteq \mathbb{R}$  is a number  $b$  such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say  $S$  is **bounded above** by  $b$ .

If  $b \in S$  also, then  $b$  is called a **maximal element** of  $S$

## Examples:

- 34 is an upper bound of  $[-1, 5]$ , but not a maximal element

# Upper bounds

An **upper bound** for a set  $S \subseteq \mathbb{R}$  is a number  $b$  such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say  $S$  is **bounded above** by  $b$ .

If  $b \in S$  also, then  $b$  is called a **maximal element** of  $S$

## Examples:

- 34 is an upper bound of  $[-1, 5]$ , but not a maximal element
- 5 is a maximal element of  $[-1, 5]$



# Upper bounds

An **upper bound** for a set  $S \subseteq \mathbb{R}$  is a number  $b$  such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say  $S$  is **bounded above** by  $b$ .

If  $b \in S$  also, then  $b$  is called a **maximal element** of  $S$

## Examples:

- 34 is an upper bound of  $[-1, 5]$ , but not a maximal element
- 5 is a maximal element of  $[-1, 5]$
- 3 is an upper bound of  $[0, 3]$ , but not a maximal element

# Upper bounds

An **upper bound** for a set  $S \subseteq \mathbb{R}$  is a number  $b$  such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say  $S$  is **bounded above** by  $b$ .

If  $b \in S$  also, then  $b$  is called a **maximal element** of  $S$

## Examples:

- 34 is an upper bound of  $[-1, 5]$ , but not a maximal element
- 5 is a maximal element of  $[-1, 5]$
- 3 is an upper bound of  $[0, 3)$ , but not a maximal element
- $\mathbb{Z}_+$  has no upper bound

# Upper bounds

An **upper bound** for a set  $S \subseteq \mathbb{R}$  is a number  $b$  such that

$$x \leq b \quad \text{for all } x \in S.$$

In this case, we say  $S$  is **bounded above** by  $b$ .

If  $b \in S$  also, then  $b$  is called a **maximal element** of  $S$

## Examples:

- 34 is an upper bound of  $[-1, 5]$ , but not a maximal element
- 5 is a maximal element of  $[-1, 5]$
- 3 is an upper bound of  $[0, 3)$ , but not a maximal element
- $\mathbb{Z}_+$  has no upper bound
- $[3, 7)$  has an upper bound but no maximal element

# Challenge!

## Problem

Show that if  $S$  has a maximal element, then it is unique.

# Challenge!

## Problem

Show that if  $S$  has a maximal element, then it is unique.

**Hint: use the definition!**

# Challenge!

## Problem

Show that if  $S$  has a maximal element, then it is unique.

**Hint: use the definition!**

## Solution

Suppose that  $b_1$  and  $b_2$  are both maximal elements of  $S$ .

# Challenge!

## Problem

Show that if  $S$  has a maximal element, then it is unique.

**Hint: use the definition!**

## Solution

Suppose that  $b_1$  and  $b_2$  are both maximal elements of  $S$ .  
Then  $x \leq b_1$  and  $x \leq b_2$  for all  $x \in S$ .

# Challenge!

## Problem

Show that if  $S$  has a maximal element, then it is unique.

**Hint: use the definition!**

## Solution

Suppose that  $b_1$  and  $b_2$  are both maximal elements of  $S$ .

Then  $x \leq b_1$  and  $x \leq b_2$  for all  $x \in S$ .

Moreover,  $b_1 \in S$  and  $b_2 \in S$ .



# Challenge!

## Problem

Show that if  $S$  has a maximal element, then it is unique.

**Hint: use the definition!**

## Solution

Suppose that  $b_1$  and  $b_2$  are both maximal elements of  $S$ .

Then  $x \leq b_1$  and  $x \leq b_2$  for all  $x \in S$ .

Moreover,  $b_1 \in S$  and  $b_2 \in S$ .

Since  $b_1 \in S$  and  $b_2$  is an upper bound of  $S$ ,  $b_1 \leq b_2$ .

# Challenge!

## Problem

Show that if  $S$  has a maximal element, then it is unique.

**Hint: use the definition!**

## Solution

Suppose that  $b_1$  and  $b_2$  are both maximal elements of  $S$ .

Then  $x \leq b_1$  and  $x \leq b_2$  for all  $x \in S$ .

Moreover,  $b_1 \in S$  and  $b_2 \in S$ .

Since  $b_1 \in S$  and  $b_2$  is an upper bound of  $S$ ,  $b_1 \leq b_2$ .

Likewise, since  $b_2 \in S$  and  $b_1$  is an upper bound of  $S$ ,  $b_2 \leq b_1$ .

# Challenge!

## Problem

Show that if  $S$  has a maximal element, then it is unique.

**Hint: use the definition!**

## Solution

Suppose that  $b_1$  and  $b_2$  are both maximal elements of  $S$ .

Then  $x \leq b_1$  and  $x \leq b_2$  for all  $x \in S$ .

Moreover,  $b_1 \in S$  and  $b_2 \in S$ .

Since  $b_1 \in S$  and  $b_2$  is an upper bound of  $S$ ,  $b_1 \leq b_2$ .

Likewise, since  $b_2 \in S$  and  $b_1$  is an upper bound of  $S$ ,  $b_2 \leq b_1$ .

By the trichotomy, we find  $b_1 = b_2$ .

# Challenge!

## Problem

Show that if  $S$  has a maximal element, then it is unique.

**Hint: use the definition!**

## Solution

Suppose that  $b_1$  and  $b_2$  are both maximal elements of  $S$ .

Then  $x \leq b_1$  and  $x \leq b_2$  for all  $x \in S$ .

Moreover,  $b_1 \in S$  and  $b_2 \in S$ .

Since  $b_1 \in S$  and  $b_2$  is an upper bound of  $S$ ,  $b_1 \leq b_2$ .

Likewise, since  $b_2 \in S$  and  $b_1$  is an upper bound of  $S$ ,  $b_2 \leq b_1$ .

By the trichotomy, we find  $b_1 = b_2$ .

Now we can say *the* maximum,  $\max(S)$

# Supremum

A **supremum** of a set  $S$  of real numbers is a real number  $b \in \mathbb{R}$  such that

- $b$  is an upper bound of  $S$
- if  $b' < b$ , then  $b'$  is not an upper bound of  $S$

# Supremum

A **supremum** of a set  $S$  of real numbers is a real number  $b \in \mathbb{R}$  such that

- $b$  is an upper bound of  $S$
- if  $b' < b$ , then  $b'$  is not an upper bound of  $S$

In other words

a supremum is a **least upper bound**

# Challenge!

## Problem

Show that if  $S$  has a supremum, then it is unique.

# Challenge!

## Problem

Show that if  $S$  has a supremum, then it is unique.

**Hint: use the definition!**



# Challenge!

## Problem

Show that if  $S$  has a supremum, then it is unique.

**Hint: use the definition!**

## Solution

Suppose that  $b_1$  and  $b_2$  are both suprema of  $S$ .

# Challenge!

## Problem

Show that if  $S$  has a supremum, then it is unique.

**Hint: use the definition!**

## Solution

Suppose that  $b_1$  and  $b_2$  are both suprema of  $S$ .  
Then  $b_1$  and  $b_2$  are both upper bounds of  $S$ .

# Challenge!

## Problem

Show that if  $S$  has a supremum, then it is unique.

**Hint: use the definition!**

## Solution

Suppose that  $b_1$  and  $b_2$  are both suprema of  $S$ .  
Then  $b_1$  and  $b_2$  are both upper bounds of  $S$ .  
Since  $b_1$  is a least upper bound,  $b_1 \leq b_2$ .

# Challenge!

## Problem

Show that if  $S$  has a supremum, then it is unique.

**Hint: use the definition!**

## Solution

Suppose that  $b_1$  and  $b_2$  are both suprema of  $S$ .

Then  $b_1$  and  $b_2$  are both upper bounds of  $S$ .

Since  $b_1$  is a least upper bound,  $b_1 \leq b_2$ .

Since  $b_2$  is also a least upper bound,  $b_2 \leq b_1$ .

# Challenge!

## Problem

Show that if  $S$  has a supremum, then it is unique.

**Hint: use the definition!**

## Solution

Suppose that  $b_1$  and  $b_2$  are both suprema of  $S$ .

Then  $b_1$  and  $b_2$  are both upper bounds of  $S$ .

Since  $b_1$  is a least upper bound,  $b_1 \leq b_2$ .

Since  $b_2$  is also a least upper bound,  $b_2 \leq b_1$ .

By the trichotomy, we find  $b_1 = b_2$ .

# Challenge!

## Problem

Show that if  $S$  has a supremum, then it is unique.

**Hint: use the definition!**

## Solution

Suppose that  $b_1$  and  $b_2$  are both suprema of  $S$ .

Then  $b_1$  and  $b_2$  are both upper bounds of  $S$ .

Since  $b_1$  is a least upper bound,  $b_1 \leq b_2$ .

Since  $b_2$  is also a least upper bound,  $b_2 \leq b_1$ .

By the trichotomy, we find  $b_1 = b_2$ .

Now we can say *the* supremum,  $\sup(S)$

# Challenge!

## Problem

Show that if  $S$  has a maximum element. Then  $S$  has a supremum and  $\max(S) = \sup(S)$

# Challenge!

## Problem

Show that if  $S$  has a maximum element. Then  $S$  has a supremum and  $\max(S) = \sup(S)$

**Hint: use the definition!**



# Challenge!

## Problem

Show that if  $S$  has a maximum element. Then  $S$  has a supremum and  $\max(S) = \sup(S)$

**Hint: use the definition!**

## Solution

Let  $b = \max(S)$ .

# Challenge!

## Problem

Show that if  $S$  has a maximum element. Then  $S$  has a supremum and  $\max(S) = \sup(S)$

**Hint: use the definition!**

## Solution

Let  $b = \max(S)$ .

Then  $b$  is an upper bound of  $S$  and  $b \in S$ .

# Challenge!

## Problem

Show that if  $S$  has a maximum element. Then  $S$  has a supremum and  $\max(S) = \sup(S)$

**Hint: use the definition!**

## Solution

Let  $b = \max(S)$ .

Then  $b$  is an upper bound of  $S$  and  $b \in S$ .

If  $b'$  is another upper bound of  $S$ , then by definition  $b \leq b'$ .

# Challenge!

## Problem

Show that if  $S$  has a maximum element. Then  $S$  has a supremum and  $\max(S) = \sup(S)$

**Hint: use the definition!**

## Solution

Let  $b = \max(S)$ .

Then  $b$  is an upper bound of  $S$  and  $b \in S$ .

If  $b'$  is another upper bound of  $S$ , then by definition  $b \leq b'$ .

Thus  $b$  is the least upper bound of  $S$ .

# Completeness Axiom


Now we have the machinery in place to state the Completeness Axiom.

# Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.


# Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.

 **completeness axiom:** if  $S$  is any subset of real numbers which is bounded above, then it has a supremum  $\sup(S)$

# Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.


 **completeness axiom:** if  $S$  is any subset of real numbers which is bounded above, then it has a supremum  $\sup(S)$

**Examples:**



# Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.

 **completeness axiom:** if  $S$  is any subset of real numbers which is bounded above, then it has a supremum  $\sup(S)$

**Examples:**

- 3 is the supremum of  $(0, 3)$

# Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.

**A10 completeness axiom:** if  $S$  is any subset of real numbers which is bounded above, then it has a supremum  $\sup(S)$

## Examples:

- 3 is the supremum of  $(0, 3)$
- 1 is the supremum of

$$\left\{ \frac{n}{n+1} : n \in \mathbb{Z}_+ \right\}$$

# Completeness Axiom

Now we have the machinery in place to state the Completeness Axiom.

**completeness axiom:** if  $S$  is any subset of real numbers which is bounded above, then it has a supremum  $\sup(S)$

## Examples:

- 3 is the supremum of  $(0, 3)$
- 1 is the supremum of

$$\left\{ \frac{n}{n+1} : n \in \mathbb{Z}_+ \right\}$$

- $\pi$  is the supremum of

$$\{3, 3.1, 3.14, 3.141, 3.1415, 3.14159, 3.141592, \dots\}.$$

# Challenge

## Problem

Prove that if  $A$  is a set of integers that is bounded above, then  $A$  has a maximum.

# Challenge

## Solution

The set  $A$  is bounded above, so it has a supremum  $b = \sup(A)$ .

# Challenge

## Solution

The set  $A$  is bounded above, so it has a supremum  $b = \sup(A)$ . Since  $b - 1$  is not an upper bound, so there exists  $a \in A$  with  $b - 1 < a$ .

# Challenge

## Solution

The set  $A$  is bounded above, so it has a supremum  $b = \sup(A)$ . Since  $b - 1$  is not an upper bound, so there exists  $a \in A$  with  $b - 1 < a$ .

If  $a' \in A$ , then  $a'$  is an integer and therefore  $a' = a + k$  for  $k \in \mathbb{Z}$ .

# Challenge

## Solution

The set  $A$  is bounded above, so it has a supremum  $b = \sup(A)$ . Since  $b - 1$  is not an upper bound, so there exists  $a \in A$  with  $b - 1 < a$ .

If  $a' \in A$ , then  $a'$  is an integer and therefore  $a' = a + k$  for  $k \in \mathbb{Z}$ . Since  $b$  is an upper bound,  $b \geq a + k > b - 1 + k$ , making  $0 > k - 1$ .



# Challenge

## Solution

The set  $A$  is bounded above, so it has a supremum  $b = \sup(A)$ . Since  $b - 1$  is not an upper bound, so there exists  $a \in A$  with  $b - 1 < a$ .

If  $a' \in A$ , then  $a'$  is an integer and therefore  $a' = a + k$  for  $k \in \mathbb{Z}$ . Since  $b$  is an upper bound,  $b \geq a + k > b - 1 + k$ , making  $0 > k - 1$ .

Therefore  $k \leq 0$  and  $a' \leq a$ .

# Challenge

## Solution

The set  $A$  is bounded above, so it has a supremum  $b = \sup(A)$ . Since  $b - 1$  is not an upper bound, so there exists  $a \in A$  with  $b - 1 < a$ .

If  $a' \in A$ , then  $a'$  is an integer and therefore  $a' = a + k$  for  $k \in \mathbb{Z}$ . Since  $b$  is an upper bound,  $b \geq a + k > b - 1 + k$ , making  $0 > k - 1$ .

Therefore  $k \leq 0$  and  $a' \leq a$ .

It follows that  $a$  is an upper bound of  $A$ , and since  $a \in A$  it is a maximum.

# Lower bounds and infima

A **lower bound** for a set  $S \subseteq \mathbb{R}$  is a number  $b$  such that

$$b \leq x \quad \text{for all } x \in S.$$

# Lower bounds and infima

A **lower bound** for a set  $S \subseteq \mathbb{R}$  is a number  $b$  such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say  $S$  is **bounded below** by  $b$ .

# Lower bounds and infima

A **lower bound** for a set  $S \subseteq \mathbb{R}$  is a number  $b$  such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say  $S$  is **bounded below** by  $b$ .

If  $b \in S$  also, then  $b$  is called a **minimal element** of  $S$ .

# Lower bounds and infima

A **lower bound** for a set  $S \subseteq \mathbb{R}$  is a number  $b$  such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say  $S$  is **bounded below** by  $b$ .

If  $b \in S$  also, then  $b$  is called a **minimal element** of  $S$ . An **infimum** of a set  $S$  of real numbers is a real number  $b \in \mathbb{R}$  such that

- $b$  is a lower bound of  $S$
- if  $b < b'$ , then  $b'$  is not a lower bound of  $S$

# Lower bounds and infima

A **lower bound** for a set  $S \subseteq \mathbb{R}$  is a number  $b$  such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say  $S$  is **bounded below** by  $b$ .

If  $b \in S$  also, then  $b$  is called a **minimal element** of  $S$ . An **infimum** of a set  $S$  of real numbers is a real number  $b \in \mathbb{R}$  such that

- $b$  is a lower bound of  $S$
- if  $b < b'$ , then  $b'$  is not a lower bound of  $S$

In other words

an infimum is a **greatest lower bound**

# Properties of Suprema

The first property of suprema is that they must be arbitrarily close to elements of the set.

## Theorem (Approximation Property)

*Let  $S \subseteq \mathbb{R}$  be bounded above, and let  $b = \sup(S)$ . Then for all  $\epsilon > 0$ , there exists  $a \in S$  with*

$$b - \epsilon < a < b.$$



# Properties of Suprema

Proof.

Since  $b - \epsilon < b$ , the definition of a supremum implies  $b - \epsilon$  cannot be an upper bound.

# Properties of Suprema

Proof.

Since  $b - \epsilon < b$ , the definition of a supremum implies  $b - \epsilon$  cannot be an upper bound.

Therefore there must exist  $a \in S$  with  $a > b - \epsilon$ . □

# Important example: The number $e$

For each  $n$ , let  $s_n$  denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

# Important example: The number $e$

For each  $n$ , let  $s_n$  denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

# Important example: The number $e$

For each  $n$ , let  $s_n$  denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Consider the set

$$S = \{s_n : n \in \mathbb{Z}_+\}.$$

# Important example: The number $e$

For each  $n$ , let  $s_n$  denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Consider the set

$$S = \{s_n : n \in \mathbb{Z}_+\}.$$

For every  $n$ ,

$$s_n \leq 1 + 1 + \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \cdots + \frac{1}{2^n} = 3 - \frac{1}{2^n} < 3$$

so  $S$  is bounded above by 3.

# Important example: The number $e$

For each  $n$ , let  $s_n$  denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Consider the set

$$S = \{s_n : n \in \mathbb{Z}_+\}.$$

For every  $n$ ,

$$s_n \leq 1 + 1 + \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \cdots + \frac{1}{2^n} = 3 - \frac{1}{2^n} < 3$$

so  $S$  is bounded above by 3. Therefore  $S$  has a supremum,  $\sup(S)$ .

# Important example: The number $e$

Even if  $\epsilon$  is very small ( $\epsilon = 0.000000001$ ),  $\sup(S) - \epsilon$  is not an upper bound of  $S$ .



# Important example: The number $e$

Even if  $\epsilon$  is very small ( $\epsilon = 0.000000001$ ),  $\sup(S) - \epsilon$  is not an upper bound of  $S$ .

So there exists  $N$  with  $s_N > \sup(S) - \epsilon$ .

# Important example: The number $e$

Even if  $\epsilon$  is very small ( $\epsilon = 0.000000001$ ),  $\sup(S) - \epsilon$  is not an upper bound of  $S$ .

So there exists  $N$  with  $s_N > \sup(S) - \epsilon$ .

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

# Important example: The number $e$

Even if  $\epsilon$  is very small ( $\epsilon = 0.000000001$ ),  $\sup(S) - \epsilon$  is not an upper bound of  $S$ .

So there exists  $N$  with  $s_N > \sup(S) - \epsilon$ .

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says  $\sup(S)$  is *really* close to  $s_N$ .

# Important example: The number $e$

Even if  $\epsilon$  is very small ( $\epsilon = 0.000000001$ ),  $\sup(S) - \epsilon$  is not an upper bound of  $S$ .

So there exists  $N$  with  $s_N > \sup(S) - \epsilon$ .

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says  $\sup(S)$  is *really* close to  $s_N$ . Taking  $\epsilon$  smaller and smaller

# Important example: The number $e$

Even if  $\epsilon$  is very small ( $\epsilon = 0.000000001$ ),  $\sup(S) - \epsilon$  is not an upper bound of  $S$ .

So there exists  $N$  with  $s_N > \sup(S) - \epsilon$ .

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says  $\sup(S)$  is *really* close to  $s_N$ . Taking  $\epsilon$  smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

# Important example: The number $e$

Even if  $\epsilon$  is very small ( $\epsilon = 0.000000001$ ),  $\sup(S) - \epsilon$  is not an upper bound of  $S$ .

So there exists  $N$  with  $s_N > \sup(S) - \epsilon$ .

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says  $\sup(S)$  is *really* close to  $s_N$ . Taking  $\epsilon$  smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

If we remember Taylor series

# Important example: The number $e$

Even if  $\epsilon$  is very small ( $\epsilon = 0.000000001$ ),  $\sup(S) - \epsilon$  is not an upper bound of  $S$ .

So there exists  $N$  with  $s_N > \sup(S) - \epsilon$ .

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says  $\sup(S)$  is *really* close to  $s_N$ . Taking  $\epsilon$  smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

If we remember Taylor series

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots$$

# Important example: The number $e$

Even if  $\epsilon$  is very small ( $\epsilon = 0.000000001$ ),  $\sup(S) - \epsilon$  is not an upper bound of  $S$ .

So there exists  $N$  with  $s_N > \sup(S) - \epsilon$ .

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says  $\sup(S)$  is *really* close to  $s_N$ . Taking  $\epsilon$  smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

If we remember Taylor series

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots$$

We've just shown that  $e^1 = e$  exists.



# Properties of Suprema

Also, suprema play nicely with addition.

## Theorem (Additive Property)

*Let  $A, B \subseteq \mathbb{R}$  be bounded above set*

$$C = \{x + y : x \in A, y \in B\}.$$

*Then  $C$  is bounded above and*

$$\sup(C) = \sup(A) + \sup(B).$$

# Properties of Suprema

Proof.

Let  $a = \sup(A)$ ,  $b = \sup(B)$ , and  $c = \sup(C)$ .

# Properties of Suprema

Proof.

Let  $a = \sup(A)$ ,  $b = \sup(B)$ , and  $c = \sup(C)$ .

We will prove  $c \leq a + b$  and then  $a + b \leq c$ .

# Properties of Suprema

## Proof.

Let  $a = \sup(A)$ ,  $b = \sup(B)$ , and  $c = \sup(C)$ .

We will prove  $c \leq a + b$  and then  $a + b \leq c$ .

First, note  $a$  is an upper bound of  $A$ , so  $x \leq a$  for all  $x \in A$ .

# Properties of Suprema

## Proof.

Let  $a = \sup(A)$ ,  $b = \sup(B)$ , and  $c = \sup(C)$ .

We will prove  $c \leq a + b$  and then  $a + b \leq c$ .

First, note  $a$  is an upper bound of  $A$ , so  $x \leq a$  for all  $x \in A$ .

Also  $b$  is an upper bound of  $B$ , so  $y \leq b$  for all  $y \in B$ .

# Properties of Suprema

## Proof.

Let  $a = \sup(A)$ ,  $b = \sup(B)$ , and  $c = \sup(C)$ .

We will prove  $c \leq a + b$  and then  $a + b \leq c$ .

First, note  $a$  is an upper bound of  $A$ , so  $x \leq a$  for all  $x \in A$ .

Also  $b$  is an upper bound of  $B$ , so  $y \leq b$  for all  $y \in B$ .

If  $z \in C$ , then  $z = x + y$  for some  $x \in A$  and  $y \in B$ , and therefore  $z = x + y \leq a + b$ .

# Properties of Suprema

## Proof.

Let  $a = \sup(A)$ ,  $b = \sup(B)$ , and  $c = \sup(C)$ .

We will prove  $c \leq a + b$  and then  $a + b \leq c$ .

First, note  $a$  is an upper bound of  $A$ , so  $x \leq a$  for all  $x \in A$ .

Also  $b$  is an upper bound of  $B$ , so  $y \leq b$  for all  $y \in B$ .

If  $z \in C$ , then  $z = x + y$  for some  $x \in A$  and  $y \in B$ , and therefore  $z = x + y \leq a + b$ .

Therefore  $a + b$  is an upper bound of  $C$ .

# Properties of Suprema

## Proof.

Let  $a = \sup(A)$ ,  $b = \sup(B)$ , and  $c = \sup(C)$ .

We will prove  $c \leq a + b$  and then  $a + b \leq c$ .

First, note  $a$  is an upper bound of  $A$ , so  $x \leq a$  for all  $x \in A$ .

Also  $b$  is an upper bound of  $B$ , so  $y \leq b$  for all  $y \in B$ .

If  $z \in C$ , then  $z = x + y$  for some  $x \in A$  and  $y \in B$ , and therefore  $z = x + y \leq a + b$ .

Therefore  $a + b$  is an upper bound of  $C$ .

This means  $c \leq a + b$ .



# Properties of Suprema

## Proof.

Let  $a = \sup(A)$ ,  $b = \sup(B)$ , and  $c = \sup(C)$ .

We will prove  $c \leq a + b$  and then  $a + b \leq c$ .

First, note  $a$  is an upper bound of  $A$ , so  $x \leq a$  for all  $x \in A$ .

Also  $b$  is an upper bound of  $B$ , so  $y \leq b$  for all  $y \in B$ .

If  $z \in C$ , then  $z = x + y$  for some  $x \in A$  and  $y \in B$ , and therefore  $z = x + y \leq a + b$ .

Therefore  $a + b$  is an upper bound of  $C$ .

This means  $c \leq a + b$ .

Next, note for all  $x \in A$  and  $y \in B$  that  $x \leq c - y$ .

# Properties of Suprema

## Proof.

Let  $a = \sup(A)$ ,  $b = \sup(B)$ , and  $c = \sup(C)$ .

We will prove  $c \leq a + b$  and then  $a + b \leq c$ .

First, note  $a$  is an upper bound of  $A$ , so  $x \leq a$  for all  $x \in A$ .

Also  $b$  is an upper bound of  $B$ , so  $y \leq b$  for all  $y \in B$ .

If  $z \in C$ , then  $z = x + y$  for some  $x \in A$  and  $y \in B$ , and therefore  $z = x + y \leq a + b$ .

Therefore  $a + b$  is an upper bound of  $C$ .

This means  $c \leq a + b$ .

Next, note for all  $x \in A$  and  $y \in B$  that  $x \leq c - y$ .

Therefore  $c - y$  is an upper bound of  $A$  and  $a \leq c - y$ .

# Properties of Suprema

## Proof.

Let  $a = \sup(A)$ ,  $b = \sup(B)$ , and  $c = \sup(C)$ .

We will prove  $c \leq a + b$  and then  $a + b \leq c$ .

First, note  $a$  is an upper bound of  $A$ , so  $x \leq a$  for all  $x \in A$ .

Also  $b$  is an upper bound of  $B$ , so  $y \leq b$  for all  $y \in B$ .

If  $z \in C$ , then  $z = x + y$  for some  $x \in A$  and  $y \in B$ , and therefore  $z = x + y \leq a + b$ .

Therefore  $a + b$  is an upper bound of  $C$ .

This means  $c \leq a + b$ .

Next, note for all  $x \in A$  and  $y \in B$  that  $x \leq c - y$ .

Therefore  $c - y$  is an upper bound of  $A$  and  $a \leq c - y$ .

It follows that  $y \leq c - a$  for all  $y \in B$ .

# Properties of Suprema

## Proof.

Let  $a = \sup(A)$ ,  $b = \sup(B)$ , and  $c = \sup(C)$ .

We will prove  $c \leq a + b$  and then  $a + b \leq c$ .

First, note  $a$  is an upper bound of  $A$ , so  $x \leq a$  for all  $x \in A$ .

Also  $b$  is an upper bound of  $B$ , so  $y \leq b$  for all  $y \in B$ .

If  $z \in C$ , then  $z = x + y$  for some  $x \in A$  and  $y \in B$ , and therefore  $z = x + y \leq a + b$ .

Therefore  $a + b$  is an upper bound of  $C$ .

This means  $c \leq a + b$ .

Next, note for all  $x \in A$  and  $y \in B$  that  $x \leq c - y$ .

Therefore  $c - y$  is an upper bound of  $A$  and  $a \leq c - y$ .

It follows that  $y \leq c - a$  for all  $y \in B$ .

Thus  $c - a$  is an upper bound of  $B$ , and it follows  $b \leq c - a$ .

# Properties of Suprema

## Proof.

Let  $a = \sup(A)$ ,  $b = \sup(B)$ , and  $c = \sup(C)$ .

We will prove  $c \leq a + b$  and then  $a + b \leq c$ .

First, note  $a$  is an upper bound of  $A$ , so  $x \leq a$  for all  $x \in A$ .

Also  $b$  is an upper bound of  $B$ , so  $y \leq b$  for all  $y \in B$ .

If  $z \in C$ , then  $z = x + y$  for some  $x \in A$  and  $y \in B$ , and therefore  $z = x + y \leq a + b$ .

Therefore  $a + b$  is an upper bound of  $C$ .

This means  $c \leq a + b$ .

Next, note for all  $x \in A$  and  $y \in B$  that  $x \leq c - y$ .

Therefore  $c - y$  is an upper bound of  $A$  and  $a \leq c - y$ .

It follows that  $y \leq c - a$  for all  $y \in B$ .

Thus  $c - a$  is an upper bound of  $B$ , and it follows  $b \leq c - a$ .

Therefore  $a + b \leq c$ . □

# Challenge

## Problem

Use the completeness axiom to prove that the  $\mathbb{Z}_+$  is not bounded above. (Apostol Theorem 1.17)

# Challenge

## Problem

Use the completeness axiom to prove that the  $\mathbb{Z}_+$  is not bounded above. (Apostol Theorem 1.17)

**Hint:** assume it is and consider  $\sup(\mathbb{Z}_+) - 1$

# Challenge

## Problem

Use the completeness axiom to prove that the  $\mathbb{Z}_+$  is not bounded above. (Apostol Theorem 1.17)

**Hint:** assume it is and consider  $\sup(\mathbb{Z}_+) - 1$

## Solution

Assume it is.



# Challenge

## Problem

Use the completeness axiom to prove that the  $\mathbb{Z}_+$  is not bounded above. (Apostol Theorem 1.17)

**Hint:** assume it is and consider  $\sup(\mathbb{Z}_+) - 1$

## Solution

Assume it is.

The completeness axiom implies that  $b = \sup(\mathbb{Z}_+)$  exists

# Challenge

## Problem

Use the completeness axiom to prove that the  $\mathbb{Z}_+$  is not bounded above. (Apostol Theorem 1.17)

**Hint:** assume it is and consider  $\sup(\mathbb{Z}_+) - 1$

## Solution

Assume it is.

The completeness axiom implies that  $b = \sup(\mathbb{Z}_+)$  exists

The number  $b$  is an upper bound and if  $b' < b$ , then  $b'$  is not.

# Challenge

## Problem

Use the completeness axiom to prove that the  $\mathbb{Z}_+$  is not bounded above. (Apostol Theorem 1.17)

**Hint:** assume it is and consider  $\sup(\mathbb{Z}_+) - 1$

## Solution

Assume it is.

The completeness axiom implies that  $b = \sup(\mathbb{Z}_+)$  exists

The number  $b$  is an upper bound and if  $b' < b$ , then  $b'$  is not.

Then  $b - 1 < b$  by Axiom 7, so  $b - 1$  cannot be an upper bound.

# Challenge

## Problem

Use the completeness axiom to prove that the  $\mathbb{Z}_+$  is not bounded above. (Apostol Theorem 1.17)

**Hint:** assume it is and consider  $\sup(\mathbb{Z}_+) - 1$

## Solution

Assume it is.

The completeness axiom implies that  $b = \sup(\mathbb{Z}_+)$  exists

The number  $b$  is an upper bound and if  $b' < b$ , then  $b'$  is not.

Then  $b - 1 < b$  by Axiom 7, so  $b - 1$  cannot be an upper bound.

This means there exists  $n \in \mathbb{Z}_+$  with  $b - 1 < n$ .

# Challenge

## Problem

Use the completeness axiom to prove that the  $\mathbb{Z}_+$  is not bounded above. (Apostol Theorem 1.17)

**Hint:** assume it is and consider  $\sup(\mathbb{Z}_+) - 1$

## Solution

Assume it is.

The completeness axiom implies that  $b = \sup(\mathbb{Z}_+)$  exists

The number  $b$  is an upper bound and if  $b' < b$ , then  $b'$  is not.

Then  $b - 1 < b$  by Axiom 7, so  $b - 1$  cannot be an upper bound.

This means there exists  $n \in \mathbb{Z}_+$  with  $b - 1 < n$ .

It follows from Axiom 7 that  $b < n + 1$ .

# Challenge

## Problem

Use the completeness axiom to prove that the  $\mathbb{Z}_+$  is not bounded above. (Apostol Theorem 1.17)

**Hint:** assume it is and consider  $\sup(\mathbb{Z}_+) - 1$

## Solution

Assume it is.

The completeness axiom implies that  $b = \sup(\mathbb{Z}_+)$  exists

The number  $b$  is an upper bound and if  $b' < b$ , then  $b'$  is not.

Then  $b - 1 < b$  by Axiom 7, so  $b - 1$  cannot be an upper bound.

This means there exists  $n \in \mathbb{Z}_+$  with  $b - 1 < n$ .

It follows from Axiom 7 that  $b < n + 1$ .

However,  $n + 1 \in \mathbb{Z}$ , so this contradicts  $b$  being an upper bound.

# Archimedian property

Theorem (Apostol Theorem 1.18)

*For every  $x \in \mathbb{R}$ , there exists  $n \in \mathbb{Z}_+$  with  $n > x$ .*

# Archimedian property

Theorem (Apostol Theorem 1.18)

*For every  $x \in \mathbb{R}$ , there exists  $n \in \mathbb{Z}_+$  with  $n > x$ .*

Proof.

If not, then  $x$  is an upper bound of  $\mathbb{Z}_+$ . □



# Archimedean property

## Theorem (Apostol Theorem 1.18)

*For every  $x \in \mathbb{R}$ , there exists  $n \in \mathbb{Z}_+$  with  $n > x$ .*

## Proof.

If not, then  $x$  is an upper bound of  $\mathbb{Z}_+$ . □

## Theorem (Archimedean Property of Reals)

*For every  $x, y \in \mathbb{R}$  with  $x > 0$ , there exists  $n \in \mathbb{Z}_+$  with  $y < nx$ .*

# Archimedean property

## Theorem (Apostol Theorem 1.18)

*For every  $x \in \mathbb{R}$ , there exists  $n \in \mathbb{Z}_+$  with  $n > x$ .*

Proof.

If not, then  $x$  is an upper bound of  $\mathbb{Z}_+$ .  $\square$

## Theorem (Archimedean Property of Reals)

*For every  $x, y \in \mathbb{R}$  with  $x > 0$ , there exists  $n \in \mathbb{Z}_+$  with  $y < nx$ .*

Proof.

Replace  $x$  with  $y/x$  in the previous theorem.  $\square$

# Outline

- 1 Real Analysis Lecture 2
  - Irrational numbers
  - Upper Bound and Supremum
  - Decimal expansions

# Decimal approximations

A **finite decimal expansion** is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \cdots + \frac{a_n}{10^n},$$

where  $a_0 \in \mathbb{Z}_+$  and  $0 \leq a_k \leq 9$  for  $1 \leq k \leq n$ .

# Decimal approximations

A **finite decimal expansion** is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \cdots + \frac{a_n}{10^n},$$

where  $a_0 \in \mathbb{Z}_+$  and  $0 \leq a_k \leq 9$  for  $1 \leq k \leq n$ .

**Notation:**

$$a_0.a_1a_2a_3\dots a_n.$$

# Decimal approximations

A **finite decimal expansion** is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \cdots + \frac{a_n}{10^n},$$

where  $a_0 \in \mathbb{Z}_+$  and  $0 \leq a_k \leq 9$  for  $1 \leq k \leq n$ .

**Notation:**

$$a_0.a_1a_2a_3\dots a_n.$$

Any positive real number  $x > 0$  can be approximated by a finite decimal expansion.

# Decimal approximations

## Theorem (Apostol Theorem 1.20)

*For any real  $x > 0$  and  $n \in \mathbb{Z}_+$ , there exists a finite decimal expansion  $r_n = a_0.a_1a_2 \dots a_n$  with*

$$r_n \leq x < r_n + \frac{1}{10^n}.$$

# Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$



# Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set  $A$  is a set of integers which is bounded above by  $x$ , so it has a maximum  $a_0$ .

# Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set  $A$  is a set of integers which is bounded above by  $x$ , so it has a maximum  $a_0$ .

Then clearly  $x_1 = x - a_0 \in [0, 1)$ .

# Decimal approximations

## Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set  $A$  is a set of integers which is bounded above by  $x$ , so it has a maximum  $a_0$ .

Then clearly  $x_1 = x - a_0 \in [0, 1)$ .

Define  $a_1, a_2, a_3, \dots$  and  $x_1, x_2, x_3, \dots$  recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and  $x_{k+1} = 10x_k - a_k$ .

# Decimal approximations

## Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set  $A$  is a set of integers which is bounded above by  $x$ , so it has a maximum  $a_0$ .

Then clearly  $x_1 = x - a_0 \in [0, 1)$ .

Define  $a_1, a_2, a_3, \dots$  and  $x_1, x_2, x_3, \dots$  recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and  $x_{k+1} = 10x_k - a_k$ . Then  $0 \leq a_k \leq 9$  for all  $k \geq 1$

# Decimal approximations

## Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set  $A$  is a set of integers which is bounded above by  $x$ , so it has a maximum  $a_0$ .

Then clearly  $x_1 = x - a_0 \in [0, 1)$ .

Define  $a_1, a_2, a_3, \dots$  and  $x_1, x_2, x_3, \dots$  recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and  $x_{k+1} = 10x_k - a_k$ . Then  $0 \leq a_k \leq 9$  for all  $k \geq 1$  and

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n + 1}{10^n}$$

# Decimal expansions

We say that  $x > 0$  has the decimal expansion  $a_0.a_1a_2a_3\dots$  and write

$$x = a_0.a_1a_2a_3a_4\dots$$

# Decimal expansions

We say that  $x > 0$  has the decimal expansion  $a_0.a_1a_2a_3\dots$  and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all  $n \in \mathbb{Z}_+$ ,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

# Decimal expansions

We say that  $x > 0$  has the decimal expansion  $a_0.a_1a_2a_3\dots$  and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all  $n \in \mathbb{Z}_+$ ,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:



# Decimal expansions

We say that  $x > 0$  has the decimal expansion  $a_0.a_1a_2a_3\dots$  and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all  $n \in \mathbb{Z}_+$ ,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- 1 we haven't defined limits

# Decimal expansions

We say that  $x > 0$  has the decimal expansion  $a_0.a_1a_2a_3\dots$  and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all  $n \in \mathbb{Z}_+$ ,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- 1 we haven't defined limits
- 2 with this definition, numbers have unique decimal expansions!

# Decimal expansions

We say that  $x > 0$  has the decimal expansion  $a_0.a_1a_2a_3\dots$  and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all  $n \in \mathbb{Z}_+$ ,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- 1 we haven't defined limits
- 2 with this definition, numbers have unique decimal expansions!

$$1 \neq 0.999999999\dots$$

# Decimal expansions

We say that  $x > 0$  has the decimal expansion  $a_0.a_1a_2a_3\dots$  and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all  $n \in \mathbb{Z}_+$ ,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- ① we haven't defined limits
- ② with this definition, numbers have unique decimal expansions!

$$1 \neq 0.999999999\dots$$

$$1 \not\leq 0 + \frac{9}{10} + \frac{9}{100} + \dots + \frac{9+1}{10^n} = 1.$$