

# MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics  
California State University Fullerton

October 14, 2024

# Outline

- 1 Real Analysis Lecture 13
  - Metric subspaces
  - Limits of sequences

# Outline

- 1 Real Analysis Lecture 13
  - Metric subspaces
  - Limits of sequences

# Metric subspace

## Definition

Let  $(M, d)$  be a metric space.

# Metric subspace

## Definition

Let  $(M, d)$  be a metric space.

If  $S \subseteq M$  is a subset, then  $d : M \times M \rightarrow \mathbb{R}$  restricts to a metric  $d : S \times S \rightarrow \mathbb{R}$ , called the **relative metric**.

# Metric subspace

## Definition

Let  $(M, d)$  be a metric space.

If  $S \subseteq M$  is a subset, then  $d : M \times M \rightarrow \mathbb{R}$  restricts to a metric  $d : S \times S \rightarrow \mathbb{R}$ , called the **relative metric**.

The space  $(S, d)$  with the relative metric is called a **metric subspace**.

# Metric subspace

## Definition

Let  $(M, d)$  be a metric space.

If  $S \subseteq M$  is a subset, then  $d : M \times M \rightarrow \mathbb{R}$  restricts to a metric  $d : S \times S \rightarrow \mathbb{R}$ , called the **relative metric**.

The space  $(S, d)$  with the relative metric is called a **metric subspace**.

Open balls in  $S$ :

$$B_S(x; r) = B_M(x; r) \cap S.$$

# Example

Let  $M = \mathbb{R}$  and  $d$  be the Euclidean metric.

Consider the metric subspace  $(S, d)$  for  $S = [-1, 1) \cup \{2\}$ .

Open balls:

$$\{2\}$$

$$[-1, s), \quad -1 < s < 1$$

$$(r, s), \quad -1 < r < s < 1$$

$$[-1, s) \cup \{2\}, \quad -1 < s < 1$$

$$(r, s) \cup \{2\}, \quad -1 < r < s < 1$$



# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

## Proof.

# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

## Proof.

$\implies$ : Assume  $V$  is an open subset of  $S$ .

# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

## Proof.

$\implies$ : Assume  $V$  is an open subset of  $S$ .

Then for all  $x \in V$ , there exists  $r_x > 0$  such that  $B_S(x; r_x) \subseteq V$ .

# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

## Proof.

$\implies$ : Assume  $V$  is an open subset of  $S$ .

Then for all  $x \in V$ , there exists  $r_x > 0$  such that  $B_S(x; r_x) \subseteq V$ .

This says  $B_M(x; r_x) \cap S \subseteq V$ .

# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

## Proof.

$\implies$ : Assume  $V$  is an open subset of  $S$ .

Then for all  $x \in V$ , there exists  $r_x > 0$  such that  $B_S(x; r_x) \subseteq V$ .

This says  $B_M(x; r_x) \cap S \subseteq V$ .

Define

$$U = \bigcup_{x \in V} B_M(x; r_x).$$

# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

## Proof.

$\implies$ : Assume  $V$  is an open subset of  $S$ .

Then for all  $x \in V$ , there exists  $r_x > 0$  such that  $B_S(x; r_x) \subseteq V$ .

This says  $B_M(x; r_x) \cap S \subseteq V$ .

Define

$$U = \bigcup_{x \in V} B_M(x; r_x).$$

This is a union of open sets, and is therefore open.

# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

## Proof.

$\implies$ : Assume  $V$  is an open subset of  $S$ .

Then for all  $x \in V$ , there exists  $r_x > 0$  such that  $B_S(x; r_x) \subseteq V$ .

This says  $B_M(x; r_x) \cap S \subseteq V$ .

Define

$$U = \bigcup_{x \in V} B_M(x; r_x).$$

This is a union of open sets, and is therefore open.

Furthermore  $U \cap S = V$ .





# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

## Proof.

# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

## Proof.

$\Leftarrow$ : Assume  $U$  is an open subset of  $M$ .

# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

## Proof.

$\Leftarrow$ : Assume  $U$  is an open subset of  $M$ .  
We need to show that  $V := U \cap S$  is open.

# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

## Proof.

$\Leftarrow$ : Assume  $U$  is an open subset of  $M$ .

We need to show that  $V := U \cap S$  is open.

If  $x \in V$ , then  $x \in U$  and there exists  $r > 0$  such that  $B_M(x; r) \subseteq U$ .

# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

## Proof.

$\Leftarrow$ : Assume  $U$  is an open subset of  $M$ .

We need to show that  $V := U \cap S$  is open.

If  $x \in V$ , then  $x \in U$  and there exists  $r > 0$  such that

$$B_M(x; r) \subseteq U.$$

It follows that  $B_S(x, r) = B_M(x, r) \cap S \subseteq U \cap S = V$ .

# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

## Proof.

$\Leftarrow$ : Assume  $U$  is an open subset of  $M$ .

We need to show that  $V := U \cap S$  is open.

If  $x \in V$ , then  $x \in U$  and there exists  $r > 0$  such that  $B_M(x; r) \subseteq U$ .

It follows that  $B_S(x, r) = B_M(x, r) \cap S \subseteq U \cap S = V$ .

This shows that  $x$  is an interior point of  $S$  in the metric subspace  $(S, d)$ .

# Open sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $V$  is an open subset of  $S$  if and only if there exists an open subset  $U$  of  $M$  with  $V = U \cap S$ .*

## Proof.

$\Leftarrow$ : Assume  $U$  is an open subset of  $M$ .

We need to show that  $V := U \cap S$  is open.

If  $x \in V$ , then  $x \in U$  and there exists  $r > 0$  such that  $B_M(x; r) \subseteq U$ .

It follows that  $B_S(x, r) = B_M(x, r) \cap S \subseteq U \cap S = V$ .

This shows that  $x$  is an interior point of  $S$  in the metric subspace  $(S, d)$ .

Since  $x \in V$  was arbitrary, this shows that  $V$  is open in  $S$ . □



# Closed sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $B$  is a closed subset of  $S$  if and only if there exists a closed subset  $A$  of  $M$  with  $B = A \cap S$ .*

# Closed sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $B$  is a closed subset of  $S$  if and only if there exists a closed subset  $A$  of  $M$  with  $B = A \cap S$ .*

## Proof.

# Closed sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $B$  is a closed subset of  $S$  if and only if there exists a closed subset  $A$  of  $M$  with  $B = A \cap S$ .*

## Proof.

Let  $B \subseteq S$ .

# Closed sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $B$  is a closed subset of  $S$  if and only if there exists a closed subset  $A$  of  $M$  with  $B = A \cap S$ .*

## Proof.

Let  $B \subseteq S$ .

$B$  is closed in  $S$  if and only if  $S \setminus B$  is open in  $B$

# Closed sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $B$  is a closed subset of  $S$  if and only if there exists a closed subset  $A$  of  $M$  with  $B = A \cap S$ .*

## Proof.

Let  $B \subseteq S$ .

$B$  is closed in  $S$  if and only if  $S \setminus B$  is open in  $B$

This is true if and only if there exists an open subset  $U$  of  $M$  with  $S \setminus B = U \cap S$ .

# Closed sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $B$  is a closed subset of  $S$  if and only if there exists a closed subset  $A$  of  $M$  with  $B = A \cap S$ .*

## Proof.

Let  $B \subseteq S$ .

$B$  is closed in  $S$  if and only if  $S \setminus B$  is open in  $B$

This is true if and only if there exists an open subset  $U$  of  $M$  with  $S \setminus B = U \cap S$ .

This is true if and only if there exists an open subset  $U$  of  $M$  with  $B = (M \setminus U) \cap S$ .

# Closed sets in subspaces

## Theorem

*Let  $(M, d)$  be a metric space and let  $(S, d)$  be a subspace. Then  $B$  is a closed subset of  $S$  if and only if there exists a closed subset  $A$  of  $M$  with  $B = A \cap S$ .*

## Proof.

Let  $B \subseteq S$ .

$B$  is closed in  $S$  if and only if  $S \setminus B$  is open in  $B$

This is true if and only if there exists an open subset  $U$  of  $M$  with  $S \setminus B = U \cap S$ .

This is true if and only if there exists an open subset  $U$  of  $M$  with  $B = (M \setminus U) \cap S$ .

this is true if and only if there exists a closed subset  $A$  of  $M$  with  $B = A \cap S$ .



# Example

Consider the metric space  $(\mathbb{R}, d)$  with  $d = d_{\text{eucl}}$  the Euclidean metric, and the metric subspace  $(S, d)$  for  $S = [0, 1]$ .



# Example

Consider the metric space  $(\mathbb{R}, d)$  with  $d = d_{\text{eucl}}$  the Euclidean metric, and the metric subspace  $(S, d)$  for  $S = [0, 1]$ .

- $[0, 1]$  is open in  $S$  and closed in  $S$ , but it is not open in  $M$

# Example

Consider the metric space  $(\mathbb{R}, d)$  with  $d = d_{\text{eucl}}$  the Euclidean metric, and the metric subspace  $(S, d)$  for  $S = [0, 1]$ .

- $[0, 1]$  is open in  $S$  and closed in  $S$ , but it is not open in  $M$
- $[0, 1/2)$  is open in  $S$ , but not in  $M$

# Example

Consider the metric space  $(\mathbb{R}, d)$  with  $d = d_{\text{eucl}}$  the Euclidean metric, and the metric subspace  $(S, d)$  for  $S = [0, 1]$ .

- $[0, 1]$  is open in  $S$  and closed in  $S$ , but it is not open in  $M$
- $[0, 1/2)$  is open in  $S$ , but not in  $M$
- $(0, 1/2)$  is open in both  $S$  and in  $M$

# Example

Consider the metric space  $(\mathbb{R}, d)$  with  $d = d_{\text{eucl}}$  the Euclidean metric, and the metric subspace  $(S, d)$  for  $S = [0, 1]$ .

- $[0, 1]$  is open in  $S$  and closed in  $S$ , but it is not open in  $M$
- $[0, 1/2)$  is open in  $S$ , but not in  $M$
- $(0, 1/2)$  is open in both  $S$  and in  $M$
- $(1/2, 1]$  is open in  $S$ , but not in  $M$

# Outline

- 1 Real Analysis Lecture 13
  - Metric subspaces
  - Limits of sequences

# Definition of a limit

# Definition of a limit

Back in Calc. I:

# Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$



# Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as  $n$  gets arbitrarily large, the value of  $x_n$  gets arbitrarily close to  $L$ .

# Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as  $n$  gets arbitrarily large, the value of  $x_n$  gets arbitrarily close to  $L$ .

**Formally:**

# Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as  $n$  gets arbitrarily large, the value of  $x_n$  gets arbitrarily close to  $L$ .

**Formally:**

$\forall \epsilon > 0$ , there exists  $N \in \mathbb{Z}_+$  such that  $|x_n - L| < \epsilon \forall n \geq N$ .

# Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as  $n$  gets arbitrarily large, the value of  $x_n$  gets arbitrarily close to  $L$ .

**Formally:**

$\forall \epsilon > 0$ , there exists  $N \in \mathbb{Z}_+$  such that  $|x_n - L| < \epsilon \forall n \geq N$ .

**Metric space version:**

# Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as  $n$  gets arbitrarily large, the value of  $x_n$  gets arbitrarily close to  $L$ .

**Formally:**

$\forall \epsilon > 0$ , there exists  $N \in \mathbb{Z}_+$  such that  $|x_n - L| < \epsilon \forall n \geq N$ .

**Metric space version:**

$\forall \epsilon > 0$ , there exists  $N \in \mathbb{Z}_+$  such that  $d(x_n, L) < \epsilon \forall n \geq N$ .

# Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as  $n$  gets arbitrarily large, the value of  $x_n$  gets arbitrarily close to  $L$ .

**Formally:**

$\forall \epsilon > 0$ , there exists  $N \in \mathbb{Z}_+$  such that  $|x_n - L| < \epsilon \forall n \geq N$ .

**Metric space version:**

$\forall \epsilon > 0$ , there exists  $N \in \mathbb{Z}_+$  such that  $d(x_n, L) < \epsilon \forall n \geq N$ .

We say  $\{x_n\}$  **converges** to  $L$  and  $L$  is the **limit of the sequence**.

# Example

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .



# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  converges to zero.

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  converges to zero.

Proof.

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  converges to zero.

**Proof.**

Let  $\epsilon > 0$ .

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  converges to zero.

**Proof.**

Let  $\epsilon > 0$ .

Choose  $N \in \mathbb{Z}_+$  with  $N > 1/\epsilon$ .

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  converges to zero.

**Proof.**

Let  $\epsilon > 0$ .

Choose  $N \in \mathbb{Z}_+$  with  $N > 1/\epsilon$ .

Then  $n > N$  implies

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  converges to zero.

**Proof.**

Let  $\epsilon > 0$ .

Choose  $N \in \mathbb{Z}_+$  with  $N > 1/\epsilon$ .

Then  $n > N$  implies

$$d(x_n, 0) = |1/n - 0| = 1/n < 1/N < \epsilon.$$

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  converges to zero.

**Proof.**

Let  $\epsilon > 0$ .

Choose  $N \in \mathbb{Z}_+$  with  $N > 1/\epsilon$ .

Then  $n > N$  implies

$$d(x_n, 0) = |1/n - 0| = 1/n < 1/N < \epsilon.$$

Since  $\epsilon > 0$  was arbitrary, this proves  $\{x_n\}$  converges to 0.

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  converges to zero.

**Proof.**

Let  $\epsilon > 0$ .

Choose  $N \in \mathbb{Z}_+$  with  $N > 1/\epsilon$ .

Then  $n > N$  implies

$$d(x_n, 0) = |1/n - 0| = 1/n < 1/N < \epsilon.$$

Since  $\epsilon > 0$  was arbitrary, this proves  $\{x_n\}$  converges to 0.  $\square$



# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = (n - 1)/n$  converges to 1.

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = (n - 1)/n$  converges to 1.

Proof.

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = (n - 1)/n$  converges to 1.

**Proof.**

Let  $\epsilon > 0$ .

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = (n - 1)/n$  converges to 1.

**Proof.**

Let  $\epsilon > 0$ .

Choose  $N \in \mathbb{Z}_+$  with  $N > 1/\epsilon$ .

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = (n - 1)/n$  converges to 1.

**Proof.**

Let  $\epsilon > 0$ .

Choose  $N \in \mathbb{Z}_+$  with  $N > 1/\epsilon$ .

Then  $n > N$  implies

$$d(x_n, 1) = |(n - 1)/n - 1| = 1/n < 1/N < \epsilon.$$

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the Euclidean metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = (n - 1)/n$  converges to 1.

**Proof.**

Let  $\epsilon > 0$ .

Choose  $N \in \mathbb{Z}_+$  with  $N > 1/\epsilon$ .

Then  $n > N$  implies

$$d(x_n, 1) = |(n - 1)/n - 1| = 1/n < 1/N < \epsilon.$$

Since  $\epsilon > 0$  was arbitrary, this proves  $\{x_n\}$  converges to 1.  $\square$

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the **discrete** metric  $d$ .



# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the **discrete** metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  does not converge.

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the **discrete** metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  does not converge.

Proof.

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the **discrete** metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  does not converge.

**Proof.**

Suppose that it converges to a number  $L$ .

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the **discrete** metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  does not converge.

**Proof.**

Suppose that it converges to a number  $L$ .

Then for all  $\epsilon > 0$ , there must exist an  $N \in \mathbb{Z}_+$  such that  $d(x_n, L) < \epsilon$  for all  $N \geq n$ .

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the **discrete** metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  does not converge.

**Proof.**

Suppose that it converges to a number  $L$ .

Then for all  $\epsilon > 0$ , there must exist an  $N \in \mathbb{Z}_+$  such that  $d(x_n, L) < \epsilon$  for all  $N \geq n$ .

Take  $\epsilon = 1$  and choose  $N \in \mathbb{Z}_+$  such that  $d(x_n, L) < 1$  for all  $N \geq n$ .

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the **discrete** metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  does not converge.

**Proof.**

Suppose that it converges to a number  $L$ .

Then for all  $\epsilon > 0$ , there must exist an  $N \in \mathbb{Z}_+$  such that  $d(x_n, L) < \epsilon$  for all  $N \geq n$ .

Take  $\epsilon = 1$  and choose  $N \in \mathbb{Z}_+$  such that  $d(x_n, L) < 1$  for all  $N \geq n$ .

With the discrete metric  $d(x_n, L) < 1$  implies  $x_n = L$ .

# Example

Consider the metric space  $(M, d)$  defined by  $M = \mathbb{R}$  with the **discrete** metric  $d$ .

The sequence  $\{x_n\}$  defined by  $x_n = 1/n$  does not converge.

**Proof.**

Suppose that it converges to a number  $L$ .

Then for all  $\epsilon > 0$ , there must exist an  $N \in \mathbb{Z}_+$  such that  $d(x_n, L) < \epsilon$  for all  $N \geq n$ .

Take  $\epsilon = 1$  and choose  $N \in \mathbb{Z}_+$  such that  $d(x_n, L) < 1$  for all  $N \geq n$ .

With the discrete metric  $d(x_n, L) < 1$  implies  $x_n = L$ .

This means  $x_n = 1/n = L$  for all  $n \geq N$ , which is a contradiction! □

# Uniqueness of limits

## Theorem (Uniqueness of limits)



# Uniqueness of limits

## Theorem (Uniqueness of limits)

*Let  $(M, d)$  be a metric space.*

# Uniqueness of limits

## Theorem (Uniqueness of limits)

*Let  $(M, d)$  be a metric space.*

*If a sequence  $\{x_n\}$  in  $M$  converges to  $L$  and converges to  $M$ , then  $L = M$ .*

# Uniqueness of limits

## Theorem (Uniqueness of limits)

*Let  $(M, d)$  be a metric space.*

*If a sequence  $\{x_n\}$  in  $M$  converges to  $L$  and converges to  $M$ , then  $L = M$ .*

Proof.

# Uniqueness of limits

## Theorem (Uniqueness of limits)

*Let  $(M, d)$  be a metric space.*

*If a sequence  $\{x_n\}$  in  $M$  converges to  $L$  and converges to  $M$ , then  $L = M$ .*

## Proof.

Suppose that  $\{x_n\}$  converges to  $L$  and  $M$ .

# Uniqueness of limits

## Theorem (Uniqueness of limits)

*Let  $(M, d)$  be a metric space.*

*If a sequence  $\{x_n\}$  in  $M$  converges to  $L$  and converges to  $M$ , then  $L = M$ .*

## Proof.

Suppose that  $\{x_n\}$  converges to  $L$  and  $M$ .

Assume further that  $L \neq M$  and let  $\epsilon = d(L, M)/2$ .

# Uniqueness of limits

## Theorem (Uniqueness of limits)

*Let  $(M, d)$  be a metric space.*

*If a sequence  $\{x_n\}$  in  $M$  converges to  $L$  and converges to  $M$ , then  $L = M$ .*

## Proof.

Suppose that  $\{x_n\}$  converges to  $L$  and  $M$ .

Assume further that  $L \neq M$  and let  $\epsilon = d(L, M)/2$ .

Then there exists  $N_1$  such that  $d(x_n, L) < \epsilon$  for  $n > N_1$ .

# Uniqueness of limits

## Theorem (Uniqueness of limits)

*Let  $(M, d)$  be a metric space.*

*If a sequence  $\{x_n\}$  in  $M$  converges to  $L$  and converges to  $M$ , then  $L = M$ .*

## Proof.

Suppose that  $\{x_n\}$  converges to  $L$  and  $M$ .

Assume further that  $L \neq M$  and let  $\epsilon = d(L, M)/2$ .

Then there exists  $N_1$  such that  $d(x_n, L) < \epsilon$  for  $n > N_1$ .

Likewise, there exists  $N_2$  such that  $d(x_n, M) < \epsilon$  for  $n > N_2$ .

# Uniqueness of limits

## Theorem (Uniqueness of limits)

*Let  $(M, d)$  be a metric space.*

*If a sequence  $\{x_n\}$  in  $M$  converges to  $L$  and converges to  $M$ , then  $L = M$ .*

## Proof.

Suppose that  $\{x_n\}$  converges to  $L$  and  $M$ .

Assume further that  $L \neq M$  and let  $\epsilon = d(L, M)/2$ .

Then there exists  $N_1$  such that  $d(x_n, L) < \epsilon$  for  $n > N_1$ .

Likewise, there exists  $N_2$  such that  $d(x_n, M) < \epsilon$  for  $n > N_2$ .

Then for  $n > \max\{N_1, N_2\}$ , the triangle inequality says



# Uniqueness of limits

## Theorem (Uniqueness of limits)

*Let  $(M, d)$  be a metric space.*

*If a sequence  $\{x_n\}$  in  $M$  converges to  $L$  and converges to  $M$ , then  $L = M$ .*

## Proof.

Suppose that  $\{x_n\}$  converges to  $L$  and  $M$ .

Assume further that  $L \neq M$  and let  $\epsilon = d(L, M)/2$ .

Then there exists  $N_1$  such that  $d(x_n, L) < \epsilon$  for  $n > N_1$ .

Likewise, there exists  $N_2$  such that  $d(x_n, M) < \epsilon$  for  $n > N_2$ .

Then for  $n > \max\{N_1, N_2\}$ , the triangle inequality says

$$d(L, M) \leq d(L, x_n) + d(x_n, M) < 2\epsilon = d(L, M).$$

# Uniqueness of limits

## Theorem (Uniqueness of limits)

*Let  $(M, d)$  be a metric space.*

*If a sequence  $\{x_n\}$  in  $M$  converges to  $L$  and converges to  $M$ , then  $L = M$ .*

## Proof.

Suppose that  $\{x_n\}$  converges to  $L$  and  $M$ .

Assume further that  $L \neq M$  and let  $\epsilon = d(L, M)/2$ .

Then there exists  $N_1$  such that  $d(x_n, L) < \epsilon$  for  $n > N_1$ .

Likewise, there exists  $N_2$  such that  $d(x_n, M) < \epsilon$  for  $n > N_2$ .

Then for  $n > \max\{N_1, N_2\}$ , the triangle inequality says

$$d(L, M) \leq d(L, x_n) + d(x_n, M) < 2\epsilon = d(L, M).$$

This is a contradiction.



# Existence of limits

## Theorem

*If a sequence  $\{x_n\}$  in a metric space  $(M, d)$  converges to a value  $L \in M$ , then the range*

$$X = \{x_1, x_2, \dots\} \subseteq M$$

*is a bounded set and  $L$  is an adherent point of  $X$ .*

# Existence of limits

## Theorem

*If a sequence  $\{x_n\}$  in a metric space  $(M, d)$  converges to a value  $L \in M$ , then the range*

$$X = \{x_1, x_2, \dots\} \subseteq M$$

*is a bounded set and  $L$  is an adherent point of  $X$ .*

When the range of a sequence is bounded, we call the sequence  $\{x_n\}$  bounded.

# Challenge!

Try to prove the previous theorem.

# Existence of limits

Sort of a converse of the previous theorem.

# Existence of limits

Sort of a converse of the previous theorem.

## Theorem

*If  $S$  is a subset of a metric space  $(M, d)$  and  $L \in M$  is an adherent point of  $S$ , then there exists a sequence  $\{x_n\}$  of elements of  $S$  which converges to  $L$ .*

# Existence of limits

Recall a **subsequence** of a sequence  $\{x_n\}$  is a sequence of the form  $\{x_{k(n)}\}$ , for  $k(1) < k(2) < \dots$  positive integers.



# Existence of limits

Recall a **subsequence** of a sequence  $\{x_n\}$  is a sequence of the form  $\{x_{k(n)}\}$ , for  $k(1) < k(2) < \dots$  positive integers.

## Theorem

*A sequence  $\{x_n\}$  in a metric space  $(M, d)$  converges to a value  $L \in M$  if and only if every subsequence  $\{x_{k(n)}\}$  also converges to  $L$ .*

# Challenge!

Try to prove the previous theorem.

# Monotone sequences

## Definition

# Monotone sequences

## Definition

A sequence  $\{x_n\}$  of real numbers is called **increasing** or **monotone increasing** if  $x_n \leq x_{n+1}$  for all  $n$ .

# Monotone sequences

## Definition

A sequence  $\{x_n\}$  of real numbers is called **increasing** or **monotone increasing** if  $x_n \leq x_{n+1}$  for all  $n$ .

It is called **strictly increasing** if  $x_n < x_{n+1}$  for all  $n$ .

# Monotone sequences

## Definition

A sequence  $\{x_n\}$  of real numbers is called **increasing** or **monotone increasing** if  $x_n \leq x_{n+1}$  for all  $n$ .

It is called **strictly increasing** if  $x_n < x_{n+1}$  for all  $n$ .

A sequence  $\{x_n\}$  of real numbers is called **decreasing** or **monotone decreasing** if  $x_n \geq x_{n+1}$  for all  $n$ .

# Monotone sequences

## Definition

A sequence  $\{x_n\}$  of real numbers is called **increasing** or **monotone increasing** if  $x_n \leq x_{n+1}$  for all  $n$ .

It is called **strictly increasing** if  $x_n < x_{n+1}$  for all  $n$ .

A sequence  $\{x_n\}$  of real numbers is called **decreasing** or **monotone decreasing** if  $x_n \geq x_{n+1}$  for all  $n$ .

It is called **strictly decreasing** if  $x_n > x_{n+1}$  for all  $n$ .

# Monotone sequences

## Definition

A sequence  $\{x_n\}$  of real numbers is called **increasing** or **monotone increasing** if  $x_n \leq x_{n+1}$  for all  $n$ .

It is called **strictly increasing** if  $x_n < x_{n+1}$  for all  $n$ .

A sequence  $\{x_n\}$  of real numbers is called **decreasing** or **monotone decreasing** if  $x_n \geq x_{n+1}$  for all  $n$ .

It is called **strictly decreasing** if  $x_n > x_{n+1}$  for all  $n$ .

In any of these cases, we call the sequence **monotone**.



# Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

# Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

Theorem (Monotone convergence theorem)

# Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

## Theorem (Monotone convergence theorem)

*Consider the metric space formed by  $\mathbb{R}$  with Euclidean distance.*

# Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

## Theorem (Monotone convergence theorem)

*Consider the metric space formed by  $\mathbb{R}$  with Euclidean distance.*

*A monotone increasing sequence  $\{x_n\}$  of real numbers which is bounded above converges to  $L = \sup\{x_1, x_2, \dots\}$ .*

# Monotone sequences

Bounded monotone sequences always converge (with Euclidean metric)!

## Theorem (Monotone convergence theorem)

*Consider the metric space formed by  $\mathbb{R}$  with Euclidean distance.*

*A monotone increasing sequence  $\{x_n\}$  of real numbers which is bounded above converges to  $L = \sup\{x_1, x_2, \dots\}$ .*

*A monotone decreasing sequence  $\{x_n\}$  which is bounded below converges to  $L = \inf\{x_1, x_2, \dots\}$ .*

# Monotone sequences

Proof.

# Monotone sequences

Proof.

Suppose that  $\{x_n\}$  is a monotone increasing sequence which is bounded above and let  $X = \{x_1, x_2, \dots\}$ .

# Monotone sequences

## Proof.

Suppose that  $\{x_n\}$  is a monotone increasing sequence which is bounded above and let  $X = \{x_1, x_2, \dots\}$ . Then by the Completeness Axiom  $L = \sup X$  exists.



# Monotone sequences

## Proof.

Suppose that  $\{x_n\}$  is a monotone increasing sequence which is bounded above and let  $X = \{x_1, x_2, \dots\}$ . Then by the Completeness Axiom  $L = \sup X$  exists.

Let  $\epsilon > 0$ .

# Monotone sequences

## Proof.

Suppose that  $\{x_n\}$  is a monotone increasing sequence which is bounded above and let  $X = \{x_1, x_2, \dots\}$ . Then by the Completeness Axiom  $L = \sup X$  exists.

Let  $\epsilon > 0$ .

Then  $L - \epsilon$  is not a supremum of  $X$ , so there exists  $x_N \in X$  such that  $x_N > L - \epsilon$ .

# Monotone sequences

## Proof.

Suppose that  $\{x_n\}$  is a monotone increasing sequence which is bounded above and let  $X = \{x_1, x_2, \dots\}$ . Then by the Completeness Axiom  $L = \sup X$  exists.

Let  $\epsilon > 0$ .

Then  $L - \epsilon$  is not a supremum of  $X$ , so there exists  $x_N \in X$  such that  $x_N > L - \epsilon$ .

Since the sequence is increasing, for any  $n > N$ , we have that  $L - \epsilon < x_N \leq x_n < L$ .

# Monotone sequences

## Proof.

Suppose that  $\{x_n\}$  is a monotone increasing sequence which is bounded above and let  $X = \{x_1, x_2, \dots\}$ . Then by the Completeness Axiom  $L = \sup X$  exists.

Let  $\epsilon > 0$ .

Then  $L - \epsilon$  is not a supremum of  $X$ , so there exists  $x_N \in X$  such that  $x_N > L - \epsilon$ .

Since the sequence is increasing, for any  $n > N$ , we have that  $L - \epsilon < x_N \leq x_n < L$ .

It follows that

$$d(x_n, L) = |x_n - L| = L - x_n < \epsilon.$$

# Monotone sequences

## Proof.

Suppose that  $\{x_n\}$  is a monotone increasing sequence which is bounded above and let  $X = \{x_1, x_2, \dots\}$ . Then by the Completeness Axiom  $L = \sup X$  exists.

Let  $\epsilon > 0$ .

Then  $L - \epsilon$  is not a supremum of  $X$ , so there exists  $x_N \in X$  such that  $x_N > L - \epsilon$ .

Since the sequence is increasing, for any  $n > N$ , we have that  $L - \epsilon < x_N \leq x_n < L$ .

It follows that

$$d(x_n, L) = |x_n - L| = L - x_n < \epsilon.$$

Since  $\epsilon > 0$  was arbitrary, this proves  $x_n$  converges to  $L$ . □