

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

September 16, 2024

Outline

1 Real Analysis Lecture 6

- Functions
- Cardinality
- Set algebra

Outline

1 Real Analysis Lecture 6

- Functions
- Cardinality
- Set algebra

Functions

A **function** from A to B is a relation \mathcal{R} from A to B with the property

Functions

A **function** from A to B is a relation \mathcal{R} from A to B with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \quad \text{implies} \quad b = c.$$

Functions

A **function** from A to B is a relation \mathcal{R} from A to B with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

If a relation \mathcal{R} is a function, we usually use a symbol like f .

Functions

A **function** from A to B is a relation \mathcal{R} from A to B with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

If a relation \mathcal{R} is a function, we usually use a symbol like f .

NOTATION: $f : A \rightarrow B$ means f is a function from A to B

NOTATION: $f(a) = b$ means $(a, b) \in f$.

Functions

A **function** from A to B is a relation \mathcal{R} from A to B with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

If a relation \mathcal{R} is a function, we usually use a symbol like f .

NOTATION: $f : A \rightarrow B$ means f is a function from A to B

NOTATION: $f(a) = b$ means $(a, b) \in f$.

The set

$$\text{img}(f) = \{f(a) : a \in A\}$$

is called the **range** or **image** of f

Challenge!

Problem

Determine all the equivalence relations on \mathbb{R} which are also functions.

Challenge!

Problem

Determine all the equivalence relations on \mathbb{R} which are also functions.

Solution

$f = \mathcal{R}$ must be reflexive, so $x\mathcal{R}x$ for all x

Challenge!

Problem

Determine all the equivalence relations on \mathbb{R} which are also functions.

Solution

$f = \mathcal{R}$ must be reflexive, so $x\mathcal{R}x$ for all x
This means $f(x) = x$ for all x .

Challenge!

Problem

Determine all the equivalence relations on \mathbb{R} which are also functions.

Solution

$f = \mathcal{R}$ must be reflexive, so $x\mathcal{R}x$ for all x

This means $f(x) = x$ for all x .

Thus the only function which is an equivalence relation is the identity function

$$f(x) = x.$$

Properties of functions

A function f from A to B is called **one-to-one** or **injective** if

Properties of functions

A function f from A to B is called **one-to-one** or **injective** if

$$f(x) = f(y) \text{ implies } x = y \text{ for all } x, y \in A.$$

Properties of functions

A function f from A to B is called **one-to-one** or **injective** if

$$f(x) = f(y) \text{ implies } x = y \text{ for all } x, y \in A.$$

It is called **onto** or **surjective** if $\text{img}(f) = B$, or equivalently

$$\text{for all } b \in B \text{ there exists } a \in A \text{ with } f(a) = b.$$

Properties of functions

A function f from A to B is called **one-to-one** or **injective** if

$$f(x) = f(y) \text{ implies } x = y \text{ for all } x, y \in A.$$

It is called **onto** or **surjective** if $\text{img}(f) = B$, or equivalently

$$\text{for all } b \in B \text{ there exists } a \in A \text{ with } f(a) = b.$$

If it satisfies both properties, it is called **bijective**.

Inverses

The **converse** of a relation \mathcal{R} from A to B is the relation

Inverses

The **converse** of a relation \mathcal{R} from A to B is the relation

$$\check{\mathcal{R}} = \{(b, a) \in B \times A : a\mathcal{R}b\}.$$

Inverses

The **converse** of a relation \mathcal{R} from A to B is the relation

$$\check{\mathcal{R}} = \{(b, a) \in B \times A : a\mathcal{R}b\}.$$

If f is function, then \check{f} may or not be a function.

Inverses

The **converse** of a relation \mathcal{R} from A to B is the relation

$$\check{\mathcal{R}} = \{(b, a) \in B \times A : a\mathcal{R}b\}.$$

If f is function, then \check{f} may or not be a function.
If it is, we call it the **inverse** of f .

Compositions

The composition of $f : A \rightarrow B$ and $g : B \rightarrow C$ is the function

Compositions

The composition of $f : A \rightarrow B$ and $g : B \rightarrow C$ is the function

$$g \circ f : A \rightarrow C$$

Compositions

The composition of $f : A \rightarrow B$ and $g : B \rightarrow C$ is the function

$$g \circ f : A \rightarrow C$$

with domain A and codomain C defined by

Compositions

The composition of $f : A \rightarrow B$ and $g : B \rightarrow C$ is the function

$$g \circ f : A \rightarrow C$$

with domain A and codomain C defined by

$$(g \circ f)(x) = g(f(x)).$$

Sequences

A **finite sequence** is a function $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$.

Sequences

A **finite sequence** is a function $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$.

An **infinite sequence** is a function $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$.

Sequences

A **finite sequence** is a function $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$.

An **infinite sequence** is a function $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$.

NOTATION: f_n indicates the value $f(n)$

Sequences

A **finite sequence** is a function $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$.

An **infinite sequence** is a function $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$.

NOTATION: f_n indicates the value $f(n)$

NOTATION: $\{f_n\}$ is another way of writing the function f

Sequences

A **finite sequence** is a function $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$.

An **infinite sequence** is a function $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$.

NOTATION: f_n indicates the value $f(n)$

NOTATION: $\{f_n\}$ is another way of writing the function f

If $k : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ is a function which is **strictly increasing**, meaning

$$m < n \quad \Rightarrow \quad k(m) < k(n),$$

Sequences

A **finite sequence** is a function $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$.

An **infinite sequence** is a function $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$.

NOTATION: f_n indicates the value $f(n)$

NOTATION: $\{f_n\}$ is another way of writing the function f

If $k : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ is a function which is **strictly increasing**, meaning

$$m < n \Rightarrow k(m) < k(n),$$

then the composition $f \circ k : \mathbb{Z}_+ \rightarrow \mathbb{R}$ forms a sequence called a **subsequence** of f .

Sequences

A **finite sequence** is a function $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$.

An **infinite sequence** is a function $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$.

NOTATION: f_n indicates the value $f(n)$

NOTATION: $\{f_n\}$ is another way of writing the function f

If $k : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ is a function which is **strictly increasing**, meaning

$$m < n \Rightarrow k(m) < k(n),$$

then the composition $f \circ k : \mathbb{Z}_+ \rightarrow \mathbb{R}$ forms a sequence called a **subsequence** of f .

NOTATION: $\{f_{k(n)}\}$ or $\{f_{k_n}\}$ both really mean $f \circ k$

Outline

1 Real Analysis Lecture 6

- Functions
- **Cardinality**
- Set algebra

Cantor's Paradise

Philosophy: sets are the same size, or **cardinality**, if there is a bijection between them.

Cantor's Paradise

Philosophy: sets are the same size, or **cardinality**, if there is a bijection between them.

- $|A|$ denotes the cardinality of A

Cantor's Paradise

Philosophy: sets are the same size, or **cardinality**, if there is a bijection between them.

- $|A|$ denotes the cardinality of A
- $|A| \leq |B|$ means there is an injection $f : A \rightarrow B$

Cantor's Paradise

Philosophy: sets are the same size, or **cardinality**, if there is a bijection between them.

- $|A|$ denotes the cardinality of A
- $|A| \leq |B|$ means there is an injection $f : A \rightarrow B$
- $|A| \geq |B|$ means there is a surjection $f : A \rightarrow B$

Cantor's Paradise

Philosophy: sets are the same size, or **cardinality**, if there is a bijection between them.

- $|A|$ denotes the cardinality of A
- $|A| \leq |B|$ means there is an injection $f : A \rightarrow B$
- $|A| \geq |B|$ means there is a surjection $f : A \rightarrow B$
- $|A| = |B|$ means there is a bijection $f : A \rightarrow B$

Cantor's Paradise

Philosophy: sets are the same size, or **cardinality**, if there is a bijection between them.

- $|A|$ denotes the cardinality of A
- $|A| \leq |B|$ means there is an injection $f : A \rightarrow B$
- $|A| \geq |B|$ means there is a surjection $f : A \rightarrow B$
- $|A| = |B|$ means there is a bijection $f : A \rightarrow B$

Cantor's Paradise

Theorem (Cantor-Schroeder-Bernstein Theorem)

The following are equivalent

- (i) $|A| \leq |B|$ and $|B| \leq |A|$
- (ii) $|A| \geq |B|$ and $|B| \geq |A|$
- (iii) $|A| \leq |B|$ and $|A| \geq |A|$
- (iv) $|A| = |B|$

Challenge

Problem

Show that \mathbb{Z}_+ and $\mathbb{Z}_+ \times \mathbb{Z}_+$ have the same cardinality.

Challenge

Problem

Show that \mathbb{Z}_+ and $\mathbb{Z}_+ \times \mathbb{Z}_+$ have the same cardinality.

Hint: consider $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$, $f(m, n) = 2^m 3^n$

Solution

Challenge

Problem

Show that \mathbb{Z}_+ and $\mathbb{Z}_+ \times \mathbb{Z}_+$ have the same cardinality.

Hint: consider $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$, $f(m, n) = 2^m 3^n$

Solution

$2^a 3^b = 2^c 3^d$ if and only if $a = c$ and $b = d$ (Fund. Thm. Arith.)

Challenge

Problem

Show that \mathbb{Z}_+ and $\mathbb{Z}_+ \times \mathbb{Z}_+$ have the same cardinality.

Hint: consider $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$, $f(m, n) = 2^m 3^n$

Solution

$2^a 3^b = 2^c 3^d$ if and only if $a = c$ and $b = d$ (Fund. Thm. Arith.)
 $f(m, n) = 2^m 3^n$ is injective

Challenge

Problem

Show that \mathbb{Z}_+ and $\mathbb{Z}_+ \times \mathbb{Z}_+$ have the same cardinality.

Hint: consider $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$, $f(m, n) = 2^m 3^n$

Solution

$2^a 3^b = 2^c 3^d$ if and only if $a = c$ and $b = d$ (Fund. Thm. Arith.)

$f(m, n) = 2^m 3^n$ is injective

$$|\mathbb{Z}_+ \times \mathbb{Z}_+| \leq |\mathbb{Z}_+|$$

Challenge

Problem

Show that \mathbb{Z}_+ and $\mathbb{Z}_+ \times \mathbb{Z}_+$ have the same cardinality.

Hint: consider $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$, $f(m, n) = 2^m 3^n$

Solution

$2^a 3^b = 2^c 3^d$ if and only if $a = c$ and $b = d$ (Fund. Thm. Arith.)

$f(m, n) = 2^m 3^n$ is injective

$$|\mathbb{Z}_+ \times \mathbb{Z}_+| \leq |\mathbb{Z}_+|$$

$g : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$, $g(m, n) = m$ is surjective

Challenge

Problem

Show that \mathbb{Z}_+ and $\mathbb{Z}_+ \times \mathbb{Z}_+$ have the same cardinality.

Hint: consider $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$, $f(m, n) = 2^m 3^n$

Solution

$2^a 3^b = 2^c 3^d$ if and only if $a = c$ and $b = d$ (Fund. Thm. Arith.)

$f(m, n) = 2^m 3^n$ is injective

$$|\mathbb{Z}_+ \times \mathbb{Z}_+| \leq |\mathbb{Z}_+|$$

$g : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$, $g(m, n) = m$ is surjective

$$|\mathbb{Z}_+ \times \mathbb{Z}_+| \geq |\mathbb{Z}_+|$$

Challenge

Problem

Show that \mathbb{Z}_+ and \mathbb{Z} have the same cardinality.

Challenge

Problem

Show that \mathbb{Z}_+ and \mathbb{Z} have the same cardinality.

Hint: consider $f : \mathbb{Z}_+ \rightarrow \mathbb{Z}$

$$f(n) = \begin{cases} n/2 & n \text{ is even} \\ -(n-1)/2 & n \text{ is odd} \end{cases}$$

Finite and infinite

Finite and infinite

Sets with finite cardinality:

$$\{1, 2, 3\}, \quad \{\mathbb{Z}, \mathbb{R}\}, \quad \{x : x \text{ is a student at CSUF}\}$$

Finite and infinite

Sets with finite cardinality:

$$\{1, 2, 3\}, \quad \{\mathbb{Z}, \mathbb{R}\}, \quad \{x : x \text{ is a student at CSUF}\}$$

Sets with infinite cardinality:

$$\mathbb{Z}_+, \quad \mathbb{Q}, \quad \mathbb{R}, \quad \mathbb{C}.$$

Finite and infinite

Sets with finite cardinality:

$$\{1, 2, 3\}, \quad \{\mathbb{Z}, \mathbb{R}\}, \quad \{x : x \text{ is a student at CSUF}\}$$

Sets with infinite cardinality:

$$\mathbb{Z}_+, \quad \mathbb{Q}, \quad \mathbb{R}, \quad \mathbb{C}.$$

Cantor's discovery: there are multiple sizes of infinity!

$$\mathbb{Z}_+, \mathbb{Z}, \mathbb{Z} \times \mathbb{Z}, \mathbb{Q}, \text{ are all the same cardinality}$$

Finite and infinite

Sets with finite cardinality:

$$\{1, 2, 3\}, \quad \{\mathbb{Z}, \mathbb{R}\}, \quad \{x : x \text{ is a student at CSUF}\}$$

Sets with infinite cardinality:

$$\mathbb{Z}_+, \quad \mathbb{Q}, \quad \mathbb{R}, \quad \mathbb{C}.$$

Cantor's discovery: there are multiple sizes of infinity!

$$\mathbb{Z}_+, \mathbb{Z}, \mathbb{Z} \times \mathbb{Z}, \mathbb{Q}, \text{ are all the same cardinality}$$

\mathbb{R} has larger cardinality than \mathbb{Z}_+ .

Cantor diagonalization

Suppose there were a bijection

$$f : \mathbb{Z}_+ \rightarrow (0, 1)$$

Cantor diagonalization

Suppose there were a bijection

$$f : \mathbb{Z}_+ \rightarrow (0, 1)$$

$$f(1) = 0.a_{11}a_{12}a_{13}a_{14}a_{15}\dots$$

$$f(2) = 0.a_{21}a_{22}a_{23}a_{24}a_{25}\dots$$

$$f(3) = 0.a_{31}a_{32}a_{33}a_{34}a_{35}\dots$$

$$f(4) = 0.a_{41}a_{42}a_{43}a_{44}a_{45}\dots$$

$$f(5) = 0.a_{51}a_{52}a_{53}a_{54}a_{55}\dots$$

$$\vdots$$

Cantor diagonalization

Suppose there were a surjection

$$f : \mathbb{Z}_+ \rightarrow (0, 1)$$

Cantor diagonalization

Suppose there were a surjection

$$f : \mathbb{Z}_+ \rightarrow (0, 1)$$

$$f(1) = 0.\textcolor{red}{a}_{11} a_{12} a_{13} a_{14} a_{15} \dots$$

$$f(2) = 0.a_{21} \textcolor{red}{a}_{22} a_{23} a_{24} a_{25} \dots$$

$$f(3) = 0.a_{31} a_{32} \textcolor{red}{a}_{33} a_{34} a_{35} \dots$$

$$f(4) = 0.a_{41} a_{42} a_{43} \textcolor{red}{a}_{44} a_{45} \dots$$

$$f(5) = 0.a_{51} a_{52} a_{53} a_{54} \textcolor{red}{a}_{55} \dots$$

$$\vdots$$

Cantor diagonalization

Now define $b_1, b_2, b_3 \cdots \in \{0, 9\}$ by

$$b_j = \begin{cases} 1, & a_{jj} = 0 \\ 0, & a_{jj} \neq 0 \end{cases}$$

Cantor diagonalization

Now define $b_1, b_2, b_3 \dots \in \{0, 9\}$ by

$$b_j = \begin{cases} 1, & a_{jj} = 0 \\ 0, & a_{jj} \neq 0 \end{cases}$$

$$b = 0.b_1 b_2 b_3 b_4 b_5 \dots$$

Cantor diagonalization

- $b \in (0, 1)$

Cantor diagonalization

- $b \in (0, 1)$
- f is surjective

Cantor diagonalization

- $b \in (0, 1)$
- f is surjective
- there exists $n \in \mathbb{Z}_+$ with $f(n) = b$

Cantor diagonalization

- $b \in (0, 1)$
- f is surjective
- there exists $n \in \mathbb{Z}_+$ with $f(n) = b$

$$0.b_1b_2b_3b_4b_5\cdots = 0.a_{n1}a_{n2}a_{n3}a_{n4}a_{n5}\cdots$$

Cantor diagonalization

- $b \in (0, 1)$
- f is surjective
- there exists $n \in \mathbb{Z}_+$ with $f(n) = b$

$$0.b_1b_2b_3b_4b_5\cdots = 0.a_{n1}a_{n2}a_{n3}a_{n4}a_{n5}\cdots$$

- $b_n = a_{nn}$

Cantor diagonalization

- $b \in (0, 1)$
- f is surjective
- there exists $n \in \mathbb{Z}_+$ with $f(n) = b$

$$0.b_1b_2b_3b_4b_5\cdots = 0.a_{n1}a_{n2}a_{n3}a_{n4}a_{n5}\cdots$$

- $b_n = a_{nn}$ CONTRADICTION!

Countable and uncountable

A set is

Countable and uncountable

A set is

- **countably infinite** if it has the same cardinality as \mathbb{Z}_+

Countable and uncountable

A set is

- **countably infinite** if it has the same cardinality as \mathbb{Z}_+
- **countable** if it is finite or countably infinite

Countable and uncountable

A set is

- **countably infinite** if it has the same cardinality as \mathbb{Z}_+
- **countable** if it is finite or countably infinite
- **uncountable** otherwise

Countable and uncountable

A set is

- **countably infinite** if it has the same cardinality as \mathbb{Z}_+
- **countable** if it is finite or countably infinite
- **uncountable** otherwise

Theorem (Cantor's Theorem)

\mathbb{R} is uncountable

Ultimate Cantor diagonalization

The **power set** $\mathcal{P}(A)$ of a set A is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

Ultimate Cantor diagonalization

The **power set** $\mathcal{P}(A)$ of a set A is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

Theorem

The cardinality of $\mathcal{P}(A)$ is strictly larger than A .

Ultimate Cantor diagonalization

The **power set** $\mathcal{P}(A)$ of a set A is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

Theorem

The cardinality of $\mathcal{P}(A)$ is strictly larger than A .

Proof.

Ultimate Cantor diagonalization

The **power set** $\mathcal{P}(A)$ of a set A is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

Theorem

The cardinality of $\mathcal{P}(A)$ is strictly larger than A .

Proof.

Suppose $f : A \rightarrow \mathcal{P}(A)$ is surjective.

Ultimate Cantor diagonalization

The **power set** $\mathcal{P}(A)$ of a set A is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

Theorem

The cardinality of $\mathcal{P}(A)$ is strictly larger than A .

Proof.

Suppose $f : A \rightarrow \mathcal{P}(A)$ is surjective.
Consider the set

$$S = \{a \in A : a \notin f(a)\}.$$



Challenge

Challenge

Since f is surjective, $S = f(x)$ for some $x \in A$.

Challenge

Since f is surjective, $S = f(x)$ for some $x \in A$.

Problem

Consider the statement $x \in S$. What can you conclude?

Outline

1 Real Analysis Lecture 6

- Functions
- Cardinality
- Set algebra

Unions of sets

Unions:

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

$$A_1 \cup A_2 \cup \cdots \cup A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

$$A_1 \cup A_2 \cup \cdots \cup A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

NOTATION:

$$\bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \cdots \cup A_n.$$

Intersections of sets

Intersections:

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

$$A_1 \cap A_2 \cap \cdots \cap A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

$$A_1 \cap A_2 \cap \cdots \cap A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

NOTATION:

$$\bigcap_{i=1}^n A_i = A_1 \cap A_2 \cap \cdots \cap A_n.$$

Complements of sets

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Theorem

De Morgan's Laws

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Theorem

De Morgan's Laws

$$B - \bigcup_{i=1}^n A_i = \bigcap_{i=1}^n (B - A_i)$$

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Theorem

De Morgan's Laws

$$B - \bigcup_{i=1}^n A_i = \bigcap_{i=1}^n (B - A_i)$$

$$B - \bigcap_{i=1}^n A_i = \bigcup_{i=1}^n (B - A_i)$$

Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- I any set, called the **index set**

Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- I any set, called the **index set**
- A_i is a set for all i

Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- I any set, called the **index set**
- A_i is a set for all i

$$\bigcup_{i \in I} A_i = \{x : x \in A_i, \text{ for some } i \in I\}$$

Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- I any set, called the **index set**
- A_i is a set for all i

$$\bigcup_{i \in I} A_i = \{x : x \in A_i, \text{ for some } i \in I\}$$

$$\bigcap_{i \in I} A_i = \{x : x \in A_i, \text{ for all } i \in I\}$$

Challenge

- index set $I = \mathbb{Z}_+$
- family of sets $\{A_i : i \in I\}$
- $A_i = [0, i)$

Problem

Determine $\bigcup_{i \in I} A_i$.

Challenge

- index set $I = \mathbb{Z}_+$
- family of sets $\{A_i : i \in I\}$
- $A_i = [0, i)$

Problem

Determine $\bigcap_{i \in I} A_i$.