

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

September 4, 2024

Outline

- 1 Real Analysis Lecture 3
 - Suprema and Infima
 - Decimal expansions
 - The Triangle Inequality

Outline

- 1 Real Analysis Lecture 3
 - Suprema and Infima
 - Decimal expansions
 - The Triangle Inequality

Completeness Axiom

Last time, we finished with the Completeness Axiom:

Completeness Axiom

Last time, we finished with the Completeness Axiom:

Completeness Axiom

Last time, we finished with the Completeness Axiom:

- Ⓐ10 **completeness axiom:** if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Completeness Axiom

Last time, we finished with the Completeness Axiom:

A10 completeness axiom: if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

Completeness Axiom

Last time, we finished with the Completeness Axiom:

- **completeness axiom:** if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

- 3 is the supremum of $(0, 3)$

Completeness Axiom

Last time, we finished with the Completeness Axiom:

A10 completeness axiom: if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

- 3 is the supremum of $(0, 3)$
- 1 is the supremum of

$$\left\{ \frac{n}{n+1} : n \in \mathbb{Z}_+ \right\}$$

Completeness Axiom

Last time, we finished with the Completeness Axiom:

- Ⓐ10 **completeness axiom:** if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

- 3 is the supremum of $(0, 3)$
- 1 is the supremum of

$$\left\{ \frac{n}{n+1} : n \in \mathbb{Z}_+ \right\}$$

- π is the supremum of

$$\{3, 3.1, 3.14, 3.141, 3.1415, 3.14159, 3.141592, \dots\}.$$

Challenge

Problem

Prove that if A is a set of integers that is bounded above, then A has a maximum.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

If $a' \in A$, then a' is an integer and therefore $a' = a + k$ for $k \in \mathbb{Z}$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

If $a' \in A$, then a' is an integer and therefore $a' = a + k$ for $k \in \mathbb{Z}$. Since b is an upper bound, $b \geq a + k > b - 1 + k$, making $0 > k - 1$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

If $a' \in A$, then a' is an integer and therefore $a' = a + k$ for $k \in \mathbb{Z}$. Since b is an upper bound, $b \geq a + k > b - 1 + k$, making $0 > k - 1$.

Therefore $k \leq 0$ and $a' \leq a$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

If $a' \in A$, then a' is an integer and therefore $a' = a + k$ for $k \in \mathbb{Z}$. Since b is an upper bound, $b \geq a + k > b - 1 + k$, making $0 > k - 1$.

Therefore $k \leq 0$ and $a' \leq a$.

It follows that a is an upper bound of A , and since $a \in A$ it is a maximum.

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

If $b \in S$ also, then b is called a **minimal element** of S .

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

If $b \in S$ also, then b is called a **minimal element** of S . An **infimum** of a set S of real numbers is a real number $b \in \mathbb{R}$ such that

- b is a lower bound of S
- if $b < b'$, then b' is not a lower bound of S

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

If $b \in S$ also, then b is called a **minimal element** of S . An **infimum** of a set S of real numbers is a real number $b \in \mathbb{R}$ such that

- b is a lower bound of S
- if $b < b'$, then b' is not a lower bound of S

In other words

an infimum is a **greatest lower bound**

Properties of Suprema

The first property of suprema is that they must be arbitrarily close to elements of the set.

Theorem (Approximation Property)

Let $S \subseteq \mathbb{R}$ be bounded above, and let $b = \sup(S)$. Then for all $\epsilon > 0$, there exists $a \in S$ with

$$b - \epsilon < a \leq b.$$

Properties of Suprema

Proof.

Since $b - \epsilon < b$, the definition of a supremum implies $b - \epsilon$ cannot be an upper bound.

Properties of Suprema

Proof.

Since $b - \epsilon < b$, the definition of a supremum implies $b - \epsilon$ cannot be an upper bound.

Therefore there must exist $a \in S$ with $a > b - \epsilon$. □

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Consider the set

$$S = \{s_n : n \in \mathbb{Z}_+\}.$$

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Consider the set

$$S = \{s_n : n \in \mathbb{Z}_+\}.$$

For every n ,

$$s_n \leq 1 + 1 + \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \cdots + \frac{1}{2^n} = 3 - \frac{1}{2^n} < 3$$

so S is bounded above by 3.

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Consider the set

$$S = \{s_n : n \in \mathbb{Z}_+\}.$$

For every n ,

$$s_n \leq 1 + 1 + \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \cdots + \frac{1}{2^n} = 3 - \frac{1}{2^n} < 3$$

so S is bounded above by 3. Therefore S has a supremum, $\sup(S)$.

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N .

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

If we remember Taylor series

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

If we remember Taylor series

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots$$

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

If we remember Taylor series

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots$$

We've just shown that $e^1 = e$ exists.

Properties of Suprema

Also, suprema play nicely with addition.

Theorem (Additive Property)

Let $A, B \subseteq \mathbb{R}$ be bounded above set

$$C = \{x + y : x \in A, y \in B\}.$$

Then C is bounded above and

$$\sup(C) = \sup(A) + \sup(B).$$

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

It follows that $y \leq c - a$ for all $y \in B$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

It follows that $y \leq c - a$ for all $y \in B$.

Thus $c - a$ is an upper bound of B , and it follows $b \leq c - a$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

It follows that $y \leq c - a$ for all $y \in B$.

Thus $c - a$ is an upper bound of B , and it follows $b \leq c - a$.

Therefore $a + b \leq c$. □

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

This means there exists $n \in \mathbb{Z}_+$ with $b - 1 < n$.

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

This means there exists $n \in \mathbb{Z}_+$ with $b - 1 < n$.

It follows from Axiom 7 that $b < n + 1$.

Challenge

Problem

Use the completeness axiom to prove that the \mathbb{Z}_+ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

This means there exists $n \in \mathbb{Z}_+$ with $b - 1 < n$.

It follows from Axiom 7 that $b < n + 1$.

However, $n + 1 \in \mathbb{Z}$, so this contradicts b being an upper bound.

Archimedian property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{Z}_+$ with $n > x$.

Archimedian property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{Z}_+$ with $n > x$.

Proof.

If not, then x is an upper bound of \mathbb{Z}_+ . □

Archimedean property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{Z}_+$ with $n > x$.

Proof.

If not, then x is an upper bound of \mathbb{Z}_+ . □

Theorem (Archimedean Property of Reals)

For every $x, y \in \mathbb{R}$ with $x > 0$, there exists $n \in \mathbb{Z}_+$ with $y < nx$.

Archimedean property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{Z}_+$ with $n > x$.

Proof.

If not, then x is an upper bound of \mathbb{Z}_+ . \square

Theorem (Archimedean Property of Reals)

For every $x, y \in \mathbb{R}$ with $x > 0$, there exists $n \in \mathbb{Z}_+$ with $y < nx$.

Proof.

Replace x with y/x in the previous theorem. \square

Outline

- 1 Real Analysis Lecture 3
 - Suprema and Infima
 - **Decimal expansions**
 - The Triangle Inequality

Decimal approximations

A **finite decimal expansion** is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \cdots + \frac{a_n}{10^n},$$

where $a_0 \in \mathbb{Z}_+$ and $0 \leq a_k \leq 9$ for $1 \leq k \leq n$.

Decimal approximations

A **finite decimal expansion** is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \cdots + \frac{a_n}{10^n},$$

where $a_0 \in \mathbb{Z}_+$ and $0 \leq a_k \leq 9$ for $1 \leq k \leq n$.

Notation:

$$a_0.a_1a_2a_3\dots a_n.$$

Decimal approximations

A **finite decimal expansion** is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \cdots + \frac{a_n}{10^n},$$

where $a_0 \in \mathbb{Z}_+$ and $0 \leq a_k \leq 9$ for $1 \leq k \leq n$.

Notation:

$$a_0.a_1a_2a_3\ldots a_n.$$

Any positive real number $x > 0$ can be approximated by a finite decimal expansion.

Decimal approximations

Theorem (Apostol Theorem 1.20)

For any real $x > 0$ and $n \in \mathbb{Z}_+$, there exists a finite decimal expansion $r_n = a_0.a_1a_2 \dots a_n$ with

$$r_n \leq x < r_n + \frac{1}{10^n}.$$

Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set A is a set of integers which is bounded above by x , so it has a maximum a_0 .

Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set A is a set of integers which is bounded above by x , so it has a maximum a_0 .

Then clearly $x_1 = x - a_0 \in [0, 1)$.

Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set A is a set of integers which is bounded above by x , so it has a maximum a_0 .

Then clearly $x_1 = x - a_0 \in [0, 1)$.

Define a_1, a_2, a_3, \dots and x_1, x_2, x_3, \dots recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and $x_{k+1} = 10x_k - a_k$.

Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set A is a set of integers which is bounded above by x , so it has a maximum a_0 .

Then clearly $x_1 = x - a_0 \in [0, 1)$.

Define a_1, a_2, a_3, \dots and x_1, x_2, x_3, \dots recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and $x_{k+1} = 10x_k - a_k$. Then $0 \leq a_k \leq 9$ for all $k \geq 1$

Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set A is a set of integers which is bounded above by x , so it has a maximum a_0 .

Then clearly $x_1 = x - a_0 \in [0, 1)$.

Define a_1, a_2, a_3, \dots and x_1, x_2, x_3, \dots recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and $x_{k+1} = 10x_k - a_k$. Then $0 \leq a_k \leq 9$ for all $k \geq 1$ and

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n + 1}{10^n}$$



Decimal expansions

We say that $x > 0$ has the decimal expansion $a_0.a_1a_2a_3\dots$ and write

$$x = a_0.a_1a_2a_3a_4\dots$$

Decimal expansions

We say that $x > 0$ has the decimal expansion $a_0.a_1a_2a_3\dots$ and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Decimal expansions

We say that $x > 0$ has the decimal expansion $a_0.a_1a_2a_3\dots$ and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

Decimal expansions

We say that $x > 0$ has the decimal expansion $a_0.a_1a_2a_3\dots$ and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- 1 we haven't defined limits

Decimal expansions

We say that $x > 0$ has the decimal expansion $a_0.a_1a_2a_3\dots$ and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- 1 we haven't defined limits
- 2 with this definition, numbers have unique decimal expansions!

Decimal expansions

We say that $x > 0$ has the decimal expansion $a_0.a_1a_2a_3\dots$ and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- 1 we haven't defined limits
- 2 with this definition, numbers have unique decimal expansions!

$$1 \neq 0.999999999\dots$$

Decimal expansions

We say that $x > 0$ has the decimal expansion $a_0.a_1a_2a_3\dots$ and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all $n \in \mathbb{Z}_+$,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- ① we haven't defined limits
- ② with this definition, numbers have unique decimal expansions!

$$1 \neq 0.999999999\dots$$

$$1 \not\leq 0 + \frac{9}{10} + \frac{9}{100} + \dots + \frac{9+1}{10^n} = 1.$$

Outline

- 1 Real Analysis Lecture 3
 - Suprema and Infima
 - Decimal expansions
 - The Triangle Inequality

Absolute value

The **absolute value** of $x \in \mathbb{R}$ is

$$|x| = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$$

In particular,

$$0 \leq |x|$$

and also

$$-|x| \leq x \leq |x|.$$

Challenge!

Problem

Prove Apostol Theorem 1.21 that if $a \geq 0$, then $|x| \leq a$ if and only if $-a \leq x \leq a$.

Solution

If $|x| \leq a$ then $-a \leq -|x|$

Challenge!

Problem

Prove Apostol Theorem 1.21 that if $a \geq 0$, then $|x| \leq a$ if and only if $-a \leq x \leq a$.

Solution

If $|x| \leq a$ then $-a \leq -|x|$ and therefore

$$-a \leq -|x| \leq x \leq |x| \leq a$$

.

Challenge!

Problem

Prove Apostol Theorem 1.21 that if $a \geq 0$, then $|x| \leq a$ if and only if $-a \leq x \leq a$.

Solution

If $|x| \leq a$ then $-a \leq -|x|$ and therefore

$$-a \leq -|x| \leq x \leq |x| \leq a$$

. Conversely, if $-a \leq x \leq a$ then

Challenge!

Problem

Prove Apostol Theorem 1.21 that if $a \geq 0$, then $|x| \leq a$ if and only if $-a \leq x \leq a$.

Solution

If $|x| \leq a$ then $-a \leq -|x|$ and therefore

$$-a \leq -|x| \leq x \leq |x| \leq a$$

. Conversely, if $-a \leq x \leq a$ then

$$x \geq 0 \Rightarrow |x| = x \leq a$$

Challenge!

Problem

Prove Apostol Theorem 1.21 that if $a \geq 0$, then $|x| \leq a$ if and only if $-a \leq x \leq a$.

Solution

If $|x| \leq a$ then $-a \leq -|x|$ and therefore

$$-a \leq -|x| \leq x \leq |x| \leq a$$

. Conversely, if $-a \leq x \leq a$ then

$$x \geq 0 \Rightarrow |x| = x \leq a$$

$$x \leq 0 \Rightarrow |x| = -x \leq -(-a) = a$$

Basic triangle inequality

Theorem (Triangle inequality)

For any real numbers $x, y \in \mathbb{R}$ we have

$$|x + y| \leq |x| + |y|.$$

Proof.

$$-|x| \leq x \leq |x| \quad \text{and} \quad -|y| \leq y \leq |y|$$

Basic triangle inequality

Theorem (Triangle inequality)

For any real numbers $x, y \in \mathbb{R}$ we have

$$|x + y| \leq |x| + |y|.$$

Proof.

$$-|x| \leq x \leq |x| \quad \text{and} \quad -|y| \leq y \leq |y|$$

Adding these together, we get

$$-(|x| + |y|) \leq x + y \leq |x| + |y|$$

Basic triangle inequality

Theorem (Triangle inequality)

For any real numbers $x, y \in \mathbb{R}$ we have

$$|x + y| \leq |x| + |y|.$$

Proof.

$$-|x| \leq x \leq |x| \quad \text{and} \quad -|y| \leq y \leq |y|$$

Adding these together, we get

$$-(|x| + |y|) \leq x + y \leq |x| + |y|$$

It follows from the previous theorem that

$$|x + y| \leq |x| + |y|$$

Advanced triangle inequality

Theorem (Triangle inequality)

For any real numbers $x_1, x_2, \dots, x_n \in \mathbb{R}$ we have

$$|x_1 + x_2 + \cdots + x_n| \leq |x_1| + |x_2| + \cdots + |x_n|.$$

Advanced triangle inequality

Theorem (Triangle inequality)

For any real numbers $x_1, x_2, \dots, x_n \in \mathbb{R}$ we have

$$|x_1 + x_2 + \dots + x_n| \leq |x_1| + |x_2| + \dots + |x_n|.$$

Proof.

Induction. □

Higher-dimensional triangle inequality

Theorem (Cauchy-Schwartz Inequality (Apostol Theorem 1.23))

If x_1, \dots, x_n and y_1, \dots, y_n are real numbers, then

$$\left(\sum_{k=1}^n x_k y_k \right)^2 \leq \left(\sum_{k=1}^n x_k^2 \right) \left(\sum_{k=1}^n y_k^2 \right)$$

If y_k isn't always zero, then equality holds if and only if there exists $t \in \mathbb{R}$ with $x_k = ty_k$ for all k .

Higher-dimensional triangle inequality

Proof.

$$\sum_{k=1}^n (x_k + ty_k)^2 \geq 0, \quad \text{with equality iff all terms zero}$$

$$\sum_{k=1}^n x_k^2 + 2t \sum_{k=1}^n x_k y_k + t^2 \sum_{k=1}^n y_k^2 \geq 0$$

Take $t = -(\sum_{k=1}^n x_k y_k) / (\sum_{k=1}^n y_k^2)$:

$$\sum_{k=1}^n x_k^2 - \frac{(\sum_{k=1}^n x_k y_k)^2}{(\sum_{k=1}^n y_k^2)} \geq 0.$$

