

# MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics  
California State University Fullerton

September 16, 2024





# Ultimate Cantor diagonalization

The **power set**  $\mathcal{P}(A)$  of a set  $A$  is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

# Ultimate Cantor diagonalization

The **power set**  $\mathcal{P}(A)$  of a set  $A$  is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

## Theorem

*The cardinality of  $\mathcal{P}(A)$  is strictly larger than  $A$ .*

# Ultimate Cantor diagonalization

The **power set**  $\mathcal{P}(A)$  of a set  $A$  is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

## Theorem

*The cardinality of  $\mathcal{P}(A)$  is strictly larger than  $A$ .*

## Proof.

# Ultimate Cantor diagonalization

The **power set**  $\mathcal{P}(A)$  of a set  $A$  is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

## Theorem

*The cardinality of  $\mathcal{P}(A)$  is strictly larger than  $A$ .*

## Proof.

Suppose  $f : A \rightarrow \mathcal{P}(A)$  is surjective.

# Ultimate Cantor diagonalization

The **power set**  $\mathcal{P}(A)$  of a set  $A$  is

$$\mathcal{P}(A) = \{S : S \subseteq A\}.$$

## Theorem

*The cardinality of  $\mathcal{P}(A)$  is strictly larger than  $A$ .*

## Proof.

Suppose  $f : A \rightarrow \mathcal{P}(A)$  is surjective.

Consider the set

$$S = \{a \in A : a \notin f(a)\}.$$





# Challenge

# Challenge

Since  $f$  is surjective,  $S = f(x)$  for some  $x \in A$ .

# Challenge

Since  $f$  is surjective,  $S = f(x)$  for some  $x \in A$ .

## Problem

Consider the statement  $x \in S$ . What can you conclude?



# Unions of sets

Unions:

# Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

# Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

# Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

$$A_1 \cup A_2 \cup \cdots \cup A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$



# Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

$$A_1 \cup A_2 \cup \cdots \cup A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

NOTATION:

$$\bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \cdots \cup A_n.$$

# Intersections of sets

Intersections:

# Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

# Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

# Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

$$A_1 \cap A_2 \cap \cdots \cap A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

# Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

$$A_1 \cap A_2 \cap \cdots \cap A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

NOTATION:

$$\bigcap_{i=1}^n A_i = A_1 \cap A_2 \cap \cdots \cap A_n.$$

# Complements of sets

# Complements of sets

The **complement** of  $A$  relative to  $B$  is

$$B - A = \{b \in B : b \notin A\}.$$



# Complements of sets

The **complement** of  $A$  relative to  $B$  is

$$B - A = \{b \in B : b \notin A\}.$$

## Theorem

*De Morgan's Laws*

# Complements of sets

The **complement** of  $A$  relative to  $B$  is

$$B - A = \{b \in B : b \notin A\}.$$

## Theorem

*De Morgan's Laws*

$$B - \bigcup_{i=1}^n A_i = \bigcap_{i=1}^n (B - A_i)$$

# Complements of sets

The **complement** of  $A$  relative to  $B$  is

$$B - A = \{b \in B : b \notin A\}.$$

## Theorem

*De Morgan's Laws*

$$B - \bigcup_{i=1}^n A_i = \bigcap_{i=1}^n (B - A_i)$$

$$B - \bigcap_{i=1}^n A_i = \bigcup_{i=1}^n (B - A_i)$$

# Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

# Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- $I$  any set, called the **index set**

# Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- $I$  any set, called the **index set**
- $A_i$  is a set for all  $i$

# Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- $I$  any set, called the **index set**
- $A_i$  is a set for all  $i$

$$\bigcup_{i \in I} A_i = \{x : x \in A_i, \text{ for some } i \in I\}$$

# Families of sets

A **family of sets** is a collection

$$\{A_i; i \in I\}$$

- $I$  any set, called the **index set**
- $A_i$  is a set for all  $i$

$$\bigcup_{i \in I} A_i = \{x : x \in A_i, \text{ for some } i \in I\}$$

$$\bigcap_{i \in I} A_i = \{x : x \in A_i, \text{ for all } i \in I\}$$



# Challenge

- index set  $I = \mathbb{Z}_+$
- family of sets  $\{A_i : i \in I\}$
- $A_i = [0, i)$

## Problem

Determine  $\bigcup_{i \in I} A_i$ .

# Challenge

- index set  $I = \mathbb{Z}_+$
- family of sets  $\{A_i : i \in I\}$
- $A_i = [0, i)$

## Problem

Determine  $\bigcap_{i \in I} A_i$ .