

# MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics  
California State University Fullerton

September 9, 2024

# Outline

- 1 Real Analysis Lecture 3
  - Decimal expansions
  - The Triangle Inequality
  - Sets, Relations, Functions

# Outline

- 1 Real Analysis Lecture 3
  - Decimal expansions
  - The Triangle Inequality
  - Sets, Relations, Functions

# Decimal approximations

A **finite decimal expansion** is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \cdots + \frac{a_n}{10^n},$$

where  $a_0 \in \mathbb{Z}_+$  and  $0 \leq a_k \leq 9$  for  $1 \leq k \leq n$ .

# Decimal approximations

A **finite decimal expansion** is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \cdots + \frac{a_n}{10^n},$$

where  $a_0 \in \mathbb{Z}_+$  and  $0 \leq a_k \leq 9$  for  $1 \leq k \leq n$ .

**Notation:**

$$a_0.a_1a_2a_3\dots a_n.$$

# Decimal approximations

A **finite decimal expansion** is an expression

$$r = a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \frac{a_3}{10^3} + \cdots + \frac{a_n}{10^n},$$

where  $a_0 \in \mathbb{Z}_+$  and  $0 \leq a_k \leq 9$  for  $1 \leq k \leq n$ .

**Notation:**

$$a_0.a_1a_2a_3 \dots a_n.$$

Any positive real number  $x > 0$  can be approximated by a finite decimal expansion.

# Decimal approximations

## Theorem (Apostol Theorem 1.20)

*For any real  $x > 0$  and  $n \in \mathbb{Z}_+$ , there exists a finite decimal expansion  $r_n = a_0.a_1a_2 \dots a_n$  with*

$$r_n \leq x < r_n + \frac{1}{10^n}.$$

# Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$



# Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set  $A$  is a set of integers which is bounded above by  $x$ , so it has a maximum  $a_0$ .

# Decimal approximations

Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set  $A$  is a set of integers which is bounded above by  $x$ , so it has a maximum  $a_0$ .

Then clearly  $x_1 = x - a_0 \in [0, 1)$ .

# Decimal approximations

## Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set  $A$  is a set of integers which is bounded above by  $x$ , so it has a maximum  $a_0$ .

Then clearly  $x_1 = x - a_0 \in [0, 1)$ .

Define  $a_1, a_2, a_3, \dots$  and  $x_1, x_2, x_3, \dots$  recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and  $x_{k+1} = 10x_k - a_k$ .

# Decimal approximations

## Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set  $A$  is a set of integers which is bounded above by  $x$ , so it has a maximum  $a_0$ .

Then clearly  $x_1 = x - a_0 \in [0, 1)$ .

Define  $a_1, a_2, a_3, \dots$  and  $x_1, x_2, x_3, \dots$  recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and  $x_{k+1} = 10x_k - a_k$ . Then  $0 \leq a_k \leq 9$  for all  $k \geq 1$

# Decimal approximations

## Proof.

Consider the set

$$A = \{a \in \mathbb{Z} : a \leq x\}.$$

The set  $A$  is a set of integers which is bounded above by  $x$ , so it has a maximum  $a_0$ .

Then clearly  $x_1 = x - a_0 \in [0, 1)$ .

Define  $a_1, a_2, a_3, \dots$  and  $x_1, x_2, x_3, \dots$  recursively by

$$a_k = \max\{a \in \mathbb{Z}_+ : a \leq 10x_k\}$$

and  $x_{k+1} = 10x_k - a_k$ . Then  $0 \leq a_k \leq 9$  for all  $k \geq 1$  and

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \cdots + \frac{a_n + 1}{10^n}$$



# Decimal expansions

We say that  $x > 0$  has the decimal expansion  $a_0.a_1a_2a_3\dots$  and write

$$x = a_0.a_1a_2a_3a_4\dots$$

# Decimal expansions

We say that  $x > 0$  has the decimal expansion  $a_0.a_1a_2a_3\dots$  and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all  $n \in \mathbb{Z}_+$ ,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

# Decimal expansions

We say that  $x > 0$  has the decimal expansion  $a_0.a_1a_2a_3\dots$  and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all  $n \in \mathbb{Z}_+$ ,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:



# Decimal expansions

We say that  $x > 0$  has the decimal expansion  $a_0.a_1a_2a_3\dots$  and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all  $n \in \mathbb{Z}_+$ ,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- 1 we haven't defined limits

# Decimal expansions

We say that  $x > 0$  has the decimal expansion  $a_0.a_1a_2a_3\dots$  and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all  $n \in \mathbb{Z}_+$ ,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- 1 we haven't defined limits
- 2 with this definition, numbers have unique decimal expansions!

# Decimal expansions

We say that  $x > 0$  has the decimal expansion  $a_0.a_1a_2a_3\dots$  and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all  $n \in \mathbb{Z}_+$ ,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- 1 we haven't defined limits
- 2 with this definition, numbers have unique decimal expansions!

$$1 \neq 0.999999999\dots$$

# Decimal expansions

We say that  $x > 0$  has the decimal expansion  $a_0.a_1a_2a_3\dots$  and write

$$x = a_0.a_1a_2a_3a_4\dots$$

if for all  $n \in \mathbb{Z}_+$ ,

$$a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n}{10^n} \leq x < a_0 + \frac{a_1}{10} + \frac{a_2}{100} + \dots + \frac{a_n + 1}{10^n}.$$

Note: this is slightly different than the usual limit meaning, for two good reasons:

- ① we haven't defined limits
- ② with this definition, numbers have unique decimal expansions!

$$1 \neq 0.999999999\dots$$

$$1 \not\leq 0 + \frac{9}{10} + \frac{9}{100} + \dots + \frac{9+1}{10^n} = 1.$$

# Challenge!

## Problem

Find the decimal expansion of  $1/7$ .

# Challenge!

## Problem

Find a rational number whose decimal expansion is

$$0.45454545\dots$$

# Challenge!

## Problem

Which kinds of numbers have decimal expansions that end?  
(Meaning that after a while, all the decimals are zero?)

# Challenge!

## Problem

Which kinds of numbers have decimal expansions that repeat?



# Outline

- 1 Real Analysis Lecture 3
  - Decimal expansions
  - The Triangle Inequality
  - Sets, Relations, Functions

# Absolute value

The **absolute value** of  $x \in \mathbb{R}$  is

$$|x| = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$$

In particular,

$$0 \leq |x|$$

and also

$$-|x| \leq x \leq |x|.$$

# Challenge!

## Problem

Prove Apostol Theorem 1.21 that if  $a \geq 0$ , then  $|x| \leq a$  if and only if  $-a \leq x \leq a$ .

# Challenge!

## Problem

Prove Apostol Theorem 1.21 that if  $a \geq 0$ , then  $|x| \leq a$  if and only if  $-a \leq x \leq a$ .

## Solution

If  $|x| \leq a$  then  $-a \leq -|x|$

# Challenge!

## Problem

Prove Apostol Theorem 1.21 that if  $a \geq 0$ , then  $|x| \leq a$  if and only if  $-a \leq x \leq a$ .

## Solution

If  $|x| \leq a$  then  $-a \leq -|x|$  and therefore

$$-a \leq -|x| \leq x \leq |x| \leq a$$

# Challenge!

## Problem

Prove Apostol Theorem 1.21 that if  $a \geq 0$ , then  $|x| \leq a$  if and only if  $-a \leq x \leq a$ .

## Solution

If  $|x| \leq a$  then  $-a \leq -|x|$  and therefore

$$-a \leq -|x| \leq x \leq |x| \leq a$$

Conversely, if  $-a \leq x \leq a$  then

# Challenge!

## Problem

Prove Apostol Theorem 1.21 that if  $a \geq 0$ , then  $|x| \leq a$  if and only if  $-a \leq x \leq a$ .

## Solution

If  $|x| \leq a$  then  $-a \leq -|x|$  and therefore

$$-a \leq -|x| \leq x \leq |x| \leq a$$

Conversely, if  $-a \leq x \leq a$  then

$$x \geq 0 \Rightarrow |x| = x \leq a$$

# Challenge!

## Problem

Prove Apostol Theorem 1.21 that if  $a \geq 0$ , then  $|x| \leq a$  if and only if  $-a \leq x \leq a$ .

## Solution

If  $|x| \leq a$  then  $-a \leq -|x|$  and therefore

$$-a \leq -|x| \leq x \leq |x| \leq a$$

Conversely, if  $-a \leq x \leq a$  then

$$x \geq 0 \Rightarrow |x| = x \leq a$$

$$x \leq 0 \Rightarrow |x| = -x \leq -(-a) = a$$



# Basic triangle inequality

## Theorem (Triangle inequality)

*For any real numbers  $x, y \in \mathbb{R}$  we have*

$$|x + y| \leq |x| + |y|.$$

# Basic triangle inequality

## Theorem (Triangle inequality)

*For any real numbers  $x, y \in \mathbb{R}$  we have*

$$|x + y| \leq |x| + |y|.$$

Proof.

$$-|x| \leq x \leq |x| \quad \text{and} \quad -|y| \leq y \leq |y|$$

# Basic triangle inequality

## Theorem (Triangle inequality)

*For any real numbers  $x, y \in \mathbb{R}$  we have*

$$|x + y| \leq |x| + |y|.$$

## Proof.

$$-|x| \leq x \leq |x| \quad \text{and} \quad -|y| \leq y \leq |y|$$

Adding these together, we get

$$-(|x| + |y|) \leq x + y \leq |x| + |y|$$

# Basic triangle inequality

## Theorem (Triangle inequality)

*For any real numbers  $x, y \in \mathbb{R}$  we have*

$$|x + y| \leq |x| + |y|.$$

## Proof.

$$-|x| \leq x \leq |x| \quad \text{and} \quad -|y| \leq y \leq |y|$$

Adding these together, we get

$$-(|x| + |y|) \leq x + y \leq |x| + |y|$$

It follows from the previous theorem that

$$|x + y| \leq |x| + |y|$$

# Advanced triangle inequality

## Theorem (Triangle inequality)

*For any real numbers  $x_1, x_2, \dots, x_n \in \mathbb{R}$  we have*

$$|x_1 + x_2 + \cdots + x_n| \leq |x_1| + |x_2| + \cdots + |x_n|.$$

# Advanced triangle inequality

## Theorem (Triangle inequality)

*For any real numbers  $x_1, x_2, \dots, x_n \in \mathbb{R}$  we have*

$$|x_1 + x_2 + \dots + x_n| \leq |x_1| + |x_2| + \dots + |x_n|.$$

Proof.

Induction. □

# Higher-dimensional triangle inequality

**Theorem (Cauchy-Schwartz Inequality (Apostol Theorem 1.23))**

*If  $x_1, \dots, x_n$  and  $y_1, \dots, y_n$  are real numbers, then*

$$\left( \sum_{k=1}^n x_k y_k \right)^2 \leq \left( \sum_{k=1}^n x_k^2 \right) \left( \sum_{k=1}^n y_k^2 \right)$$

*If  $y_k$  isn't always zero, then equality holds if and only if there exists  $t \in \mathbb{R}$  with  $x_k = ty_k$  for all  $k$ .*

# Higher-dimensional triangle inequality

## Theorem (Cauchy-Schwartz Inequality (Apostol Theorem 1.23))

*If  $x_1, \dots, x_n$  and  $y_1, \dots, y_n$  are real numbers, then*

$$\left( \sum_{k=1}^n x_k y_k \right)^2 \leq \left( \sum_{k=1}^n x_k^2 \right) \left( \sum_{k=1}^n y_k^2 \right)$$

*If  $y_k$  isn't always zero, then equality holds if and only if there exists  $t \in \mathbb{R}$  with  $x_k = ty_k$  for all  $k$ .*

Vector version:

$$(\vec{x} \cdot \vec{y})^2 \leq |\vec{x}|^2 |\vec{y}|^2.$$



# Higher-dimensional triangle inequality

Proof.

# Higher-dimensional triangle inequality

Proof.

$$\sum_{k=1}^n (x_k + ty_k)^2 \geq 0, \quad \text{with equality iff all terms zero}$$

# Higher-dimensional triangle inequality

Proof.

$$\sum_{k=1}^n (x_k + ty_k)^2 \geq 0, \quad \text{with equality iff all terms zero}$$

$$\sum_{k=1}^n x_k^2 + 2t \sum_{k=1}^n x_k y_k + t^2 \sum_{k=1}^n y_k^2 \geq 0$$

# Higher-dimensional triangle inequality

Proof.

$$\sum_{k=1}^n (x_k + ty_k)^2 \geq 0, \quad \text{with equality iff all terms zero}$$

$$\sum_{k=1}^n x_k^2 + 2t \sum_{k=1}^n x_k y_k + t^2 \sum_{k=1}^n y_k^2 \geq 0$$

Take  $t = -(\sum_{k=1}^n x_k y_k) / (\sum_{k=1}^n y_k^2)$  :

# Higher-dimensional triangle inequality

Proof.

$$\sum_{k=1}^n (x_k + ty_k)^2 \geq 0, \quad \text{with equality iff all terms zero}$$

$$\sum_{k=1}^n x_k^2 + 2t \sum_{k=1}^n x_k y_k + t^2 \sum_{k=1}^n y_k^2 \geq 0$$

Take  $t = -(\sum_{k=1}^n x_k y_k) / (\sum_{k=1}^n y_k^2)$  :

$$\sum_{k=1}^n x_k^2 - \frac{(\sum_{k=1}^n x_k y_k)^2}{(\sum_{k=1}^n y_k^2)} \geq 0.$$



# Higher-dimensional triangle inequality

## Theorem (Minkowski inequality)

*For any real numbers  $x_1, x_2, \dots, x_n \in \mathbb{R}$  and  $y_1, y_2, \dots, y_n \in \mathbb{R}$  we have*

$$\left( \sum_{k=1}^n (x_k + y_k)^2 \right)^{1/2} \leq \left( \sum_{k=1}^n x_k^2 \right)^{1/2} + \left( \sum_{k=1}^n y_k^2 \right)^{1/2}$$

# Higher-dimensional triangle inequality

## Theorem (Minkowski inequality)

For any real numbers  $x_1, x_2, \dots, x_n \in \mathbb{R}$  and  $y_1, y_2, \dots, y_n \in \mathbb{R}$  we have

$$\left( \sum_{k=1}^n (x_k + y_k)^2 \right)^{1/2} \leq \left( \sum_{k=1}^n x_k^2 \right)^{1/2} + \left( \sum_{k=1}^n y_k^2 \right)^{1/2}$$

Vector version:

$$|\vec{x} + \vec{y}| \leq |\vec{x}| + |\vec{y}|.$$

# Higher-dimensional triangle inequality

## Theorem (Minkowski inequality)

For any real numbers  $x_1, x_2, \dots, x_n \in \mathbb{R}$  and  $y_1, y_2, \dots, y_n \in \mathbb{R}$  we have

$$\left( \sum_{k=1}^n (x_k + y_k)^2 \right)^{1/2} \leq \left( \sum_{k=1}^n x_k^2 \right)^{1/2} + \left( \sum_{k=1}^n y_k^2 \right)^{1/2}$$

Vector version:

$$|\vec{x} + \vec{y}| \leq |\vec{x}| + |\vec{y}|.$$

A higher dimensional triangle inequality!



# Higher-dimensional triangle inequality

Proof.

By the triangle inequality:

# Higher-dimensional triangle inequality

Proof.

By the triangle inequality:

$$\begin{aligned}\sum_{k=1}^n (x_k + y_k)^2 &\leq \sum_{k=1}^n (|x_k| + |y_k|)|x_k + y_k| \\ &= \sum_{k=1}^n |x_k| \cdot |x_k + y_k| + \sum_{k=1}^n |y_k| \cdot |x_k + y_k|\end{aligned}$$

# Higher-dimensional triangle inequality

Proof.

By the triangle inequality:

$$\begin{aligned}\sum_{k=1}^n (x_k + y_k)^2 &\leq \sum_{k=1}^n (|x_k| + |y_k|)|x_k + y_k| \\ &= \sum_{k=1}^n |x_k| \cdot |x_k + y_k| + \sum_{k=1}^n |y_k| \cdot |x_k + y_k|\end{aligned}$$

Now applying the Cauchy-Schwartz inequality:

# Higher-dimensional triangle inequality

Proof.

By the triangle inequality:

$$\begin{aligned}\sum_{k=1}^n (x_k + y_k)^2 &\leq \sum_{k=1}^n (|x_k| + |y_k|) |x_k + y_k| \\ &= \sum_{k=1}^n |x_k| \cdot |x_k + y_k| + \sum_{k=1}^n |y_k| \cdot |x_k + y_k|\end{aligned}$$

Now applying the Cauchy-Schwartz inequality:

$$\begin{aligned}&\leq \left( \sum_{k=1}^n |x_k|^2 \right)^{1/2} \left( \sum_{k=1}^n |x_k + y_k|^2 \right)^{1/2} + \left( \sum_{k=1}^n |y_k|^2 \right)^{1/2} \left( \sum_{k=1}^n |x_k + y_k|^2 \right)^{1/2} \\ &= \left[ \left( \sum_{k=1}^n |x_k|^2 \right)^{1/2} + \left( \sum_{k=1}^n |y_k|^2 \right)^{1/2} \right] \left( \sum_{k=1}^n |x_k + y_k|^2 \right)^{1/2}\end{aligned}$$



# Higher-dimensional triangle inequality

Proof.

By the triangle inequality:

$$\begin{aligned}\sum_{k=1}^n (x_k + y_k)^2 &\leq \sum_{k=1}^n (|x_k| + |y_k|) |x_k + y_k| \\ &= \sum_{k=1}^n |x_k| \cdot |x_k + y_k| + \sum_{k=1}^n |y_k| \cdot |x_k + y_k|\end{aligned}$$

Now applying the Cauchy-Schwartz inequality:

$$\begin{aligned}&\leq \left( \sum_{k=1}^n |x_k|^2 \right)^{1/2} \left( \sum_{k=1}^n |x_k + y_k|^2 \right)^{1/2} + \left( \sum_{k=1}^n |y_k|^2 \right)^{1/2} \left( \sum_{k=1}^n |x_k + y_k|^2 \right)^{1/2} \\ &= \left[ \left( \sum_{k=1}^n |x_k|^2 \right)^{1/2} + \left( \sum_{k=1}^n |y_k|^2 \right)^{1/2} \right] \left( \sum_{k=1}^n |x_k + y_k|^2 \right)^{1/2}\end{aligned}$$



# Outline

- 1 Real Analysis Lecture 3
  - Decimal expansions
  - The Triangle Inequality
  - Sets, Relations, Functions

# Set basics

Intuitively, a **set**  $A$  is a "collection of things" which we call the **elements** of  $A$ .

# Set basics

Intuitively, a **set**  $A$  is a "collection of things" which we call the **elements** of  $A$ .

- in practice, this is a bad definition (Russell's Paradox)



# Set basics

Intuitively, a **set**  $A$  is a "collection of things" which we call the **elements** of  $A$ .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

# Set basics

Intuitively, a **set**  $A$  is a "collection of things" which we call the **elements** of  $A$ .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

# Set basics

Intuitively, a **set**  $A$  is a "collection of things" which we call the **elements** of  $A$ .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$

# Set basics

Intuitively, a **set**  $A$  is a "collection of things" which we call the **elements** of  $A$ .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$

# Set basics

Intuitively, a **set**  $A$  is a "collection of things" which we call the **elements** of  $A$ .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$
- empty set  $\emptyset$

# Set basics

Intuitively, a **set**  $A$  is a "collection of things" which we call the **elements** of  $A$ .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$
- empty set  $\emptyset$
- $\{\heartsuit, \text{Fall}, \{\emptyset\}\}$

# Set basics

Intuitively, a **set**  $A$  is a "collection of things" which we call the **elements** of  $A$ .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$
- empty set  $\emptyset$
- $\{\heartsuit, \text{Fall}, \{\emptyset\}\}$
- $\{n \in \mathbb{Z} : n \text{ is prime}\}$

# Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.



# Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

# Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

# Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

- ordered pairs

# Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

- ordered pairs

$$(a, b) = \{a, \{a, b\}\}$$

# Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

- ordered pairs

$$(a, b) = \{a, \{a, b\}\}$$

- even relations and functions are sets!

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

$$(a, b) = (c, d) \text{ if and only if } \{a, \{a, b\}\} = \{c, \{c, d\}\}$$

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

$(a, b) = (c, d)$  if and only if  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if  $a = c$  and  $b = d$ , then  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$



# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

$(a, b) = (c, d)$  if and only if  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if  $a = c$  and  $b = d$ , then  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

$(a, b) = (c, d)$  if and only if  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if  $a = c$  and  $b = d$ , then  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

Suppose  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$ .

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

$(a, b) = (c, d)$  if and only if  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if  $a = c$  and  $b = d$ , then  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

Suppose  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$ .

Two possible cases:

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

$(a, b) = (c, d)$  if and only if  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if  $a = c$  and  $b = d$ , then  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

Suppose  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$ .

Two possible cases:

Case I:  $a = c$  and  $\{a, b\} = \{c, d\}$

Case II:  $a = \{c, d\}$  and  $\{a, b\} = c$

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

$(a, b) = (c, d)$  if and only if  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if  $a = c$  and  $b = d$ , then  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

Suppose  $\{a, \{a, b\}\} = \{c, \{c, d\}\}$ .

Two possible cases:

Case I:  $a = c$  and  $\{a, b\} = \{c, d\}$

Case II:  $a = \{c, d\}$  and  $\{a, b\} = c$

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

$$\text{Case I: } a = c \text{ and } \{a, b\} = \{c, d\}$$

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

Case I:  $a = c$  and  $\{a, b\} = \{c, d\}$

Since  $\{a, b\} = \{c, d\}$ , we know  $b \in \{c, d\}$ .

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

Case I:  $a = c$  and  $\{a, b\} = \{c, d\}$

Since  $\{a, b\} = \{c, d\}$ , we know  $b \in \{c, d\}$ .

Therefore  $b = c$  or  $b = d$ .

If  $b = d$ , we're done!



# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

$$\text{Case I: } a = c \text{ and } \{a, b\} = \{c, d\}$$

Since  $\{a, b\} = \{c, d\}$ , we know  $b \in \{c, d\}$ .

Therefore  $b = c$  or  $b = d$ .

If  $b = d$ , we're done! ... so assume instead that  $b = c$

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

Case I:  $a = c$  and  $\{a, b\} = \{c, d\}$

Since  $\{a, b\} = \{c, d\}$ , we know  $b \in \{c, d\}$ .

Therefore  $b = c$  or  $b = d$ .

If  $b = d$ , we're done! ... so assume instead that  $b = c$

Then  $a = c$  implies  $a = b$ .

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

Case I:  $a = c$  and  $\{a, b\} = \{c, d\}$

Since  $\{a, b\} = \{c, d\}$ , we know  $b \in \{c, d\}$ .

Therefore  $b = c$  or  $b = d$ .

If  $b = d$ , we're done! ... so assume instead that  $b = c$

Then  $a = c$  implies  $a = b$ .

Therefore  $\{c, d\} = \{a, b\} = \{a, a\} = \{a\}$ .

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

Case I:  $a = c$  and  $\{a, b\} = \{c, d\}$

Since  $\{a, b\} = \{c, d\}$ , we know  $b \in \{c, d\}$ .

Therefore  $b = c$  or  $b = d$ .

If  $b = d$ , we're done! ... so assume instead that  $b = c$

Then  $a = c$  implies  $a = b$ .

Therefore  $\{c, d\} = \{a, b\} = \{a, a\} = \{a\}$ .

It follows that  $d = c = b = a$ .

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

$$\text{Case II: } a = \{c, d\} \text{ and } \{a, b\} = c$$

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

$$\text{Case II: } a = \{c, d\} \text{ and } \{a, b\} = c$$

This would imply that  $c \in a$  and  $a \in c$ .

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

$$\text{Case II: } a = \{c, d\} \text{ and } \{a, b\} = c$$

This would imply that  $c \in a$  and  $a \in c$ .

This can be shown to contradict the ZF Axioms of Set Theory.

# Challenge!

## Problem

Prove that for ordered pairs  $(a, b)$  and  $(c, d)$  that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

## Solution

$$\text{Case II: } a = \{c, d\} \text{ and } \{a, b\} = c$$

This would imply that  $c \in a$  and  $a \in c$ .

This can be shown to contradict the ZF Axioms of Set Theory.  
Specifically the regularity axiom for the set  $\{a, c\}$ ...



# Relations

The **Cartesian product** of  $A$  and  $B$  is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

# Relations

The **Cartesian product** of  $A$  and  $B$  is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation**  $\mathcal{R}$  from  $A$  to  $B$  is a subset of  $A \times B$ .

# Relations

The **Cartesian product** of  $A$  and  $B$  is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation**  $\mathcal{R}$  from  $A$  to  $B$  is a subset of  $A \times B$ .

**NOTATION:**  $a\mathcal{R}b$  means  $(a, b) \in \mathcal{R}$ .

# Relations

The **Cartesian product** of  $A$  and  $B$  is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation**  $\mathcal{R}$  from  $A$  to  $B$  is a subset of  $A \times B$ .

**NOTATION:**  $a\mathcal{R}b$  means  $(a, b) \in \mathcal{R}$ .

Domain and codomain:

# Relations

The **Cartesian product** of  $A$  and  $B$  is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation**  $\mathcal{R}$  from  $A$  to  $B$  is a subset of  $A \times B$ .

**NOTATION:**  $a\mathcal{R}b$  means  $(a, b) \in \mathcal{R}$ .

Domain and codomain:

$$\begin{aligned}\text{dom}(\mathcal{R}) &= \{a \in A : \exists b \in B, a\mathcal{R}b\} \\ \text{codom}(\mathcal{R}) &= \{b \in B : \exists a \in A, a\mathcal{R}b\}\end{aligned}$$

# Relations

The **Cartesian product** of  $A$  and  $B$  is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation**  $\mathcal{R}$  from  $A$  to  $B$  is a subset of  $A \times B$ .

**NOTATION:**  $a\mathcal{R}b$  means  $(a, b) \in \mathcal{R}$ .

Domain and codomain:

$$\begin{aligned}\text{dom}(\mathcal{R}) &= \{a \in A : \exists b \in B, a\mathcal{R}b\} \\ \text{codom}(\mathcal{R}) &= \{b \in B : \exists a \in A, a\mathcal{R}b\}\end{aligned}$$

# Relations

A relation  $\mathcal{R}$  from  $A$  to  $A$  is called a **relation on  $A$** . A relation on  $A$  is

# Relations

A relation  $\mathcal{R}$  from  $A$  to  $A$  is called a **relation on  $A$** . A relation on  $A$  is

- **reflexive** if  $a\mathcal{R}a$  for all  $a \in A$



# Relations

A relation  $\mathcal{R}$  from  $A$  to  $A$  is called a **relation on  $A$** . A relation on  $A$  is

- **reflexive** if  $a\mathcal{R}a$  for all  $a \in A$
- **symmetric** if  $a\mathcal{R}b$  implies  $b\mathcal{R}a$  for all  $a, b \in A$

# Relations

A relation  $\mathcal{R}$  from  $A$  to  $A$  is called a **relation on  $A$** . A relation on  $A$  is

- **reflexive** if  $a\mathcal{R}a$  for all  $a \in A$
- **symmetric** if  $a\mathcal{R}b$  implies  $b\mathcal{R}a$  for all  $a, b \in A$
- **transitive** if  $a\mathcal{R}b$  and  $b\mathcal{R}c$  implies  $a\mathcal{R}c$  for all  $a, b, c \in A$

# Relations

A relation  $\mathcal{R}$  from  $A$  to  $A$  is called a **relation on  $A$** . A relation on  $A$  is

- **reflexive** if  $a\mathcal{R}a$  for all  $a \in A$
- **symmetric** if  $a\mathcal{R}b$  implies  $b\mathcal{R}a$  for all  $a, b \in A$
- **transitive** if  $a\mathcal{R}b$  and  $b\mathcal{R}c$  implies  $a\mathcal{R}c$  for all  $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

# Relations

A relation  $\mathcal{R}$  from  $A$  to  $A$  is called a **relation on  $A$** . A relation on  $A$  is

- **reflexive** if  $a\mathcal{R}a$  for all  $a \in A$
- **symmetric** if  $a\mathcal{R}b$  implies  $b\mathcal{R}a$  for all  $a, b \in A$
- **transitive** if  $a\mathcal{R}b$  and  $b\mathcal{R}c$  implies  $a\mathcal{R}c$  for all  $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Examples:

# Relations

A relation  $\mathcal{R}$  from  $A$  to  $A$  is called a **relation on  $A$** . A relation on  $A$  is

- **reflexive** if  $a\mathcal{R}a$  for all  $a \in A$
- **symmetric** if  $a\mathcal{R}b$  implies  $b\mathcal{R}a$  for all  $a, b \in A$
- **transitive** if  $a\mathcal{R}b$  and  $b\mathcal{R}c$  implies  $a\mathcal{R}c$  for all  $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Examples:

# Relations

A relation  $\mathcal{R}$  from  $A$  to  $A$  is called a **relation on  $A$** . A relation on  $A$  is

- **reflexive** if  $a\mathcal{R}a$  for all  $a \in A$
- **symmetric** if  $a\mathcal{R}b$  implies  $b\mathcal{R}a$  for all  $a, b \in A$
- **transitive** if  $a\mathcal{R}b$  and  $b\mathcal{R}c$  implies  $a\mathcal{R}c$  for all  $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Examples:

- $< = \{(x, y) : y - x \in (0, \infty)\}$  is transitive but not reflexive or symmetric on  $\mathbb{R}$

# Relations

A relation  $\mathcal{R}$  from  $A$  to  $A$  is called a **relation on  $A$** . A relation on  $A$  is

- **reflexive** if  $a\mathcal{R}a$  for all  $a \in A$
- **symmetric** if  $a\mathcal{R}b$  implies  $b\mathcal{R}a$  for all  $a, b \in A$
- **transitive** if  $a\mathcal{R}b$  and  $b\mathcal{R}c$  implies  $a\mathcal{R}c$  for all  $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Examples:

- $< = \{(x, y) : y - x \in (0, \infty)\}$  is transitive but not reflexive or symmetric on  $\mathbb{R}$
- $\leq = \{(x, y) : y - x \in [0, \infty)\}$  is reflexive and transitive but not symmetric on  $\mathbb{R}$

# Challenge!

## Problem

Give an example of a relation on  $\mathbb{R}$  which is symmetric and transitive but not reflexive.



# Challenge!

## Problem

Give an example of a relation on  $\mathbb{R}$  which is reflexive and symmetric but not transitive.

# Functions

A **function** from  $A$  to  $B$  is a relation  $\mathcal{R}$  from  $A$  to  $B$  with the property

# Functions

A **function** from  $A$  to  $B$  is a relation  $\mathcal{R}$  from  $A$  to  $B$  with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \quad \text{implies} \quad b = c.$$

# Functions

A **function** from  $A$  to  $B$  is a relation  $\mathcal{R}$  from  $A$  to  $B$  with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

If a relation  $\mathcal{R}$  is a function, we usually use a symbol like  $f$ .

# Functions

A **function** from  $A$  to  $B$  is a relation  $\mathcal{R}$  from  $A$  to  $B$  with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

If a relation  $\mathcal{R}$  is a function, we usually use a symbol like  $f$ .

**NOTATION:**  $f : A \rightarrow B$  means  $f$  is a function from  $A$  to  $B$

**NOTATION:**  $f(a) = b$  means  $(a, b) \in f$ .

# Functions

A **function** from  $A$  to  $B$  is a relation  $\mathcal{R}$  from  $A$  to  $B$  with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

If a relation  $\mathcal{R}$  is a function, we usually use a symbol like  $f$ .

**NOTATION:**  $f : A \rightarrow B$  means  $f$  is a function from  $A$  to  $B$

**NOTATION:**  $f(a) = b$  means  $(a, b) \in f$ .

The set

$$\text{img}(f) = \{f(a) : a \in A\}$$

is called the **range** or **image** of  $f$

# Challenge!

## Problem

Determine all the equivalence relations on  $\mathbb{R}$  which are also functions.

# Challenge!

## Problem

Determine all the equivalence relations on  $\mathbb{R}$  which are also functions.

## Solution

$f = \mathcal{R}$  must be reflexive, so  $x\mathcal{R}x$  for all  $x$



# Challenge!

## Problem

Determine all the equivalence relations on  $\mathbb{R}$  which are also functions.

## Solution

$f = \mathcal{R}$  must be reflexive, so  $x\mathcal{R}x$  for all  $x$   
This means  $f(x) = x$  for all  $x$ .

# Challenge!

## Problem

Determine all the equivalence relations on  $\mathbb{R}$  which are also functions.

## Solution

$f = \mathcal{R}$  must be reflexive, so  $x\mathcal{R}x$  for all  $x$

This means  $f(x) = x$  for all  $x$ .

Thus the only function which is an equivalence relation is the identity function

$$f(x) = x.$$

# Properties of functions

A function  $f$  from  $A$  to  $B$  is called **one-to-one** or **injective** if

# Properties of functions

A function  $f$  from  $A$  to  $B$  is called **one-to-one** or **injective** if

$$f(x) = f(y) \text{ implies } x = y \text{ for all } x, y \in A.$$

# Properties of functions

A function  $f$  from  $A$  to  $B$  is called **one-to-one** or **injective** if

$$f(x) = f(y) \text{ implies } x = y \text{ for all } x, y \in A.$$

It is called **onto** or **surjective** if  $\text{img}(f) = B$ , or equivalently

$$\text{for all } b \in B \text{ there exists } a \in A \text{ with } f(a) = b.$$

# Properties of functions

A function  $f$  from  $A$  to  $B$  is called **one-to-one** or **injective** if

$$f(x) = f(y) \text{ implies } x = y \text{ for all } x, y \in A.$$

It is called **onto** or **surjective** if  $\text{img}(f) = B$ , or equivalently

$$\text{for all } b \in B \text{ there exists } a \in A \text{ with } f(a) = b.$$

If it satisfies both properties, it is called **bijective**.

# Inverses

The **converse** of a relation  $\mathcal{R}$  from  $A$  to  $B$  is the relation

# Inverses

The **converse** of a relation  $\mathcal{R}$  from  $A$  to  $B$  is the relation

$$\check{\mathcal{R}} = \{(b, a) \in B \times A : a\mathcal{R}b\}.$$



# Inverses

The **converse** of a relation  $\mathcal{R}$  from  $A$  to  $B$  is the relation

$$\check{\mathcal{R}} = \{(b, a) \in B \times A : a\mathcal{R}b\}.$$

If  $f$  is function, then  $\check{f}$  may or not be a function.

# Inverses

The **converse** of a relation  $\mathcal{R}$  from  $A$  to  $B$  is the relation

$$\check{\mathcal{R}} = \{(b, a) \in B \times A : a\mathcal{R}b\}.$$

If  $f$  is function, then  $\check{f}$  may or not be a function.  
If it is, we call it the **inverse** of  $f$ .

# Compositions

The composition of  $f : A \rightarrow B$  and  $g : B \rightarrow C$  is the function

# Compositions

The composition of  $f : A \rightarrow B$  and  $g : B \rightarrow C$  is the function

$$g \circ f : A \rightarrow C$$

# Compositions

The composition of  $f : A \rightarrow B$  and  $g : B \rightarrow C$  is the function

$$g \circ f : A \rightarrow C$$

with domain  $A$  and codomain  $C$  defined by

# Compositions

The composition of  $f : A \rightarrow B$  and  $g : B \rightarrow C$  is the function

$$g \circ f : A \rightarrow C$$

with domain  $A$  and codomain  $C$  defined by

$$(g \circ f)(x) = g(f(x)).$$