

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

September 3, 2025

Outline

- 1 Real Analysis Lecture 3
 - Set basics
 - Set algebra

Outline

- 1 Real Analysis Lecture 3
 - Set basics
 - Set algebra

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$
- empty set $\emptyset$

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$
- empty set $\emptyset$
- $\{\heartsuit, \text{Fall}, \{\emptyset\}\}$

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{Z}_+, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$
- empty set $\emptyset$
- $\{\heartsuit, \text{Fall}, \{\emptyset\}\}$
- $\{n \in \mathbb{Z} : n \text{ is prime}\}$

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

- ordered pairs

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

- ordered pairs

$$(a, b) = \{a, \{a, b\}\}$$

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

- ordered pairs

$$(a, b) = \{a, \{a, b\}\}$$

- even relations and functions are sets!

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$$(a, b) = (c, d) \text{ if and only if } \{a, \{a, b\}\} = \{c, \{c, d\}\}$$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$(a, b) = (c, d)$ if and only if $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if $a = c$ and $b = d$, then $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$(a, b) = (c, d)$ if and only if $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if $a = c$ and $b = d$, then $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$(a, b) = (c, d)$ if and only if $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if $a = c$ and $b = d$, then $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

Suppose $\{a, \{a, b\}\} = \{c, \{c, d\}\}$.

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$(a, b) = (c, d)$ if and only if $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if $a = c$ and $b = d$, then $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

Suppose $\{a, \{a, b\}\} = \{c, \{c, d\}\}$.

Two possible cases:

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$(a, b) = (c, d)$ if and only if $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if $a = c$ and $b = d$, then $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

Suppose $\{a, \{a, b\}\} = \{c, \{c, d\}\}$.

Two possible cases:

Case I: $a = c$ and $\{a, b\} = \{c, d\}$

Case II: $a = \{c, d\}$ and $\{a, b\} = c$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$(a, b) = (c, d)$ if and only if $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

Clearly, if $a = c$ and $b = d$, then $\{a, \{a, b\}\} = \{c, \{c, d\}\}$

The tough part is the opposite direction!

Suppose $\{a, \{a, b\}\} = \{c, \{c, d\}\}$.

Two possible cases:

Case I: $a = c$ and $\{a, b\} = \{c, d\}$

Case II: $a = \{c, d\}$ and $\{a, b\} = c$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$$\text{Case I: } a = c \text{ and } \{a, b\} = \{c, d\}$$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

Case I: $a = c$ and $\{a, b\} = \{c, d\}$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

Case I: $a = c$ and $\{a, b\} = \{c, d\}$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Therefore $b = c$ or $b = d$.

If $b = d$, we're done!

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

Case I: $a = c$ and $\{a, b\} = \{c, d\}$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Therefore $b = c$ or $b = d$.

If $b = d$, we're done! ... so assume instead that $b = c$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

Case I: $a = c$ and $\{a, b\} = \{c, d\}$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Therefore $b = c$ or $b = d$.

If $b = d$, we're done! ... so assume instead that $b = c$

Then $a = c$ implies $a = b$.

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

Case I: $a = c$ and $\{a, b\} = \{c, d\}$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Therefore $b = c$ or $b = d$.

If $b = d$, we're done! ... so assume instead that $b = c$

Then $a = c$ implies $a = b$.

Therefore $\{c, d\} = \{a, b\} = \{a, a\} = \{a\}$.

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

Case I: $a = c$ and $\{a, b\} = \{c, d\}$

Since $\{a, b\} = \{c, d\}$, we know $b \in \{c, d\}$.

Therefore $b = c$ or $b = d$.

If $b = d$, we're done! ... so assume instead that $b = c$

Then $a = c$ implies $a = b$.

Therefore $\{c, d\} = \{a, b\} = \{a, a\} = \{a\}$.

It follows that $d = c = b = a$.

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$$\text{Case II: } a = \{c, d\} \text{ and } \{a, b\} = c$$

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

Case II: $a = \{c, d\}$ and $\{a, b\} = c$

This would imply that $c \in a$ and $a \in c$.

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$$\text{Case II: } a = \{c, d\} \text{ and } \{a, b\} = c$$

This would imply that $c \in a$ and $a \in c$.

This can be shown to contradict the ZF Axioms of Set Theory.

Challenge!

Problem

Prove that for ordered pairs (a, b) and (c, d) that

$$(a, b) = (c, d) \quad \text{if and only if} \quad a = c \text{ and } b = d$$

Solution

$$\text{Case II: } a = \{c, d\} \text{ and } \{a, b\} = c$$

This would imply that $c \in a$ and $a \in c$.

This can be shown to contradict the ZF Axioms of Set Theory.
Specifically the regularity axiom for the set $\{a, c\}$...

Outline

- 1 Real Analysis Lecture 3
 - Set basics
 - Set algebra

Unions of sets

Unions:

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

$$A_1 \cup A_2 \cup \cdots \cup A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

$$A_1 \cup A_2 \cup \cdots \cup A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

NOTATION:

$$\bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \cdots \cup A_n.$$

Intersections of sets

Intersections:

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

$$A_1 \cap A_2 \cap \cdots \cap A_n = \{x : x \in A_i \text{ for all } 1 \leq i \leq n\}$$

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

$$A_1 \cap A_2 \cap \cdots \cap A_n = \{x : x \in A_i \text{ for all } 1 \leq i \leq n\}$$

NOTATION:

$$\bigcap_{i=1}^n A_i = A_1 \cap A_2 \cap \cdots \cap A_n.$$

Complements of sets

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Theorem

De Morgan's Laws

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Theorem

De Morgan's Laws

$$B - \bigcup_{i=1}^n A_i = \bigcap_{i=1}^n (B - A_i)$$

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Theorem

De Morgan's Laws

$$B - \bigcup_{i=1}^n A_i = \bigcap_{i=1}^n (B - A_i)$$

$$B - \bigcap_{i=1}^n A_i = \bigcup_{i=1}^n (B - A_i)$$

Families of sets

A **family of sets** is a collection

$$\{A_i : i \in I\}$$

Families of sets

A **family of sets** is a collection

$$\{A_i : i \in I\}$$

- I any set, called the **index set**

Families of sets

A **family of sets** is a collection

$$\{A_i : i \in I\}$$

- I any set, called the **index set**
- A_i is a set for all i

Families of sets

A **family of sets** is a collection

$$\{A_i : i \in I\}$$

- I any set, called the **index set**
- A_i is a set for all i

$$\bigcup_{i \in I} A_i = \{x : x \in A_i, \text{ for some } i \in I\}$$

Families of sets

A **family of sets** is a collection

$$\{A_i : i \in I\}$$

- I any set, called the **index set**
- A_i is a set for all i

$$\bigcup_{i \in I} A_i = \{x : x \in A_i, \text{ for some } i \in I\}$$

$$\bigcap_{i \in I} A_i = \{x : x \in A_i, \text{ for all } i \in I\}$$

Challenge

- index set $I = (1, 2)$
- family of sets $\{A_i : i \in I\}$
- $A_i = [0, i]$

Problem

Determine $\bigcup_{i \in I} A_i$.

Challenge

- index set $I = \mathbb{Z}_+$
- family of sets $\{A_i : i \in I\}$
- $A_i = [0, i)$

Problem

Determine $\bigcap_{i \in I} A_i$.

Challenge

Problem

Prove that De Morgan's Laws apply to families of sets too, ie.

$$B - \bigcup_{i \in I} A_i = \bigcap_{i \in I} (B - A_i)$$

Challenge

Problem

Prove that De Morgan's Laws apply to families of sets too, ie.

$$B - \bigcup_{i \in I} A_i = \bigcap_{i \in I} (B - A_i)$$

$$B - \bigcap_{i \in I} A_i = \bigcup_{i \in I} (B - A_i)$$

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** $\mathcal{R}$ from A to B is a subset of $A \times B$.

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** $\mathcal{R}$ from A to B is a subset of $A \times B$.

NOTATION: $a\mathcal{R}b$ means $(a, b) \in \mathcal{R}$.

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** $\mathcal{R}$ from A to B is a subset of $A \times B$.

NOTATION: $a\mathcal{R}b$ means $(a, b) \in \mathcal{R}$.

Domain and codomain:

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** $\mathcal{R}$ from A to B is a subset of $A \times B$.

NOTATION: $a\mathcal{R}b$ means $(a, b) \in \mathcal{R}$.

Domain and codomain:

$$\begin{aligned}\text{dom}(\mathcal{R}) &= \{a \in A : \exists b \in B, a\mathcal{R}b\} \\ \text{codom}(\mathcal{R}) &= \{b \in B : \exists a \in A, a\mathcal{R}b\}\end{aligned}$$

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** $\mathcal{R}$ from A to B is a subset of $A \times B$.

NOTATION: $a\mathcal{R}b$ means $(a, b) \in \mathcal{R}$.

Domain and codomain:

$$\begin{aligned}\text{dom}(\mathcal{R}) &= \{a \in A : \exists b \in B, a\mathcal{R}b\} \\ \text{codom}(\mathcal{R}) &= \{b \in B : \exists a \in A, a\mathcal{R}b\}\end{aligned}$$

Relations

A relation $\mathcal{R}$ from A to A is called a **relation on A** . A relation on A is

Relations

A relation $\mathcal{R}$ from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$

Relations

A relation $\mathcal{R}$ from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$
- **symmetric** if $a\mathcal{R}b$ implies $b\mathcal{R}a$ for all $a, b \in A$

Relations

A relation $\mathcal{R}$ from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$
- **symmetric** if $a\mathcal{R}b$ implies $b\mathcal{R}a$ for all $a, b \in A$
- **transitive** if $a\mathcal{R}b$ and $b\mathcal{R}c$ implies $a\mathcal{R}c$ for all $a, b, c \in A$

Relations

A relation $\mathcal{R}$ from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$
- **symmetric** if $a\mathcal{R}b$ implies $b\mathcal{R}a$ for all $a, b \in A$
- **transitive** if $a\mathcal{R}b$ and $b\mathcal{R}c$ implies $a\mathcal{R}c$ for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Relations

A relation $\mathcal{R}$ from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$
- **symmetric** if $a\mathcal{R}b$ implies $b\mathcal{R}a$ for all $a, b \in A$
- **transitive** if $a\mathcal{R}b$ and $b\mathcal{R}c$ implies $a\mathcal{R}c$ for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Examples:

Relations

A relation $\mathcal{R}$ from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$
- **symmetric** if $a\mathcal{R}b$ implies $b\mathcal{R}a$ for all $a, b \in A$
- **transitive** if $a\mathcal{R}b$ and $b\mathcal{R}c$ implies $a\mathcal{R}c$ for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Examples:

Relations

A relation $\mathcal{R}$ from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$
- **symmetric** if $a\mathcal{R}b$ implies $b\mathcal{R}a$ for all $a, b \in A$
- **transitive** if $a\mathcal{R}b$ and $b\mathcal{R}c$ implies $a\mathcal{R}c$ for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Examples:

- $< = \{(x, y) : y - x \in (0, \infty)\}$ is transitive but not reflexive or symmetric on $\mathbb{R}$

Relations

A relation $\mathcal{R}$ from A to A is called a **relation on A** . A relation on A is

- **reflexive** if $a\mathcal{R}a$ for all $a \in A$
- **symmetric** if $a\mathcal{R}b$ implies $b\mathcal{R}a$ for all $a, b \in A$
- **transitive** if $a\mathcal{R}b$ and $b\mathcal{R}c$ implies $a\mathcal{R}c$ for all $a, b, c \in A$

An **equivalence relation** satisfies all three properties.

Examples:

- $< = \{(x, y) : y - x \in (0, \infty)\}$ is transitive but not reflexive or symmetric on $\mathbb{R}$
- $\leq = \{(x, y) : y - x \in [0, \infty)\}$ is reflexive and transitive but not symmetric on $\mathbb{R}$

Challenge!

Problem

Give an example of a relation on $\mathbb{R}$ which is symmetric and transitive but not reflexive.

Challenge!

Problem

Give an example of a relation on $\mathbb{R}$ which is reflexive and symmetric but not transitive.

Functions

A **function** from A to B is a relation $\mathcal{R}$ from A to B with the property

Functions

A **function** from A to B is a relation $\mathcal{R}$ from A to B with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

Functions

A **function** from A to B is a relation $\mathcal{R}$ from A to B with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

If a relation $\mathcal{R}$ is a function, we usually use a symbol like f .

Functions

A **function** from A to B is a relation $\mathcal{R}$ from A to B with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

If a relation $\mathcal{R}$ is a function, we usually use a symbol like f .

NOTATION: $f : A \rightarrow B$ means f is a function from A to B

NOTATION: $f(a) = b$ means $(a, b) \in f$.

Functions

A **function** from A to B is a relation $\mathcal{R}$ from A to B with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

If a relation $\mathcal{R}$ is a function, we usually use a symbol like f .

NOTATION: $f : A \rightarrow B$ means f is a function from A to B

NOTATION: $f(a) = b$ means $(a, b) \in f$.

The set

$$\text{img}(f) = \{f(a) : a \in A\}$$

is called the **range** or **image** of f

Challenge!

Problem

Determine all the equivalence relations on $\mathbb{R}$ which are also functions.

Challenge!

Problem

Determine all the equivalence relations on $\mathbb{R}$ which are also functions.

Solution

$f = \mathcal{R}$ must be reflexive, so $x\mathcal{R}x$ for all x

Challenge!

Problem

Determine all the equivalence relations on $\mathbb{R}$ which are also functions.

Solution

$f = \mathcal{R}$ must be reflexive, so $x\mathcal{R}x$ for all x
This means $f(x) = x$ for all x .

Challenge!

Problem

Determine all the equivalence relations on $\mathbb{R}$ which are also functions.

Solution

$f = \mathcal{R}$ must be reflexive, so $x\mathcal{R}x$ for all x

This means $f(x) = x$ for all x .

Thus the only function which is an equivalence relation is the identity function

$$f(x) = x.$$

Properties of functions

A function f from A to B is called **one-to-one** or **injective** if

Properties of functions

A function f from A to B is called **one-to-one** or **injective** if

$$f(x) = f(y) \text{ implies } x = y \text{ for all } x, y \in A.$$

Properties of functions

A function f from A to B is called **one-to-one** or **injective** if

$$f(x) = f(y) \text{ implies } x = y \text{ for all } x, y \in A.$$

It is called **onto** or **surjective** if $\text{img}(f) = B$, or equivalently

$$\text{for all } b \in B \text{ there exists } a \in A \text{ with } f(a) = b.$$

Properties of functions

A function f from A to B is called **one-to-one** or **injective** if

$$f(x) = f(y) \text{ implies } x = y \text{ for all } x, y \in A.$$

It is called **onto** or **surjective** if $\text{img}(f) = B$, or equivalently

$$\text{for all } b \in B \text{ there exists } a \in A \text{ with } f(a) = b.$$

If it satisfies both properties, it is called **bijective**.

Inverses

The **converse** of a relation $\mathcal{R}$ from A to B is the relation

Inverses

The **converse** of a relation $\mathcal{R}$ from A to B is the relation

$$\check{\mathcal{R}} = \{(b, a) \in B \times A : a\mathcal{R}b\}.$$

Inverses

The **converse** of a relation $\mathcal{R}$ from A to B is the relation

$$\check{\mathcal{R}} = \{(b, a) \in B \times A : a\mathcal{R}b\}.$$

If f is function, then $\check{f}$ may or not be a function.

Inverses

The **converse** of a relation $\mathcal{R}$ from A to B is the relation

$$\check{\mathcal{R}} = \{(b, a) \in B \times A : a\mathcal{R}b\}.$$

If f is function, then $\check{f}$ may or not be a function.
If it is, we call it the **inverse** of f .

Compositions

The composition of $f : A \rightarrow B$ and $g : B \rightarrow C$ is the function

Compositions

The composition of $f : A \rightarrow B$ and $g : B \rightarrow C$ is the function

$$g \circ f : A \rightarrow C$$

Compositions

The composition of $f : A \rightarrow B$ and $g : B \rightarrow C$ is the function

$$g \circ f : A \rightarrow C$$

with domain A and codomain C defined by

Compositions

The composition of $f : A \rightarrow B$ and $g : B \rightarrow C$ is the function

$$g \circ f : A \rightarrow C$$

with domain A and codomain C defined by

$$(g \circ f)(x) = g(f(x)).$$

Sequences

A **finite sequence** is a function $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$.

Sequences

A **finite sequence** is a function $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$.

An **infinite sequence** is a function $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$.

Sequences

A **finite sequence** is a function $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$.

An **infinite sequence** is a function $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$.

NOTATION: f_n indicates the value $f(n)$

Sequences

A **finite sequence** is a function $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$.

An **infinite sequence** is a function $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$.

NOTATION: f_n indicates the value $f(n)$

NOTATION: $\{f_n\}$ is another way of writing the function f

Sequences

A **finite sequence** is a function $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$.

An **infinite sequence** is a function $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$.

NOTATION: f_n indicates the value $f(n)$

NOTATION: $\{f_n\}$ is another way of writing the function f

If $k : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ is a function which is **strictly increasing**,
meaning

$$m < n \quad \Rightarrow \quad k(m) < k(n),$$

Sequences

A **finite sequence** is a function $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$.

An **infinite sequence** is a function $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$.

NOTATION: f_n indicates the value $f(n)$

NOTATION: $\{f_n\}$ is another way of writing the function f

If $k : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ is a function which is **strictly increasing**,
meaning

$$m < n \quad \Rightarrow \quad k(m) < k(n),$$

then the composition $f \circ k : \mathbb{Z}_+ \rightarrow \mathbb{R}$ forms a sequence called a
subsequence of f .

Sequences

A **finite sequence** is a function $f : \{1, 2, \dots, n\} \rightarrow \mathbb{R}$.

An **infinite sequence** is a function $f : \mathbb{Z}_+ \rightarrow \mathbb{R}$.

NOTATION: f_n indicates the value $f(n)$

NOTATION: $\{f_n\}$ is another way of writing the function f

If $k : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ is a function which is **strictly increasing**, meaning

$$m < n \quad \Rightarrow \quad k(m) < k(n),$$

then the composition $f \circ k : \mathbb{Z}_+ \rightarrow \mathbb{R}$ forms a sequence called a **subsequence** of f .

NOTATION: $\{f_{k(n)}\}$ or $\{f_{k_n}\}$ both really mean $f \circ k$