

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

August 23, 2025

Outline

- 1 Real Analysis Lecture 2
 - Types of real numbers
 - Suprema and Infima

Outline

- 1 Real Analysis Lecture 2
 - Types of real numbers
 - Suprema and Infima

Inductive sets

A subset S of $\mathbb{R}$ is called a **inductive set** if

- $1 \in S$
- if $x \in S$ then $x + 1 \in S$

Inductive sets

A subset S of $\mathbb{R}$ is called a **inductive set** if

- $1 \in S$
- if $x \in S$ then $x + 1 \in S$

Examples:

$$\mathbb{R}, \quad \mathbb{Q}, \quad (0, \infty), \quad \mathbb{Z}, \quad \mathbb{N}, \dots$$

Integers and rationals

How do we reverse-engineer the integers from the reals?

Integers and rationals

How do we reverse-engineer the integers from the reals?

- the set of **positive integers** is

$$\mathbb{Z}_+ = \{x : x \text{ belongs to every inductive set}\}$$

Integers and rationals

How do we reverse-engineer the integers from the reals?

- the set of **positive integers** is

$$\mathbb{Z}_+ = \{x : x \text{ belongs to every inductive set}\}$$

- integers are

$$\mathbb{Z} = \{x : x \text{ is zero or } \pm x \text{ is a positive integer}\}$$

Integers and rationals

How do we reverse-engineer the integers from the reals?

- the set of **positive integers** is

$$\mathbb{Z}_+ = \{x : x \text{ belongs to every inductive set}\}$$

- integers are

$$\mathbb{Z} = \{x : x \text{ is zero or } \pm x \text{ is a positive integer}\}$$

- rationals are

$$\mathbb{Q} = \{a/b : a, b \in \mathbb{Z}, b \neq 0\}.$$

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof:

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof:

Let

$$S = \{n \in \mathbb{Z}_+ : 2|n(n+1)\}$$

.

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof:

Let

$$S = \{n \in \mathbb{Z}_+ : 2|n(n+1)\}$$

. We see that $1 \in S$ because 2 divides $1(1+1)$.

Induction

Principle of Induction:

If S is an inductive set, then $\mathbb{Z}_+ \subseteq S$

Question

Waaaaait...How is this this induction?!

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof:

Let

$$S = \{n \in \mathbb{Z}_+ : 2|n(n+1)\}$$

. We see that $1 \in S$ because 2 divides $1(1+1)$.

Now suppose $x \in S$. (This is our usual inductive assumption).

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.
Since $x+1$ was arbitrary, this shows that S is an inductive set.

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.
Since $x+1$ was arbitrary, this shows that S is an inductive set.
By the Principle of Induction, this means $\mathbb{Z}_+ \subseteq S$.

Induction

Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.
Since $x+1$ was arbitrary, this shows that S is an inductive set.
By the Principle of Induction, this means $\mathbb{Z}_+ \subseteq S$.
In other words $n \in S$ for every positive integer n .

Induction


Example: Let's prove $n(n+1)$ is divisible by 2 for all positive integers n , using the Principle of Induction.

Proof continued:

Then 2 divides $x(x+1)$, so $x(x+1) = 2k$ for some integer k .
This means $(x+1)(x+2) = x(x+1) + 2(x+1) = 2(k+x+1)$.
Thus 2 divides $(x+1)(x+2)$, showing that $x+1 \in S$.
Since $x+1$ was arbitrary, this shows that S is an inductive set.
By the Principle of Induction, this means $\mathbb{Z}_+ \subseteq S$.
In other words $n \in S$ for every positive integer n .
Hence 2 divides $n(n+1)$ for every positive integers n .

Types of numbers


Transcendental Numbers



Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Discovered by Hippasus, a pythagorean (a student of Pythagoras)



Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery

Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- dramatic reenactment, found online:

Irrational numbers


Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- dramatic reenactment, found online:

Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- dramatic reenactment, found online:



Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- anime style:

Irrational numbers

Numbers which are not of the form a/b with $a, b \in \mathbb{Z}$ are called **irrational**.

- Pythagoras drowned Hippasus at sea for his discovery
- anime style:



Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then $\sqrt{n}$ is irrational? (Apostol Theorem 1.10)

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then $\sqrt{n}$ is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then $\sqrt{n}$ is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b .

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then $\sqrt{n}$ is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then $\sqrt{n}$ is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then $\sqrt{n}$ is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then $\sqrt{n}$ is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$. It follows that $b^2 n' = 1$, so b^2 divides 1.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then $\sqrt{n}$ is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$. It follows that $b^2 n' = 1$, so b^2 divides 1. This implies n' divides 1, so $n' = \pm 1$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then $\sqrt{n}$ is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$. It follows that $b^2 n' = 1$, so b^2 divides 1. This implies n' divides 1, so $n' = \pm 1$. Since $n > 0$, $n' > 0$ so $n' = 1$.

Challenge!

Problem

Can you prove that if $n \in \mathbb{Z}_+$ is not a perfect square, then $\sqrt{n}$ is irrational? (Apostol Theorem 1.10)

Solution

We prove by contradiction. Assume $\sqrt{n} = a/b$ for integers a, b . Without loss of generality, we may assume $\gcd(a, b) = 1$. Then $b^2 n = a^2$. Since $\gcd(a, b) = 1$, a^2 must divide n , ie. $n = n' a^2$. It follows that $b^2 n' = 1$, so b^2 divides 1. This implies n' divides 1, so $n' = \pm 1$. Since $n > 0$, $n' > 0$ so $n' = 1$. Thus $n = a^2$, which is a contradiction.

Algebraic and Transcendental

Irrational numbers can be further divided into to categories

Algebraic and Transcendental

Irrational numbers can be further divided into to categories

- 1 **algebraic numbers:** numbers which are roots of polynomials with integer coefficients, like

$$\sqrt{2}, \quad \sqrt{3}, \quad \text{and} \quad \sqrt[3]{\sqrt{5} + \sqrt{7}}.$$

Algebraic and Transcendental

Irrational numbers can be further divided into to categories

- 1 **algebraic numbers:** numbers which are roots of polynomials with integer coefficients, like

$$\sqrt{2}, \quad \sqrt{3}, \quad \text{and} \quad \sqrt[3]{\sqrt{5} + \sqrt{7}}.$$

- 2 **transcendental numbers:** numbers which are not algebraic, like

$$\pi, \quad e, \quad e^{\pi}, \quad \text{and maybe} \quad \pi + e?$$

Algebraic and Transcendental

Irrational numbers can be further divided into to categories

- 1 **algebraic numbers:** numbers which are roots of polynomials with integer coefficients, like

$$\sqrt{2}, \quad \sqrt{3}, \quad \text{and} \quad \sqrt[3]{\sqrt{5} + \sqrt{7}}.$$

- 2 **transcendental numbers:** numbers which are not algebraic, like

$$\pi, \quad e, \quad e^{\pi}, \quad \text{and maybe} \quad \pi + e?$$

- 3 transcendentals are mysterious ... but most real numbers are transcendental!

Outline

- 1 Real Analysis Lecture 2
 - Types of real numbers
 - Suprema and Infima

Completeness Axiom

Last time, we finished with the Completeness Axiom:

Completeness Axiom

Last time, we finished with the Completeness Axiom:

Completeness Axiom

Last time, we finished with the Completeness Axiom:

- Ⓐ10 **completeness axiom:** if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Completeness Axiom

Last time, we finished with the Completeness Axiom:

A10 completeness axiom: if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

Completeness Axiom

Last time, we finished with the Completeness Axiom:

- Ⓐ10 **completeness axiom:** if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

- 3 is the supremum of $(0, 3)$

Completeness Axiom

Last time, we finished with the Completeness Axiom:

- Ⓐ10 **completeness axiom:** if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

- 3 is the supremum of $(0, 3)$
- 1 is the supremum of

$$\left\{ \frac{n}{n+1} : n \in \mathbb{Z}_+ \right\}$$

Completeness Axiom

Last time, we finished with the Completeness Axiom:

- **completeness axiom:** if S is any subset of real numbers which is bounded above, then it has a supremum $\sup(S)$

Examples:

- 3 is the supremum of $(0, 3)$
- 1 is the supremum of

$$\left\{ \frac{n}{n+1} : n \in \mathbb{Z}_+ \right\}$$

- π is the supremum of

$$\{3, 3.1, 3.14, 3.141, 3.1415, 3.14159, 3.141592, \dots\}.$$

Challenge

Problem

Prove that if A is a set of integers that is bounded above, then A has a maximum.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

If $a' \in A$, then a' is an integer and therefore $a' = a + k$ for $k \in \mathbb{Z}$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

If $a' \in A$, then a' is an integer and therefore $a' = a + k$ for $k \in \mathbb{Z}$. Since b is an upper bound, $b \geq a + k > b - 1 + k$, making $0 > k - 1$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

If $a' \in A$, then a' is an integer and therefore $a' = a + k$ for $k \in \mathbb{Z}$. Since b is an upper bound, $b \geq a + k > b - 1 + k$, making $0 > k - 1$.

Therefore $k \leq 0$ and $a' \leq a$.

Challenge

Solution

The set A is bounded above, so it has a supremum $b = \sup(A)$. Since $b - 1$ is not an upper bound, so there exists $a \in A$ with $b - 1 < a$.

If $a' \in A$, then a' is an integer and therefore $a' = a + k$ for $k \in \mathbb{Z}$. Since b is an upper bound, $b \geq a + k > b - 1 + k$, making $0 > k - 1$.

Therefore $k \leq 0$ and $a' \leq a$.

It follows that a is an upper bound of A , and since $a \in A$ it is a maximum.

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

If $b \in S$ also, then b is called a **minimal element** of S .

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

If $b \in S$ also, then b is called a **minimal element** of S . An **infimum** of a set S of real numbers is a real number $b \in \mathbb{R}$ such that

- b is a lower bound of S
- if $b < b'$, then b' is not a lower bound of S

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

If $b \in S$ also, then b is called a **minimal element** of S . An **infimum** of a set S of real numbers is a real number $b \in \mathbb{R}$ such that

- b is a lower bound of S
- if $b < b'$, then b' is not a lower bound of S

In other words

an infimum is a **greatest lower bound**

Properties of Suprema

The first property of suprema is that they must be arbitrarily close to elements of the set.

Theorem (Approximation Property)

Let $S \subseteq \mathbb{R}$ be bounded above, and let $b = \sup(S)$. Then for all $\epsilon > 0$, there exists $a \in S$ with

$$b - \epsilon < a \leq b.$$

Properties of Suprema

Proof.

Since $b - \epsilon < b$, the definition of a supremum implies $b - \epsilon$ cannot be an upper bound.

Properties of Suprema

Proof.

Since $b - \epsilon < b$, the definition of a supremum implies $b - \epsilon$ cannot be an upper bound.

Therefore there must exist $a \in S$ with $a > b - \epsilon$. □

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Consider the set

$$S = \{s_n : n \in \mathbb{Z}_+\}.$$

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Consider the set

$$S = \{s_n : n \in \mathbb{Z}_+\}.$$

For every n ,

$$s_n \leq 1 + 1 + \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \cdots + \frac{1}{2^n} = 3 - \frac{1}{2^n} < 3$$

so S is bounded above by 3.

Important example: The number e

For each n , let s_n denote the sum

$$s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Notice that

$$s_1 < s_2 < s_3 < s_4 < \dots$$

Consider the set

$$S = \{s_n : n \in \mathbb{Z}_+\}.$$

For every n ,

$$s_n \leq 1 + 1 + \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \cdots + \frac{1}{2^n} = 3 - \frac{1}{2^n} < 3$$

so S is bounded above by 3. Therefore S has a supremum, $\sup(S)$.

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N .

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

If we remember Taylor series

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

If we remember Taylor series

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots$$

Important example: The number e

Even if ϵ is very small ($\epsilon = 0.000000001$), $\sup(S) - \epsilon$ is not an upper bound of S .

So there exists N with $s_N > \sup(S) - \epsilon$.

$$s_1 < s_2 < s_3 < s_4 < \cdots < \sup(S) - \epsilon < s_N < \cdots < \sup(S).$$

This says $\sup(S)$ is *really* close to s_N . Taking ϵ smaller and smaller

$$\sup(S) = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

If we remember Taylor series

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots$$

We've just shown that $e^1 = e$ exists.

Properties of Suprema

Also, suprema play nicely with addition.

Theorem (Additive Property)

Let $A, B \subseteq \mathbb{R}$ be bounded above set

$$C = \{x + y : x \in A, y \in B\}.$$

Then C is bounded above and

$$\sup(C) = \sup(A) + \sup(B).$$

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.
We will prove $c \leq a + b$ and then $a + b \leq c$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

It follows that $y \leq c - a$ for all $y \in B$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

It follows that $y \leq c - a$ for all $y \in B$.

Thus $c - a$ is an upper bound of B , and it follows $b \leq c - a$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

It follows that $y \leq c - a$ for all $y \in B$.

Thus $c - a$ is an upper bound of B , and it follows $b \leq c - a$.

Therefore $a + b \leq c$. □

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{Z}_+$ is not bounded above. (Apostol Theorem 1.17)

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{Z}_+$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{Z}_+$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{Z}_+$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{Z}_+$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{Z}_+$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{Z}_+$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

This means there exists $n \in \mathbb{Z}_+$ with $b - 1 < n$.

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{Z}_+$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

This means there exists $n \in \mathbb{Z}_+$ with $b - 1 < n$.

It follows from Axiom 7 that $b < n + 1$.

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{Z}_+$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{Z}_+) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{Z}_+)$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

This means there exists $n \in \mathbb{Z}_+$ with $b - 1 < n$.

It follows from Axiom 7 that $b < n + 1$.

However, $n + 1 \in \mathbb{Z}$, so this contradicts b being an upper bound.

Archimedian property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{Z}_+$ with $n > x$.

Archimedean property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{Z}_+$ with $n > x$.

Proof.

If not, then x is an upper bound of $\mathbb{Z}_+$. □

Archimedean property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{Z}_+$ with $n > x$.

Proof.

If not, then x is an upper bound of $\mathbb{Z}_+$. □

Theorem (Archimedean Property of Reals)

For every $x, y \in \mathbb{R}$ with $x > 0$, there exists $n \in \mathbb{Z}_+$ with $y < nx$.

Archimedean property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{Z}_+$ with $n > x$.

Proof.

If not, then x is an upper bound of $\mathbb{Z}_+$. □

Theorem (Archimedean Property of Reals)

For every $x, y \in \mathbb{R}$ with $x > 0$, there exists $n \in \mathbb{Z}_+$ with $y < nx$.

Proof.

Replace x with y/x in the previous theorem. □