

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

September 17, 2025

Outline

- 1 Real Analysis Lecture 7
 - More on Closed Sets

Outline

- 1 Real Analysis Lecture 7
 - More on Closed Sets

Adherent and Accumulation Points

Definition

A point $\vec{x} \in \mathbb{R}^n$ is called an **adherent point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A .

Adherent and Accumulation Points

Definition

A point $\vec{x} \in \mathbb{R}^n$ is called an **adherent point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A .

It is called an **accumulation point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A **different from $\vec{x}$** .

Adherent and Accumulation Points

Definition

A point $\vec{x} \in \mathbb{R}^n$ is called an **adherent point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A .

It is called an **accumulation point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A **different from $\vec{x}$** .

Examples:

Adherent and Accumulation Points

Definition

A point $\vec{x} \in \mathbb{R}^n$ is called an **adherent point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A .

It is called an **accumulation point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A **different from $\vec{x}$** .

Examples:

- every point in A is adherent

Adherent and Accumulation Points

Definition

A point $\vec{x} \in \mathbb{R}^n$ is called an **adherent point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A .

It is called an **accumulation point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A **different from $\vec{x}$** .

Examples:

- every point in A is adherent
- accumulation points are adherent points

Adherent and Accumulation Points

Definition

A point $\vec{x} \in \mathbb{R}^n$ is called an **adherent point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A .

It is called an **accumulation point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A **different from $\vec{x}$** .

Examples:

- every point in A is adherent
- accumulation points are adherent points
- -1 is an accumulation point of $(-1, 1)$.

Adherent and Accumulation Points

Definition

A point $\vec{x} \in \mathbb{R}^n$ is called an **adherent point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A .

It is called an **accumulation point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A **different from $\vec{x}$** .

Examples:

- every point in A is adherent
- accumulation points are adherent points
- -1 is an accumulation point of $(-1, 1)$.
- suprema and infima are accumulation points!

Adherent and Accumulation Points

Definition

A point $\vec{x} \in \mathbb{R}^n$ is called an **adherent point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A .

It is called an **accumulation point** if for all $r > 0$ the ball $B(\vec{x}; r)$ contains at least one element of A **different from $\vec{x}$** .

Examples:

- every point in A is adherent
- accumulation points are adherent points
- -1 is an accumulation point of $(-1, 1)$.
- suprema and infima are accumulation points!
- 0 is an accumulation point of $\{1/1, 1/2, 1/3, \dots\}$

Characterizing accumulation points

Theorem (Apostol Theorem 3.17)

A point $\vec{x}$ is an accumulation point of A if and only if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A .

Characterizing accumulation points

Theorem (Apostol Theorem 3.17)

A point $\vec{x}$ is an accumulation point of A if and only if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A .

Proof.

Characterizing accumulation points

Theorem (Apostol Theorem 3.17)

A point $\vec{x}$ is an accumulation point of A if and only if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A .

Proof.

Obviously, if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A , then it contains at least one point of A different from $\vec{x}$, so it's an accumulation point.

Characterizing accumulation points

Theorem (Apostol Theorem 3.17)

A point $\vec{x}$ is an accumulation point of A if and only if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A .

Proof.

Obviously, if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A , then it contains at least one point of A different from $\vec{x}$, so it's an accumulation point.

Suppose instead that $\vec{x}$ is an accumulation point and let $r > 0$.

Characterizing accumulation points

Theorem (Apostol Theorem 3.17)

A point $\vec{x}$ is an accumulation point of A if and only if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A .

Proof.

Obviously, if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A , then it contains at least one point of A different from $\vec{x}$, so it's an accumulation point.

Suppose instead that $\vec{x}$ is an accumulation point and let $r > 0$. It suffices to show that the set $C = (B(\vec{x}; r) \setminus \{\vec{x}\}) \cap A$ is infinite.



Characterizing accumulation points

Theorem (Apostol Theorem 3.17)

A point $\vec{x}$ is an accumulation point of A if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A .

Characterizing accumulation points

Theorem (Apostol Theorem 3.17)

A point $\vec{x}$ is an accumulation point of A if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A .

Proof.

Characterizing accumulation points

Theorem (Apostol Theorem 3.17)

A point $\vec{x}$ is an accumulation point of A if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A .

Proof.

Suppose that C is finite.

Characterizing accumulation points

Theorem (Apostol Theorem 3.17)

A point $\vec{x}$ is an accumulation point of A if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A .

Proof.

Suppose that C is finite.

Then the set $\{|\vec{y} - \vec{x}| : \vec{y} \in C\}$ is also finite.

Characterizing accumulation points

Theorem (Apostol Theorem 3.17)

A point $\vec{x}$ is an accumulation point of A if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A .

Proof.

Suppose that C is finite.

Then the set $\{|\vec{y} - \vec{x}| : \vec{y} \in C\}$ is also finite.

It is also nonempty and has only positive values.

Characterizing accumulation points

Theorem (Apostol Theorem 3.17)

A point $\vec{x}$ is an accumulation point of A if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A .

Proof.

Suppose that C is finite.

Then the set $\{|\vec{y} - \vec{x}| : \vec{y} \in C\}$ is also finite.

It is also nonempty and has only positive values. This means that it has a minimum value $s > 0$.

Characterizing accumulation points

Theorem (Apostol Theorem 3.17)

A point $\vec{x}$ is an accumulation point of A if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A .

Proof.

Suppose that C is finite.

Then the set $\{|\vec{y} - \vec{x}| : \vec{y} \in C\}$ is also finite.

It is also nonempty and has only positive values. This means that it has a minimum value $s > 0$.

Since $\vec{x}$ is an accumulation point, $B(\vec{x}; s) \cap A$ has an element $\vec{y}$ different from $\vec{x}$.

Characterizing accumulation points

Theorem (Apostol Theorem 3.17)

A point $\vec{x}$ is an accumulation point of A if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A .

Proof.

Suppose that C is finite.

Then the set $\{|\vec{y} - \vec{x}| : \vec{y} \in C\}$ is also finite.

It is also nonempty and has only positive values. This means that it has a minimum value $s > 0$.

Since $\vec{x}$ is an accumulation point, $B(\vec{x}; s) \cap A$ has an element $\vec{y}$ different from $\vec{x}$.

However, $\vec{y} \in C$ and $|\vec{y} - \vec{x}| < s$, contradicting the minimality of s .

Characterizing accumulation points

Theorem (Apostol Theorem 3.17)

A point $\vec{x}$ is an accumulation point of A if for all $r > 0$, the ball $B(\vec{x}; r)$ contains infinitely many points of A .

Proof.

Suppose that C is finite.

Then the set $\{|\vec{y} - \vec{x}| : \vec{y} \in C\}$ is also finite.

It is also nonempty and has only positive values. This means that it has a minimum value $s > 0$.

Since $\vec{x}$ is an accumulation point, $B(\vec{x}; s) \cap A$ has an element $\vec{y}$ different from $\vec{x}$.

However, $\vec{y} \in C$ and $|\vec{y} - \vec{x}| < s$, contradicting the minimality of s .

We conclude that C is infinite. □

Closure of a set

Definition

The **closure** of a set A is the set $\bar{A}$ of all adherent points of A

Closure of a set

Definition

The **closure** of a set A is the set $\bar{A}$ of all adherent points of A

Theorem (Apostol Theorem 3.18, 3.20, 3.22)

A set is closed if and only if it contains all of its adherent points (ie. $A = \bar{A}$), or equivalently if and only if A contains all of its accumulation points.

Closure of a set

Proof.

Suppose that A is closed.

Closure of a set

Proof.

Suppose that A is closed.

Then $\mathbb{R}^n \setminus A$ is open.

Closure of a set

Proof.

Suppose that A is closed.

Then $\mathbb{R}^n \setminus A$ is open.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then there exist $r > 0$ such that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

Closure of a set

Proof.

Suppose that A is closed.

Then $\mathbb{R}^n \setminus A$ is open.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then there exist $r > 0$ such that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

This means that $B(\vec{x}; r) \cap A = \emptyset$.

Closure of a set

Proof.

Suppose that A is closed.

Then $\mathbb{R}^n \setminus A$ is open.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then there exist $r > 0$ such that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

This means that $B(\vec{x}; r) \cap A = \emptyset$.

Thus $\vec{x}$ is not an adherent point of A .

Closure of a set

Proof.

Suppose that A is closed.

Then $\mathbb{R}^n \setminus A$ is open.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then there exist $r > 0$ such that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

This means that $B(\vec{x}; r) \cap A = \emptyset$.

Thus $\vec{x}$ is not an adherent point of A .

Thus every adherent point of A is an element of A .

Closure of a set

Proof.

Suppose that A is closed.

Then $\mathbb{R}^n \setminus A$ is open.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then there exist $r > 0$ such that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

This means that $B(\vec{x}; r) \cap A = \emptyset$.

Thus $\vec{x}$ is not an adherent point of A .

Thus every adherent point of A is an element of A .

Conversely, suppose that A contains all of its accumulation points.

Closure of a set

Proof.

Suppose that A is closed.

Then $\mathbb{R}^n \setminus A$ is open.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then there exist $r > 0$ such that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

This means that $B(\vec{x}; r) \cap A = \emptyset$.

Thus $\vec{x}$ is not an adherent point of A .

Thus every adherent point of A is an element of A .

Conversely, suppose that A contains all of its accumulation points.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then $\vec{x}$ is not an accumulation point.

Closure of a set

Proof.

Suppose that A is closed.

Then $\mathbb{R}^n \setminus A$ is open.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then there exist $r > 0$ such that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

This means that $B(\vec{x}; r) \cap A = \emptyset$.

Thus $\vec{x}$ is not an adherent point of A .

Thus every adherent point of A is an element of A .

Conversely, suppose that A contains all of its accumulation points.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then $\vec{x}$ is not an accumulation point.

Therefore there exists $r > 0$ with $B(\vec{x}; r) \cap A = \emptyset$.

Closure of a set

Proof.

Suppose that A is closed.

Then $\mathbb{R}^n \setminus A$ is open.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then there exist $r > 0$ such that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

This means that $B(\vec{x}; r) \cap A = \emptyset$.

Thus $\vec{x}$ is not an adherent point of A .

Thus every adherent point of A is an element of A .

Conversely, suppose that A contains all of its accumulation points.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then $\vec{x}$ is not an accumulation point.

Therefore there exists $r > 0$ with $B(\vec{x}; r) \cap A = \emptyset$.

This implies that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

Closure of a set

Proof.

Suppose that A is closed.

Then $\mathbb{R}^n \setminus A$ is open.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then there exist $r > 0$ such that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

This means that $B(\vec{x}; r) \cap A = \emptyset$.

Thus $\vec{x}$ is not an adherent point of A .

Thus every adherent point of A is an element of A .

Conversely, suppose that A contains all of its accumulation points.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then $\vec{x}$ is not an accumulation point.

Therefore there exists $r > 0$ with $B(\vec{x}; r) \cap A = \emptyset$.

This implies that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

Thus $\vec{x}$ is an interior point of $\mathbb{R}^n \setminus A$.

Closure of a set

Proof.

Suppose that A is closed.

Then $\mathbb{R}^n \setminus A$ is open.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then there exist $r > 0$ such that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

This means that $B(\vec{x}; r) \cap A = \emptyset$.

Thus $\vec{x}$ is not an adherent point of A .

Thus every adherent point of A is an element of A .

Conversely, suppose that A contains all of its accumulation points.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then $\vec{x}$ is not an accumulation point.

Therefore there exists $r > 0$ with $B(\vec{x}; r) \cap A = \emptyset$.

This implies that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

Thus $\vec{x}$ is an interior point of $\mathbb{R}^n \setminus A$.

Since $\vec{x}$ was an arbitrary element of $\mathbb{R}^n \setminus A$, this proves $\mathbb{R}^n \setminus A$ is open.

Closure of a set

Proof.

Suppose that A is closed.

Then $\mathbb{R}^n \setminus A$ is open.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then there exist $r > 0$ such that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

This means that $B(\vec{x}; r) \cap A = \emptyset$.

Thus $\vec{x}$ is not an adherent point of A .

Thus every adherent point of A is an element of A .

Conversely, suppose that A contains all of its accumulation points.

If $\vec{x} \in \mathbb{R}^n \setminus A$, then $\vec{x}$ is not an accumulation point.

Therefore there exists $r > 0$ with $B(\vec{x}; r) \cap A = \emptyset$.

This implies that $B(\vec{x}; r) \subseteq \mathbb{R}^n \setminus A$.

Thus $\vec{x}$ is an interior point of $\mathbb{R}^n \setminus A$.

Since $\vec{x}$ was an arbitrary element of $\mathbb{R}^n \setminus A$, this proves $\mathbb{R}^n \setminus A$ is open.

Therefore A is closed.



Closures are closed

Theorem

If $A \subseteq \mathbb{R}^n$ is any set, then $\overline{A}$ is closed.

Proof.

Exercise!



Outline

- 1 Real Analysis Lecture 7
 - More on Closed Sets

Warm-up Challenge

Problem

Show that the closed ball

$$\overline{B}(\vec{x}; r) = \{\vec{y} \in \mathbb{R}^n : |\vec{x} - \vec{y}| \leq r\}$$

is a closed set.

Point set topology

We have been studying the beginnings of **point set topology**.

Point set topology

We have been studying the beginnings of **point set topology**. In topology, we start with a set X called the **space** and a collection of special subsets called **open sets**.

Point set topology

We have been studying the beginnings of **point set topology**. In topology, we start with a set X called the **space** and a collection of special subsets called **open sets**.

Fundamental questions:

Point set topology

We have been studying the beginnings of **point set topology**. In topology, we start with a set X called the **space** and a collection of special subsets called **open sets**.

Fundamental questions:

- kinds of points: interior points, boundary points, accumulation points, adherent points

Point set topology

We have been studying the beginnings of **point set topology**. In topology, we start with a set X called the **space** and a collection of special subsets called **open sets**.

Fundamental questions:

- kinds of points: interior points, boundary points, accumulation points, adherent points
- kinds of sets: open, closed, connected, component, compact

Point set topology

We have been studying the beginnings of **point set topology**. In topology, we start with a set X called the **space** and a collection of special subsets called **open sets**.

Fundamental questions:

- kinds of points: interior points, boundary points, accumulation points, adherent points
- kinds of sets: open, closed, connected, component, compact
- kinds of functions: continuous, homeomorphism

Point set topology

We have been studying the beginnings of **point set topology**. In topology, we start with a set X called the **space** and a collection of special subsets called **open sets**.

Fundamental questions:

- kinds of points: interior points, boundary points, accumulation points, adherent points
- kinds of sets: open, closed, connected, component, compact
- kinds of functions: continuous, homeomorphism

Answers must be in terms of open sets

Open covers

Definition

A **open cover** of a set $A \subseteq \mathbb{R}^n$ is a family $\{U_i : i \in I\}$ of open sets with $A \subseteq \bigcup_{i \in I} U_i$.

Open covers

Definition

A **open cover** of a set $A \subseteq \mathbb{R}^n$ is a family $\{U_i : i \in I\}$ of open sets with $A \subseteq \bigcup_{i \in I} U_i$.

A **subcover** is a family $\{U_j : j \in J\}$ with $J \subseteq I$ which still covers A , ie. $A \subseteq \bigcup_{j \in J} U_j$.

Open covers

Definition

A **open cover** of a set $A \subseteq \mathbb{R}^n$ is a family $\{U_i : i \in I\}$ of open sets with $A \subseteq \bigcup_{i \in I} U_i$.

A **subcover** is a family $\{U_j : j \in J\}$ with $J \subseteq I$ which still covers A , ie. $A \subseteq \bigcup_{j \in J} U_j$.

Examples:

Open covers

Definition

A **open cover** of a set $A \subseteq \mathbb{R}^n$ is a family $\{U_i : i \in I\}$ of open sets with $A \subseteq \bigcup_{i \in I} U_i$.

A **subcover** is a family $\{U_j : j \in J\}$ with $J \subseteq I$ which still covers A , ie. $A \subseteq \bigcup_{j \in J} U_j$.

Examples:

- $\{\mathbb{R}\}$ is an open cover of the interval $(0, 1]$

Open covers

Definition

A **open cover** of a set $A \subseteq \mathbb{R}^n$ is a family $\{U_i : i \in I\}$ of open sets with $A \subseteq \bigcup_{i \in I} U_i$.

A **subcover** is a family $\{U_j : j \in J\}$ with $J \subseteq I$ which still covers A , ie. $A \subseteq \bigcup_{j \in J} U_j$.

Examples:

- $\{\mathbb{R}\}$ is an open cover of the interval $(0, 1]$
- $\{(0, 1/2), (1/2, 2)\}$ is an open cover of the interval $(0, 1]$

Open covers

Definition

A **open cover** of a set $A \subseteq \mathbb{R}^n$ is a family $\{U_i : i \in I\}$ of open sets with $A \subseteq \bigcup_{i \in I} U_i$.

A **subcover** is a family $\{U_j : j \in J\}$ with $J \subseteq I$ which still covers A , ie. $A \subseteq \bigcup_{j \in J} U_j$.

Examples:

- $\{\mathbb{R}\}$ is an open cover of the interval $(0, 1]$
- $\{(0, 1/2), (1/2, 2)\}$ is an open cover of the interval $(0, 1]$
- $\{(\frac{1}{n}, \frac{n+1}{n}) : n \in \mathbb{Z}_+\}$ is an open cover of $(0, 1]$

Lindelöf Covering Theorem

Theorem (Lindelöf Covering Theorem)

Let $A \subseteq \mathbb{R}^n$ be a set and suppose $\{U_i : i \in I\}$ is an open covering of A . Then there exists a countable subcover $\{U_j : j \in J\}$.

Lindelöf Covering Theorem

Theorem (Lindelöf Covering Theorem)

Let $A \subseteq \mathbb{R}^n$ be a set and suppose $\{U_i : i \in I\}$ is an open covering of A . Then there exists a countable subcover $\{U_j : j \in J\}$.

Question

Can we do better than this?

Lindelöf Covering Theorem

Theorem (Lindelöf Covering Theorem)

Let $A \subseteq \mathbb{R}^n$ be a set and suppose $\{U_i : i \in I\}$ is an open covering of A . Then there exists a countable subcover $\{U_j : j \in J\}$.

Question

Can we do better than this?

Definition

A set $A \subseteq \mathbb{R}^n$ is called **compact** if every open cover of A has a *finite* subcover.

Challenge

Problem

Show that a singleton set $A = \{\vec{x}\}$ is compact.

Challenge

Problem

Show that a singleton set $A = \{\vec{x}\}$ is compact.

Solution

Challenge

Problem

Show that a singleton set $A = \{\vec{x}\}$ is compact.

Solution

Suppose that $\{U_i : i \in I\}$ is an open cover of X .

Challenge

Problem

Show that a singleton set $A = \{\vec{x}\}$ is compact.

Solution

Suppose that $\{U_i : i \in I\}$ is an open cover of X .
Then $A \subseteq \bigcup_{i \in I} U_i$.

Challenge

Problem

Show that a singleton set $A = \{\vec{x}\}$ is compact.

Solution

Suppose that $\{U_i : i \in I\}$ is an open cover of X .

Then $A \subseteq \bigcup_{i \in I} U_i$.

This means that $\vec{x} \in \bigcup_{i \in I} U_i$.

Challenge

Problem

Show that a singleton set $A = \{\vec{x}\}$ is compact.

Solution

Suppose that $\{U_i : i \in I\}$ is an open cover of X .

Then $A \subseteq \bigcup_{i \in I} U_i$.

This means that $\vec{x} \in \bigcup_{i \in I} U_i$.

Therefore $\vec{x} \in U_j$ for some $j \in I$.

Challenge

Problem

Show that a singleton set $A = \{\vec{x}\}$ is compact.

Solution

Suppose that $\{U_i : i \in I\}$ is an open cover of X .

Then $A \subseteq \bigcup_{i \in I} U_i$.

This means that $\vec{x} \in \bigcup_{i \in I} U_i$.

Therefore $\vec{x} \in U_j$ for some $j \in I$.

It follows that $\{U_j\}$ is a subcover of A consisting of a single set.

Challenge

Problem

Show that a singleton set $A = \{\vec{x}\}$ is compact.

Solution

Suppose that $\{U_i : i \in I\}$ is an open cover of X .

Then $A \subseteq \bigcup_{i \in I} U_i$.

This means that $\vec{x} \in \bigcup_{i \in I} U_i$.

Therefore $\vec{x} \in U_j$ for some $j \in I$.

It follows that $\{U_j\}$ is a subcover of A consisting of a single set.

Thus every open cover has a finite subcover, making A compact!

Bounded sets

Bounded sets

We will show that the compact subsets of $\mathbb{R}^n$ are exactly the closed sets that are bounded.

Bounded sets

We will show that the compact subsets of $\mathbb{R}^n$ are exactly the closed sets that are bounded.

Definition

A subset $A \subseteq \mathbb{R}^n$ is called **bounded** if there exists $\vec{x} \in \mathbb{R}^n$ and $r > 0$ such that $A \subseteq B(\vec{x}; r)$.

Bounded sets

We will show that the compact subsets of $\mathbb{R}^n$ are exactly the closed sets that are bounded.

Definition

A subset $A \subseteq \mathbb{R}^n$ is called **bounded** if there exists $\vec{x} \in \mathbb{R}^n$ and $r > 0$ such that $A \subseteq B(\vec{x}; r)$.

Theorem

If A is a compact set, then A must be bounded.

Bounded sets

We will show that the compact subsets of $\mathbb{R}^n$ are exactly the closed sets that are bounded.

Definition

A subset $A \subseteq \mathbb{R}^n$ is called **bounded** if there exists $\vec{x} \in \mathbb{R}^n$ and $r > 0$ such that $A \subseteq B(\vec{x}; r)$.

Theorem

If A is a compact set, then A must be bounded.

Proof.

Suppose A is a compact set.

Bounded sets

We will show that the compact subsets of $\mathbb{R}^n$ are exactly the closed sets that are bounded.

Definition

A subset $A \subseteq \mathbb{R}^n$ is called **bounded** if there exists $\vec{x} \in \mathbb{R}^n$ and $r > 0$ such that $A \subseteq B(\vec{x}; r)$.

Theorem

If A is a compact set, then A must be bounded.

Proof.

Suppose A is a compact set.

Consider the family of sets $\{U_i : i \in I\}$ where $I = \mathbb{Z}_+$ and $U_i = B(\vec{0}; i)$.



Challenge

Problem

Show that $\{U_i : i \in I\}$ is an open cover of A .

Challenge

Problem

Show that $\{U_i : i \in I\}$ is an open cover of A .

Solution

Note that U_i is an open subset for all i , so we need only show $A \subseteq \bigcup_{i \in I} U_i$.

Challenge

Problem

Show that $\{U_i : i \in I\}$ is an open cover of A .

Solution

Note that U_i is an open subset for all i , so we need only show $A \subseteq \bigcup_{i \in I} U_i$.

$$\bigcup_{i \in I} U_i = \bigcup_{i=1}^{\infty} B(\vec{0}; i) \subseteq \mathbb{R}^n.$$

Challenge

Problem

Show that $\{U_i : i \in I\}$ is an open cover of A .

Solution

Note that U_i is an open subset for all i , so we need only show $A \subseteq \bigcup_{i \in I} U_i$.

$$\bigcup_{i \in I} U_i = \bigcup_{i=1}^{\infty} B(\vec{0}; i) \subseteq \mathbb{R}^n.$$

Moreover, if $\vec{x} \in \mathbb{R}^n$, then we can choose $N \in \mathbb{Z}_+$ such that $N > |\vec{x}|$.

Challenge

Problem

Show that $\{U_i : i \in I\}$ is an open cover of A .

Solution

Note that U_i is an open subset for all i , so we need only show $A \subseteq \bigcup_{i \in I} U_i$.

$$\bigcup_{i \in I} U_i = \bigcup_{i=1}^{\infty} B(\vec{0}; i) \subseteq \mathbb{R}^n.$$

Moreover, if $\vec{x} \in \mathbb{R}^n$, then we can choose $N \in \mathbb{Z}_+$ such that $N > |\vec{x}|$.

Therefore $\vec{x} \in B(\vec{0}; N)$ and it follows $\vec{x} \in \bigcup_{i \in I} U_i$.

Challenge

Problem

Show that $\{U_i : i \in I\}$ is an open cover of A .

Solution

Note that U_i is an open subset for all i , so we need only show $A \subseteq \bigcup_{i \in I} U_i$.

$$\bigcup_{i \in I} U_i = \bigcup_{i=1}^{\infty} B(\vec{0}; i) \subseteq \mathbb{R}^n.$$

Moreover, if $\vec{x} \in \mathbb{R}^n$, then we can choose $N \in \mathbb{Z}_+$ such that $N > |\vec{x}|$.

Therefore $\vec{x} \in B(\vec{0}; N)$ and it follows $\vec{x} \in \bigcup_{i \in I} U_i$.

Thus $\mathbb{R}^n = \bigcup_{i \in I} U_i$, and since $A \subseteq \mathbb{R}^n$, we see $A \subseteq \bigcup_{i \in I} U_i$.

Challenge

Problem

Explain what it means that $\{U_i : i \in I\}$ has a finite subcover.

Challenge

Problem

Explain what it means that $\{U_i : i \in I\}$ has a finite subcover.

Solution

There is a finite subset $\{i_1, i_2, \dots, i_m\} \subseteq I$ with $A \subseteq \bigcup_{k=1}^m U_{i_k}$.

Challenge

Problem

Show that $\bigcup_{k=1}^m U_{i_k}$ is an open ball, and thus A is bounded.

Challenge

Problem

Show that $\bigcup_{k=1}^m U_{i_k}$ is an open ball, and thus A is bounded.

Solution

Without loss of generality, $i_1 < i_2 < \dots i_m$.

Challenge

Problem

Show that $\bigcup_{k=1}^m U_{i_k}$ is an open ball, and thus A is bounded.

Solution

Without loss of generality, $i_1 < i_2 < \dots < i_m$. Then

$$B(\vec{0}; i_1) \subseteq B(\vec{0}; i_2) \subseteq \dots \subseteq B(\vec{0}; i_m).$$

Challenge

Problem

Show that $\bigcup_{k=1}^m U_{i_k}$ is an open ball, and thus A is bounded.

Solution

Without loss of generality, $i_1 < i_2 < \dots i_m$. Then

$$B(\vec{0}; i_1) \subseteq B(\vec{0}; i_2) \subseteq \dots \subseteq B(\vec{0}; i_m).$$

This means

$$A \subseteq \bigcup_{k=1}^m A_{i_k} = \bigcup_{k=1}^m B(\vec{0}; i_k) = B(\vec{0}, i_m).$$

Challenge

Problem

Show that $\bigcup_{k=1}^m U_{i_k}$ is an open ball, and thus A is bounded.

Solution

Without loss of generality, $i_1 < i_2 < \dots < i_m$. Then

$$B(\vec{0}; i_1) \subseteq B(\vec{0}; i_2) \subseteq \dots \subseteq B(\vec{0}; i_m).$$

This means

$$A \subseteq \bigcup_{k=1}^m A_{i_k} = \bigcup_{k=1}^m B(\vec{0}; i_k) = B(\vec{0}, i_m).$$

This shows A is bounded.

Compact implies closed

Theorem

Suppose that $A \subseteq \mathbb{R}^n$ is compact. Then A is closed.

Compact implies closed

Theorem

Suppose that $A \subseteq \mathbb{R}^n$ is compact. Then A is closed.

Proof.

Compact implies closed

Theorem

Suppose that $A \subseteq \mathbb{R}^n$ is compact. Then A is closed.

Proof.

Suppose that A is compact and let $\vec{x} \in \mathbb{R}^n \setminus A$.

Compact implies closed

Theorem

Suppose that $A \subseteq \mathbb{R}^n$ is compact. Then A is closed.

Proof.

Suppose that A is compact and let $\vec{x} \in \mathbb{R}^n \setminus A$.

We will show $\vec{x}$ cannot be an adherent point of A and therefore A contains all of its adherent points.

Compact implies closed

Theorem

Suppose that $A \subseteq \mathbb{R}^n$ is compact. Then A is closed.

Proof.

Suppose that A is compact and let $\vec{x} \in \mathbb{R}^n \setminus A$.

We will show $\vec{x}$ cannot be an adherent point of A and therefore A contains all of its adherent points.

Consider the family of sets $\{U_i : i \in I\}$ where $I = \mathbb{Z}_+$ and $U_i = \mathbb{R}^n \setminus \overline{B}(\vec{x}; 1/i)$. □

Challenge

Problem

Prove that $\bigcup_{i \in I} U_i = \mathbb{R}^n \setminus \{\vec{x}\}$.

Challenge

Problem

Explain why $\{U_i : i \in I\}$ is an open cover of A .

Challenge

Problem

Show that if $\{U_i : i \in I\}$ has a finite subcover, then $\vec{x}$ can't be an adherent point of $\vec{A}$.

Bolzano-Weierstrass Theorem

So far, we have seen compact sets are closed and bounded.

Bolzano-Weierstrass Theorem

So far, we have seen compact sets are closed and bounded.
We want to prove the opposite is true too!

Bolzano-Weierstrass Theorem

So far, we have seen compact sets are closed and bounded.
We want to prove the opposite is true too!
This will require obtaining some other fundamental results
about real numbers.

Bolzano-Weierstrass Theorem

So far, we have seen compact sets are closed and bounded.
We want to prove the opposite is true too!
This will require obtaining some other fundamental results
about real numbers.

Theorem (Bolzano-Weierstrass Theorem)

A bounded set $A \subseteq \mathbb{R}^n$ with infinitely many points must have an accumulation point.

Cantor Intersection Theorem

Theorem (Cantor Intersection Theorem)

Suppose that $C_1, C_2, C_3, \dots$ are non-empty closed, bounded sets with $C_{i+1} \subseteq C_i$ for all $i \in I = \mathbb{Z}_+$. Then $\bigcap_{i \in I} C_i$ is also non-empty.

Cantor Intersection Theorem

Theorem (Cantor Intersection Theorem)

Suppose that $C_1, C_2, C_3, \dots$ are non-empty closed, bounded sets with $C_{i+1} \subseteq C_i$ for all $i \in I = \mathbb{Z}_+$. Then $\bigcap_{i \in I} C_i$ is also non-empty.

Proof.

Cantor Intersection Theorem

Theorem (Cantor Intersection Theorem)

Suppose that $C_1, C_2, C_3, \dots$ are non-empty closed, bounded sets with $C_{i+1} \subseteq C_i$ for all $i \in I = \mathbb{Z}_+$. Then $\bigcap_{i \in I} C_i$ is also non-empty.

Proof.

If C_i is finite for some i , then the proof is simple. Assume otherwise.

Cantor Intersection Theorem

Theorem (Cantor Intersection Theorem)

Suppose that $C_1, C_2, C_3, \dots$ are non-empty closed, bounded sets with $C_{i+1} \subseteq C_i$ for all $i \in I = \mathbb{Z}_+$. Then $\bigcap_{i \in I} C_i$ is also non-empty.

Proof.

If C_i is finite for some i , then the proof is simple. Assume otherwise.

Then choose $\vec{x}_i \in C_i$ for all $i \in I$ all distinct.

Cantor Intersection Theorem

Theorem (Cantor Intersection Theorem)

Suppose that $C_1, C_2, C_3, \dots$ are non-empty closed, bounded sets with $C_{i+1} \subseteq C_i$ for all $i \in I = \mathbb{Z}_+$. Then $\bigcap_{i \in I} C_i$ is also non-empty.

Proof.

If C_i is finite for some i , then the proof is simple. Assume otherwise.

Then choose $\vec{x}_i \in C_i$ for all $i \in I$ all distinct.

The set $A = \{\vec{x}_i : i \in \mathbb{Z}_+\}$ is infinite and bounded (because it is contained in C_1).

Cantor Intersection Theorem

Theorem (Cantor Intersection Theorem)

Suppose that $C_1, C_2, C_3, \dots$ are non-empty closed, bounded sets with $C_{i+1} \subseteq C_i$ for all $i \in I = \mathbb{Z}_+$. Then $\bigcap_{i \in I} C_i$ is also non-empty.

Proof.

If C_i is finite for some i , then the proof is simple. Assume otherwise.

Then choose $\vec{x}_i \in C_i$ for all $i \in I$ all distinct.

The set $A = \{\vec{x}_i : i \in \mathbb{Z}_+\}$ is infinite and bounded (because it is contained in C_1).

Therefore the Bolzano-Weierstrass Theorem tells us it has an accumulation point $\vec{x}$.



Cantor Intersection Theorem

Theorem (Cantor Intersection Theorem)

Suppose that $C_1, C_2, C_3, \dots$ are non-empty closed, bounded sets with $C_{i+1} \subseteq C_i$ for all $i \in I = \mathbb{Z}_+$. Then $\bigcap_{i \in I} C_i$ is also non-empty.

Cantor Intersection Theorem

Theorem (Cantor Intersection Theorem)

Suppose that $C_1, C_2, C_3, \dots$ are non-empty closed, bounded sets with $C_{i+1} \subseteq C_i$ for all $i \in I = \mathbb{Z}_+$. Then $\bigcap_{i \in I} C_i$ is also non-empty.

Proof by Contradiction.

Cantor Intersection Theorem

Theorem (Cantor Intersection Theorem)

Suppose that $C_1, C_2, C_3, \dots$ are non-empty closed, bounded sets with $C_{i+1} \subseteq C_i$ for all $i \in I = \mathbb{Z}_+$. Then $\bigcap_{i \in I} C_i$ is also non-empty.

Proof by Contradiction.

In fact, if we let $A_m = \{\vec{x}_i : i \in \mathbb{Z}_+, i \geq m\}$, then $\vec{x}$ is an accumulation point of A_m for all m .

Cantor Intersection Theorem

Theorem (Cantor Intersection Theorem)

Suppose that $C_1, C_2, C_3, \dots$ are non-empty closed, bounded sets with $C_{i+1} \subseteq C_i$ for all $i \in I = \mathbb{Z}_+$. Then $\bigcap_{i \in I} C_i$ is also non-empty.

Proof by Contradiction.

In fact, if we let $A_m = \{\vec{x}_i : i \in \mathbb{Z}_+, i \geq m\}$, then $\vec{x}$ is an accumulation point of A_m for all m .

Moreover, $A_m \subseteq C_m$ so $\vec{x}$ is an accumulation point of C_m for all m .

Cantor Intersection Theorem

Theorem (Cantor Intersection Theorem)

Suppose that $C_1, C_2, C_3, \dots$ are non-empty closed, bounded sets with $C_{i+1} \subseteq C_i$ for all $i \in I = \mathbb{Z}_+$. Then $\bigcap_{i \in I} C_i$ is also non-empty.

Proof by Contradiction.

In fact, if we let $A_m = \{\vec{x}_i : i \in \mathbb{Z}_+, i \geq m\}$, then $\vec{x}$ is an accumulation point of A_m for all m .

Moreover, $A_m \subseteq C_m$ so $\vec{x}$ is an accumulation point of C_m for all m .

Since C_m is closed, it follows $\vec{x} \in C_m$, and thus $\vec{x} \in \bigcap_{i \in I} C_i$.

Cantor Intersection Theorem

Theorem (Cantor Intersection Theorem)

Suppose that $C_1, C_2, C_3, \dots$ are non-empty closed, bounded sets with $C_{i+1} \subseteq C_i$ for all $i \in I = \mathbb{Z}_+$. Then $\bigcap_{i \in I} C_i$ is also non-empty.

Proof by Contradiction.

In fact, if we let $A_m = \{\vec{x}_i : i \in \mathbb{Z}_+, i \geq m\}$, then $\vec{x}$ is an accumulation point of A_m for all m .

Moreover, $A_m \subseteq C_m$ so $\vec{x}$ is an accumulation point of C_m for all m .

Since C_m is closed, it follows $\vec{x} \in C_m$, and thus $\vec{x} \in \bigcap_{i \in I} C_i$. In particular the intersection is non-empty! □

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

We already proved compact implies closed and bounded.

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

We already proved compact implies closed and bounded.
Assume A is closed and bounded.

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

We already proved compact implies closed and bounded.

Assume A is closed and bounded.

Take an open cover $\{U_i : i \in I\}$.

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

We already proved compact implies closed and bounded.

Assume A is closed and bounded.

Take an open cover $\{U_i : i \in I\}$.

Then by Lindelöf's Theorem, we can take a countable subcover.

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

We already proved compact implies closed and bounded.

Assume A is closed and bounded.

Take an open cover $\{U_i : i \in I\}$.

Then by Lindelöf's Theorem, we can take a countable subcover.

Therefore without loss of generality, assume $I = \mathbb{Z}_+$.

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

We already proved compact implies closed and bounded.

Assume A is closed and bounded.

Take an open cover $\{U_i : i \in I\}$.

Then by Lindelöf's Theorem, we can take a countable subcover.

Therefore without loss of generality, assume $I = \mathbb{Z}_+$.

Define closed sets $C_1, C_2, C_3, \dots$ by

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

We already proved compact implies closed and bounded.

Assume A is closed and bounded.

Take an open cover $\{U_i : i \in I\}$.

Then by Lindelöf's Theorem, we can take a countable subcover.

Therefore without loss of generality, assume $I = \mathbb{Z}_+$.

Define closed sets $C_1, C_2, C_3, \dots$ by

$$C_m = A \cap \left(\mathbb{R}^n \setminus \bigcup_{i=1}^m U_i \right).$$

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

If C_m is empty for some m , then $\{U_1, \dots, U_m\}$ covers A .

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

If C_m is empty for some m , then $\{U_1, \dots, U_m\}$ covers A .
To prove this must happen, we assume otherwise.

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

If C_m is empty for some m , then $\{U_1, \dots, U_m\}$ covers A .
To prove this must happen, we assume otherwise.
Notice that $C_{m+1} \subseteq C_m$ for all i .

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

If C_m is empty for some m , then $\{U_1, \dots, U_m\}$ covers A .

To prove this must happen, we assume otherwise.

Notice that $C_{m+1} \subseteq C_m$ for all i .

Since they are non-empty, closed and bounded, the Cantor Intersection Theorem says $\bigcap_{m=1}^{\infty} C_m$ is non-empty.

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

If C_m is empty for some m , then $\{U_1, \dots, U_m\}$ covers A .

To prove this must happen, we assume otherwise.

Notice that $C_{m+1} \subseteq C_m$ for all i .

Since they are non-empty, closed and bounded, the Cantor Intersection Theorem says $\bigcap_{m=1}^{\infty} C_m$ is non-empty.

Let $\vec{x} \in \bigcap_{m=1}^{\infty} C_m$.

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

If C_m is empty for some m , then $\{U_1, \dots, U_m\}$ covers A .

To prove this must happen, we assume otherwise.

Notice that $C_{m+1} \subseteq C_m$ for all i .

Since they are non-empty, closed and bounded, the Cantor Intersection Theorem says $\bigcap_{m=1}^{\infty} C_m$ is non-empty.

Let $\vec{x} \in \bigcap_{m=1}^{\infty} C_m$.

Since $\vec{x} \in A \subseteq \bigcup_{i \in I} U_i$, we have that $\vec{x} \in U_k$ for some $k \in I$, meaning $\vec{x} \notin C_k$.

Heine-Borel Theorem

Theorem (Heine-Borel Theorem)

A subset $A \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof.

If C_m is empty for some m , then $\{U_1, \dots, U_m\}$ covers A .

To prove this must happen, we assume otherwise.

Notice that $C_{m+1} \subseteq C_m$ for all i .

Since they are non-empty, closed and bounded, the Cantor Intersection Theorem says $\bigcap_{m=1}^{\infty} C_m$ is non-empty.

Let $\vec{x} \in \bigcap_{m=1}^{\infty} C_m$.

Since $\vec{x} \in A \subseteq \bigcup_{i \in I} U_i$, we have that $\vec{x} \in U_k$ for some $k \in I$, meaning $\vec{x} \notin C_k$.

This is a contradiction.

