

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

November 12, 2024

Outline

- 1 Real Analysis Lecture 20
 - More on differentiation
 - Riemann-Stieltjes integrals

Outline

- 1 Real Analysis Lecture 20
 - More on differentiation
 - Riemann-Stieltjes integrals

Local maxima and minima

Definition

Let (M, d) be a metric space and f be a real-valued function on a subset $S \subseteq M$.

Local maxima and minima

Definition

Let (M, d) be a metric space and f be a real-valued function on a subset $S \subseteq M$. Then f has an **absolute (or global) maximum at** $c \in S$ if

$$f(x) \leq f(c) \quad \text{for all } x \in S.$$

Local maxima and minima

Definition

Let (M, d) be a metric space and f be a real-valued function on a subset $S \subseteq M$. Then f has an **absolute (or global) maximum at** $c \in S$ if

$$f(x) \leq f(c) \quad \text{for all } x \in S.$$

It has **local maximum at** $a \in S$ if there exists an $r > 0$ such that

$$f(x) \leq f(c) \quad \text{for all } x \in B_M(c; r) \cap S.$$

Local maxima and minima

Definition

Let (M, d) be a metric space and f be a real-valued function on a subset $S \subseteq M$. Then f has an **absolute (or global) maximum at** $c \in S$ if

$$f(x) \leq f(c) \quad \text{for all } x \in S.$$

It has **local maximum at** $a \in S$ if there exists an $r > 0$ such that

$$f(x) \leq f(c) \quad \text{for all } x \in B_M(c; r) \cap S.$$

Absolute and local minima are defined similarly.

Critical points

Definition

Let f be a real-valued function on an open interval (a, b) .

Critical points

Definition

Let f be a real-valued function on an open interval (a, b) . A point $c \in (a, b)$ is called a **critical point** of f if either f is not differentiable at c or else $f'(c) = 0$.

Critical points

Definition

Let f be a real-valued function on an open interval (a, b) . A point $c \in (a, b)$ is called a **critical point** of f if either f is not differentiable at c or else $f'(c) = 0$.

Theorem

Let f be a real-valued function on an open interval (a, b) and suppose that f has a local maximum or local minimum at $c \in (a, b)$. Then c is a critical point of f .

Critical points

Proof.

Critical points

Proof.

We will prove this in the case c is a local maximum. The proof in the case of a local minimum is similar.

Critical points

Proof.

We will prove this in the case c is a local maximum. The proof in the case of a local minimum is similar.

If f is not differentiable at c , then we're done!

Critical points

Proof.

We will prove this in the case c is a local maximum. The proof in the case of a local minimum is similar.

If f is not differentiable at c , then we're done!

Therefore assume that it is differentiable.

Critical points

Proof.

We will prove this in the case c is a local maximum. The proof in the case of a local minimum is similar.

If f is not differentiable at c , then we're done!

Therefore assume that it is differentiable.

By definition, there exists $r > 0$ such that

$$f(x) \leq f(c) \quad \text{for all } x \in (c - r, c + r) \cap (a, b).$$

Critical points

Proof.

We will prove this in the case c is a local maximum. The proof in the case of a local minimum is similar.

If f is not differentiable at c , then we're done!

Therefore assume that it is differentiable.

By definition, there exists $r > 0$ such that

$$f(x) \leq f(c) \quad \text{for all } x \in (c - r, c + r) \cap (a, b).$$

To obtain a contradiction, suppose that $f'(c) \neq 0$.

Critical points

Proof.

We will prove this in the case c is a local maximum. The proof in the case of a local minimum is similar.

If f is not differentiable at c , then we're done!

Therefore assume that it is differentiable.

By definition, there exists $r > 0$ such that

$$f(x) \leq f(c) \quad \text{for all } x \in (c - r, c + r) \cap (a, b).$$

To obtain a contradiction, suppose that $f'(c) \neq 0$.

Then by the Helper Theorem, there is a function $f^*(x)$ defined on (a, b) and continuous at $c \in (a, b)$ with

$$f(x) - f(c) = (x - c)f^*(x).$$



Critical points

Proof.

Moreover, $f^*(c) = f'(c) \neq 0$.

Critical points

Proof.

Moreover, $f^*(c) = f'(c) \neq 0$.

Since f is continuous, we may choose $\delta > 0$ such that

$$|x - c| < \delta \Rightarrow |f^*(x) - f^*(c)| < |f^*(c)|/2.$$

Critical points

Proof.

Moreover, $f^*(c) = f'(c) \neq 0$.

Since f is continuous, we may choose $\delta > 0$ such that

$$|x - c| < \delta \Rightarrow |f^*(x) - f^*(c)| < |f^*(c)|/2.$$

Moreover, we may assume that $\delta < r$.

Critical points

Proof.

Moreover, $f^*(c) = f'(c) \neq 0$.

Since f is continuous, we may choose $\delta > 0$ such that

$$|x - c| < \delta \Rightarrow |f^*(x) - f^*(c)| < |f^*(c)|/2.$$

Moreover, we may assume that $\delta < r$.

It follows that $f^*(x) > |f^*(c)|/2$ for all $x \in (a, b)$ with $|x - c| < \delta$.

Critical points

Proof.

Moreover, $f^*(c) = f'(c) \neq 0$.

Since f is continuous, we may choose $\delta > 0$ such that

$$|x - c| < \delta \Rightarrow |f^*(x) - f^*(c)| < |f^*(c)|/2.$$

Moreover, we may assume that $\delta < r$.

It follows that $f^*(x) > |f^*(c)|/2$ for all $x \in (a, b)$ with $|x - c| < \delta$.

However,

$$f(c \pm \delta/2) = f(c) \pm \frac{1}{2}\delta f^*(c \pm \delta/2),$$

one of these will be larger than $f(c)$.

Critical points

Proof.

Moreover, $f^*(c) = f'(c) \neq 0$.

Since f is continuous, we may choose $\delta > 0$ such that

$$|x - c| < \delta \Rightarrow |f^*(x) - f^*(c)| < |f^*(c)|/2.$$

Moreover, we may assume that $\delta < r$.

It follows that $f^*(x) > |f^*(c)|/2$ for all $x \in (a, b)$ with $|x - c| < \delta$.

However,

$$f(c \pm \delta/2) = f(c) \pm \frac{1}{2} \delta f^*(c \pm \delta/2),$$

one of these will be larger than $f(c)$. Contradiction! □

Rolle's Theorem

Theorem

Rolle's Theorem Suppose that $f(x)$ is a real-valued function on $[a, b]$ which is

Rolle's Theorem

Theorem

Rolle's Theorem Suppose that $f(x)$ is a real-valued function on $[a, b]$ which is

- *continuous on $[a, b]$;*

Rolle's Theorem

Theorem

Rolle's Theorem Suppose that $f(x)$ is a real-valued function on $[a, b]$ which is

- *continuous on $[a, b]$;*
- *differentiable at each point in (a, b) ;*

Rolle's Theorem

Theorem

Rolle's Theorem Suppose that $f(x)$ is a real-valued function on $[a, b]$ which is

- *continuous on $[a, b]$;*
- *differentiable at each point in (a, b) ;*
- *and which satisfies $f(a) = f(b)$.*

Rolle's Theorem

Theorem

Rolle's Theorem Suppose that $f(x)$ is a real-valued function on $[a, b]$ which is

- *continuous on $[a, b]$;*
- *differentiable at each point in (a, b) ;*
- *and which satisfies $f(a) = f(b)$.*

Then there exists $c \in (a, b)$ with $f'(c) = 0$.

Rolle's Theorem

Proof.

Suppose otherwise.

Rolle's Theorem

Proof.

Suppose otherwise.

By the Extreme Value Theorem, $f(x)$ has an absolute maximum on $[a, b]$, as well as an absolute minimum on $[a, b]$.

Rolle's Theorem

Proof.

Suppose otherwise.

By the Extreme Value Theorem, $f(x)$ has an absolute maximum on $[a, b]$, as well as an absolute minimum on $[a, b]$. If both of these occur at the endpoints, then the absolute max is the absolute min, forcing f to be constant.

Rolle's Theorem

Proof.

Suppose otherwise.

By the Extreme Value Theorem, $f(x)$ has an absolute maximum on $[a, b]$, as well as an absolute minimum on $[a, b]$.

If both of these occur at the endpoints, then the absolute max is the absolute min, forcing f to be constant.

Then $f'(c) = 0$ for every $c \in (a, b)$.

Rolle's Theorem

Proof.

Suppose otherwise.

By the Extreme Value Theorem, $f(x)$ has an absolute maximum on $[a, b]$, as well as an absolute minimum on $[a, b]$. If both of these occur at the endpoints, then the absolute max is the absolute min, forcing f to be constant.

Then $f'(c) = 0$ for every $c \in (a, b)$.

Otherwise at least one of the points happens inside, so $f(x)$ has a local max or local min at a value c inside (a, b) .

Rolle's Theorem

Proof.

Suppose otherwise.

By the Extreme Value Theorem, $f(x)$ has an absolute maximum on $[a, b]$, as well as an absolute minimum on $[a, b]$. If both of these occur at the endpoints, then the absolute max is the absolute min, forcing f to be constant.

Then $f'(c) = 0$ for every $c \in (a, b)$.

Otherwise at least one of the points happens inside, so $f(x)$ has a local max or local min at a value c inside (a, b) .

The point c is a critical point, and since f is differentiable at c , we must have $f'(c) = 0$. □

Mean Value Theorem

Theorem

Rolle's Theorem Suppose that $f(x)$ is a real-valued function on $[a, b]$ which is

Mean Value Theorem

Theorem

Rolle's Theorem Suppose that $f(x)$ is a real-valued function on $[a, b]$ which is

- *continuous on $[a, b]$;*

Mean Value Theorem

Theorem

Rolle's Theorem Suppose that $f(x)$ is a real-valued function on $[a, b]$ which is

- *continuous on $[a, b]$;*
- *and differentiable at each point in (a, b) .*

Mean Value Theorem

Theorem

Rolle's Theorem Suppose that $f(x)$ is a real-valued function on $[a, b]$ which is

- *continuous on $[a, b]$;*
- *and differentiable at each point in (a, b) .*

Then there exists $c \in (a, b)$ with

$$f'(c) = \frac{f(b) - f(a)}{b - a}.$$

Mean Value Theorem

Proof.

Consider the function

$$g(x) = f(x) - \left(\frac{f(b) - f(a)}{b - a} \right) (x - a) + f(a).$$

Mean Value Theorem

Proof.

Consider the function

$$g(x) = f(x) - \left(\frac{f(b) - f(a)}{b - a} \right) (x - a) + f(a).$$

Then $g(x)$ is

Mean Value Theorem

Proof.

Consider the function

$$g(x) = f(x) - \left(\frac{f(b) - f(a)}{b - a} \right) (x - a) + f(a).$$

Then $g(x)$ is

- continuous on $[a, b]$;

Mean Value Theorem

Proof.

Consider the function

$$g(x) = f(x) - \left(\frac{f(b) - f(a)}{b - a} \right) (x - a) + f(a).$$

Then $g(x)$ is

- continuous on $[a, b]$;
- differentiable at each point in (a, b) ;

Mean Value Theorem

Proof.

Consider the function

$$g(x) = f(x) - \left(\frac{f(b) - f(a)}{b - a} \right) (x - a) + f(a).$$

Then $g(x)$ is

- continuous on $[a, b]$;
- differentiable at each point in (a, b) ;
- and satisfies $g(a) = g(b)$.

Mean Value Theorem

Proof.

Consider the function

$$g(x) = f(x) - \left(\frac{f(b) - f(a)}{b - a} \right) (x - a) + f(a).$$

Then $g(x)$ is

- continuous on $[a, b]$;
- differentiable at each point in (a, b) ;
- and satisfies $g(a) = g(b)$.

By Rolle's Theorem, there exists $c \in (a, b)$ with

$$0 = g'(c) = f'(c) - \frac{f(b) - f(a)}{b - a}.$$

Outline

- 1 Real Analysis Lecture 20
 - More on differentiation
 - Riemann-Stieltjes integrals

Integrals

Back in Calculus, we learned about **integrals**.

Integrals

Back in Calculus, we learned about **integrals**.
It turns out that there are many types of integrals:

Integrals

Back in Calculus, we learned about **integrals**.
It turns out that there are many types of integrals:

- the Riemann integral (our familiar friend)

Integrals

Back in Calculus, we learned about **integrals**.

It turns out that there are many types of integrals:

- the Riemann integral (our familiar friend)
- the Riemann-Stieltjes integral

Integrals

Back in Calculus, we learned about **integrals**.

It turns out that there are many types of integrals:

- the Riemann integral (our familiar friend)
- the Riemann-Stieltjes integral
- the Lebesgue integral ...

Integrals

Back in Calculus, we learned about **integrals**.

It turns out that there are many types of integrals:

- the Riemann integral (our familiar friend)
- the Riemann-Stieltjes integral
- the Lebesgue integral ...

Most of these generalize the Riemann integral in some sense.

Partitions

Definition

A **partition** P of $[a, b]$ is a set of points

$$\{x_0, x_1, \dots, x_n\}$$

with

$$a = x_0 < x_1 < x_2 < \dots < x_n = b.$$

Partitions

Definition

A **partition** P of $[a, b]$ is a set of points

$$\{x_0, x_1, \dots, x_n\}$$

with

$$a = x_0 < x_1 < x_2 < \dots < x_n = b.$$

A set of **sample points** for P is a set $\{t_1, \dots, t_n\}$ with $t_k \in [x_{k-1}, x_k]$ for all $1 \leq k \leq n$.

Partitions

Definition

A **partition** P of $[a, b]$ is a set of points

$$\{x_0, x_1, \dots, x_n\}$$

with

$$a = x_0 < x_1 < x_2 < \dots < x_n = b.$$

A set of **sample points** for P is a set $\{t_1, \dots, t_n\}$ with $t_k \in [x_{k-1}, x_k]$ for all $1 \leq k \leq n$. A partition P' is a **refinement** of P if $P \subseteq P'$.

Riemann-Stieltjes sums

Notation: given a real-valued function α on $[a, b]$

$$\Delta\alpha_k(P) = \alpha(x_k) - \alpha(x_{k-1}), \quad k = 1, 2, \dots, n.$$

Riemann-Stieltjes sums

Notation: given a real-valued function α on $[a, b]$

$$\Delta\alpha_k(P) = \alpha(x_k) - \alpha(x_{k-1}), \quad k = 1, 2, \dots, n.$$

Definition

Riemann-Stieltjes sums

Notation: given a real-valued function α on $[a, b]$

$$\Delta\alpha_k(P) = \alpha(x_k) - \alpha(x_{k-1}), \quad k = 1, 2, \dots, n.$$

Definition

Let f and α be real-valued functions on $[a, b]$, and let $P = \{x_0, x_1, \dots, x_n\}$ be a partition of $[a, b]$.

Riemann-Stieltjes sums

Notation: given a real-valued function α on $[a, b]$

$$\Delta\alpha_k(P) = \alpha(x_k) - \alpha(x_{k-1}), \quad k = 1, 2, \dots, n.$$

Definition

Let f and α be real-valued functions on $[a, b]$, and let $P = \{x_0, x_1, \dots, x_n\}$ be a partition of $[a, b]$. A sum of the form

$$S(P, f, \alpha, \{t_k\}) = \sum_{k=1}^n f(t_k) \Delta\alpha_k(P),$$

for some set of sample points $\{t_k\}$ of P .

Riemann-Stieltjes integral

Definition

Riemann-Stieltjes integral

Definition

Let f and α be real-valued functions on $[a, b]$.

Riemann-Stieltjes integral

Definition

Let f and α be real-valued functions on $[a, b]$. We say f is **Riemann integrable** with respect to α on $[a, b]$ if there exists $A \in \mathbb{R}$ with the following property:

Riemann-Stieltjes integral

Definition

Let f and α be real-valued functions on $[a, b]$. We say f is **Riemann integrable** with respect to α on $[a, b]$ if there exists $A \in \mathbb{R}$ with the following property: for all $\epsilon > 0$, there exists a partition P_ϵ of $[a, b]$ such that for all refinements P of P_ϵ

$$|S(P, f, \alpha, \{t_k\}) - A| < \epsilon,$$

for all sets of sample points $\{t_k\}$ of P .

Riemann-Stieltjes integral

Definition

Let f and α be real-valued functions on $[a, b]$. We say f is **Riemann integrable** with respect to α on $[a, b]$ if there exists $A \in \mathbb{R}$ with the following property: for all $\epsilon > 0$, there exists a partition P_ϵ of $[a, b]$ such that for all refinements P of P_ϵ

$$|S(P, f, \alpha, \{t_k\}) - A| < \epsilon,$$

for all sets of sample points $\{t_k\}$ of P .

Notation: in this case A is denoted by $\int_a^b f d\alpha$ or $\int_a^b f(x) d\alpha(x)$

Linearity of integration

Theorem

Suppose that f and g are Riemann integrable with respect to α on $[a, b]$.

Linearity of integration

Theorem

Suppose that f and g are Riemann integrable with respect to α on $[a, b]$.

Then for any $c_1, c_2 \in \mathbb{R}$, $c_1 f + c_2 g$ is Riemann integrable with respect to α and

$$\int_a^b (c_1 f + c_2 g) d\alpha = c_1 \int_a^b f d\alpha + c_2 \int_a^b g d\alpha.$$

Linearity of integration

Proof.

Linearity of integration

Proof.

Let $\epsilon > 0$ and WLOG assume $c_1, c_2 \neq 0$.

Linearity of integration

Proof.

Let $\epsilon > 0$ and WLOG assume $c_1, c_2 \neq 0$.

Choose a partition P'_ϵ such that

$$|S(P, f, \alpha, \{t_k\}) - \int_a^b f d\alpha| < \epsilon/2|c_1|,$$

for all refinements P of P'_ϵ and sets of sample points $\{t_k\}$ of P .

Linearity of integration

Proof.

Let $\epsilon > 0$ and WLOG assume $c_1, c_2 \neq 0$.

Choose a partition P'_ϵ such that

$$|S(P, f, \alpha, \{t_k\}) - \int_a^b f d\alpha| < \epsilon/2|c_1|,$$

for all refinements P of P'_ϵ and sets of sample points $\{t_k\}$ of P .

Also, choose a partition P''_ϵ such that

$$|S(P, g, \alpha, \{t_k\}) - \int_a^b g d\alpha| < \epsilon/2|c_2|,$$

for all refinements P of P''_ϵ and sets of sample points $\{t_k\}$ of P .

Linearity of integration

Proof.

Let $\epsilon > 0$ and WLOG assume $c_1, c_2 \neq 0$.

Choose a partition P'_ϵ such that

$$|S(P, f, \alpha, \{t_k\}) - \int_a^b f d\alpha| < \epsilon/2|c_1|,$$

for all refinements P of P'_ϵ and sets of sample points $\{t_k\}$ of P .

Also, choose a partition P''_ϵ such that

$$|S(P, g, \alpha, \{t_k\}) - \int_a^b g d\alpha| < \epsilon/2|c_2|,$$

for all refinements P of P''_ϵ and sets of sample points $\{t_k\}$ of P .

Let $P_\epsilon = P'_\epsilon \cup P''_\epsilon$.

Linearity of integration

Proof.

Let $\epsilon > 0$ and WLOG assume $c_1, c_2 \neq 0$.

Choose a partition P'_ϵ such that

$$|S(P, f, \alpha, \{t_k\}) - \int_a^b f d\alpha| < \epsilon/2|c_1|,$$

for all refinements P of P'_ϵ and sets of sample points $\{t_k\}$ of P .

Also, choose a partition P''_ϵ such that

$$|S(P, g, \alpha, \{t_k\}) - \int_a^b g d\alpha| < \epsilon/2|c_2|,$$

for all refinements P of P''_ϵ and sets of sample points $\{t_k\}$ of P .
Let $P_\epsilon = P'_\epsilon \cup P''_\epsilon$. Then any refinement P of P_ϵ , is a refinement of P'_ϵ and P''_ϵ .



Linearity of integration

Proof.

Therefore if P is a refinement of P_ϵ , and $\{t_k\}$ is a set of sample points of P , then by the triangle inequality

Linearity of integration

Proof.

Therefore if P is a refinement of P_ϵ , and $\{t_k\}$ is a set of sample points of P , then by the triangle inequality

$$\begin{aligned}
 & |S(P, c_1 f + c_2 g, \alpha, \{t_k\}) - (c_1 \int_a^b f d\alpha + c_2 \int_a^b g d\alpha)| \\
 &= |c_1 S(P, f, \alpha, \{t_k\}) + c_2 S(P, g, \alpha, \{t_k\}) - (c_1 \int_a^b f d\alpha + c_2 \int_a^b g d\alpha)| \\
 &= |c_1 S(P, f, \alpha, \{t_k\}) - c_1 \int_a^b f d\alpha + c_2 S(P, g, \alpha, \{t_k\}) - c_2 \int_a^b g d\alpha| \\
 &\leq |c_1| \cdot |S(P, f, \alpha, \{t_k\}) - \int_a^b f d\alpha| + |c_2| \cdot |S(P, g, \alpha, \{t_k\}) - \int_a^b g d\alpha| \\
 &\qquad\qquad\qquad < \epsilon/2 + \epsilon/2 = \epsilon.
 \end{aligned}$$

Linearity of integration

Proof.

Therefore if P is a refinement of P_ϵ , and $\{t_k\}$ is a set of sample points of P , then by the triangle inequality

$$\begin{aligned}
 & |S(P, c_1 f + c_2 g, \alpha, \{t_k\}) - (c_1 \int_a^b f d\alpha + c_2 \int_a^b g d\alpha)| \\
 &= |c_1 S(P, f, \alpha, \{t_k\}) + c_2 S(P, g, \alpha, \{t_k\}) - (c_1 \int_a^b f d\alpha + c_2 \int_a^b g d\alpha)| \\
 &= |c_1 S(P, f, \alpha, \{t_k\}) - c_1 \int_a^b f d\alpha + c_2 S(P, g, \alpha, \{t_k\}) - c_2 \int_a^b g d\alpha| \\
 &\leq |c_1| \cdot |S(P, f, \alpha, \{t_k\}) - \int_a^b f d\alpha| + |c_2| \cdot |S(P, g, \alpha, \{t_k\}) - \int_a^b g d\alpha| \\
 &\qquad\qquad\qquad < \epsilon/2 + \epsilon/2 = \epsilon.
 \end{aligned}$$

Since $\epsilon > 0$ was arbitrary, this proves the theorem. □

Linearity of integration

Theorem

Suppose that f is Riemann integrable with respect to both α and β on $[a, b]$.

Linearity of integration

Theorem

Suppose that f is Riemann integrable with respect to both α and β on $[a, b]$.

Then for any $c_1, c_2 \in \mathbb{R}$, f is Riemann integrable with respect to $c_1\alpha + c_2\beta$ and

$$\int_a^b f d(c_1\alpha + c_2\beta) = c_1 \int_a^b f d\alpha + c_2 \int_a^b f d\beta.$$

Challenge!

Problem

Prove the last theorem!

Linearity of integration

Theorem

Suppose that f is a real-valued function on $[a, b]$ and that $c \in (a, b)$.

Linearity of integration

Theorem

Suppose that f is a real-valued function on $[a, b]$ and that $c \in (a, b)$. If $\int_a^c f d\alpha$ and $\int_c^b f d\alpha$ exist, then $\int_a^b f d\alpha$ exists and

$$\int_a^c f d\alpha + \int_c^b f d\alpha = \int_a^b f d\alpha.$$

Linearity of integration

Theorem

Suppose that f is a real-valued function on $[a, b]$ and that $c \in (a, b)$. If $\int_a^c f d\alpha$ and $\int_c^b f d\alpha$ exist, then $\int_a^b f d\alpha$ exists and

$$\int_a^c f d\alpha + \int_c^b f d\alpha = \int_a^b f d\alpha.$$

Conversely, if $\int_a^b f d\alpha$ exists then $\int_a^c f d\alpha$ and $\int_c^b f d\alpha$ exist and

$$\int_a^c f d\alpha + \int_c^b f d\alpha = \int_a^b f d\alpha.$$