

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

October 20, 2025

Outline

- 1 Real Analysis Lecture 16
 - Continuity
 - Connectedness

Outline

- 1 Real Analysis Lecture 16
 - Continuity
 - Connectedness

Continuity

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

Continuity

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

Definition

Continuity

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

Definition

Then f is continuous at a point $p \in S$ if for all $\epsilon > 0$ there exists $\delta > 0$ such that

$$d_S(x, p) < \delta \Rightarrow d_T(f(x), f(p)) < \epsilon.$$

Continuity

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

Definition

Then f is continuous at a point $p \in S$ if for all $\epsilon > 0$ there exists $\delta > 0$ such that

$$d_S(x, p) < \delta \Rightarrow d_T(f(x), f(p)) < \epsilon.$$

Equivalently, $f(B_S(p; \delta)) \subseteq B_T(f(p); \epsilon)$.

Continuity

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

Definition

Then f is continuous at a point $p \in S$ if for all $\epsilon > 0$ there exists $\delta > 0$ such that

$$d_S(x, p) < \delta \Rightarrow d_T(f(x), f(p)) < \epsilon.$$

Equivalently, $f(B_S(p; \delta)) \subseteq B_T(f(p); \epsilon)$. We say f is continuous on $A \subseteq S$ if $f(x)$ is continuous at every point $p \in A$.

Continuity

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

Definition

Then f is continuous at a point $p \in S$ if for all $\epsilon > 0$ there exists $\delta > 0$ such that

$$d_S(x, p) < \delta \Rightarrow d_T(f(x), f(p)) < \epsilon.$$

Equivalently, $f(B_S(p; \delta)) \subseteq B_T(f(p); \epsilon)$. We say f is continuous on $A \subseteq S$ if $f(x)$ is continuous at every point $p \in A$. If f is continuous on S , we simply say f is continuous.

Sequence version

Sequence version

Theorem

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

Sequence version

Theorem

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function. Then f is continuous at a point p in S if for all sequences $\{x_n\}$ of values in S which converge to p , we have

Sequence version

Theorem

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function. Then f is continuous at a point p in S if for all sequences $\{x_n\}$ of values in S which converge to p , we have

$$\lim_{n \rightarrow \infty} f(x_n) = f(p).$$

Sequence version

Theorem

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function. Then f is continuous at a point p in S if for all sequences $\{x_n\}$ of values in S which converge to p , we have

$$\lim_{n \rightarrow \infty} f(x_n) = f(p).$$

Intuitively, continuity means

$$\lim_{n \rightarrow \infty} f(x_n) = f\left(\lim_{n \rightarrow \infty} x_n\right).$$

Compositions

Theorem

Suppose (S, d_S) , (T, d_T) , and (U, d_U) are metric spaces and $f : S \rightarrow T$ and $g : T \rightarrow U$ are functions.

Compositions

Theorem

Suppose (S, d_S) , (T, d_T) , and (U, d_U) are metric spaces and $f : S \rightarrow T$ and $g : T \rightarrow U$ are functions.

If f is continuous at p and g is continuous at $f(p)$, then the composition $h = g \circ f : S \rightarrow U$ defined by $h(x) = g(f(x))$ is continuous at p .

Challenge!

Problem

Prove the previous theorem!

Preimages

Definition

Suppose $f : S \rightarrow T$ is a function and that $Y \subseteq T$.

Preimages

Definition

Suppose $f : S \rightarrow T$ is a function and that $Y \subseteq T$.
The **inverse image** or **preimage** of Y is the set

$$f^{-1}(Y) = \{x : x \in S, \text{ and } f(x) \in Y\}.$$

Challenge!

Problem

Suppose that $f : S \rightarrow T$ and $Y \subseteq Z \subseteq T$.
Prove that

$$f^{-1}(Y) \subseteq f^{-1}(Z).$$

Open sets

Theorem

Suppose (S, d_S) and (T, d_T) are metric spaces and $f : S \rightarrow T$ is a function.

Open sets

Theorem

Suppose (S, d_S) and (T, d_T) are metric spaces and $f : S \rightarrow T$ is a function.

Then f is continuous if and only if $f^{-1}(V)$ is open for all open subsets $V \subseteq T$.

Closed sets

Theorem

Suppose (S, d_S) and (T, d_T) are metric spaces and $f : S \rightarrow T$ is a function.

Closed sets

Theorem

Suppose (S, d_S) and (T, d_T) are metric spaces and $f : S \rightarrow T$ is a function.

Then f is continuous if and only if $f^{-1}(A)$ is closed for all closed subsets $A \subseteq T$.

Open sets

Theorem

Suppose (S, d_S) and (T, d_T) are metric spaces and $f : S \rightarrow T$ is a function.

Open sets

Theorem

Suppose (S, d_S) and (T, d_T) are metric spaces and $f : S \rightarrow T$ is a function.

Then f is continuous if and only if $f^{-1}(V)$ is open for all open subsets $V \subseteq T$.

Open sets

Proof.

Open sets

Proof.

$(\Rightarrow)$

Open sets

Proof.

($\Rightarrow$)

Suppose $V \subseteq T$ is open.

Open sets

Proof.

$(\Rightarrow)$

Suppose $V \subseteq T$ is open.

Then for all $x \in f^{-1}(V)$, we have $x \in S$ and $f(x) \in V$.

Open sets

Proof.

($\Rightarrow$)

Suppose $V \subseteq T$ is open.

Then for all $x \in f^{-1}(V)$, we have $x \in S$ and $f(x) \in V$.

Since V is open, there exists $\epsilon > 0$ such that $B_T(f(x); \epsilon) \subseteq V$.

Open sets

Proof.

($\Rightarrow$)

Suppose $V \subseteq T$ is open.

Then for all $x \in f^{-1}(V)$, we have $x \in S$ and $f(x) \in V$.

Since V is open, there exists $\epsilon > 0$ such that $B_T(f(x); \epsilon) \subseteq V$.

Since f is continuous at x , there exists $\delta > 0$ such that

$f(B_S(x; \delta)) \subseteq B_T(f(x); \epsilon) \subseteq V$.

Open sets

Proof.

($\Rightarrow$)

Suppose $V \subseteq T$ is open.

Then for all $x \in f^{-1}(V)$, we have $x \in S$ and $f(x) \in V$.

Since V is open, there exists $\epsilon > 0$ such that $B_T(f(x); \epsilon) \subseteq V$.

Since f is continuous at x , there exists $\delta > 0$ such that

$f(B_S(x; \delta)) \subseteq B_T(f(x); \epsilon) \subseteq V$.

Consequently $B_S(x; \delta) \subseteq f^{-1}(V)$.

Open sets

Proof.

($\Rightarrow$)

Suppose $V \subseteq T$ is open.

Then for all $x \in f^{-1}(V)$, we have $x \in S$ and $f(x) \in V$.

Since V is open, there exists $\epsilon > 0$ such that $B_T(f(x); \epsilon) \subseteq V$.

Since f is continuous at x , there exists $\delta > 0$ such that

$f(B_S(x; \delta)) \subseteq B_T(f(x); \epsilon) \subseteq V$.

Consequently $B_S(x; \delta) \subseteq f^{-1}(V)$.

This shows that x is an interior point of $f^{-1}(V)$.

Open sets

Proof.

($\Rightarrow$)

Suppose $V \subseteq T$ is open.

Then for all $x \in f^{-1}(V)$, we have $x \in S$ and $f(x) \in V$.

Since V is open, there exists $\epsilon > 0$ such that $B_T(f(x); \epsilon) \subseteq V$.

Since f is continuous at x , there exists $\delta > 0$ such that

$f(B_S(x; \delta)) \subseteq B_T(f(x); \epsilon) \subseteq V$.

Consequently $B_S(x; \delta) \subseteq f^{-1}(V)$.

This shows that x is an interior point of $f^{-1}(V)$.

Since x was arbitrary, this proves $f^{-1}(V)$ is open. □

Open sets

Proof.

Open sets

Proof.

($\Leftarrow$)

Open sets

Proof.

($\Leftarrow$)

Suppose that for all open subsets $V \subseteq T$, the preimage $f^{-1}(V)$ is open.

Open sets

Proof.

($\Leftarrow$)

Suppose that for all open subsets $V \subseteq T$, the preimage $f^{-1}(V)$ is open.

Let $x \in S$.

Open sets

Proof.

($\Leftarrow$)

Suppose that for all open subsets $V \subseteq T$, the preimage $f^{-1}(V)$ is open.

Let $x \in S$.

Then for all $\epsilon > 0$, the ball $B_T(f(x); \epsilon)$ is open.

Open sets

Proof.

($\Leftarrow$)

Suppose that for all open subsets $V \subseteq T$, the preimage $f^{-1}(V)$ is open.

Let $x \in S$.

Then for all $\epsilon > 0$, the ball $B_T(f(x); \epsilon)$ is open.

Therefore the preimage $f^{-1}(B_T(f(x); \epsilon))$ is open.

Open sets

Proof.

($\Leftarrow$)

Suppose that for all open subsets $V \subseteq T$, the preimage $f^{-1}(V)$ is open.

Let $x \in S$.

Then for all $\epsilon > 0$, the ball $B_T(f(x); \epsilon)$ is open.

Therefore the preimage $f^{-1}(B_T(f(x); \epsilon))$ is open.

It contains the point x , which must be an interior point.

Open sets

Proof.

($\Leftarrow$)

Suppose that for all open subsets $V \subseteq T$, the preimage $f^{-1}(V)$ is open.

Let $x \in S$.

Then for all $\epsilon > 0$, the ball $B_T(f(x); \epsilon)$ is open.

Therefore the preimage $f^{-1}(B_T(f(x); \epsilon))$ is open.

It contains the point x , which must be an interior point.

This means that there exists $\delta > 0$ such that

$$B_S(x; \delta) \subseteq f^{-1}(B_T(f(x); \epsilon)).$$

Open sets

Proof.

($\Leftarrow$)

Suppose that for all open subsets $V \subseteq T$, the preimage $f^{-1}(V)$ is open.

Let $x \in S$.

Then for all $\epsilon > 0$, the ball $B_T(f(x); \epsilon)$ is open.

Therefore the preimage $f^{-1}(B_T(f(x); \epsilon))$ is open.

It contains the point x , which must be an interior point.

This means that there exists $\delta > 0$ such that

$$B_S(x; \delta) \subseteq f^{-1}(B_T(f(x); \epsilon)).$$

$$\text{Thus } f(B_S(x; \delta)) \subseteq B_T(f(x); \epsilon).$$



Relatively open sets

Theorem

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

If f is continuous on a subset $A \subseteq S$, then for any open subset $V \subseteq T$ there exists an open subset $U \subseteq S$ with

$$f^{-1}(V) \cap A = U \cap A.$$

Relatively open sets

Proof.

Relatively open sets

Proof.

Consider the subspace $(A, d_A = d_S)$ of S .

Relatively open sets

Proof.

Consider the subspace $(A, d_A = d_S)$ of S .

The function f restricts to a continuous function $f|_A : A \rightarrow T$.

Relatively open sets

Proof.

Consider the subspace $(A, d_A = d_S)$ of S .

The function f restricts to a continuous function $f|_A : A \rightarrow T$.

Therefore if $V \subseteq T$ is open, the preimage

$$f|_A^{-1}(V) = \{x : x \in A, \text{ and } f(x) \in V\} = f^{-1}(V) \cap A$$

Relatively open sets

Proof.

Consider the subspace $(A, d_A = d_S)$ of S .

The function f restricts to a continuous function $f|_A : A \rightarrow T$.

Therefore if $V \subseteq T$ is open, the preimage

$$f|_A^{-1}(V) = \{x : x \in A, \text{ and } f(x) \in V\} = f^{-1}(V) \cap A$$

is open in the subspace A of S .

Relatively open sets

Proof.

Consider the subspace $(A, d_A = d_S)$ of S .

The function f restricts to a continuous function $f|_A : A \rightarrow T$.

Therefore if $V \subseteq T$ is open, the preimage

$$f|_A^{-1}(V) = \{x : x \in A, \text{ and } f(x) \in V\} = f^{-1}(V) \cap A$$

is open in the subspace A of S .

This is the same thing as

$$f^{-1}(V) \cap A = U \cap A$$

for some open subset

$$U \subseteq S$$



Compact sets

Theorem

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

If f is continuous on a compact subset $A \subseteq S$, then $f(A)$ is compact.

Compact sets

Theorem

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

If f is continuous on a compact subset $A \subseteq S$, then $f(A)$ is compact.

Proof.

Compact sets

Theorem

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

If f is continuous on a compact subset $A \subseteq S$, then $f(A)$ is compact.

Proof.

Suppose that $\{V_i : i \in I\}$ is an open cover of $f(A)$.

Compact sets

Theorem

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

If f is continuous on a compact subset $A \subseteq S$, then $f(A)$ is compact.

Proof.

Suppose that $\{V_i : i \in I\}$ is an open cover of $f(A)$.

Then $f^{-1}(V_i) \cap A = U_i \cap A$ for some open subset $U_i \subseteq S$.

Compact sets

Theorem

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

If f is continuous on a compact subset $A \subseteq S$, then $f(A)$ is compact.

Proof.

Suppose that $\{V_i : i \in I\}$ is an open cover of $f(A)$.

Then $f^{-1}(V_i) \cap A = U_i \cap A$ for some open subset $U_i \subseteq S$.

The open subsets $\{U_i : i \in I\}$ cover A , and since A is compact there is a finite subcover $\{U_1, \dots, U_n\}$.

Compact sets

Theorem

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

If f is continuous on a compact subset $A \subseteq S$, then $f(A)$ is compact.

Proof.

Suppose that $\{V_i : i \in I\}$ is an open cover of $f(A)$.

Then $f^{-1}(V_i) \cap A = U_i \cap A$ for some open subset $U_i \subseteq S$.

The open subsets $\{U_i : i \in I\}$ cover A , and since A is compact there is a finite subcover $\{U_1, \dots, U_n\}$.

This means $A \subseteq \bigcup_{i=1}^n U_i$ and therefore

Compact sets

Theorem

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

If f is continuous on a compact subset $A \subseteq S$, then $f(A)$ is compact.

Proof.

Suppose that $\{V_i : i \in I\}$ is an open cover of $f(A)$.

Then $f^{-1}(V_i) \cap A = U_i \cap A$ for some open subset $U_i \subseteq S$.

The open subsets $\{U_i : i \in I\}$ cover A , and since A is compact there is a finite subcover $\{U_1, \dots, U_n\}$.

This means $A \subseteq \bigcup_{i=1}^n U_i$ and therefore

$$f(A) \subseteq \bigcup_{i=1}^n V_i.$$

Compact sets

Theorem

Let (S, d_S) and (T, d_T) be metric spaces and $f : S \rightarrow T$ a function.

If f is continuous on a compact subset $A \subseteq S$, then $f(A)$ is compact.

Proof.

Suppose that $\{V_i : i \in I\}$ is an open cover of $f(A)$.

Then $f^{-1}(V_i) \cap A = U_i \cap A$ for some open subset $U_i \subseteq S$.

The open subsets $\{U_i : i \in I\}$ cover A , and since A is compact there is a finite subcover $\{U_1, \dots, U_n\}$.

This means $A \subseteq \bigcup_{i=1}^n U_i$ and therefore

$$f(A) \subseteq \bigcup_{i=1}^n V_i.$$

Extreme value theorem

Theorem

Suppose (S, d_S) is a metric space and $f : S \rightarrow \mathbb{R}$ is a function (to Euclidean space).

If f is continuous on a compact subset $A \subseteq S$, then there exists $a, b \in A$ such that

$$f(a) = \inf f(A), \quad \text{and} \quad f(b) = \sup f(A).$$

Extreme value theorem

Extreme value theorem

Proof.

Extreme value theorem

Proof.

The set $f(A)$ is compact in $\mathbb{R}$, so it is closed and bounded.

Extreme value theorem

Proof.

The set $f(A)$ is compact in $\mathbb{R}$, so it is closed and bounded. The Completeness Axiom therefore tells us that it has an infimum $y = \inf(f(A))$ and supremum $z = \sup(f(A))$.

Extreme value theorem

Proof.

The set $f(A)$ is compact in $\mathbb{R}$, so it is closed and bounded.

The Completeness Axiom therefore tells us that it has an infimum $y = \inf(f(A))$ and supremum $z = \sup(f(A))$.

The infimum and supremum of a set are adherent points of that set.

Extreme value theorem

Proof.

The set $f(A)$ is compact in $\mathbb{R}$, so it is closed and bounded.

The Completeness Axiom therefore tells us that it has an infimum $y = \inf(f(A))$ and supremum $z = \sup(f(A))$.

The infimum and supremum of a set are adherent points of that set.

Therefore, since $f(A)$ is closed $y, z \in f(A)$.

Extreme value theorem

Proof.

The set $f(A)$ is compact in $\mathbb{R}$, so it is closed and bounded.

The Completeness Axiom therefore tells us that it has an infimum $y = \inf(f(A))$ and supremum $z = \sup(f(A))$.

The infimum and supremum of a set are adherent points of that set.

Therefore, since $f(A)$ is closed $y, z \in f(A)$.

This means that there exists $a, b \in A$ such that $y = f(a)$ and $z = f(b)$. □

Outline

- 1 Real Analysis Lecture 16
 - Continuity
 - Connectedness

Characterization of intervals

Theorem

If $I \subseteq \mathbb{R}$ is an interval, then there does not exist two nonempty, disjoint open sets $U, V \subseteq \mathbb{R}$ (Euclidean space), with

$$I \subseteq U \cup V.$$

Connected spaces

Definition

A metric space (S, d) is called **disconnected** if there exists two nonempty, disjoint open subsets $U, V \subseteq S$ with $S = U \cup V$.

Connected spaces

Definition

A metric space (S, d) is called **disconnected** if there exists two nonempty, disjoint open subsets $U, V \subseteq S$ with $S = U \cup V$. We call S **connected** if it is not disconnected.

Connected spaces

Definition

A metric space (S, d) is called **disconnected** if there exists two nonempty, disjoint open subsets $U, V \subseteq S$ with $S = U \cup V$.

We call S **connected** if it is not disconnected.

A subset $A \subseteq S$ is connected if the corresponding subspace (A, d) is connected.

Connected spaces

Definition

A metric space (S, d) is called **disconnected** if there exists two nonempty, disjoint open subsets $U, V \subseteq S$ with $S = U \cup V$.

We call S **connected** if it is not disconnected.

A subset $A \subseteq S$ is connected if the corresponding subspace (A, d) is connected.

Examples:

Connected spaces

Definition

A metric space (S, d) is called **disconnected** if there exists two nonempty, disjoint open subsets $U, V \subseteq S$ with $S = U \cup V$.

We call S **connected** if it is not disconnected.

A subset $A \subseteq S$ is connected if the corresponding subspace (A, d) is connected.

Examples:

- the real line with the Euclidean metric $(\mathbb{R}, d_{eucl})$ is connected

Connected spaces

Definition

A metric space (S, d) is called **disconnected** if there exists two nonempty, disjoint open subsets $U, V \subseteq S$ with $S = U \cup V$.

We call S **connected** if it is not disconnected.

A subset $A \subseteq S$ is connected if the corresponding subspace (A, d) is connected.

Examples:

- the real line with the Euclidean metric $(\mathbb{R}, d_{eucl})$ is connected
- the real line with the discrete metric $(\mathbb{R}, d_{disc})$ is connected

Connected spaces

Definition

A metric space (S, d) is called **disconnected** if there exists two nonempty, disjoint open subsets $U, V \subseteq S$ with $S = U \cup V$.

We call S **connected** if it is not disconnected.

A subset $A \subseteq S$ is connected if the corresponding subspace (A, d) is connected.

Examples:

- the real line with the Euclidean metric $(\mathbb{R}, d_{eucl})$ is connected
- the real line with the discrete metric $(\mathbb{R}, d_{disc})$ is connected
- euclidean space $\mathbb{R}^n$ is connected

Connected spaces

Definition

A metric space (S, d) is called **disconnected** if there exists two nonempty, disjoint open subsets $U, V \subseteq S$ with $S = U \cup V$.

We call S **connected** if it is not disconnected.

A subset $A \subseteq S$ is connected if the corresponding subspace (A, d) is connected.

Examples:

- the real line with the Euclidean metric $(\mathbb{R}, d_{eucl})$ is connected
- the real line with the discrete metric $(\mathbb{R}, d_{disc})$ is connected
- euclidean space $\mathbb{R}^n$ is connected
- an interval $I \subseteq \mathbb{R}$ with the Euclidean metric is connected

Intermediate value theorem

Theorem (Intermediate Value Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function with the Euclidean metric.

Then for any value y between $f(a)$ and $f(b)$, there exists $c \in [a, b]$ with $f(c) = y$.