

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

September 30, 2026

Outline

- 1 Real Analysis Lecture 11
 - Limits of sequences

Outline

- 1 Real Analysis Lecture 11
 - Limits of sequences

Warm-up

Definition

The **Oxford dictionary metric space** is the set M of all words in the Oxford English dictionary, where the metric is the discrete metric.

Example:

$$d(\text{fish}, \text{bird}) = 1, \quad d(\text{apple}, \text{apple}) = 0$$

Prove that for any word $w \in M$

$$d(\text{fire}, \text{ice}) \leq d(\text{fire}, w) + d(w, \text{ice})$$

Sequences

Definition

A **sequence** of real numbers is a function $x : \mathbb{N} \rightarrow \mathbb{R}$.

A **sequence** of points in $\mathbb{R}^n$ is a function $x : \mathbb{N} \rightarrow \mathbb{R}^n$

A **sequence** of points in a metric space (M, d) is a function $x : \mathbb{N} \rightarrow M$

Notation:

$$x_n = \text{formula}, \quad x(n) = \text{formula}, \quad x = (a, b, c, \dots)$$

Examples:

- $x_n = 1/n$ is a sequence in $\mathbb{R}$
- $x_n = (1/n, 1/n^2)$ is a sequence in $\mathbb{R}^2$
- $x = (\text{cat}, \text{dog}, \text{cat}, \text{dog}, \dots)$ is a sequence in the Oxford dictionary metric space

Definition of a limit

Definition of a limit

Back in Calc. I:

Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as n gets arbitrarily large, the value of x_n gets arbitrarily close to L .

Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as n gets arbitrarily large, the value of x_n gets arbitrarily close to L .

Formally:

Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as n gets arbitrarily large, the value of x_n gets arbitrarily close to L .

Formally:

$\forall \epsilon > 0$, there exists $N \in \mathbb{Z}_+$ such that $|x_n - L| < \epsilon \quad \forall n \geq N$.

Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as n gets arbitrarily large, the value of x_n gets arbitrarily close to L .

Formally:

$\forall \epsilon > 0$, there exists $N \in \mathbb{Z}_+$ such that $|x_n - L| < \epsilon \quad \forall n \geq N$.

Metric space version:

Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as n gets arbitrarily large, the value of x_n gets arbitrarily close to L .

Formally:

$\forall \epsilon > 0$, there exists $N \in \mathbb{Z}_+$ such that $|x_n - L| < \epsilon \quad \forall n \geq N$.

Metric space version:

$\forall \epsilon > 0$, there exists $N \in \mathbb{Z}_+$ such that $d(x_n, L) < \epsilon \quad \forall n \geq N$.

Definition of a limit

Back in Calc. I:

$$\lim_{n \rightarrow \infty} x_n = L \quad \text{means}$$

that as n gets arbitrarily large, the value of x_n gets arbitrarily close to L .

Formally:

$$\forall \epsilon > 0, \quad \text{there exists } N \in \mathbb{Z}_+ \text{ such that } |x_n - L| < \epsilon \quad \forall n \geq N.$$

Metric space version:

$$\forall \epsilon > 0, \quad \text{there exists } N \in \mathbb{Z}_+ \text{ such that } d(x_n, L) < \epsilon \quad \forall n \geq N.$$

We say $\{x_n\}$ **converges** to L and L is the **limit of the sequence**.

Example

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Proof.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Proof.

Let $\epsilon > 0$.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Then $n > N$ implies

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Then $n > N$ implies

$$d(x_n, 0) = |1/n - 0| = 1/n < 1/N < \epsilon.$$

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Then $n > N$ implies

$$d(x_n, 0) = |1/n - 0| = 1/n < 1/N < \epsilon.$$

Since $\epsilon > 0$ was arbitrary, this proves $\{x_n\}$ converges to 0.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ converges to zero.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Then $n > N$ implies

$$d(x_n, 0) = |1/n - 0| = 1/n < 1/N < \epsilon.$$

Since $\epsilon > 0$ was arbitrary, this proves $\{x_n\}$ converges to 0. $\square$

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = (n - 1)/n$ converges to 1.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = (n - 1)/n$ converges to 1.

Proof.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = (n - 1)/n$ converges to 1.

Proof.

Let $\epsilon > 0$.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = (n - 1)/n$ converges to 1.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = (n - 1)/n$ converges to 1.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Then $n > N$ implies

$$d(x_n, 1) = |(n - 1)/n - 1| = 1/n < 1/N < \epsilon.$$

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the Euclidean metric d .

The sequence $\{x_n\}$ defined by $x_n = (n - 1)/n$ converges to 1.

Proof.

Let $\epsilon > 0$.

Choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Then $n > N$ implies

$$d(x_n, 1) = |(n - 1)/n - 1| = 1/n < 1/N < \epsilon.$$

Since $\epsilon > 0$ was arbitrary, this proves $\{x_n\}$ converges to 1. $\square$

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ does not converge.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ does not converge.

Proof.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ does not converge.

Proof.

Suppose that it converges to a number L .

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ does not converge.

Proof.

Suppose that it converges to a number L .

Then for all $\epsilon > 0$, there must exist an $N \in \mathbb{Z}_+$ such that $d(x_n, L) < \epsilon$ for all $n \geq N$.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ does not converge.

Proof.

Suppose that it converges to a number L .

Then for all $\epsilon > 0$, there must exist an $N \in \mathbb{Z}_+$ such that $d(x_n, L) < \epsilon$ for all $n \geq N$.

Take $\epsilon = 1$ and choose $N \in \mathbb{Z}_+$ such that $d(x_n, L) < 1$ for all $n \geq N$.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ does not converge.

Proof.

Suppose that it converges to a number L .

Then for all $\epsilon > 0$, there must exist an $N \in \mathbb{Z}_+$ such that $d(x_n, L) < \epsilon$ for all $n \geq N$.

Take $\epsilon = 1$ and choose $N \in \mathbb{Z}_+$ such that $d(x_n, L) < 1$ for all $n \geq N$.

With the discrete metric $d(x_n, L) < 1$ implies $x_n = L$.

Example

Consider the metric space (M, d) defined by $M = \mathbb{R}$ with the **discrete** metric d .

The sequence $\{x_n\}$ defined by $x_n = 1/n$ does not converge.

Proof.

Suppose that it converges to a number L .

Then for all $\epsilon > 0$, there must exist an $N \in \mathbb{Z}_+$ such that $d(x_n, L) < \epsilon$ for all $n \geq N$.

Take $\epsilon = 1$ and choose $N \in \mathbb{Z}_+$ such that $d(x_n, L) < 1$ for all $n \geq N$.

With the discrete metric $d(x_n, L) < 1$ implies $x_n = L$.

This means $x_n = 1/n = L$ for all $n \geq N$, which is a contradiction!



Uniqueness of limits

Theorem (Uniqueness of limits)

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to T , then $L = T$.

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to T , then $L = T$.

Proof.

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to T , then $L = T$.

Proof.

Suppose that $\{x_n\}$ converges to L and T .

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to T , then $L = T$.

Proof.

Suppose that $\{x_n\}$ converges to L and T .

Assume further that $L \neq T$ and let $\epsilon = d(L, T)/2$.

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to T , then $L = T$.

Proof.

Suppose that $\{x_n\}$ converges to L and T .

Assume further that $L \neq T$ and let $\epsilon = d(L, T)/2$.

Then there exists N_1 such that $d(x_n, L) < \epsilon$ for $n \geq N_1$.

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to T , then $L = T$.

Proof.

Suppose that $\{x_n\}$ converges to L and T .

Assume further that $L \neq T$ and let $\epsilon = d(L, T)/2$.

Then there exists N_1 such that $d(x_n, L) < \epsilon$ for $n \geq N_1$.

Likewise, there exists N_2 such that $d(x_n, T) < \epsilon$ for $n \geq N_2$.

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to T , then $L = T$.

Proof.

Suppose that $\{x_n\}$ converges to L and T .

Assume further that $L \neq T$ and let $\epsilon = d(L, T)/2$.

Then there exists N_1 such that $d(x_n, L) < \epsilon$ for $n \geq N_1$.

Likewise, there exists N_2 such that $d(x_n, T) < \epsilon$ for $n \geq N_2$.

Then for $n > \max\{N_1, N_2\}$, the triangle inequality says

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to T , then $L = T$.

Proof.

Suppose that $\{x_n\}$ converges to L and T .

Assume further that $L \neq T$ and let $\epsilon = d(L, T)/2$.

Then there exists N_1 such that $d(x_n, L) < \epsilon$ for $n \geq N_1$.

Likewise, there exists N_2 such that $d(x_n, T) < \epsilon$ for $n \geq N_2$.

Then for $n > \max\{N_1, N_2\}$, the triangle inequality says

$$d(L, T) \leq d(L, x_n) + d(x_n, T) < 2\epsilon = d(L, T).$$

Uniqueness of limits

Theorem (Uniqueness of limits)

Let (M, d) be a metric space.

If a sequence $\{x_n\}$ in M converges to L and converges to T , then $L = T$.

Proof.

Suppose that $\{x_n\}$ converges to L and T .

Assume further that $L \neq T$ and let $\epsilon = d(L, T)/2$.

Then there exists N_1 such that $d(x_n, L) < \epsilon$ for $n \geq N_1$.

Likewise, there exists N_2 such that $d(x_n, T) < \epsilon$ for $n \geq N_2$.

Then for $n > \max\{N_1, N_2\}$, the triangle inequality says

$$d(L, T) \leq d(L, x_n) + d(x_n, T) < 2\epsilon = d(L, T).$$

This is a contradiction.



Existence of limits

Theorem

If a sequence $\{x_n\}$ in a metric space (M, d) converges to a value $L \in M$, then the range

$$X = \{x_1, x_2, \dots\} \subseteq M$$

is a bounded set and L is an adherent point of X .

Existence of limits

Theorem

If a sequence $\{x_n\}$ in a metric space (M, d) converges to a value $L \in M$, then the range

$$X = \{x_1, x_2, \dots\} \subseteq M$$

is a bounded set and L is an adherent point of X .

When the range of a sequence is bounded, we call the sequence $\{x_n\}$ bounded.

Existence of limits

Proof.

Suppose that $\lim_{n \rightarrow \infty} x_n = L$.

Existence of limits

Proof.

Suppose that $\lim_{n \rightarrow \infty} x_n = L$.

Let $\epsilon = 2024$.

Existence of limits

Proof.

Suppose that $\lim_{n \rightarrow \infty} x_n = L$.

Let $\epsilon = 2024$.

Then there exists $N \in \mathbb{Z}_+$ with $d(x_n, L) \leq 2024$ for all $n \geq N$.

Existence of limits

Proof.

Suppose that $\lim_{n \rightarrow \infty} x_n = L$.

Let $\epsilon = 2024$.

Then there exists $N \in \mathbb{Z}_+$ with $d(x_n, L) \leq 2024$ for all $n \geq N$.

If we take

$$R = \max\{d(x_1, L), d(x_2, L), \dots, d(x_{N-1}, L), 2024\},$$

Existence of limits

Proof.

Suppose that $\lim_{n \rightarrow \infty} x_n = L$.

Let $\epsilon = 2024$.

Then there exists $N \in \mathbb{Z}_+$ with $d(x_n, L) \leq 2024$ for all $n \geq N$.

If we take

$$R = \max\{d(x_1, L), d(x_2, L), \dots, d(x_{N-1}, L), 2024\},$$

then $d(x_n, L) \leq R$ for all N .

Existence of limits

Proof.

Suppose that $\lim_{n \rightarrow \infty} x_n = L$.

Let $\epsilon = 2024$.

Then there exists $N \in \mathbb{Z}_+$ with $d(x_n, L) \leq 2024$ for all $n \geq N$.

If we take

$$R = \max\{d(x_1, L), d(x_2, L), \dots, d(x_{N-1}, L), 2024\},$$

then $d(x_n, L) \leq R$ for all N .

Therefore $X \subseteq B_M(L, R)$ and in particular X is bounded. □

Existence of limits

Sort of a converse of the previous theorem.

Existence of limits

Sort of a converse of the previous theorem.

Theorem

If S is a subset of a metric space (M, d) and $L \in M$ is an adherent point of S , then there exists a sequence $\{x_n\}$ of elements of S which converges to L .

Existence of limits

Proof.

Suppose L is an adherent point of S .

Existence of limits

Proof.

Suppose L is an adherent point of S .

Then for all $n \in \mathbb{Z}_+$, the ball $B_M(x, \frac{1}{n})$ contains at least one element of S .

Existence of limits

Proof.

Suppose L is an adherent point of S .

Then for all $n \in \mathbb{Z}_+$, the ball $B_M(x, \frac{1}{n})$ contains at least one element of S .

For each $n \in \mathbb{Z}_+$, choose $x_n \in S \cap B_M(x, \frac{1}{n})$.

Existence of limits

Proof.

Suppose L is an adherent point of S .

Then for all $n \in \mathbb{Z}_+$, the ball $B_M(x, \frac{1}{n})$ contains at least one element of S .

For each $n \in \mathbb{Z}_+$, choose $x_n \in S \cap B_M(x, \frac{1}{n})$.

We claim $\lim x_n = L$.

Existence of limits

Proof.

Suppose L is an adherent point of S .

Then for all $n \in \mathbb{Z}_+$, the ball $B_M(x, \frac{1}{n})$ contains at least one element of S .

For each $n \in \mathbb{Z}_+$, choose $x_n \in S \cap B_M(x, \frac{1}{n})$.

We claim $\lim x_n = L$.

To see this, let $\epsilon > 0$.

Existence of limits

Proof.

Suppose L is an adherent point of S .

Then for all $n \in \mathbb{Z}_+$, the ball $B_M(x, \frac{1}{n})$ contains at least one element of S .

For each $n \in \mathbb{Z}_+$, choose $x_n \in S \cap B_M(x, \frac{1}{n})$.

We claim $\lim x_n = L$.

To see this, let $\epsilon > 0$.

Then choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

Existence of limits

Proof.

Suppose L is an adherent point of S .

Then for all $n \in \mathbb{Z}_+$, the ball $B_M(x, \frac{1}{n})$ contains at least one element of S .

For each $n \in \mathbb{Z}_+$, choose $x_n \in S \cap B_M(x, \frac{1}{n})$.

We claim $\lim x_n = L$.

To see this, let $\epsilon > 0$.

Then choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

It follows that for all $n \geq N$

$$d(x_n, L) < 1/n < 1/N < \epsilon.$$

Existence of limits

Proof.

Suppose L is an adherent point of S .

Then for all $n \in \mathbb{Z}_+$, the ball $B_M(x, \frac{1}{n})$ contains at least one element of S .

For each $n \in \mathbb{Z}_+$, choose $x_n \in S \cap B_M(x, \frac{1}{n})$.

We claim $\lim x_n = L$.

To see this, let $\epsilon > 0$.

Then choose $N \in \mathbb{Z}_+$ with $N > 1/\epsilon$.

It follows that for all $n \geq N$

$$d(x_n, L) < 1/n < 1/N < \epsilon.$$

Since $\epsilon > 0$ was arbitrary, this proves convergence. □