

MATH 350-2 Advanced Calculus

W.R. Casper

Department of Mathematics
California State University Fullerton

September 2, 2026

Outline

- 1 Real Analysis Lecture 3
 - Completeness Axiom (continued)
 - Set basics
 - Set algebra

Outline

1

Real Analysis Lecture 3

- Completeness Axiom (continued)
- Set basics
- Set algebra

Completeness Axiom

Completeness Axiom

Definition

An **upper bound** of a set A is a number b with $x \leq b$ for all $x \in A$. If A has an upper bound, it is called **bounded above**.

Completeness Axiom

Definition

An **upper bound** of a set A is a number b with $x \leq b$ for all $x \in A$. If A has an upper bound, it is called **bounded above**.

Definition

A **supremum** of a set A is a number s with two properties:

Completeness Axiom

Definition

An **upper bound** of a set A is a number b with $x \leq b$ for all $x \in A$. If A has an upper bound, it is called **bounded above**.

Definition

A **supremum** of a set A is a number s with two properties:

- s is an upper bound of A , and

Completeness Axiom

Definition

An **upper bound** of a set A is a number b with $x \leq b$ for all $x \in A$. If A has an upper bound, it is called **bounded above**.

Definition

A **supremum** of a set A is a number s with two properties:

- s is an upper bound of A , and
- if $t < s$ then t is NOT an upper bound of A

Completeness Axiom

Definition

An **upper bound** of a set A is a number b with $x \leq b$ for all $x \in A$. If A has an upper bound, it is called **bounded above**.

Definition

A **supremum** of a set A is a number s with two properties:

- s is an upper bound of A , and
- if $t < s$ then t is NOT an upper bound of A

Completeness Axiom: every nonempty set of real numbers which is bounded above has a supremum.

Decimals

The completeness axiom allows us to define **decimals**.

Decimals

The completeness axiom allows us to define **decimals**. Let $0 \leq n$ be an integer and $d_1, d_2, d_3, \dots$ be a sequence of integers with $0 \leq d_k \leq 9$ for all $k \in \mathbb{N}$.

Decimals

The completeness axiom allows us to define **decimals**. Let $0 \leq n$ be an integer and $d_1, d_2, d_3, \dots$ be a sequence of integers with $0 \leq d_k \leq 9$ for all $k \in \mathbb{N}$.

Definition

The k 'th **decimal approximation** of $n.d_1d_2d_3\dots$ is

$$s_k = n + \frac{d_1}{10} + \frac{d_2}{10^2} + \frac{d_3}{10^3} + \dots + \frac{d_k}{10^k}.$$

Decimals

The completeness axiom allows us to define **decimals**. Let $0 \leq n$ be an integer and $d_1, d_2, d_3, \dots$ be a sequence of integers with $0 \leq d_k \leq 9$ for all $k \in \mathbb{N}$.

Definition

The k 'th **decimal approximation** of $n.d_0d_1d_2d_3\dots$ is

$$s_k = n + \frac{d_1}{10} + \frac{d_2}{10^2} + \frac{d_3}{10^3} + \dots + \frac{d_k}{10^k}.$$

Definition

The **decimal** $n.d_0d_1d_2d_3\dots$ is the real number defined as the supremum of decimal approximations

$$n.d_0d_1d_2d_3\dots = \sup\{s_k : k \in \mathbb{N}\}.$$

Examples of decimals

- Liouville's constant

0.11000100000000000000000010...

Examples of decimals

- Liouville's constant

0.1100010000000000000000010...

- π

3.1415926535...

Examples of decimals

- Liouville's constant

0.1100010000000000000000010...

- π

3.1415926535...

- 1

0.9999999999...

Examples of decimals

- Liouville's constant

0.1100010000000000000000010...

- π

3.1415926535...

- 1

0.9999999999...

- 1

1.0000000000...

Examples of decimals

- Liouville's constant

0.11000100000000000000000010...

- π

3.1415926535...

- 1

0.9999999999...

- 1

1.0000000000...

- later we will show *any* positive number has at least one decimal expansion

Decimal expansions of rationals

Theorem

A decimal expansion

$$n.d_0d_1d_2d_3\dots$$

is rational if and only if it is eventually periodic, ie. there exists N and p such that $d_{k+p} = d_p$ for all $k \geq N$.

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

If $b \in S$ also, then b is called a **minimal element** of S .

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

If $b \in S$ also, then b is called a **minimal element** of S . An **infimum** of a set S of real numbers is a real number $b \in \mathbb{R}$ such that

- b is a lower bound of S
- if $b < b'$, then b' is not a lower bound of S

Lower bounds and infima

A **lower bound** for a set $S \subseteq \mathbb{R}$ is a number b such that

$$b \leq x \quad \text{for all } x \in S.$$

In this case, we say S is **bounded below** by b .

If $b \in S$ also, then b is called a **minimal element** of S . An **infimum** of a set S of real numbers is a real number $b \in \mathbb{R}$ such that

- b is a lower bound of S
- if $b < b'$, then b' is not a lower bound of S

In other words

an infimum is a **greatest lower bound**

Properties of Suprema

The first property of suprema is that they must be arbitrarily close to elements of the set.

Theorem (Approximation Property)

Let $S \subseteq \mathbb{R}$ be nonempty and bounded above, and let $b = \sup(S)$. Then for all $\epsilon > 0$, there exists $a \in S$ with

$$b - \epsilon < a \leq b.$$

Properties of Suprema

Proof.

Since $b - \epsilon < b$, the definition of a supremum implies $b - \epsilon$ cannot be an upper bound.

Properties of Suprema

Proof.

Since $b - \epsilon < b$, the definition of a supremum implies $b - \epsilon$ cannot be an upper bound.

Therefore there must exist $a \in S$ with $a > b - \epsilon$. □

Properties of Suprema

Also, suprema play nicely with addition.

Theorem (Additive Property)

Let $A, B \subseteq \mathbb{R}$ be nonempty, bounded above sets.

$$C = \{x + y : x \in A, y \in B\}.$$

Then C is bounded above and

$$\sup(C) = \sup(A) + \sup(B).$$

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of b , so $y \leq b$ for all $y \in B$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

It follows that $y \leq c - a$ for all $y \in B$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of B , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \in B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

It follows that $y \leq c - a$ for all $y \in B$.

Thus $c - a$ is an upper bound of B , and it follows $b \leq c - a$.

Properties of Suprema

Proof.

Let $a = \sup(A)$, $b = \sup(B)$, and $c = \sup(C)$.

We will prove $c \leq a + b$ and then $a + b \leq c$.

First, note a is an upper bound of A , so $x \leq a$ for all $x \in A$.

Also b is an upper bound of b , so $y \leq b$ for all $y \in B$.

If $z \in C$, then $z = x + y$ for some $x \in A$ and $y \in B$, and therefore $z = x + y \leq a + b$.

Therefore $a + b$ is an upper bound of C .

This means $c \leq a + b$.

Next, note for all $x \in A$ and $y \leq B$ that $x \leq c - y$.

Therefore $c - y$ is an upper bound of A and $a \leq c - y$.

It follows that $y \leq c - a$ for all $y \in B$.

Thus $c - a$ is an upper bound of B , and it follows $b \leq c - a$.

Therefore $a + b \leq c$. □

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{N}$ is not bounded above. (Apostol Theorem 1.17)

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{N}$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{N}) - 1$

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{N}$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{N}) - 1$

Solution

Assume it is.

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{N}$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{N}) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{N})$ exists

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{N}$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{N}) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{N})$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{N}$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{N}) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{N})$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{N}$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{N}) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{N})$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

This means there exists $n \in \mathbb{N}$ with $b - 1 < n$.

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{N}$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{N}) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{N})$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

This means there exists $n \in \mathbb{N}$ with $b - 1 < n$.

It follows from Axiom 7 that $b < n + 1$.

Challenge

Problem

Use the completeness axiom to prove that the $\mathbb{N}$ is not bounded above. (Apostol Theorem 1.17)

Hint: assume it is and consider $\sup(\mathbb{N}) - 1$

Solution

Assume it is.

The completeness axiom implies that $b = \sup(\mathbb{N})$ exists

The number b is an upper bound and if $b' < b$, then b' is not.

Then $b - 1 < b$ by Axiom 7, so $b - 1$ cannot be an upper bound.

This means there exists $n \in \mathbb{N}$ with $b - 1 < n$.

It follows from Axiom 7 that $b < n + 1$.

However, $n + 1 \in \mathbb{Z}$, so this contradicts b being an upper bound.

Archimedean property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{N}$ with $n > x$.

Archimedean property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{N}$ with $n > x$.

Proof.

If not, then x is an upper bound of $\mathbb{N}$. □

Archimedean property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{N}$ with $n > x$.

Proof.

If not, then x is an upper bound of $\mathbb{N}$. □

Theorem (Archimedean Property of Reals)

For every $x, y \in \mathbb{R}$ with $x > 0$, there exists $n \in \mathbb{N}$ with $y < nx$.

Archimedean property

Theorem (Apostol Theorem 1.18)

For every $x \in \mathbb{R}$, there exists $n \in \mathbb{N}$ with $n > x$.

Proof.

If not, then x is an upper bound of $\mathbb{N}$. □

Theorem (Archimedean Property of Reals)

For every $x, y \in \mathbb{R}$ with $x > 0$, there exists $n \in \mathbb{N}$ with $y < nx$.

Proof.

Replace x with y/x in the previous theorem. □

Outline

1

Real Analysis Lecture 3

- Completeness Axiom (continued)
- Set basics
- Set algebra

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{N}, \mathbb{Z}, \mathbb{Q}$

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{N}, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{N}, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$
- empty set $\emptyset$

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{N}, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$
- empty set $\emptyset$
- $\{\heartsuit, \text{Fall}, \{\emptyset\}\}$

Set basics

Intuitively, a **set** A is a "collection of things" which we call the **elements** of A .

- in practice, this is a bad definition (Russell's Paradox)
- true set formulation: Zermelo-Frankel Axioms

Examples:

- $\mathbb{R}, \mathbb{N}, \mathbb{Z}, \mathbb{Q}$
- $(1, 5], (0, \infty)$
- empty set $\emptyset$
- $\{\heartsuit, \text{Fall}, \{\emptyset\}\}$
- $\{n \in \mathbb{Z} : n \text{ is prime}\}$

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

- ordered pairs

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

- ordered pairs

$$(a, b) = \{a, \{a, b\}\}$$

Everything is a set

In the true minimalistic philosophy of mathematics, we want everything to be a set.

- integers

$$0 = \emptyset, 1 = \{\emptyset\}, 2 = \{\emptyset, \{\emptyset\}\}, \dots$$

- ordered pairs

$$(a, b) = \{a, \{a, b\}\}$$

- even relations and functions are sets!

Outline

1

Real Analysis Lecture 3

- Completeness Axiom (continued)
- Set basics
- Set algebra

Unions of sets

Unions:

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

$$A_1 \cup A_2 \cup \cdots \cup A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

Unions of sets

Unions:

$$A \cup B = \{x : x \in A \text{ or } x \in B\},$$

$$A \cup B \cup C = \{x : x \in A \text{ or } x \in B \text{ or } x \in C\},$$

$$A_1 \cup A_2 \cup \cdots \cup A_n = \{x : x \in A_i \text{ for some } 1 \leq i \leq n\}$$

NOTATION:

$$\bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \cdots \cup A_n.$$

Intersections of sets

Intersections:

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

$$A_1 \cap A_2 \cap \cdots \cap A_n = \{x : x \in A_i \text{ for all } 1 \leq i \leq n\}$$

Intersections of sets

Intersections:

$$A \cap B = \{x : x \in A \text{ and } x \in B\},$$

$$A \cap B \cap C = \{x : x \in A \text{ and } x \in B \text{ and } x \in C\},$$

$$A_1 \cap A_2 \cap \cdots \cap A_n = \{x : x \in A_i \text{ for all } 1 \leq i \leq n\}$$

NOTATION:

$$\bigcap_{i=1}^n A_i = A_1 \cap A_2 \cap \cdots \cap A_n.$$

Complements of sets

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Theorem

De Morgan's Laws

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Theorem

De Morgan's Laws

$$B - \bigcup_{i=1}^n A_i = \bigcap_{i=1}^n (B - A_i)$$

Complements of sets

The **complement** of A relative to B is

$$B - A = \{b \in B : b \notin A\}.$$

Theorem

De Morgan's Laws

$$B - \bigcup_{i=1}^n A_i = \bigcap_{i=1}^n (B - A_i)$$

$$B - \bigcap_{i=1}^n A_i = \bigcup_{i=1}^n (B - A_i)$$

Families of sets

A **family of sets** is a collection

$$\{A_i : i \in I\}$$

Families of sets

A **family of sets** is a collection

$$\{A_i : i \in I\}$$

- I any set, called the **index set**

Families of sets

A **family of sets** is a collection

$$\{A_i : i \in I\}$$

- I any set, called the **index set**
- A_i is a set for all i

Families of sets

A **family of sets** is a collection

$$\{A_i : i \in I\}$$

- I any set, called the **index set**
- A_i is a set for all i

$$\bigcup_{i \in I} A_i = \{x : x \in A_i, \text{ for some } i \in I\}$$

Families of sets

A **family of sets** is a collection

$$\{A_i : i \in I\}$$

- I any set, called the **index set**
- A_i is a set for all i

$$\bigcup_{i \in I} A_i = \{x : x \in A_i, \text{ for some } i \in I\}$$

$$\bigcap_{i \in I} A_i = \{x : x \in A_i, \text{ for all } i \in I\}$$

Challenge

- index set $I = (1, 2)$
- family of sets $\{A_i : i \in I\}$
- $A_i = [0, i]$

Problem

Determine $\bigcup_{i \in I} A_i$.

Challenge

- index set $I = \mathbb{N}$
- family of sets $\{A_i : i \in I\}$
- $A_i = [0, 1/i)$

Problem

Determine $\bigcap_{i \in I} A_i$.

Challenge

Problem

Prove that De Morgan's Laws apply to families of sets too, ie.

$$B - \bigcup_{i \in I} A_i = \bigcap_{i \in I} (B - A_i)$$

Challenge

Problem

Prove that De Morgan's Laws apply to families of sets too, ie.

$$B - \bigcup_{i \in I} A_i = \bigcap_{i \in I} (B - A_i)$$

$$B - \bigcap_{i \in I} A_i = \bigcup_{i \in I} (B - A_i)$$

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** $\mathcal{R}$ from A to B is a subset of $A \times B$.

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** $\mathcal{R}$ from A to B is a subset of $A \times B$.

NOTATION: $a\mathcal{R}b$ means $(a, b) \in \mathcal{R}$.

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** $\mathcal{R}$ from A to B is a subset of $A \times B$.

NOTATION: $a\mathcal{R}b$ means $(a, b) \in \mathcal{R}$.

Domain and codomain:

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** $\mathcal{R}$ from A to B is a subset of $A \times B$.

NOTATION: $a\mathcal{R}b$ means $(a, b) \in \mathcal{R}$.

Domain and codomain:

$$\text{dom}(\mathcal{R}) = \{a \in A : \exists b \in B, a\mathcal{R}b\}$$

$$\text{codom}(\mathcal{R}) = \{b \in B : \exists a \in A, a\mathcal{R}b\}$$

Relations

The **Cartesian product** of A and B is

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

A **relation** $\mathcal{R}$ from A to B is a subset of $A \times B$.

NOTATION: $a\mathcal{R}b$ means $(a, b) \in \mathcal{R}$.

Domain and codomain:

$$\begin{aligned}\text{dom}(\mathcal{R}) &= \{a \in A : \exists b \in B, a\mathcal{R}b\} \\ \text{codom}(\mathcal{R}) &= \{b \in B : \exists a \in A, a\mathcal{R}b\}\end{aligned}$$

Functions

A **function** from A to B is a relation $\mathcal{R}$ from A to B with the property

Functions

A **function** from A to B is a relation $\mathcal{R}$ from A to B with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \quad \text{implies} \quad b = c.$$

Functions

A **function** from A to B is a relation $\mathcal{R}$ from A to B with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \quad \text{implies} \quad b = c.$$

If a relation $\mathcal{R}$ is a function, we usually use a symbol like f .

Functions

A **function** from A to B is a relation $\mathcal{R}$ from A to B with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

If a relation $\mathcal{R}$ is a function, we usually use a symbol like f .

NOTATION: $f : A \rightarrow B$ means f is a function from A to B

NOTATION: $f(a) = b$ means $(a, b) \in f$.

Functions

A **function** from A to B is a relation $\mathcal{R}$ from A to B with the property

$$a\mathcal{R}b \text{ and } a\mathcal{R}c \text{ implies } b = c.$$

If a relation $\mathcal{R}$ is a function, we usually use a symbol like f .

NOTATION: $f : A \rightarrow B$ means f is a function from A to B

NOTATION: $f(a) = b$ means $(a, b) \in f$.

The set

$$\text{img}(f) = \{f(a) : a \in A\}$$

is called the **range** or **image** of f