

ABSTRACT

DAWN, WILLIAM C. Sodium Cooled Fast Reactor Simulations with the Finite Element Method. (Under the direction of Scott P. Palmtag and David J. Kropaczek.)

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis.

Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

© Copyright 2018 by William C. Dawn

All Rights Reserved

Sodium Cooled Fast Reactor Simulations with the Finite Element Method

by
William C. Dawn

A thesis submitted to the Graduate Faculty of
North Carolina State University
in partial fulfillment of the
requirements for the Degree of
Master of Science

Nuclear Engineering

Raleigh, North Carolina

2018

APPROVED BY:

Joseph M. Doster

Zhilin Li

Scott P. Palmtag
Co-chair of Advisory Committee

David J. Kropaczek
Co-chair of Advisory Committee

DEDICATION

To the future of clean energy.

BIOGRAPHY

The author was born in a small town ...

ACKNOWLEDGEMENTS

I would like to thank my advisor for his help.

TABLE OF CONTENTS

LIST OF TABLES	vii
LIST OF FIGURES	viii
Chapter 1 INTRODUCTION	1
1.1 Math environments	1
1.1.1 Equations	1
Chapter 2 FINITE ELEMENT NEUTRON DIFFUSION	4
2.1 Introduction	4
2.2 Formulation	6
2.2.1 Derivation	6
2.2.2 Matrix Quantities	10
2.2.3 Quadratures	11
2.3 Implementation	13
2.3.1 Algorithm	13
2.3.2 Memory and Storage	14
2.3.3 Linear System Solution	14
2.4 Reference Results	14
2.4.1 Triangular Element Manufactured Solutions	14
2.4.2 Two Dimension Reactors	14
2.4.3 Wedge Element Manufactured Solution Finite Cylinder	14
2.4.4 Three Dimension Reactors	14
Chapter 3 THERMAL HYDRAULICS	15
3.1 Axial Convection Model	15
3.2 Radial Conduction Model	15
3.2.1 Derivation	15
3.2.2 Relations	15
3.3 Cross Section Treatment	15
3.4 Results	15
Chapter 4 THERMAL EXPANSION	16
4.1 Necessity of Modeling	16
4.2 Model Details	16
4.3 Implementation	16
4.4 Results	16
Chapter 5 CONCLUSIONS	17
5.1 Results Discussion	17
5.2 Code Enhancements	17
5.3 Further Investigations	17

Bibliography	18
APPENDIX	19
Appendix A LOREM IPSUM	20
A.1 A First Section	20
A.2 A Second Section	22

LIST OF TABLES

Table 2.1	Jacobian Determinants	13
Table A.1	A table in the appendix.	22

LIST OF FIGURES

CHAPTER 1

INTRODUCTION

Let's start with a few paragraph basics, here is how to make **bold**, and *italics*, and *emphasized*. Let's say you need to cite something in your references, simply type `\cite{key}`, which produces [ER35]. Some other references are [GVL96] and [LK74]. Some \LaTeX compilers require a second compilation for citations and references to be sorted and matched properly in the resulting document.

Here is a quotation:

Alice, Bob and Carol are boring. Who would even want to know their secret?

Let's say we need to make a list, try this on for size

1. NCSU is great
2. I like NCSU
3. I really hope I can find a job when I graduate!

1.1 Math environments

1.1.1 Equations

There are many different ways to write equations, for example we could put $a^2 + b^2 = c^2$ directly into a sentence. Or we could use the equation environment and do

$$a^2 + b^2 = c^2. \tag{1.1}$$

And from here we can later reference it by simply doing typing `\ref{label}`, which gives 1.1. However, defining and using equation and figure reference macros will ensure that the equation references are consistent, instead of having Eq. (1), Equation 3, Eqn 4 scattered through the thesis. This template file defines `\eref` and `\fref` for this purpose. You can modify the macros to your liking in the `YourName-thesis.tex` file. For example, the command `\eref{label}` gives Eq. 1.1.

If you don't need to reference an equation you may simply do this

$$a^2 + b^2 = c^2.$$

For Greek letters you must go to the math environments, for example α , β , and γ . Let's look at equations that cover multiple lines, none of these equations may be true or mean anything, but so that the reader can get some ideas. In addition I will use some other useful notations like subscripts, superscripts, fractions, etc. One important item of note is that one uses the "ampersand" symbol to line up equations (also look at how I used quotations).

$$\gamma_1 = \alpha^\beta + \psi_0 \frac{\psi_1}{\psi_2 + \psi_3} \tag{1.2}$$

$$= \beta_1 + \beta_2 + \dots + \beta_k$$

$$\rightarrow E(\gamma_2) \tag{1.3}$$

Alternatively, one can specify a slightly different environment if none of the equations need to be numbered. Remember that if you are planning on referring to them later on, you must use a "label" statement.

$$\gamma_1 = n^{-1/2} \sum_{i=1}^n [h(X_i, \beta_0) - E\{h(X_i, \beta_0)\}]$$

$$\rightarrow \hat{q} \pm \frac{\partial \gamma_2}{\partial \beta}.$$

Lastly there may be times in which you want to use a non-italicized word your formula, such as an indicator function that may look like this $I\{\mu_i(1, \beta) > \mu_i(0, \beta)\}$, if so just use the "mbox" statement.

You could use a multiline equation for long equations. The environment is `multline`. Insert `\\` for line breaks.

$$\vec{\cdot} \cdot \vec{\nabla} \psi(\vec{r}, \cdot, E) + \Sigma_t(\vec{r}, E) \psi(\vec{r}, \cdot, E) = \int_{4\pi} d\vec{\cdot}' \int_0^\infty dE' \Sigma_s(\vec{r}, \cdot' \rightarrow \cdot, E' \rightarrow E) \psi(\vec{r}, \cdot', E') + Q(\vec{r}, \cdot, E),$$

we operate with $\int_0^\infty (\cdot) dE$ to obtain

$$\vec{\cdot} \cdot \vec{\nabla} \tilde{\psi}(\vec{r}, \cdot) + \Sigma_t(\vec{r}) \tilde{\psi}(\vec{r}, \cdot) = \int_{4\pi} d\vec{\cdot}' \int_0^\infty dE' \psi(\vec{r}, \cdot', E') \left[\int_0^\infty dE \Sigma_s(\vec{r}, \cdot' \rightarrow \cdot, E' \rightarrow E) \right] + \tilde{Q}(\vec{r}, \cdot),$$

CHAPTER 2

FINITE ELEMENT NEUTRON DIFFUSION

2.1 Introduction

For typical nuclear reactor applications, diffusion theory well approximates the neutron distribution within the reactor. The neutron diffusion equation is a second order partial differential equation in space and energy. In standard notation, the continuous neutron diffusion equation is presented as

$$-\nabla \cdot (D(\mathbf{r}, E) \nabla \phi(\mathbf{r}, E)) + \Sigma_t(\mathbf{r}, E) \phi(\mathbf{r}, E) = \frac{\chi(\mathbf{r}, E)}{k_{eff}} \int_0^\infty \nu_f(\mathbf{r}, E') \Sigma_f(\mathbf{r}, E') \phi(\mathbf{r}, E') dE' + \int_0^\infty \Sigma_s(\mathbf{r}, E' \rightarrow E) \phi(\mathbf{r}, E') dE' \quad (2.1)$$

Where D is the diffusion constant, ϕ is the scalar neutron flux, Σ_t is the total cross section, χ is the fission neutron spectrum, k_{eff} is the effective neutron multiplication factor, Σ_f is the fission cross section, ν_f is the neutron yield per fission, and $\Sigma_s(\mathbf{r}, E' \rightarrow E)$ is the scattering cross section for neutrons at position \mathbf{r} scattering from energy E' to E .

The neutron diffusion equation must be discretized in space and energy to be solved numerically. Energy discretization is relatively straight-forward and is performed using the multigroup method. Spatial discretization requires more attention and will be done with the Finite Element Method (FEM). This method is selected for several reasons. It allows for easily increasing the order of the method by increasing the order of the elements with no changing of the mesh required. Coordinates of nodes can be easily updated to reflect physical phenomena such as thermal expansion (Chapter

4). Additionally, material properties are calculated on an element basis allowing for fine detail updates to the material properties during the calculation.

For energy discretization, An energy structure is described as $\{E_g\}$ for $g = 1, 2, \dots, G$ and by convention

$$E_G > E_{G-1} > \dots > E_2 > E_1$$

Then, multigroup constants can be calculated based on the energy group structure and the known cross sections. Multigroup constants are calculated to preserve the number of neutrons produced or destroyed. That is, the calculation preserves reaction rates where the reaction rate for reaction x is defined as $R_x = \Sigma_x \phi$. Multigroup constants are then calculated. A formal derivation is given in [DH76] and the results are presented below.

$$D_g(\mathbf{r}) = \frac{\int_{E_g}^{E_{g-1}} D(\mathbf{r}, E') \nabla \phi(\mathbf{r}, E) dE}{\int_{E_g}^{E_{g-1}} \nabla \phi(\mathbf{r}, E) dE} \quad (2.2)$$

$$\Sigma_{t,g}(\mathbf{r}) = \frac{\int_{E_g}^{E_{g-1}} \Sigma_t(\mathbf{r}, E) \phi(\mathbf{r}, E) dE}{\int_{E_g}^{E_{g-1}} \phi(\mathbf{r}, E) dE} \quad (2.3)$$

$$\nu \Sigma_{f,g}(\mathbf{r}) = \frac{\int_{E_g}^{E_{g-1}} \nu_f(\mathbf{r}, E) \Sigma_f(\mathbf{r}, E) \phi(\mathbf{r}, E) dE}{\int_{E_g}^{E_{g-1}} \phi(\mathbf{r}, E) dE} \quad (2.4)$$

$$\Sigma_{s,g \rightarrow g'}(\mathbf{r}) = \frac{\int_{E_g'}^{E_{g'-1}} \int_{E_g}^{E_{g-1}} \Sigma_s(\mathbf{r}, E' \rightarrow E) \phi(\mathbf{r}, E') dE dE'}{\int_{E_g}^{E_{g-1}} \phi(\mathbf{r}, E) dE} \quad (2.5)$$

$$\chi_g(\mathbf{r}) = \int_{E_g}^{E_{g-1}} \chi(\mathbf{r}, E) dE \quad (2.6)$$

$$\phi_g(\mathbf{r}) = \int_{E_g}^{E_{g-1}} \phi(\mathbf{r}, E) dE \quad (2.7)$$

Note that cross sections $\nu_f(\mathbf{r}, E)$ and $\Sigma_f(\mathbf{r}, E)$ have been combined. This is necessary and a mathematical result of the group collapse. Then, Eq. 2.1 can be discretized in energy as

$$-\nabla \cdot (D_g(\mathbf{r}) \nabla \phi_g(\mathbf{r})) + \Sigma_{t,g}(\mathbf{r}) \phi_g(\mathbf{r}) = \frac{\chi_g(\mathbf{r})}{k_{eff}} \sum_{g'=1}^G \nu \Sigma_{f,g'}(\mathbf{r}) \phi_{g'}(\mathbf{r}) + \sum_{g'=1}^G \Sigma_{s,g' \rightarrow g}(\mathbf{r}) \phi_{g'}(\mathbf{r}) \quad (2.8)$$

The neutron diffusion equation has now been discretized in energy. Spatial discretization will be based on the Finite Element Method (FEM) and will be discussed in Sec. 2.2.1.

The total neutron cross section includes the contribution due to self-scattering. That is, due to $\Sigma_{s,g \rightarrow g}$. This can be removed from Eq. 2.8 for simplicity.

$$-\nabla \cdot (D_g(\mathbf{r}) \nabla \phi_g(\mathbf{r})) + \Sigma_{r,g}(\mathbf{r}) \phi_g(\mathbf{r}) = \frac{\chi_g(\mathbf{r})}{k_{eff}} \sum_{g'=1}^G \nu \Sigma_{f,g'}(\mathbf{r}) \phi_{g'}(\mathbf{r}) + \sum_{g'=1, g' \neq g}^G \Sigma_{s,g' \rightarrow g}(\mathbf{r}) \phi_{g'}(\mathbf{r}) \quad (2.9)$$

Where $\Sigma_{r,g}$ is the removal cross section and $\Sigma_{r,g}(\mathbf{r}) = \Sigma_{t,g}(\mathbf{r}) - \Sigma_{s,g \rightarrow g}(\mathbf{r})$. For simplicity, the neutron sources in Eq. 2.9 can be combined into a single term.

$$-\nabla \cdot (D_g(\mathbf{r}) \nabla \phi_g(\mathbf{r})) + \Sigma_{r,g}(\mathbf{r}) \phi_g(\mathbf{r}) = q_g(\mathbf{r}) \quad (2.10)$$

Where $q_g(\mathbf{r})$ is the combined neutron source at position \mathbf{r} . q can then be further divided into contributions due to fission (q_{fiss}), up-scattering (q_{up}) when a neutron increases in energy, and down-scattering when a neutron decreases in energy (q_{down}).

$$q_g(\mathbf{r}) = q_{fiss,g}(\mathbf{r}) + q_{up,g}(\mathbf{r}) + q_{down,g}(\mathbf{r}) \quad (2.11)$$

$$q_{fiss,g}(\mathbf{r}) = \frac{\chi_g(\mathbf{r})}{k_{eff}} \sum_{g'=1}^G \nu \Sigma_{f,g'}(\mathbf{r}) \phi_{g'}(\mathbf{r}) \quad (2.12)$$

$$q_{up,g}(\mathbf{r}) = \sum_{g'=g+1}^G \Sigma_{s,g' \rightarrow g}(\mathbf{r}) \phi_{g'}(\mathbf{r}) \quad (2.13)$$

$$q_{down,g}(\mathbf{r}) = \sum_{g'=1}^g \Sigma_{s,g' \rightarrow g}(\mathbf{r}) \phi_{g'}(\mathbf{r}) \quad (2.14)$$

Where the difference between q_{up} and q_{down} are the limits of the summation. This form allows for operator splitting of the neutron source term. In an iterative scheme, it will be necessary for fission and up-scatter sources to use a different flux iterate than down-scatter so this division will prove useful.

2.2 Formulation

2.2.1 Derivation

The only remaining continuous variable in the problem is the spatial variable \mathbf{r} . This will be discretized according to the Finite Element Method (FEM). The form of the diffusion equation to be discretized is Eq. 2.10. The problem is solved in a finite domain $\mathbf{r} \in \Omega$ where $\partial\Omega$ represents the boundary of the domain where some boundary condition is specified. Boundary condition options

provided include

1. Mirror. $\nabla \phi_g(\mathbf{r}) = 0$ for $\mathbf{r} \in \partial\Omega$.
2. Albedo. $D_g(\mathbf{r})\nabla \phi_g(\mathbf{r}) + \aleph \phi_g(\mathbf{r}) = 0$ for $\mathbf{r} \in \partial\Omega$ where \aleph is a real constant specified by the user. For vacuum conditions, $\alpha = \frac{1}{2}$.
3. Zero Flux. $\phi_g(\mathbf{r}) = 0$ for $\mathbf{r} \in \partial\Omega$.

(Note: the order of the above list corresponds to the order of boundary condition precedent in code with the greater the integer, the greater the precedent.)

Finite Element derivation begins with Eq. 2.10. The equation is multiplied by a testing function $v(\mathbf{r}) \in H_1(\Omega)$ Where H is the Sobolev Space. Then, the equation is integrated over the problem domain. This yields the Weak Form or Variational Form of the problem.

$$-\nabla \cdot (D_g(\mathbf{r})\nabla \phi_g(\mathbf{r})) + \Sigma_{r,g}(\mathbf{r})\phi_g(\mathbf{r}) = q_g(\mathbf{r}) \quad (2.15)$$

$$-\nabla \cdot (D_g(\mathbf{r})\nabla \phi_g(\mathbf{r}))v(\mathbf{r}) + \Sigma_{r,g}(\mathbf{r})\phi_g(\mathbf{r})v(\mathbf{r}) = q_g(\mathbf{r})v(\mathbf{r}) \quad (2.16)$$

$$-\int_{\Omega} \nabla \cdot (D_g(\mathbf{r})\nabla \phi_g(\mathbf{r}))v(\mathbf{r}) d\Omega + \int_{\Omega} \Sigma_{r,g}(\mathbf{r})\phi_g(\mathbf{r})v(\mathbf{r}) d\Omega = \int_{\Omega} q_g(\mathbf{r})v(\mathbf{r}) d\Omega \quad (2.17)$$

For the purposes of this application, material cross sections and the neutron source are assumed to be constant within an element. Then, the integral can be partitioned into a sum of integrals over the elements in the domain assuming the set of elements $\{\Omega_e\} = \Omega$ for $e = 1, 2, \dots, E$ where E is the total number of elements.

$$-\sum_{e=1}^E D_{g,e} \int_{\Omega_e} \nabla \cdot \nabla \phi_g(\mathbf{r})v(\mathbf{r}) d\Omega_e + \sum_{e=1}^E \Sigma_{r,g,e} \int_{\Omega_e} \phi_g(\mathbf{r})v(\mathbf{r}) d\Omega_e = \sum_{e=1}^E q_{g,e} \int_{\Omega_e} v(\mathbf{r}) d\Omega_e \quad (2.18)$$

The Second Green's Theorem is used to simplify the integral in the first term. A proof can be found in [Li18] in Theorem 9.2.

$$-\int_{\Omega_e} \nabla \cdot \nabla \phi_g(\mathbf{r})v(\mathbf{r}) d\Omega_e = -\int_{\partial\Omega_e} \frac{\partial \phi_g(\mathbf{r})}{\partial \mathbf{n}} v(\mathbf{r}) ds + \int_{\Omega_e} \nabla \phi_g(\mathbf{r}) \cdot \nabla v(\mathbf{r}) d\Omega_e \quad (2.19)$$

Where $\frac{\partial \phi_g(\mathbf{r})}{\partial \mathbf{n}}$ is the outward normal derivative and the integral ds is a line integral in two dimensions or a surface integral in three dimensions. Recognizing that this quantity will only be relevant on the boundary of the problem, the value of the outward normal derivative may be specified in a boundary condition. Specifically, the Albedo boundary condition which has the following form for

$\mathbf{r} \in \partial\Omega$.

$$D_g(\mathbf{r})\nabla\phi_g(\mathbf{r}) + \aleph\phi_g(\mathbf{r}) = 0 \quad (2.20)$$

$$\nabla\phi_g(\mathbf{r}) = \frac{-\aleph\phi_g(\mathbf{r})}{D_g} \quad (2.21)$$

Substituting Eq. 2.19 into Eq. 2.18 and assuming the outward normal derivative is specified in the form of an Albedo boundary condition.

$$\begin{aligned} -\sum_{e=1}^E D_{g,e} \int_{\partial\Omega_e} v(\mathbf{r}) \frac{\partial\phi_g(\mathbf{r})}{\partial\mathbf{n}} ds + \sum_{e=1}^E D_{g,e} \int_{\Omega_e} \nabla\phi_g(\mathbf{r}) \cdot \nabla v(\mathbf{r}) d\Omega_e + \\ \sum_{e=1}^E \Sigma_{r,g,e} \int_{\Omega_e} \phi_g(\mathbf{r}) v(\mathbf{r}) d\Omega_e = \sum_{e=1}^E q_{g,e} \int_{\Omega_e} v(\mathbf{r}) d\Omega_e \end{aligned} \quad (2.22)$$

$$\begin{aligned} \sum_{e=1}^E \aleph \int_{\partial\Omega_e} v(\mathbf{r}) \phi_g(\mathbf{r}) ds + \sum_{e=1}^E D_{g,e} \int_{\Omega_e} \nabla\phi_g(\mathbf{r}) \cdot \nabla v(\mathbf{r}) d\Omega_e + \\ \sum_{e=1}^E \Sigma_{r,g,e} \int_{\Omega_e} \phi_g(\mathbf{r}) v(\mathbf{r}) d\Omega_e = \sum_{e=1}^E q_{g,e} \int_{\Omega_e} v(\mathbf{r}) d\Omega_e \end{aligned} \quad (2.23)$$

Now the function of interest $\phi_g(\mathbf{r})$ is assumed to be a linear combination of chosen basis functions $\{N_i\}$.

$$\phi_g(\mathbf{r}) = \sum_{i=1}^N \alpha_{g,i} N_i(\mathbf{r}) \quad (2.24)$$

Where coefficients $\{\alpha_i\}$ are unknown and will be determined. Typically, these basis functions have unit magnitude and are centered at the node points so the coefficients α_i are the approximated solution at the nodes. Basis functions are typically polynomials of arbitrary magnitude. Linear and quadratic polynomials are common but for the application presented here, only linear basis functions are explored. The test function $v(\mathbf{r})$ is also chosen as a linear combination of the basis functions.

$$v(\mathbf{r}) = \sum_{j=1}^N N_j(\mathbf{r}) \quad (2.25)$$

The testing function is arbitrary so the magnitude is fixed to the magnitude of the basis function.

Eq. 2.24 and Eq. 2.25 are plugged into Eq. 2.23. This yields a linear system of equations.

$$\sum_{e=1}^E \mathfrak{K} \sum_{i=1}^N \alpha_{i,g} \int_{\partial\Omega_e} N_i(\mathbf{r}) N_j(\mathbf{r}) ds + \sum_{e=1}^E D_{g,e} \sum_{i=1}^N \alpha_{i,g} \int_{\Omega_e} \nabla N_i(\mathbf{r}) \cdot \nabla N_j(\mathbf{r}) d\Omega_e + \sum_{e=1}^E \Sigma_{r,g,e} \sum_{i=1}^N \alpha_{i,g} \int_{\Omega_e} N_i(\mathbf{r}) N_j(\mathbf{r}) d\Omega_e = \sum_{e=1}^E q_{g,e} \sum_{i=1}^N \int_{\Omega_e} N_i(\mathbf{r}) d\Omega_e \quad (2.26)$$

$$\sum_{i=1}^N \alpha_{i,g} \sum_{j=1}^N \left(\sum_{e=1}^E \mathfrak{K} \int_{\partial\Omega_e} N_i(\mathbf{r}) N_j(\mathbf{r}) ds + \sum_{e=1}^E D_{g,e} \int_{\Omega_e} \nabla N_i(\mathbf{r}) \cdot \nabla N_j(\mathbf{r}) d\Omega_e + \sum_{e=1}^E \Sigma_{r,g,e} \int_{\Omega_e} N_i(\mathbf{r}) N_j(\mathbf{r}) d\Omega_e \right) = \sum_{i=1}^N \left(\sum_{e=1}^E q_{g,e} \int_{\Omega_e} N_i(\mathbf{r}) d\Omega_e \right) \quad (2.27)$$

$$a(N_i, N_j) = f(N_i) \quad (2.28)$$

$$\mathbf{A}\mathbf{u} = \mathbf{f} \quad (2.29)$$

$a(N_i, N_j)$ is the bilinear form and $f(N_i)$ is the linear form of the finite element system. In matrix notation, \mathbf{A} is described by the integral quantities (more to follow), $\mathbf{u} = \{\alpha_i\}$, and \mathbf{f} is described by the source integral quantity.

Inspecting the matrix \mathbf{A} and the vector \mathbf{f} reveals the following.

$$A_{i,j,g,e} = \mathfrak{K} \int_{\partial\Omega_e} N_i(\mathbf{r}) N_j(\mathbf{r}) ds + D_{g,e} \int_{\Omega_e} \nabla N_i(\mathbf{r}) \cdot \nabla N_j(\mathbf{r}) d\Omega_e + \Sigma_{r,g,e} \int_{\Omega_e} N_i(\mathbf{r}) N_j(\mathbf{r}) d\Omega_e \quad (2.30)$$

$$f_{i,g,e} = q_{g,e} \int_{\Omega_e} N_i(\mathbf{r}) d\Omega_e \quad (2.31)$$

Then,

$$A_{i,j,g} = \sum_{e=1}^E A_{i,j,g,e} \quad (2.32)$$

$$f_{i,g} = \sum_{e=1}^E f_{i,g,e} \quad (2.33)$$

which leads to the natural population of the matrix \mathbf{A} on an element-by-element basis. That is, the matrix \mathbf{A} is assembled by looping through all of the elements and summing their contribution to the

matrix. Note that the contribution due to the surface integral will be zero in elements not on the boundary and may also be zero for problems with select boundary conditions. The population of the vector \mathbf{f} is done similarly. Then, the matrix \mathbf{A} and the vector \mathbf{f} are known for each energy group. The equations are solved on group at a time and ϕ_g is calculated and stored.

Though the notation may be obtuse, the above reduces to a linear system of equations. These equations are constructed from the integral quantities specified by the FEM and the coefficients given by the cross sections and fixed source regions. The integral quantities themselves are expressed explicitly in the next section.

2.2.2 Matrix Quantities

For certain simple elements, the integral quantities described in Eq. 2.30 and Eq. 2.31 have exact analytic forms. The for this application, linear triangles and linear wedges are investigated and many of the integrals have exact expressions. If these quantities cannot be expressed exactly or doing so would be computationally difficult, quadratures can be used and for certain problems, these quadratures can express the integrals exactly. This will be discussed in Sec. 2.2.3.

2.2.2.1 Linear Triangles

Linear triangles are common to two dimensional finite element methods and have been investigated in many applications [Hos17] [HV13] [HSD15]. This is a triangle defined by three corner coordinates with basis functions located on each corner [Sch06].

Originally proposed in [Whi85], there are simple expressions for the quantities. The expression for the line integral is found in [Eng05]. For a triangle with corners $\{x_i, y_i\}$.

$$\int_{\Omega_e} \nabla N_i(\mathbf{r}) \cdot \nabla N_j(\mathbf{r}) d\Omega_e = \frac{1}{4A_e} ((x_{i+1} - x_{i+2})(x_{j+1} - x_{j+2}) + (y_{i+1} - y_{i+2})(y_{j+1} - y_{j+2})) \quad (2.34)$$

$$\int_{\Omega_e} N_i(\mathbf{r}) N_j(\mathbf{r}) d\Omega_e = \frac{A_e}{12} (1 + \delta_{ij}) \quad (2.35)$$

$$\int_{\Omega_e} N_i(\mathbf{r}) d\Omega_e = \frac{A_e}{3} \quad (2.36)$$

$$\int_{\partial\Omega_e} N_i(\mathbf{r}) N_j(\mathbf{r}) ds = \frac{L_e}{6} (1 + \delta_{ij}) \quad (2.37)$$

$$(2.38)$$

Where A_e is the area of the triangular element L_e is the length of the edge between node i and

node j , and δ_{ij} is the Kronecker delta such that

$$\delta_{ij} = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases} \quad (2.39)$$

For higher order triangular elements, it will be necessary to employ a quadrature.

2.2.2.2 Linear Wedges

Wedge elements have not reached the same commonality as triangular elements but are extruded triangles. Geometrically, the shape is a pentahedron or a triangular prism. However, their exact geometric relation is not fixed and the nodes are free to expand. These shapes are unique because three of the faces are quadrilateral and two of the faces are triangular. The values presented herein are not yet found in literature and are calculated by there author and published here for the first time.

$$\int_{\Omega_e} N_i(\mathbf{r}) N_j(\mathbf{r}) d\Omega_e = \frac{V_e}{2} \begin{pmatrix} \frac{1}{18} & \frac{1}{36} & \frac{1}{36} & \frac{1}{36} & \frac{1}{72} & \frac{1}{72} \\ \frac{1}{36} & \frac{1}{18} & \frac{1}{36} & \frac{1}{72} & \frac{1}{36} & \frac{1}{72} \\ \frac{1}{36} & \frac{1}{36} & \frac{1}{18} & \frac{1}{72} & \frac{1}{72} & \frac{1}{36} \\ \frac{1}{36} & \frac{1}{72} & \frac{1}{72} & \frac{1}{18} & \frac{1}{36} & \frac{1}{36} \\ \frac{1}{72} & \frac{1}{36} & \frac{1}{72} & \frac{1}{36} & \frac{1}{18} & \frac{1}{36} \\ \frac{1}{72} & \frac{1}{72} & \frac{1}{36} & \frac{1}{36} & \frac{1}{36} & \frac{1}{18} \end{pmatrix} \quad (2.40)$$

$$\int_{\Omega_e} N_i(\mathbf{r}) d\Omega_e = \frac{V_e}{12} \quad (2.41)$$

$$\int_{\partial\Omega_e} N_i(\mathbf{r}) N_j(\mathbf{r}) d\partial\Omega_e = \begin{cases} \frac{A_\Delta}{12} (1 + \delta_{ij}) & \text{if triangle} \\ \frac{A_\square}{36} (1 + \delta_{ij}) (1 - \frac{1}{2} \delta_{i,(5-j)}) & \text{if quadrilateral} \end{cases} \quad (2.42)$$

Where V_e is the volume of the element. This matrix is indexed A_{ij} and is presented as a matrix because of its irregular form. Notice the integral containing the gradient operator has been omitted because if it could be computed analytically, it would be less computationally efficient than using a quadrature.

2.2.3 Quadratures

Quadratures are sets of coordinates and weights which allow for the exact integration of polynomials of given order. For a given set of weights $\{w_i\}$ and a set of coordinates $\{\mathbf{x}\}$, the quadrature can be

expressed as follows.

$$\int_{\Omega} f(\mathbf{x}) d\Omega \approx \sum_{i=1}^N w_i f(\mathbf{x}_i) \quad (2.43)$$

where Ω is an arbitrary domain described by $\{\mathbf{x}_i\}$. The above quadrature will exactly integrate a polynomial of the order of the quadrature. It is not necessarily true that N be the order of the quadrature.

For one dimensional integrals, the Gaussian quadrature is common and the most compact quadrature. The Gaussian quadrature exactly integrates a polynomial of order n using exactly n points. Weights and coordinates for this quadrature set are presented in [Kam11]. For this quadrature, $n = N$.

Two dimensional and three dimensional quadratures are necessary for the solution of this problem. Triangular quadratures are not as simply derived as line quadratures and the number of points need not equal the order of the polynomial integrated. The triangular quadrature as implemented here is symmetric and open. That is, there are no points on the boundary of the triangle. The structure for this triangular quadrature is found in [Den10].

Quadrilateral quadrature sets are simply tensor products of two line Gaussian quadratures. For an order N polynomial, now n^2 points are required.

Wedge quadrature sets are simply tensor products of a line Gaussian quadrature and a triangular quadrature.

All of the quadratures described here are tabulated for a reference element be it a line, a triangle, a quadrilateral, or a wedge. Integration in the FEM is performed on an arbitrary element in space. Therefore, it is necessary to perform a coordinate transform when using a quadrature set.

$$\int_{\Omega} f(\mathbf{x}) d\Omega = \int_{\Omega_{ref}} f(\mathbf{x}) |\mathbf{J}| d\Omega \approx \sum_{i=1}^N w_i f(\mathbf{x}_i) |\mathbf{J}_i| \quad (2.44)$$

where \mathbf{J} is the Jacobian matrix, \mathbf{J}_i is the Jacobian matrix at quadrature coordinate \mathbf{x}_i , and $|\cdot|$ represents the matrix determinant. Notationally, $J = |\mathbf{J}|$ and is termed the Jacobi.

For simple elements, the Jacobi is constant over the element and can be precalculated to save from allocating, populating, and taking the determinant of a matrix for each integration point. For the elements of concern, these values are presented below as found in [Fel04].

Table 2.1 Jacobian Determinants

Element	Jacobi	J
Triangle	$J = \frac{1}{2} \mathbf{J} $	A_e
Quadrilateral	$J = \mathbf{J} $	$\frac{1}{4} A_e$
Wedge	$J = \frac{1}{2} \mathbf{J} $	$\frac{1}{2} V_e$

2.3 Implementation

2.3.1 Algorithm

Algorithm 1 General Iteration Scheme

- 1: Initialize $\phi^{(0)}$
 - 2: Order the nodes of the problem into RCM order.
 - 3: Calculate Σ_s , Σ_t , and $\nu\Sigma_f$ for each element
 - 4: Calculate Stiffness Matrix A for each group. Store this.
 - 5: **while** Power Iteration **do**
 - 6: $s = s + 1$
 - 7: Update q_{fiss} and q_{up} from previous data $\phi^{(s-1)}$
 - 8: Update χ in each element using previous data
 - 9: **for** $g = 1, G$ **do**
 - 10: Update q_{down} from current data $\phi^{(s)}$
 - 11: Calculate total effective source in each element
 - 12: Update Load Vector with new source
 - 13: Solve $\mathbf{A}\mathbf{u} = \mathbf{f}$ using an iterative technique
 - 14: Parse \mathbf{u} for ϕ nodal solution
 - 15: Calculate element-average $\bar{\phi}$
 - 16: **end for**
 - 17: Update k_{eff}
 - 18: Check convergence
 - 19: **end while**
-

2.3.2 Memory and Storage

The Finite Element matrix **A** is large and sparse so a sparse storage implementation is required.

2.3.3 Linear System Solution

2.4 Reference Results

2.4.1 Triangular Element Manufactured Solutions

2.4.1.1 One Dimension, One-Group, Fixed Source

2.4.1.2 One Dimension, One-Group, Criticality

2.4.1.3 One Dimension, Two-Group, Criticality

2.4.1.4 One Dimension, One-Group, Two-Region, Criticality

2.4.1.5 Two Dimension, One-Group, Criticality

2.4.2 Two Dimension Reactors

2.4.2.1 VVER440

2.4.2.2 SNR

2.4.2.3 HWR

2.4.2.4 IAEA

2.4.3 Wedge Element Manufactured Solution Finite Cylinder

2.4.4 Three Dimension Reactors

2.4.4.1 MONJU

2.4.4.2 KNK

CHAPTER 3

THERMAL HYDRAULICS

3.1 Axial Convection Model

3.2 Radial Conduction Model

3.2.1 Derivation

3.2.2 Relations

3.3 Cross Section Treatment

3.4 Results

CHAPTER 4

THERMAL EXPANSION

4.1 Necessity of Modeling

4.2 Model Details

4.3 Implementation

4.4 Results

CHAPTER 5

CONCLUSIONS

5.1 Results Discussion

5.2 Code Enhancements

5.3 Further Investigations

BIBLIOGRAPHY

- [Den10] Deng, S. *Quadrature Formulas in Two Dimensions*. 2010.
- [DH76] Duderstadt, J. J. & Hamilton, L. J. *Nuclear Reactor Analysis*. New York: Wiley, 1976.
- [ER35] Einstein, A. & Rosen, N. "The particle problem in the general theory of relativity". *Physical Review* **48.1** (1935), pp. 73–77.
- [Eng05] Engblom, S. *The Finite Element Method*. 2005.
- [Fel04] Felippa, C. A. *Advanced Finite Element Methods*. Boulder, CO: University of Colorado at Boulder, 2004.
- [GVL96] Golub, G. & Van Loan, C. *Matrix computations*. Johns Hopkins Univ Pr, 1996.
- [Hos17] Hosseini, S. A. "Sensitivity Analysis of the Galerkin Finite Element Method Neutron Diffusion Solver to the Shape of the Elements". *Nuclear Engineering and Technology* **49.1** (2017), pp. 29–42.
- [HSD15] Hosseini, S. A. & Saadatian-Derakhshandeh, F. "Galerkin and Generalized Least Squares finite element: A comparative study for multi-group diffusion solvers". *Progress in Nuclear Energy* **85** (2015), pp. 473–490.
- [HV13] Hosseini, S. A. & Vosoughi, N. "Development of two-dimensional, multigroup neutron diffusion computer code based on GFEM with unstructured triangle elements". *Annals of Nuclear Energy* **51** (2013), pp. 213–226.
- [Kam11] Kamermans, M. "Gaussian Quadrature Weights and Abcissae" (2011).
- [LK74] Larsen, E. & Keller, J. "Asymptotic solution of neutron transport problems for small mean free paths". *Journal of Mathematical Physics* **15** (1974), p. 75.
- [Li18] Li, Z. et al. *Numerical Solution of Differential Equations*. Cambridge: Cambridge, 2018.
- [Sch06] Schroeder, W. et al. *The Visualization Toolkit*. 4th ed. Kitware, 2006.
- [Whi85] White, R. E. *An Introduction of the Finite Element Method with Applications to Nonlinear Problems*. New York: Wiley, 1985.

APPENDIX

APPENDIX A

LOREM IPSUM

A.1 A First Section

A.1.0.0.1 Filler Text

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique,

libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus.

Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Donec odio elit, dictum in, hendrerit sit amet, egestas sed, leo. Praesent feugiat sapien aliquet odio. Integer vitae justo. Aliquam vestibulum fringilla lorem. Sed neque lectus, consectetur at,

Table A.1 A table in the appendix.

System	Author
T _E X	Donald Knuth
L ^A T _E X	Leslie Lamport

consectetuer sed, eleifend ac, lectus. Nulla facilisi. Pellentesque eget lectus. Proin eu metus. Sed porttitor. In hac habitasse platea dictumst. Suspendisse eu lectus. Ut mi mi, lacinia sit amet, placerat et, mollis vitae, dui. Sed ante tellus, tristique ut, iaculis eu, malesuada ac, dui. Mauris nibh leo, facilisis non, adipiscing quis, ultrices a, dui.

Morbi luctus, wisi viverra faucibus pretium, nibh est placerat odio, nec commodo wisi enim eget quam. Quisque libero justo, consectetuer a, feugiat vitae, porttitor eu, libero. Suspendisse sed mauris vitae elit sollicitudin malesuada. Maecenas ultricies eros sit amet ante. Ut venenatis velit. Maecenas sed mi eget dui varius euismod. Phasellus aliquet volutpat odio. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Pellentesque sit amet pede ac sem eleifend consectetuer. Nullam elementum, urna vel imperdiet sodales, elit ipsum pharetra ligula, ac pretium ante justo a nulla. Curabitur tristique arcu eu metus. Vestibulum lectus. Proin mauris. Proin eu nunc eu urna hendrerit faucibus. Aliquam auctor, pede consequat laoreet varius, eros tellus scelerisque quam, pellentesque hendrerit ipsum dolor sed augue. Nulla nec lacus.

Suspendisse vitae elit. Aliquam arcu neque, ornare in, ullamcorper quis, commodo eu, libero. Fusce sagittis erat at erat tristique mollis. Maecenas sapien libero, molestie et, lobortis in, sodales eget, dui. Morbi ultrices rutrum lorem. Nam elementum ullamcorper leo. Morbi dui. Aliquam sagittis. Nunc placerat. Pellentesque tristique sodales est. Maecenas imperdiet lacinia velit. Cras non urna. Morbi eros pede, suscipit ac, varius vel, egestas non, eros. Praesent malesuada, diam id pretium elementum, eros sem dictum tortor, vel consectetuer odio sem sed wisi.

A.2 A Second Section

Etiam ac leo a risus tristique nonummy. Donec dignissim tincidunt nulla. Vestibulum rhoncus molestie odio. Sed lobortis, justo et pretium lobortis, mauris turpis condimentum augue, nec ultricies nibh arcu pretium enim. Nunc purus neque, placerat id, imperdiet sed, pellentesque nec, nisl. Vestibulum imperdiet neque non sem accumsan laoreet. In hac habitasse platea dictumst. Etiam condimentum facilisis libero. Suspendisse in elit quis nisl aliquam dapibus. Pellentesque

auctor sapien. Sed egestas sapien nec lectus. Pellentesque vel dui vel neque bibendum viverra. Aliquam porttitor nisl nec pede. Proin mattis libero vel turpis. Donec rutrum mauris et libero. Proin euismod porta felis. Nam lobortis, metus quis elementum commodo, nunc lectus elementum mauris, eget vulputate ligula tellus eu neque. Vivamus eu dolor.

Nulla in ipsum. Praesent eros nulla, congue vitae, euismod ut, commodo a, wisi. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Aenean nonummy magna non leo. Sed felis erat, ullamcorper in, dictum non, ultricies ut, lectus. Proin vel arcu a odio lobortis euismod. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Proin ut est. Aliquam odio. Pellentesque massa turpis, cursus eu, euismod nec, tempor congue, nulla. Duis viverra gravida mauris. Cras tincidunt. Curabitur eros ligula, varius ut, pulvinar in, cursus faucibus, augue.