

Coding of space and time for memory function.

Michael E. Hasselmo, Jennifer C. Robinson, Patrick A. LaChance, Lucius Kelton Wilmerding,
G. William Chapman

Center for Systems Neuroscience, Boston University
610 Commonwealth Ave., Boston, MA 02215

Corresponding author:

Prof. Michael Hasselmo, e-mail: hasselmo@bu.edu, twitter: @HasselmoMichael

Chapter contribution to: Time, space and memory, Oxford University Press, Edited by Sara
Aronowitz and Lynn Nadel

Acknowledgements: This work supported by the National Institutes of Health, grant numbers
R01 MH120073, R01 MH60013, R01 MH052090 and by the Office of Naval Research MURI
N00014-16-1-2832 and MURI N00014-19-1-2571 and DURIP N00014-17-1-2304. The authors
have no conflict of interest.

Manuscript sent to: memoryspacetime@gmail.com.

24 **Abstract**

25 Data from rodents shows neuronal activity that may code specific intervals of space and time, as
26 revealed by recordings in structures implicated in episodic memory function. These data are
27 relevant to modeling of episodic memories as spatiotemporal trajectories. The review will
28 include data on neuronal activity that disambiguates overlapping spatiotemporal trajectories, as
29 well as neurons that change firing rate and rhythmicity based on running speed and head
30 direction. These data support the model of episodic memory as a trajectory that includes speed
31 and direction, in contrast to Tulving's initial definition of episodic memory as a series of
32 snapshots. Spatial behavior requires coding in multiple coordinate frames, and recent data
33 supports the idea that space could involve neural coding of environmental boundaries in
34 egocentric coordinates, as well as neural coding of transformations between egocentric input and
35 memory that could be mediated by neurons coding allocentric location. These data will be
36 discussed in the context of a more recent model of the transformation between coordinate
37 systems. The egocentric input can be used both for coding the location along spatiotemporal
38 trajectories and for retrieving specific viewpoints of the environment. The chapter will also
39 include a brief discussion of the importance of mathematical models to move beyond verbal
40 definitions.

41

Introduction

Multiple cortical structures are implicated in episodic memory function. This review will present data on the neuronal activity that codes space and time in cortical structures. These data have been used in models of episodic memory that involve the encoding and retrieval of spatiotemporal trajectories. This review will also present data and modeling of the transformation between egocentric and allocentric coordinate systems.

Episodic memory was defined as memory for events that occur at a specific place and a specific time (Tulving, 1984; Eichenbaum et al., 1999). Episodic memory was described as “snapshots whose orderly succession can create the mnemonic illusion of the flow of past time” (Tulving, 1984). A neural model of episodic memory (Hasselmo, 2009; Hasselmo, 2012) proposed that episodic memory must contain more than a sequence of snapshots, but instead include a continuous representation of time and space as a spatiotemporal trajectory, that includes the speed and direction of movement of an agent or other objects (Hasselmo et al., 2010; Hasselmo, 2012), the agent’s viewpoint of an event (Conway, 2009), and the coding of prior context for disambiguation of memories (Hasselmo and Eichenbaum, 2005; Hasselmo, 2009). This chapter will review relevant neural data relevant to modeling of episodic memory, as well as briefly reviewing models and emphasizing approaches that include modeling of the transformation between egocentric and allocentric coordinate systems. Finally, there will be a section emphasizing the importance of mathematical models to move beyond verbal definitions.

Coding of time (time cells)

Episodic memory can be succinctly defined as “What did you do at time T in place P” (Tulving, 1984). The neural coding of time and space is important for this definition, and many cortical neurons code both time and space (Pastalkova et al., 2008; MacDonald et al., 2011; Kraus et al., 2013; Kraus et al., 2015). Neurophysiological data shows neurons, termed “time cells,” that code time intervals relative to task events, such as the onset of a delay period. In many tasks used to study time cells, a rodent runs around a loop of elevated track, and then remains in a consistent single location and direction while running for 10-20 seconds on a running wheel (Pastalkova et al., 2008) or a treadmill (Kraus et al., 2013; Kraus et al., 2015). During this delay period of running in one location, individual neurons fire at specific intervals after the onset of the delay. Running is not essential as time cells can also appear during delay periods of a delayed matching task performed without running (MacDonald et al., 2011) or even during a delay period in stationary head fixed animals (MacDonald et al., 2013; Heys and Dombeck, 2018). Time cell responses have been shown in a wide range of different structures, including hippocampal region CA1 (Pastalkova et al., 2008; Kraus et al., 2013; Mau et al., 2018), hippocampal region CA3 (Salz et al., 2016) as well as the entorhinal cortex (Kraus et al., 2015; Tsao et al., 2018; Heys and Dombeck, 2018). These data show that time cell responses can be important for disambiguating events that occur at one time point versus another on a time scale of seconds.

More recent data shows that neurons also show a difference in calcium activity across the trials within a given day (Mau et al., 2018), and show consistencies across days supporting coding of time on the time scale of minutes (MacDonald et al., 2011; Liu et al., 2022). This supports a model of multiscale coding of time developed by Marc Howard and colleagues (Howard et al., 2014). Calcium imaging of the same population of hippocampal neurons over several days (Mau

et al., 2018) also shows that time cells drop out or appear slowly over days, resulting in a change in correlation across the population on time scales of days (Mau et al., 2018). This slow drift in ensemble membership could provide a differential coding of memories on different days (Howard et al., 2014; Mankin et al., 2015; Rubin et al., 2015; Cai et al., 2016; Rule et al., 2019; Kinsky et al., 2020; Ziv et al., 2013; Levy et al., 2021). Models show that changes in the correlation of a population of cells on multiple longer temporal scales are essential for the capacity to differentiate episodic memories occurring at different time points on the scale of seconds, minutes, and hours (Howard et al., 2014; Liu et al., 2019). This multi-scale representation provides an efficient representation for memory on different scales, but it is also possible that time coding involves different mechanisms at different scales (Phillips, 2014), particularly for long scales on the order of weeks, months or years. The focus on disambiguating different time points motivated the use of the term “time cell” (Howard and Eichenbaum, 2013; Eichenbaum, 2014; Kraus et al., 2013; MacDonald et al., 2011). However, these cells could be coding sequences of associations of internal and external features making up the events in an episode (Buzsaki and Tingley, 2018) consistent with the initial description of these neurons as ‘episode cells’ (Pastalkova et al., 2008).

One experiment compared time versus running distance by recording as a rat ran on a treadmill at different speeds during different delay periods (Kraus et al., 2013). This experiment showed that neurons could respond on the basis of either time or running distance during the delay (Kraus et al., 2013) as predicted by previous models (Hasselmo, 2008; Burgess et al., 2007). Neurons that fire as time cells also code other dimensions related to episodic memory, consistent with evidence of mixed selectivity in other regions (Rigotti et al., 2013). Some cells that fire as

time cells during running on the treadmill might also fire as place cells during running off the treadmill on the return arms (Kraus et al., 2013; Mau et al., 2018) indicating that these cells do not only code time. Coding of both space and time was also shown for single grid cells in the entorhinal cortex (Kraus et al., 2015). These grid cells fired in an array of spatial locations when animals foraged in a two-dimensional environment, but also fired as time cells at different time points during a 16 second delay as the rat ran in a single location on the treadmill. Interestingly, a recent study showed that the overall population of space-encoding neurons and neurons coding time during immobility in entorhinal cortex form anatomically distinct sub-populations (Heys and Dombeck, 2018; Heys et al., 2014). The coding of episodes as spatiotemporal trajectories only requires a population code containing both space and time, so this model can function whether the representation is shared across individual neurons or appears in different populations (Hasselmo, 2012). Human imaging data shows neural activity associated with disambiguating the retrieval of overlapping trajectories in virtual mazes (Brown and Stern, 2014; Brown et al., 2010; Brown et al., 2014).

Time cells code time not only by overall firing rate, but also by the phase of firing of cells relative to rhythmic oscillations at theta frequency in the local field potential. Time cells show theta phase precession within their firing fields (Pastalkova et al., 2008; Terada et al., 2017; Ning et al., 2022). As a time cell starts firing, its spikes appear at late phases of theta, and as time evolves the spiking shifts to earlier phases of theta cycle before spiking ends. The coding of space by place cells shows a similar shift in phase as an animal runs through a place field. This phase coding appears important for the temporal specificity of time cell firing as time cells are lost during inactivation of the medial septum (Wang et al., 2015), which reduces theta rhythm in

the hippocampus (Brandon et al., 2014; Rawlins et al., 1979). The same inactivation of medial septum also removes spatial specificity of grid cell firing (Brandon et al., 2011; Koenig et al., 2011). Theta phase coding has the advantage that it could allow a single neuron to code a continuous dimension of time or space, which might allow a broader range of transformations on the level of single neurons that might be difficult to implement across a full population.

Coding of spatial location

A range of studies show that damage to the hippocampus causes impairments in two-dimensional spatial navigation tasks such as the Morris water maze, in which the animal learns a specific goal platform location and must then navigate to that location from a range of different starting locations (Morris et al., 1982; Eichenbaum et al., 1990). Models have addressed how different neural subtypes could underlie planning of spatiotemporal trajectories to generate the correct trajectory to the goal location from a new start location (Erdem and Hasselmo, 2014; Erdem and Hasselmo, 2012; Redish and Touretzky, 1998). Early models focused on the role of place cells and head direction cells (Redish and Touretzky, 1998), but those models would require dense representation of spatial locations by place cells. Later models address the additional role of grid cells and speed cells in planning trajectories to a goal location without needing to form a place cell code for each location (Erdem and Hasselmo, 2014; Erdem and Hasselmo, 2012; Kubie and Fenton, 2012). Impairments of goal-finding in the Morris water maze are also observed after lesions of the entorhinal cortex (Steffenach et al., 2005) or the dorsal presubiculum (Taube et al., 1992), consistent with a role of different neuronal subtypes such as head direction cells and grid cells from these regions. Further data shows neural coding of position along a trajectory in structures such as the retrosplenial cortex (Alexander and Nitz, 2017; Alexander and Nitz, 2015),

and human imaging data shows coding of arc length along a trajectory in addition to coding of euclidean distance, translation and rotation (Chrastil et al., 2016; Chrastil et al., 2015). This section will briefly review some data on coding of spatial location.

Place cells in hippocampus

Neurophysiological recording in the hippocampus revealed place cells that fire based on spatial location (O'Keefe, 1976; O'Keefe and Dostrovsky, 1971). The relationship of these place cell responses to philosophical questions about the a priori representation of space were addressed extensively in the influential book by O'Keefe and Nadel on coding of space as a cognitive map (O'Keefe and Nadel, 1978). Further studies showed place cell responses when animals were in specific locations during foraging in open field environments (Muller et al., 1987; O'Keefe and Burgess, 1996; Huxter et al., 2008; Lever et al., 2002), and in local areas of linear tracks (O'Keefe and Recce, 1993), the 8-arm radial maze (McNaughton et al., 1983), or a spatial alternation task (Ainge et al., 2007; Wood et al., 2000; Kinsky et al., 2020). The firing of place cells can vary dependent on many factors including direction through the place field (Fenton and Muller, 1998) and cells can have more than one firing field (Fenton et al., 2008). The position of an animal can be effectively decoded from the firing activity of hippocampal place cells (Brown et al., 1998), supporting their role in guiding behavior in spatial memory tasks. In more recent studies, hippocampal cell firing has been shown to code information about the direction and distance of a goal location (Ormond and O'Keefe, 2022).

In addition to coding location by firing rate, place cells also code location by their phase of firing in a phenomenon called theta phase precession (O'Keefe and Recce, 1993). This discovery

preceded the finding of phase precession by time cells described above. As the animal enters the firing field of a place cell, spiking occurs at late phases of theta and then shifts to earlier phases as the animal runs through the firing field and exits (O'Keefe and Recce, 1993; Skaggs et al., 1996; Maurer et al., 2006; Schmidt et al., 2009; Zugaro et al., 2005). Theta phase precession is associated with sequential spiking of neurons coding sequential places on different phases (Foster and Wilson, 2007), but theta phase precession appears on the first trial of running on a novel linear track, whereas theta sequences only appear on later trials (Feng et al., 2015). These data further emphasize the potential role of spike timing and phase for the coding of space and time.

Grid cells in entorhinal cortex

Neurophysiological recording in the entorhinal cortex demonstrates different types of coding of spatial dimensions. The most striking form of coding involves the response of entorhinal grid cells, which respond when a foraging animal visits an array of spatial locations in the environment that appear in a hexagonal pattern (Hafting et al., 2005). Different grid cells fire with different size and spacing between firing fields, allowing a population of grid cells to code a single location (Sargolini et al., 2006; Barry et al., 2007; Stensola et al., 2012; Heys et al., 2014). Many grid cells code both for the animal's location and the current head direction of the animal (Sargolini et al., 2006).

Entorhinal grid cells also exhibit phase coding in the form of theta phase precession as an animal runs on a linear track (Hafting et al., 2008), or as an animal forages in two-dimensions in an open field (Climer et al., 2013; Jeewajee et al., 2014). Consistent with this, the intrinsic rhythmicity of

entorhinal neurons differs with spatial scale (Jeewajee et al., 2008) and shifts with running speed (Hinman et al., 2016; Jeewajee et al., 2008). The potential role of theta rhythm in generation of grid cell responses is supported by the fact that inactivation of the medial septum causes both a dramatic reduction of theta rhythm in the entorhinal cortex (Mitchell et al., 1982; Jeffery et al., 1995), and a loss of the spatial selectivity of firing of grid cells (Brandon et al., 2011; Koenig et al., 2011). Inactivation of the medial septum or lesions of the fornix also cause impairments in the memory of goal locations (O'Keefe et al., 1975; Chrobak et al., 1989; Brioni et al., 1990). The specific population of medial septal neurons involved in regulating grid cell firing has not yet been demonstrated. However, recent studies show that inactivation of glutamatergic neurons causes a decreased specificity in grid cell firing activity, and inactivation of GABAergic neurons in the medial septum results in the loss of grid cell spatial firing along with a reduction in theta rhythm oscillations (Robinson et al., 2019).

Coding of prior context

The performance of tasks such as delayed spatial alternation or delayed non-match to position requires the capacity to distinguish (disambiguate) spatial location on different trials, as described in previous modeling work (Levy, 1996; Hasselmo and Eichenbaum, 2005; Hasselmo, 2009). Lesions of the hippocampus cause impairments in these types of tasks (Ainge et al., 2007; Aggleton et al., 1986; Aggleton et al., 1995; Hallock et al., 2013) that require memory of both spatial location and the specific time of the prior trial to disambiguate it from other previous trials. Neurophysiological data in these tasks show context-dependent activity appropriate for this behavioral disambiguation based on memory. For example, when a rat runs on the stem of spatial alternation task, individual neurons will fire selectively based on the past or future turning

response. These “splitter” neurons have been observed in the hippocampus (Wood et al., 2000; Ferbinteanu and Shapiro, 2003; Kinsky et al., 2020; Levy et al., 2021) and entorhinal cortex (Frank et al., 2000; Lipton et al., 2007; O'Neill et al., 2017). The separation and disambiguation of overlapping spatiotemporal trajectories is a challenge for any model of episodic memory, particularly when considering the large number of memories that can be generated in a single familiar environment (Robins, 2015). This raises questions of the relative overlap of representations, which could range from the extreme case of an index model, in which a small number of non-overlapping neurons code each memory, to the alternate case of a broadly distributed representation in which most neurons are involved in every memory, which is more amenable to cueing of memories, but less amenable to prevention of interference (Robins, 2015). The nature of the amount of overlap and capacity to cue less overlapping memories remains an empirical question for both simulations and neurophysiology (Hasselmo, 2015).

The context-dependent neuronal responses in a spatial alternation task can appear at specific times during training and are more stable than place cell responses in the task, possibly because the splitter responses are more necessary for accurate task performance (Kinsky et al., 2020). The left-right discriminability of splitter cell responses correlates significantly with accurate behavioral performance (Kinsky et al., 2020). Context-dependent activity can also distinguish the sample versus test trials in delayed non-match to position (Griffin et al., 2007; Levy et al., 2021), and during the course of learning the task shows a gradual shift from coding both turn direction and task phase, to showing more coding of turn direction or task phase alone (Levy et al., 2021). The separation of representation could also include the dentate gyrus, as neural

activity associated with turning to one side of the maze differs from the representation associated with turning both directions (Wilmerding et al., 2023).

Both the guidance of behavior and the learning-dependent shift in context-dependent representations over time may depend upon mechanisms of sequence retrieval during theta rhythm. Theta sequences appear to reflect planning of future trajectories, as sequences appear at choice points (Johnson and Redish, 2007; Kay et al., 2020), and the length of theta sequences increases with greater distance of future goals (Wikenheiser and Redish, 2015). The phase of firing relative to theta rhythm also appears to shift based on the novelty of individual cues (Manns et al., 2007) or the novelty of the environment (Wells et al., 2013; Douchamps et al., 2013), consistent with proposals for encoding and retrieval on different phases of theta rhythm cycles (Hasselmo et al., 2002; Hasselmo, 2006). If encoding and retrieval processes occur on different phases of theta, then the ability to discriminate a retrieved memory from current sensory input could depend upon an intact theta rhythm (Hasselmo, 2005). Loss of this phase coding could result in confabulation of imaginary memory with real memory, which has been shown to occur after damage to the medial septum (DeLuca and Cicerone, 1991). The different phases of encoding and retrieval could include a neural signal associated with phase that distinguishes the memory of a real event from the memory of an imagined event (Boyle, 2021).

Coding of trajectory speed and direction

In contrast to the model of episodic memory as a series of snapshots, the model of episodic memory as a continuous spatiotemporal trajectory includes dimensions beyond spatial location and time interval (Hasselmo, 2012; Hasselmo, 2009). The fact that one can remember a specific

viewpoint of a scene (Conway, 2009), or changes in speed of movement indicate that direction and speed are available in episodic memories.

In addition to the importance of speed and direction for episodic memory, many models of grid cell firing use path integration of self-motion to code location. This standard mechanism could function in parallel with coding of location by transformation of sensory input. Grid cells retain their spatial firing pattern in darkness, suggesting a role for path integration of self-motion in the absence of visual cues (Hafting et al., 2005; Dannenberg et al., 2020). However, sensory input is important as grid cells rotate with visual cues in a circular environment (Hafting et al., 2005), and spatial coding by grid cells is lost when all sensory cues are lost or obscured including visual, auditory, somatosensory and olfactory input (Chen et al., 2016; Pérez-Escobar et al., 2016). The role of sensory input is further supported by evidence that grid cells lose spatial coding during inactivation of regions providing head direction input (Winter et al., 2015). The subsequent sections will review data on the potential role of memory for spatial location based on path integration versus the transformation of sensory input.

Coding of speed

The running speed of animals has been shown to be coded by neurons in the hippocampus (O'Keefe et al., 1998; McNaughton et al., 1983) and medial entorhinal cortex (Sargolini et al., 2006; Wills et al., 2012; Buetfering et al., 2014; Kropff et al., 2015; Hinman et al., 2016). Some cells appear to selectively code running speed (Kropff et al., 2015), but others show mixed selectivity as grid cells and head direction cells that also code running speed (Sargolini et al., 2006; Wills et al., 2012; Buetfering et al., 2014; Jeewajee et al., 2008; Hinman et al., 2016).

Coding of running speed also appears in retrosplenial and parietal cortex (McNaughton et al., 1994; Alexander et al., 2020; Clancy et al., 2019; Carstensen et al., 2021) and sensory responses are modulated by running speed in visual cortex (Niell and Stryker, 2010) and auditory cortex (Nelson and Mooney, 2016). The direct coding of speed could occur without requiring the computation of distance over time, as neurons also show responses to acceleration guided by the vestibular system, such that integration of acceleration could generate a speed response independent of the ongoing perception of distance or time.

This speed coding could be important for episodic memory of spatiotemporal trajectories. In contrast, the data on speed coding does not uniformly support the use of speed for path integration of 2-dimensional spatial location. Models of path integration require linear coding of running speed by firing rate, as found in many speed tuning curves, but many speed-modulated cells show non-linear responses that saturate at moderate speeds (Hinman et al., 2016; Dannenberg et al., 2019). Surprisingly, neurons represent speed by firing rate over intervals of several seconds, but over shorter periods than a second, the firing rate code is too inaccurate for effective path integration (Dannenberg et al., 2019). This calls into question the use of coding of location based on integration of a firing rate code for running speed.

Coding of direction

In addition to coding of running speed, both models of episodic memory as spatiotemporal trajectories and path integration models of grid cells require coding of movement direction. Many place cells show sensitivity to movement direction on a one-dimensional linear track (McNaughton et al., 1983; Huxter et al., 2003), and hippocampal cells also appear to code the

direction of a goal on a honeycomb maze (Ormond and O'Keefe, 2022). Outside the hippocampus, there are many neurons that show responses to the current allocentric direction of an animal's head (Taube et al., 1990). These head direction cells do not depend on current location or movement direction. A systematic analysis of neurons in entorhinal cortex during periods when movement direction differed from head direction demonstrated numerous head direction cells but no cells that exclusively code movement direction (Raudies et al., 2015). Researchers have proposed that movement direction could instead be coded by rhythmic firing of theta cells (Welday et al., 2011) or spiking in theta sequences (Zutshi et al., 2017). However, the broad evidence for head direction cells in the absence of movement direction cells suggest that path integration of self-motion might be less important than the coding of sensory feature angle provided by head direction cells (Raudies et al., 2015).

Head direction cells could instead be vital for accurate transformation of egocentric coordinates into allocentric coordinates, as allocentric head direction is needed if one is to transform egocentric coordinates of sensory feature angle into allocentric location (Byrne et al., 2007; Touretzky and Redish, 1996; Bicanski and Burgess, 2018). Head direction cells have been found in a range of structures including dorsal presubiculum (Taube et al., 1990), anterior thalamus (Taube, 1995) and entorhinal cortex (Sargolini et al., 2006; Brandon et al., 2013; Brandon et al., 2011; Giocomo et al., 2014). Lesions of the dorsal presubiculum or anterior thalamic nucleus, which both provide head direction input to cortex, cause destabilization of hippocampal place cells (Goodridge and Taube, 1997) and loss of spatial coding by entorhinal grid cells (Winter et al., 2015). Head direction cells usually do not show theta rhythmicity, but in the entorhinal cortex show theta rhythmic firing that falls on alternate cycles of the theta rhythm (Brandon et

al., 2013), consistent with place cell readout of trajectories on alternate theta cycles in the hippocampus (Kay et al., 2020).

Coding based of different coordinate systems

The above data indicates that path integration of self-motion may not be the most important mechanisms for updating the memory of spatial location. In contrast, the influence of sensory feature angle could be used to update the memory of spatial location, as supported by the influence of visual cue rotation on the firing of place cells (Muller and Kubie, 1987), and grid cells (Hafting et al., 2005) and the loss of grid cell firing when darkness is combined with removal of auditory and somatosensory cues (Chen et al., 2016; Pérez-Escobar et al., 2016). Understanding the influence of sensory input on place cell firing requires understanding of coordinate transformations. The coding of spatial location by place cells is commonly described in allocentric coordinates (i.e. allocentric coordinates describe the position of an animal relative to environment boundaries). This requires a coordinate transformation from the egocentric coordinates of sensory input such as visual feature angle (i.e. egocentric coordinates describe the position of a feature relative to an animal). This may correspond to the philosophical description of a perspectival view of an object compared to a constant representation of an object (Green and Schellenberg, 2017). Studies of neural activity in the human brain have demonstrated differences in neural activity associated with viewing a navigation task from a first person, egocentric perspective, compared to performing the task from a third-person overhead perspective (Sherrill et al., 2013; Sherrill et al., 2015). The following sections will review further data relevant to this topic.

Allocentric boundary cells

The important influence of sensory cues for location coding was shown in experiments in which the distance of environmental barriers relative to other barriers (i.e. allocentric position) was changed (i.e. changing a 1x1 meter square environment to a 1x2 meter rectangle) and this was shown to alter the position of the firing fields of place cells (O'Keefe and Burgess, 1996). This data motivated the theoretical proposal of boundary vector cells that code animal position relative to boundaries (Burgess et al., 2000; Hartley et al., 2014; Hartley et al., 2000). This theoretical prediction was supported by data demonstrating boundary vector cells that fire when boundaries are at a specific distance and allocentric angle in allocentric coordinates (Solstad et al., 2008; Lever et al., 2009; Savelli et al., 2008; Barry et al., 2006). These boundary responses can occur at a distance from the boundary (Lever et al., 2009), indicating a role of visual sensory cues, and they show responses to insertion of new barriers in the environment (Lever et al., 2009). A related population of neurons, termed object vector cells, respond when the animal occupies specific allocentric angles and distances from non-boundary objects (Hoydal et al., 2019; Deshmukh and Knierim, 2011; Deshmukh and Knierim, 2013). Grid cells also respond to the allocentric positions of environmental boundaries, showing compression or expansion of the distance between firing fields (Barry et al., 2007; Stensola et al., 2012; Munn et al., 2020), and changes in coding of velocity with wall movement (Munn et al., 2020).

Models have demonstrated how allocentric spatial location can be generated from egocentric visual coding of boundaries (Byrne et al., 2007; Bicanski and Burgess, 2018; LaChance and Taube, 2023; O'Keefe, 1990). Models have demonstrated how allocentric boundary vector cells could be generated from egocentric sensory coding of environment boundaries combined with

head direction (Burgess et al., 2000; Hartley et al., 2000; Byrne et al., 2007; Bicanski and Burgess, 2018). These latter models predicted the existence of neurons that code the egocentric position of boundaries. Influences of boundary location on grid cells and allocentric boundary cells has also been modeled based on the angle and optic flow of visual features (Raudies and Hasselmo, 2015; Sherrill et al., 2015).

Egocentric boundary cells

The prediction that allocentric boundary cells could be generated from egocentric boundary cells was supported by data showing egocentric boundary cells in a range of structures, including the retrosplenial cortex (Alexander et al., 2020; van Wijngaarden et al., 2020), the postrhinal cortex (LaChance et al., 2019; Gofman et al., 2019), the entorhinal cortex (Wang et al., 2018; Wang et al., 2020) and structures receiving output from these regions such as the dorsomedial striatum (Hinman et al., 2019) and parietal cortex (Alexander et al., 2020). As animals forage in an open field environment, egocentric boundary cells fire selectively when the barriers or boundaries of the environment are at a specific angle and distance to the animal (Hinman et al., 2019; Alexander et al., 2020). They are therefore most efficiently described by plotting the position of the barrier for each spike in egocentric polar coordinates. The sum over all spikes shows how neurons respond to a barrier at a specific distance of a few to many tens of centimeters. Different neurons respond selectively for barriers at a specific range of angles and distances relative to the animal itself, with most right hemisphere neurons responding to barriers directly to the left of the animal, and most left hemisphere neurons responding to barriers directly to the right (Alexander et al., 2020). Other neurons respond to angles to the front or behind the animal. Many egocentric boundary vector responses are invariant to the appearance of environmental

boundaries indicating that the response property is not driven by high-level visual features (Hinman et al., 2019). While some papers focus on the egocentric response to barriers, others focus on the coding of position relative to the center of the environment or specific objects or goals (Wang et al., 2018; Wang et al., 2020). The neurons show tuning to barriers at a number of distances, including distances well outside the range of whisker contact, as well as at a number of angles, including positions behind the animal.

Egocentric coding of the environments has also been shown in several other cortical regions, including posterior parietal cortex, secondary motor cortex, and postrhinal cortex (LaChance et al., 2019; Gofman et al., 2019; Alexander et al., 2020). In the postrhinal cortex egocentric boundary responses persist in darkness (LaChance et al., 2019), supporting the computational theory of their generation by some mechanism of path integration based on prior contact with the barrier. In the postrhinal cortex, egocentric bearing was found to be anchored to the center of the environment rather than the boundaries (LaChance et al., 2019).

The theta phase coding shown for place cells and grid cells reviewed above suggests that theta phase coding might occur for egocentric and allocentric boundary cells. Retrosplenial neurons show phasic firing relative to hippocampal theta rhythmicity (Alexander et al., 2018) and some egocentric boundary cells show theta phase locking (Alexander et al., 2020). Consistent with the proposed separation of encoding and retrieval on different theta phases in the hippocampus (Hasselmo et al., 2002), allocentric boundary cells fire on different phases of theta during direct experience of boundaries versus trace responses to boundaries that are no longer present (Poulter et al., 2021). Theta phase coding could provide a component of spatial representation that can

contribute to the transformation from egocentric coordinates to allocentric coordinates and the encoding of egocentric information in spatiotemporal trajectories for episodic memory.

Brief review of hippocampal models

The above sections reviewed some of the data on neural representations for time and space that are relevant to episodic memory. This section will provide a brief review of some existing models of episodic memory and the internal representations used in those models. Historically, both psychological and neural models of episodic memory have focused on the representation of the world as a set of vectors, usually with an arbitrary mapping of environmental features (such as words in a verbal memory task) to individual vectors. This mapping was used in many memory models in mathematical psychology (Murdock, 2005), and the vector representation carried over into neural network models of episodic memory (McNaughton and Morris, 1987) that focused on encoding of an array of vectors representing individual memories (McNaughton and Morris, 1987). These memories were proposed to undergo orthogonalization (later called pattern separation) in the dentate gyrus (McNaughton and Morris, 1987; Hasselmo and Wyble, 1997; Treves and Rolls, 1994), and then to be stored in a recurrent auto-associative memory in region CA3 that could mediate pattern completion (McNaughton and Morris, 1987; Hasselmo and Wyble, 1997; Treves and Rolls, 1994). Finally, region CA1 would map the stored patterns back to the input.

One major criticism of these early models is their relatively small capacity relative to the number of neurons, and also the lack of a representational structure to match space and time in the world. Another concerns the simple nature of the arbitrary vector representation, which does not reflect

any hierarchical representation of information, though that could be feasible in a vector representations such as those generated by convolution (Eliasmith et al., 2012; Plate, 1995). These models also lack many of the elements of neural dynamics, such as network and cellular oscillations and dendritic subthreshold dynamics.

Other neural models of memory focus on the role of neurons with specific functional properties, such as place cells, grid cells and head direction cells. These models have address the potential functional role of these subtypes in representing a spatial environment. Models of this type have focused on a number of functions, including: 1.) guidance of behavior to a specific goal location from a variety of start locations using place cells (Redish and Touretzky, 1998; Burgess et al., 1997; Arleo et al., 2004), or using grid cells (Erdem et al., 2015; Erdem and Hasselmo, 2014; Erdem and Hasselmo, 2012; Kubie and Fenton, 2012), 2.) the encoding and retrieval of previously encoded trajectories in episodic memory (Hasselmo and Eichenbaum, 2005; Hasselmo, 2012; Hasselmo, 2009), or 3.) the transformation between egocentric and allocentric representations of environmental features (Byrne et al., 2007; Bicanski and Burgess, 2018; Alexander et al., 2023; Sheynikhovich et al., 2009). Merging these different functions in a single model would be desirable.

Models that contain functional subtypes will automatically resemble those properties of the biological data, but their function for practical behavior tends to be somewhat limited. In particular, many of these models require input that has already been coded in terms of specific aspects of spatial location or velocity, though a few have been designed to respond on the basis of direct egocentric sensory input (Sheynikhovich et al., 2009; Arleo et al., 2004). The model of

episodic memory as a spatiotemporal trajectory effectively encodes and retrieves continuous trajectories through an environment and associated features of events at different positions along the trajectory. However, this model must be modified to account for later data, as the initial version of the model used grid cell input to generate place cells and also used speed-modulated direction cells that depend on movement direction rather than head direction (Hasselmo, 2012; Hasselmo, 2009).

The models that start with functional cell types such as place cells and grid cells often do not account for detailed biophysical dynamics of individual neuronal conductances. More biophysically detailed models simulate the membrane conductances and single cell membrane potential dynamics of neurons (Traub et al., 2022; Kelley et al., 2021; Wallenstein and Hasselmo, 1997; Traub et al., 1992; Sutton and Ascoli, 2021). However, the computational demands of these biophysical models make it very difficult to simulate a wide range of functional cell types or to simulate the guidance of complex behaviors.

A different class of models uses large multi-layer systems of neuron-like elements with an implementation of an error-correcting learning rule such backpropagation of error in deep learning (Banino et al., 2018) or contrastive Hebbian learning (O'Reilly and Munakata, 2000; Naud and Sprekeler, 2018). These models have the advantage of not using an a priori representations of functional cell types, but instead generating these functional cell types within the model (Banino et al., 2018). These models also have the advantage of using a more realistic egocentric input of visual input from the environment, rather than a pre-coded neural representation. Because these models generate behavioral output that usually exceeds the

capabilities of most hand-wired models, this approach is considered to have great potential for advancing the understanding of neural representations. However, so far the internal dynamics of these models have not been easily interpretable to yield guidance concerning the internal dynamics that allow for the function of the models. The distributed code is difficult to decipher and does not seem to exhibit features of symbolic representations such as compositionality and productivity, that would allow breaking down their functional processes into interpretable rules and elements (Do and Hasselmo, 2021).

In summary, the models that best simulate behavior using error-correcting learning rules do not provide interpretable internal dynamics, the models that directly simulate a range of functional cell types are more limited in their behavioral function and still do not address the complex dynamics of membrane potentials and membrane currents, whereas the models containing the biophysical detail of membrane currents have not been used to guide behavior or to generate simulations of most functional cell types.

Need to explore broader variety of models

There is a need for better models of neural circuits. Given the limited ability of existing models to demonstrate the computational relevance of many aspects of biophysical data, one can consider the question of what would a more effective model look like? What modifications could result in a revolution in neural memory models. The dimensions of possible models are actually enormous, and existing models have only explored a tiny fraction of this space (Hasselmo et al., 2021). Some ideas are presented here for exploring different areas of model space.

1. Phase coding. The use of temporal coding by the phase of spiking activity could provide advances over the dominant use of firing rate as the neuronal code. Most of the models described above use vectors with a continuous change in value that represents a continuous change in firing rate. These models do not directly account for the temporal coding of space and time that appears in multiple studies as theta phase precession (O'Keefe and Recce, 1993; Skaggs et al., 1996; Hafting et al., 2008; Climer et al., 2013), nor do they account for the rhythmic coding of running speed (Hinman et al., 2016; Dannenberg et al., 2020). Place cells and grid cells have been effectively modeled with phase coding based on oscillatory interference (Burgess, 2008; O'Keefe and Recce, 1993), but these models have not been the focus for most researchers.

2. Dendritic processing. Another underexplored area concerns the dynamics of dendritic processing, including the potential role of phase shifts within dendritic compartments (Vaidya and Johnston, 2013; Kelley et al., 2021), which could be modulated by synaptic activation of metabotropic receptors. Previous models of dendritic H current have focused on how these currents normalize timing of synaptic potential peaks (Magee, 1999; Vaidya and Johnston, 2013), but these currents could instead play a role in generating heterogeneity of phase based on interaction with other phase shifting inputs and matching to other dendritic inputs (Alexander et al., 2023). For example, one set of phase shifts could represent the relation between two features within an object (angle and distance) and an external phase shift could represent current object angle and current viewing angle.

3. Dendritic coding of transformations. Dendritic processing via phase shifts could allow transformations to be coded at a single neuron level rather than a network level. Many models of coordinate transformations use gain modulation (Bicanski and Burgess, 2018), in which different

populations of neurons code different transformations and a gating input (such as head direction) modulates the selection of different populations, requiring large numbers of neurons. In contrast, coding of transformations by individual neurons could allow coding of a broader range of transformations, and more flexible modulation of individual neuron transformations (Alexander et al., 2023).

4. Transmission of memory representations. Another appealing feature would be to have memory representations that can be easily transmitted between neurons, so that instead of requiring a long-term synaptic modification they could instead be rapidly transmitted. One possible way of doing this would be to have neural phase codes that can be transmitted between different neurons to activate cellular mechanisms that regulate phase within individual neurons. For example, neural activity could phosphorylate the H current or potassium currents in such a manner to change their time constant (Chen et al., 2001). The H current has been shown to shift the frequency and the phase of neural activity in dendrites (Vaidya and Johnston, 2013; Kelley et al., 2021). Changes in these currents could underlie alterations in neuronal responses such as those contributing to place cell firing (Bittner et al., 2015).

5. Analogy with computer animation. Beyond the details of cellular representations, there could be more focus on the nature of neural representations for flexibly representing all elements of an existing world. There could be inspiration from the framework used in computer animation. For example, the use of matrix implementations of two dimensional bezier surfaces (Sederberg, 2012) or non-uniform rational beta-splines (NURBS) (Liu and Wang, 2002) to generate the elements of an allocentric scene and the projection transformation into an egocentric view.

Words versus equations: verbal hypotheses versus computational models

This chapter has focused on a review of neural data and computational models. However, inspired by the interaction of neuroscientists and philosophers at the meeting in Tucson that prompted this edited volume, this section will venture out of the neuroscience expertise to discuss verbal hypotheses versus computational models.

The conference in Tucson was titled Time, Space and Memory, similar to elements of many other conferences and the title of this chapter. However, one could argue that neuroscience must eventually outgrow verbal terms such as memory and associated terms such as episodic memory, semantic memory, and working memory, and to develop more sophisticated representations for words such as space and time.

Science is ultimately a quest to create new symbolic languages of description that transcend prior verbal descriptions. Mature scientific fields such as Physics have a mathematical language of theories that completely supplant and transcend verbal descriptions. Words can be used to communicate and teach about physical theories, but they are in no way a component of the theory itself. The accepted scientific theories are identical regardless of the inexact words or even variations in mathematical notation being used to describe it. Multiple theories ranging from Newtonian mechanics to the Schrodinger equations for atomic structures can be described entirely in equations that accurately guide. In most cases, the attempt to describe these theories in words alone usually introduces inaccuracies or misunderstandings. Similarly, the structure of molecules ranging from simple compounds to functional proteins to the genetic code in DNA can be described with a sequence of letters that need no words from natural language. The sequence and quantitative characteristics of each the sequence of amino acids in a protein and their

structure and interaction with other proteins or small molecules can be functionally modeled,
again without words.

At this point, systems neuroscience and psychology are still too immature as fields to have many
accepted theories that transcend words. Theories at the single cell level can do this, using the
mathematical framework of Rall and others to describe the membrane potential interactions
across the structure of dendrites and axons, and using the Hodgkin-Huxley framework to
describe the dynamics of voltage-sensitive and calcium-sensitive channels. Cellular
neurophysiology is a mature science with theories that transcend words. However, systems
neuroscience and behavioral neuroscience still suffer from the promiscuity of words to have
multiple overlapping or nonoverlapping meanings.

In an immature field such as neuroscience, words can describe experimental results for which
there is not yet a quantitative theoretical framework. And the names of researchers can provide
another type of symbolic representation for forming an interlocking web of collaborations and
dates that give some structure to the broad mass of neuroscience data. But sometimes scientific
questions become overshadowed by disagreements over definitions of terms.

In the history of science, many domains of inquiry were initially domains of natural philosophy.
Then, as empirical scientific inquiry grew more sophisticated within individual fields, the
empirical components of individual fields of inquiry sequentially separated from philosophy and
became independent disciplines. In some cases, the split seems to occur when the empirical data
from experiments and the scientific theories about this data took a form that was independent

from the verbal description of questions in the field. The philosophical questions remain in many areas of science, but the empirical theories in physics, chemistry and molecular biology have moved beyond verbal descriptions.

Replacing words

How do we replace words in systems neuroscience? Based on examples from other fields such as physics, astronomy, chemistry and engineering, the answer is to develop mathematical theories that quantitatively account for the full range of data being addressed. What we currently inaccurately call episodic memory and semantic memory and working memory could eventually be addressed with a broad continuum model with multiple time scales and spatial scales that cannot be differentiated into these primitive terms. Or conversely, a theoretical framework might allow quantization of the dynamics of neural substrates that results in precise mathematical definitions of a specific number of memory elements that have no clear relationship to our current primitive terms. Terms like episodic, semantic and working memory could be like the terms of air, water, fire and earth used in early science, and could be replaced by an orderly periodic table of neural memory dynamics that predicts and guides quantitative research on memory.

How do we defined the structure of knowledge addressed in neuroscience? Unfortunately, this is a daunting task as the scope of neuroscience is not just the empirical data, but also the scope of knowledge represented in the human brain. These internal representations include everything that we can understand about the universe. In the simplest terms, the field addresses how the nervous system guides behavior. On the one hand, we can at least focus on a single system in the body,

the nervous system, so we have a physically contained substrate. On the behavioral side, we could focus like a behaviorist on only the physical manifestations of behavior – our movements in the world. However, this immediately expands into a description of everything, because we can utter words and write mathematical equations that attempt to describe our understanding of everything in the known universe. Thus the domain of neuroscience is not everything in the universe, but the domain does include our human understanding of everything. In order to model how the nervous system guides behavior, we need to model how the nervous system represents everything an individual can know about the universe, including a representation of the conscious self as a discrete entity interacting with the external world and planning future behavior based on past memories (Hasselmo, 2010). Ultimately, a model of memory function in cortical neural circuits must provide a framework for understanding how neurons can encode a memory representation of everything that a human can think about and remember. This is clearly a challenge, but an exciting challenge.

Bibliography

- Aggleton JP, Hunt PR and Rawlins JN. (1986) The effects of hippocampal lesions upon spatial and non-spatial tests of working memory. *Behav Brain Res* 19(2): 133-146.
- Aggleton JP, Neave N, Nagle S and Hunt PR. (1995) A comparison of the effects of anterior thalamic, mamillary body and fornix lesions on reinforced spatial alternation. *Behav Brain Res* 68(1): 91-101.
- Ainge JA, van der Meer MA, Langston RF and Wood ER. (2007) Exploring the role of context-dependent hippocampal activity in spatial alternation behavior. *Hippocampus* 17(10): 988-1002.
- Alexander AS, Carstensen LC, Hinman JR, Raudies F, Chapman GW and Hasselmo ME. (2020) Egocentric boundary vector tuning of the retrosplenial cortex. *Sci Adv* 6(8): eaaz2322.
- Alexander AS and Nitz DA. (2015) Retrosplenial cortex maps the conjunction of internal and external spaces. *Nature Neuroscience* 18: 1143-1151.
- Alexander AS and Nitz DA. (2017) Spatially Periodic Activation Patterns of Retrosplenial Cortex Encode Route Sub-spaces and Distance Traveled. *Current biology: CB* 27: 1551-1560.e1554.
- Alexander AS, Rangel LM, Tingley D and Nitz DA. (2018) Neurophysiological signatures of temporal coordination between retrosplenial cortex and the hippocampal formation. *Behav Neurosci* 132(5): 453-468.
- Alexander AS, Robinson JC, Stern CE and Hasselmo ME. (2023) Gated transformations from egocentric to allocentric reference frames involving retrosplenial cortex, entorhinal cortex, and hippocampus. *Hippocampus* 33(5): 465-487.
- Arleo A, Smeraldi F and Gerstner W. (2004) Cognitive navigation based on nonuniform Gabor space sampling, unsupervised growing networks, and reinforcement learning. *IEEE Trans Neural Netw* 15(3): 639-652.
- Banino A, Barry C, Uria B, Blundell C, Lillicrap T, Mirowski P, Pritzel A, et al. (2018) Vector-based navigation using grid-like representations in artificial agents. *Nature* 557(7705): 429-433.
- Barry C, Hayman R, Burgess N and Jeffery KJ. (2007) Experience-dependent rescaling of entorhinal grids. *Nat Neurosci* 10(6): 682-684.
- Barry C, Lever C, Hayman R, Hartley T, Burton S, O'Keefe J, Jeffery K, et al. (2006) The boundary vector cell model of place cell firing and spatial memory. *Rev Neurosci* 17(1-2): 71-97.
- Bicanski A and Burgess N. (2018) A neural-level model of spatial memory and imagery. *Elife* 7: e33752.
- Bittner KC, Grienberger C, Vaidya SP, Milstein AD, Macklin JJ, Suh J, Tonegawa S, et al. (2015) Conjunctive input processing drives feature selectivity in hippocampal CA1 neurons. *Nat Neurosci* 18(8): 1133-1142.
- Boyle A. (2021) Remembering events and representing time. *Synthese* 199: 2505-2524.
- Brandon MP, Bogaard AR, Libby CP, Connerney MA, Gupta K and Hasselmo ME. (2011) Reduction of theta rhythm dissociates grid cell spatial periodicity from directional tuning. *Science* 332(6029): 595-599.
- Brandon MP, Bogaard AR, Schultheiss NW and Hasselmo ME. (2013) Segregation of cortical head direction cell assemblies on alternating theta cycles. *Nat Neurosci* 16(6): 739-748.

- Brandon MP, Koenig J, Leutgeb JK and Leutgeb S. (2014) New and distinct hippocampal place codes are generated in a new environment during septal inactivation. *Neuron* 82(4): 789-796.
- Brioni JD, Decker MW, Gamboa LP, Izquierdo I and McGaugh JL. (1990) Muscimol injections in the medial septum impair spatial learning. *Brain Res* 522(2): 227-234.
- Brown EN, Frank LM, Tang D, Quirk MC and Wilson MA. (1998) A statistical paradigm for neural spike train decoding applied to position prediction from ensemble firing patterns of rat hippocampal place cells. *J Neurosci* 18(18): 7411-7425.
- Brown TI, Hasselmo ME and Stern CE. (2014) A high-resolution study of hippocampal and medial temporal lobe correlates of spatial context and prospective overlapping route memory. *Hippocampus* 24(7): 819-839.
- Brown TI, Ross RS, Keller JB, Hasselmo ME and Stern CE. (2010) Which way was I going? Contextual retrieval supports the disambiguation of well learned overlapping navigational routes. *J Neurosci* 30(21): 7414-7422.
- Brown TI and Stern CE. (2014) Contributions of medial temporal lobe and striatal memory systems to learning and retrieving overlapping spatial memories. *Cereb Cortex* 24(7): 1906-1922.
- Buetfering C, Allen K and Monyer H. (2014) Parvalbumin interneurons provide grid cell-driven recurrent inhibition in the medial entorhinal cortex. *Nat Neurosci* 17(5): 710-718.
- Burgess N. (2008) Grid cells and theta as oscillatory interference: theory and predictions. *Hippocampus* 18(12): 1157-1174.
- Burgess N, Barry C and O'Keefe J. (2007) An oscillatory interference model of grid cell firing. *Hippocampus* 17(9): 801-812.
- Burgess N, Donnett JG, Jeffery KJ and O'Keefe J. (1997) Robotic and neuronal simulation of the hippocampus and rat navigation. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 352(1360): 1535-1543.
- Burgess N, Jackson A, Hartley T and O'Keefe J. (2000) Predictions derived from modelling the hippocampal role in navigation. *Biol Cybern* 83(3): 301-312.
- Buzsaki G and Tingley D. (2018) Space and Time: The Hippocampus as a Sequence Generator. *Trends Cogn Sci* 22(10): 853-869.
- Byrne P, Becker S and Burgess N. (2007) Remembering the past and imagining the future: a neural model of spatial memory and imagery. *Psychol Rev* 114(2): 340-375.
- Cai DJ, Aharoni D, Shuman T, Shobe J, Biane J, Song W, Wei B, et al. (2016) A shared neural ensemble links distinct contextual memories encoded close in time. *Nature* 534(7605): 115-118.
- Carstensen LC, Alexander AS, Chapman GW, Lee A, J. and Hasselmo ME. (2021) Neural responses in retrosplenial cortex associated with environmental alterations. *iScience* 24: 103377.
- Chen G, Manson D, Cacucci F and Wills TJ. (2016) Absence of Visual Input Results in the Disruption of Grid Cell Firing in the Mouse. *Current Biology* 26(17): 2335-2342.
- Chen S, Wang J and Siegelbaum SA. (2001) Properties of hyperpolarization-activated pacemaker current defined by coassembly of HCN1 and HCN2 subunits and basal modulation by cyclic nucleotide. *J Gen Physiol* 117(5): 491-504.
- Chrastil ER, Sherrill KR, Hasselmo ME and Stern CE. (2015) There and Back Again: Hippocampus and Retrosplenial Cortex Track Homing Distance during Human Path

- Integration. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 35(46): 15442-15452.
- Chrastil ER, Sherrill KR, Hasselmo ME and Stern CE. (2016) Which way and how far? Tracking of translation and rotation information for human path integration. *Human Brain Mapping* 37: 3636-3655.
- Chrobak JJ, Stackman RW and Walsh TJ. (1989) Intraseptal administration of muscimol produces dose-dependent memory impairments in the rat. *Behavioral and Neural Biology* 52: 357-369.
- Clancy KB, Orsolic I and Mrosovsky N. (2019) Locomotion-dependent remapping of distributed cortical networks. *Nat Neurosci* 22(5): 778-786.
- Climmer JR, Newman EL and Hasselmo ME. (2013) Phase coding by grid cells in unconstrained environments: two-dimensional phase precession. *Eur J Neurosci* 38(4): 2526-2541.
- Conway MA. (2009) Episodic memories. *Neuropsychologia* 47(11): 2305-2313.
- Dannenberg H, Kelley C, Hoyland A, Monaghan CK and Hasselmo ME. (2019) The Firing Rate Speed Code of Entorhinal Speed Cells Differs across Behaviorally Relevant Time Scales and Does Not Depend on Medial Septum Inputs. *J Neurosci* 39(18): 3434-3453.
- Dannenberg H, Lazaro H, Nambiar P, Hoyland A and Hasselmo ME. (2020) Effects of visual inputs on neural dynamics for coding of location and running speed in medial entorhinal cortex. *Elife* 9: e62500.
- DeLuca J and Cicerone KD. (1991) Confabulation following aneurysm of the anterior communicating artery. *Cortex* 27(3): 417-423.
- Deshmukh SS and Knierim JJ. (2011) Representation of non-spatial and spatial information in the lateral entorhinal cortex. *Front Behav Neurosci* 5: 69.
- Deshmukh SS and Knierim JJ. (2013) Influence of local objects on hippocampal representations: Landmark vectors and memory. *Hippocampus*.
- Do Q and Hasselmo ME. (2021) Neural circuits and symbolic processing. *Neurobiol Learn Mem* 186: 107552.
- Douchamps V, Jeewajee A, Blundell P, Burgess N and Lever C. (2013) Evidence for encoding versus retrieval scheduling in the hippocampus by theta phase and acetylcholine. *J Neurosci* 33(20): 8689-8704.
- Eichenbaum H. (2014) Time cells in the hippocampus: a new dimension for mapping memories. *Nat Rev Neurosci* 15(11): 732-744.
- Eichenbaum H, Dudchenko P, Wood E, Shapiro M and Tanila H. (1999) The hippocampus, memory, and place cells: is it spatial memory or a memory space? *Neuron* 23(2): 209-226.
- Eichenbaum H, Stewart C and Morris RG. (1990) Hippocampal representation in place learning. *J Neurosci* 10(11): 3531-3542.
- Eliasmith C, Stewart TC, Choo X, Bekolay T, DeWolf T, Tang Y and Rasmussen D. (2012) A large-scale model of the functioning brain. *Science* 338(6111): 1202-1205.
- Erdem UM and Hasselmo M. (2012) A goal-directed spatial navigation model using forward trajectory planning based on grid cells. *Eur J Neurosci* 35(6): 916-931.
- Erdem UM and Hasselmo ME. (2014) A biologically inspired hierarchical goal directed navigation model. *J Physiol Paris* 108(1): 28-37.
- Erdem UM, Milford MJ and Hasselmo ME. (2015) A hierarchical model of goal directed navigation selects trajectories in a visual environment. *Neurobiol Learn Mem* 117: 109-121.

- 785 Feng T, Silva D and Foster DJ. (2015) Dissociation between the experience-dependent
786 development of hippocampal theta sequences and single-trial phase precession. *J*
787 *Neurosci* 35(12): 4890-4902.
- 788 Fenton AA, Kao HY, Neymotin SA, Olypher A, Vayntrub Y, Lytton WW and Ludvig N. (2008)
789 Unmasking the CA1 ensemble place code by exposures to small and large environments:
790 more place cells and multiple, irregularly arranged, and expanded place fields in the
791 larger space. *J Neurosci* 28(44): 11250-11262.
- 792 Fenton AA and Muller RU. (1998) Place cell discharge is extremely variable during individual
793 passes of the rat through the firing field. *Proc. Natl. Acad. Sci. USA* 95(6): 3182-3187.
- 794 Ferbinteanu J and Shapiro ML. (2003) Prospective and retrospective memory coding in the
795 hippocampus. *Neuron* 40(6): 1227-1239.
- 796 Foster DJ and Wilson MA. (2007) Hippocampal theta sequences. *Hippocampus* 17(11): 1093-
797 1099.
- 798 Frank LM, Brown EN and Wilson M. (2000) Trajectory encoding in the hippocampus and
799 entorhinal cortex. *Neuron* 27(1): 169-178.
- 800 Giocomo LM, Stensola T, Bonnevie T, Van Cauter T, Moser MB and Moser EI. (2014)
801 Topography of head direction cells in medial entorhinal cortex. *Curr Biol* 24(3): 252-262.
- 802 Gofman X, Tocker G, Weiss S, Boccara CN, Lu L, Moser MB, Moser EI, et al. (2019)
803 Dissociation between Postrhinal Cortex and Downstream Parahippocampal Regions in
804 the Representation of Egocentric Boundaries. *Curr Biol* 29(16): 2751-2757 e2754.
- 805 Goodridge JP and Taube JS. (1997) Interaction between the postsubiculum and anterior thalamus
806 in the generation of head direction cell activity. *J Neurosci* 17(23): 9315-9330.
- 807 Green EJ and Schellenberg S. (2017) Spatial perception: The perspectival aspect of perception.
808 *Philosophy Compass* e12472.
- 809 Griffin AL, Eichenbaum H and Hasselmo ME. (2007) Spatial representations of hippocampal
810 CA1 neurons are modulated by behavioral context in a hippocampus-dependent memory
811 task. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*
812 27(9): 2416-2423.
- 813 Hafting T, Fyhn M, Bonnevie T, Moser MB and Moser EI. (2008) Hippocampus-independent
814 phase precession in entorhinal grid cells. *Nature* 453(7199): 1248-1252.
- 815 Hafting T, Fyhn M, Molden S, Moser M-B and Moser EI. (2005) Microstructure of a spatial map
816 in the entorhinal cortex. *Nature* 436: 801-806.
- 817 Hallock HL, Arreola AC, Shaw CL and Griffin AL. (2013) Dissociable roles of the dorsal
818 striatum and dorsal hippocampus in conditional discrimination and spatial alternation T-
819 maze tasks. *Neurobiology of Learning and Memory* 100: 108-116.
- 820 Hartley T, Burgess N, Lever C, Cacucci F and O'Keefe J. (2000) Modeling place fields in terms
821 of the cortical inputs to the hippocampus. *Hippocampus* 10(4): 369-379.
- 822 Hartley T, Lever C, Burgess N and O'Keefe J. (2014) Space in the brain: how the hippocampal
823 formation supports spatial cognition. *Philos Trans R Soc Lond B Biol Sci* 369(1635):
824 20120510.
- 825 Hasselmo ME. (2005) What is the function of hippocampal theta rhythm?--Linking behavioral
826 data to phasic properties of field potential and unit recording data. *Hippocampus* 15(7):
827 936-949.
- 828 Hasselmo ME. (2006) The role of acetylcholine in learning and memory. *Curr Opin Neurobiol*
829 16(6): 710-715.

- 830 Hasselmo ME. (2008) Grid cell mechanisms and function: contributions of entorhinal persistent
 831 spiking and phase resetting. *Hippocampus* 18(12): 1213-1229.
- 832 Hasselmo ME. (2009) A model of episodic memory: mental time travel along encoded
 833 trajectories using grid cells. *Neurobiol Learn Mem* 92(4): 559-573.
- 834 Hasselmo ME. (2010) Consciousness and neural time travel. In: Perry E, Collerton D, LeBeau
 835 FE, et al. (eds) *New Horizons in the Neuroscience of Consciousness*. Philadelphia, PA:
 836 John Benhamins Publishing Company, 73-80.
- 837 Hasselmo ME. (2012) *How we remember: Brain mechanisms of episodic memory*, Cambridge,
 838 MA: MIT Press.
- 839 Hasselmo ME. (2015) Remembering by index and content: Response to Sarah Robins. .
 840 *Philosophical Psychology* 26(8): 916-919.
- 841 Hasselmo ME, Alexander AS, Hoyland A, Robinson JC, Bezair MJ, Chapman GW,
 842 Saudargiene A, et al. (2021) The Unexplored Territory of Neural Models: Potential
 843 Guides for Exploring the Function of Metabotropic Neuromodulation. *Neuroscience* 456:
 844 143-158.
- 845 Hasselmo ME, Bodelón C and Wyble BP. (2002) A proposed function for hippocampal theta
 846 rhythm: separate phases of encoding and retrieval enhance reversal of prior learning.
 847 *Neural computation* 14(4): 793-817.
- 848 Hasselmo ME and Eichenbaum H. (2005) Hippocampal mechanisms for the context-dependent
 849 retrieval of episodes. *Neural Netw* 18(9): 1172-1190.
- 850 Hasselmo ME, Giocomo LM, Brandon MP and Yoshida M. (2010) Cellular dynamical
 851 mechanisms for encoding the time and place of events along spatiotemporal trajectories
 852 in episodic memory. *Behav Brain Res* 215(2): 261-274.
- 853 Hasselmo ME and Wyble BP. (1997) Free recall and recognition in a network model of the
 854 hippocampus: simulating effects of scopolamine on human memory function. *Behav*
 855 *Brain Res* 89(1-2): 1-34.
- 856 Heys JG and Dombeck DA. (2018) Evidence for a subcircuit in medial entorhinal cortex
 857 representing elapsed time during immobility. *Nat Neurosci* 21(11): 1574-1582.
- 858 Heys JG, Rangarajan KV and Dombeck DA. (2014) The functional micro-organization of grid
 859 cells revealed by cellular-resolution imaging. *Neuron* 84(5): 1079-1090.
- 860 Hinman JR, Brandon MP, Climer JR, Chapman GW and Hasselmo ME. (2016) Multiple
 861 Running Speed Signals in Medial Entorhinal Cortex. *Neuron* 91(3): 666-679.
- 862 Hinman JR, Chapman GW and Hasselmo ME. (2019) Neuronal representation of environmental
 863 boundaries in egocentric coordinates. *Nat Commun* 10(1): 2772.
- 864 Howard MW and Eichenbaum H. (2013) The hippocampus, time, and memory across scales. *J*
 865 *Exp Psychol Gen* 142(4): 1211-1230.
- 866 Howard MW, MacDonald CJ, Tiganj Z, Shankar KH, Du Q, Hasselmo ME and Eichenbaum H.
 867 (2014) A unified mathematical framework for coding time, space, and sequences in the
 868 hippocampal region. *J Neurosci* 34(13): 4692-4707.
- 869 Hoydal OA, Skytoen ER, Andersson SO, Moser MB and Moser EI. (2019) Object-vector coding
 870 in the medial entorhinal cortex. *Nature* 568(7752): 400-404.
- 871 Huxter J, Burgess N and O'Keefe J. (2003) Independent rate and temporal coding in
 872 hippocampal pyramidal cells. *Nature* 425(6960): 828-832.
- 873 Huxter JR, Senior TJ, Allen K and Csicsvari J. (2008) Theta phase-specific codes for two-
 874 dimensional position, trajectory and heading in the hippocampus. *Nat Neurosci* 11(5):
 875 587-594.

- 876 Jeewajee A, Barry C, Douchamps V, Manson D, Lever C and Burgess N. (2014) Theta phase
877 precession of grid and place cell firing in open environments. *Philos Trans R Soc Lond B*
878 *Biol Sci* 369(1635): 20120532.
- 879 Jeewajee A, Barry C, O'Keefe J and Burgess N. (2008) Grid cells and theta as oscillatory
880 interference: electrophysiological data from freely moving rats. *Hippocampus* 18(12):
881 1175-1185.
- 882 Jeffery KJ, Donnett JG and O'Keefe J. (1995) Medial septal control of theta-correlated unit firing
883 in the entorhinal cortex of awake rats. *Neuroreport* 6(16): 2166-2170.
- 884 Johnson A and Redish AD. (2007) Neural ensembles in CA3 transiently encode paths forward of
885 the animal at a decision point. *J Neurosci* 27(45): 12176-12189.
- 886 Kay K, Chung JE, Sosa M, Schor JS, Karlsson MP, Larkin MC, Liu DF, et al. (2020) Constant
887 Sub-second Cycling between Representations of Possible Futures in the Hippocampus.
888 *Cell* 180(3): 552-567 e525.
- 889 Kelley C, Dura-Bernal S, Neymotin SA, Antic SD, Carnevale NT, Migliore M and Lytton WW.
890 (2021) Effects of I(h) and TASK-like shunting current on dendritic impedance in layer 5
891 pyramidal-tract neurons. *J Neurophysiol* 125(4): 1501-1516.
- 892 Kinsky NR, Mau W, Sullivan DW, Levy SJ, Ruesch EA and Hasselmo ME. (2020) Trajectory-
893 modulated hippocampal neurons persist throughout memory-guided navigation. *Nat*
894 *Commun* 11(1): 2443.
- 895 Koenig J, Linder AN, Leutgeb JK and Leutgeb S. (2011) The spatial periodicity of grid cells is
896 not sustained during reduced theta oscillations. *Science* 332(6029): 592-595.
- 897 Kraus BJ, Brandon MP, Robinson RJ, 2nd, Connerney MA, Hasselmo ME and Eichenbaum H.
898 (2015) During Running in Place, Grid Cells Integrate Elapsed Time and Distance Run.
899 *Neuron* 88(3): 578-589.
- 900 Kraus BJ, Robinson RJ, 2nd, White JA, Eichenbaum H and Hasselmo ME. (2013) Hippocampal
901 "Time Cells": Time versus Path Integration. *Neuron* 78(6): 1090-1101.
- 902 Kropff E, Carmichael JE, Moser MB and Moser EI. (2015) Speed cells in the medial entorhinal
903 cortex. *Nature* 523(7561): 419-424.
- 904 Kubie JL and Fenton AA. (2012) Linear look-ahead in conjunctive cells: an entorhinal
905 mechanism for vector-based navigation. *Front Neural Circuits* 6: 20.
- 906 LaChance PA and Taube JS. (2023) A model for transforming egocentric views into goal-
907 directed behavior. *Hippocampus* In press (this issue).
- 908 LaChance PA, Todd TP and Taube JS. (2019) A sense of space in postrhinal cortex. *Science*
909 365(6449): eaax4192.
- 910 Lever C, Burton S, Jeewajee A, O'Keefe J and Burgess N. (2009) Boundary vector cells in the
911 subiculum of the hippocampal formation. *The Journal of Neuroscience* 29: 9771-9777.
- 912 Lever C, Wills T, Cacucci F, Burgess N and O'Keefe J. (2002) Long-term plasticity in
913 hippocampal place-cell representation of environmental geometry. *Nature* 416(6876): 90-
914 94.
- 915 Levy SJ, Kinsky NR, Mau W, Sullivan DW and Hasselmo ME. (2021) Hippocampal spatial
916 memory representations in mice are heterogeneously stable. *Hippocampus* 31(3): 244-
917 260.
- 918 Levy WB. (1996) A sequence predicting CA3 is a flexible associator that learns and uses context
919 to solve hippocampal-like tasks. *Hippocampus* 6(6): 579-590.
- 920 Lipton PA, White JA and Eichenbaum H. (2007) Disambiguation of overlapping experiences by
921 neurons in the medial entorhinal cortex. *J Neurosci* 27(21): 5787-5795.

- 922 Liu L and Wang G. (2002) Explicit matrix representation for NURBS curves and surfaces.
 923 *Computer Aided Geometric Design* 19: 409-419.
- 924 Liu Y, Levy S, Mau W, Geva N, Rubin A, Ziv Y, Hasselmo M, et al. (2022) Consistent
 925 population activity on the scale of minutes in the mouse hippocampus. *Hippocampus*
 926 32(5): 359-372.
- 927 Liu Y, Tiganj Z, Hasselmo ME and Howard MW. (2019) A neural microcircuit model for a
 928 scalable scale-invariant representation of time. *Hippocampus* 29(3): 260-274.
- 929 MacDonald CJ, Carrow S, Place R and Eichenbaum H. (2013) Distinct hippocampal time cell
 930 sequences represent odor memories in immobilized rats. *J Neurosci* 33(36): 14607-
 931 14616.
- 932 MacDonald CJ, Lepage KQ, Eden UT and Eichenbaum H. (2011) Hippocampal "time cells"
 933 bridge the gap in memory for discontinuous events. *Neuron* 71(4): 737-749.
- 934 Magee JC. (1999) Dendritic Ih normalizes temporal summation in hippocampal CA1 neurons.
 935 *Nat Neurosci* 2(6): 508-514.
- 936 Mankin EA, Diehl GW, Sparks FT, Leutgeb S and Leutgeb JK. (2015) Hippocampal CA2
 937 activity patterns change over time to a larger extent than between spatial contexts.
 938 *Neuron* 85(1): 190-201.
- 939 Manns JR, Zilli EA, Ong KC, Hasselmo ME and Eichenbaum H. (2007) Hippocampal CA1
 940 spiking during encoding and retrieval: relation to theta phase. *Neurobiol Learn Mem*
 941 87(1): 9-20.
- 942 Mau W, Sullivan DW, Kinsky NR, Hasselmo ME, Howard MW and Eichenbaum H. (2018) The
 943 Same Hippocampal CA1 Population Simultaneously Codes Temporal Information over
 944 Multiple Timescales. *Current Biology* 28(10): 1499-1508.e1494.
- 945 Maurer AP, Cowen SL, Burke SN, Barnes CA and McNaughton BL. (2006) Phase precession in
 946 hippocampal interneurons showing strong functional coupling to individual pyramidal
 947 cells. *J Neurosci* 26(52): 13485-13492.
- 948 McNaughton BL, Barnes CA and O'Keefe J. (1983) The contributions of position, direction, and
 949 velocity to single unit-activity in the hippocampus of freely-moving rats. *Experimental*
 950 *Brain Research* 52(1): 41-49.
- 951 McNaughton BL, Mizumori SJ, Barnes CA, Leonard BJ, Marquis M and Green EJ. (1994)
 952 Cortical representation of motion during unrestrained spatial navigation in the rat. *Cereb*
 953 *Cortex* 4(1): 27-39.
- 954 McNaughton BL and Morris RGM. (1987) Hippocampal synaptic enhancement and information
 955 storage within a distributed memory system. *Trends Neurosci.* 10: 408-415.
- 956 Mitchell SJ, Rawlins JN, Steward O and Olton DS. (1982) Medial septal area lesions disrupt
 957 theta rhythm and cholinergic staining in medial entorhinal cortex and produce impaired
 958 radial arm maze behavior in rats. *J Neurosci* 2(3): 292-302.
- 959 Morris RG, Garrud P, Rawlins JN and O'Keefe J. (1982) Place navigation impaired in rats with
 960 hippocampal lesions. *Nature* 297(5868): 681-683.
- 961 Muller RU and Kubie JL. (1987) The effects of changes in the environment on the spatial firing
 962 of hippocampal complex-spike cells. *J Neurosci* 7(7): 1951-1968.
- 963 Muller RU, Kubie JL and Ranck JB, Jr. (1987) Spatial firing patterns of hippocampal complex-
 964 spike cells in a fixed environment. *J Neurosci* 7(7): 1935-1950.
- 965 Munn RGK, Mallory CS, Hardcastle K, Chetkovich DM and Giocomo LM. (2020) Entorhinal
 966 velocity signals reflect environmental geometry. *Nat Neurosci* 23(2): 239-251.
- 967 Murdock B. (2005) Storage and retrieval of serial-order information. *Memory* 13(3-4): 259-266.

- 968 Naud R and Sprekeler H. (2018) Sparse bursts optimize information transmission in a
 969 multiplexed neural code. *Proc Natl Acad Sci U S A* 115(27): E6329-E6338.
- 970 Nelson A and Mooney R. (2016) The Basal Forebrain and Motor Cortex Provide Convergent yet
 971 Distinct Movement-Related Inputs to the Auditory Cortex. *Neuron* 90(3): 635-648.
- 972 Niell CM and Stryker MP. (2010) Modulation of visual responses by behavioral state in mouse
 973 visual cortex. *Neuron* 65: 472-479.
- 974 Ning W, Bladon JH and Hasselmo ME. (2022) Complementary representations of time in the
 975 prefrontal cortex and hippocampus. *Hippocampus* 32(8): 577-596.
- 976 O'Keefe J. (1976) Place units in the hippocampus of the freely moving rat. *Exp Neurol* 51(1): 78-
 977 109.
- 978 O'Keefe J. (1990) A computational theory of the hippocampal cognitive map. *Prog. Brain Res.*
 979 83: 301-312.
- 980 O'Keefe J and Burgess N. (1996) Geometric determinants of the place fields of hippocampal
 981 neurons. *Nature* 381: 425-428.
- 982 O'Keefe J, Burgess N, Donnett JG, Jeffery KJ and Maguire EA. (1998) Place cells, navigational
 983 accuracy, and the human hippocampus. *Philos Trans R Soc Lond B Biol Sci* 353(1373):
 984 1333-1340.
- 985 O'Keefe J and Dostrovsky J. (1971) The hippocampus as a spatial map. Preliminary evidence
 986 from unit activity in the freely-moving rat. *Brain Res* 34: 171-175.
- 987 O'Keefe J and Nadel L. (1978) *The Hippocampus as a Cognitive Map*, Oxford : New York:
 988 Oxford University Press.
- 989 O'Keefe J, Nadel L, Keightley S and Kill D. (1975) Fornix lesions selectively abolish place
 990 learning in the rat. *Exp Neurol* 48(1): 152-166.
- 991 O'Keefe J and Recce ML. (1993) Phase relationship between hippocampal place units and the
 992 EEG theta rhythm. *Hippocampus* 3: 317-330.
- 993 O'Neill J, Boccara CN, Stella F, Schoenenberger P and Csicsvari J. (2017) Superficial layers of
 994 the medial entorhinal cortex replay independently of the hippocampus. *Science*
 995 355(6321): 184-188.
- 996 O'Reilly RC and Munakata Y. (2000) *Computational explorations in cognitive neuroscience:*
 997 *Understanding the mind by simulating the brain.*, Cambridge, MA: MIT Press.
- 998 Ormond J and O'Keefe J. (2022) Hippocampal place cells have goal-oriented vector fields during
 999 navigation. *Nature* 607(7920): 741-746.
- 1000 Pastalkova E, Itskov V, Amarasingham A and Buzsaki G. (2008) Internally generated cell
 1001 assembly sequences in the rat hippocampus. *Science* 321(5894): 1322-1327.
- 1002 Pérez-Escobar JA, Kornienko O, Latuske P, Kohler L and Allen K. (2016) Visual landmarks
 1003 sharpen grid cell metric and confer context specificity to neurons of the medial entorhinal
 1004 cortex. *Elife* 5: e16937.
- 1005 Phillips I. (2014) Experience of and in Time. *Philosophy Compass* 9(2): 131-144.
- 1006 Plate TA. (1995) Holographic reduced representations. *IEEE Trans Neural Netw* 6(3): 623-641.
- 1007 Poulter S, Lee SA, Dachtler J, Willis TJ and Lever C. (2021) Vector trace cells in the subiculum
 1008 of the hippocampal formation. *Nat. Neurosci.* 24(2): 266-275.
- 1009 Raudies F, Brandon MP, Chapman GW and Hasselmo ME. (2015) Head direction is coded more
 1010 strongly than movement direction in a population of entorhinal neurons. *Brain Res* 1621:
 1011 355-367.

- Raudies F and Hasselmo ME. (2015) Differences in Visual-Spatial Input May Underlie Different Compression Properties of Firing Fields for Grid Cell Modules in Medial Entorhinal Cortex. *PLoS Comput Biol* 11(11): e1004596.
- Rawlins JN, Feldon J and Gray JA. (1979) Septo-hippocampal connections and the hippocampal theta rhythm. *Exp Brain Res* 37(1): 49-63.
- Redish AD and Touretzky DS. (1998) The role of the hippocampus in solving the Morris water maze. *Neural Comput.* 10(1): 73-111.
- Rigotti M, Barak O, Warden MR, Wang X-J, Daw ND, Miller EK and Fusi S. (2013) The importance of mixed selectivity in complex cognitive tasks. *Nature* 497: 585-590.
- Robins SK. (2015) A Mechanism for Mental Time Travel? A Critical Review of Michael Hasselmo's How We Remember: Brain Mechanisms of Episodic Memory. . *Philosophical Psychology* 28: 903-915.
- Robinson JC, Brandon MP and Hasselmo ME. (2019) Identifying the contribution of medial septum cell types in generating the entorhinal grid cell code. *Soc. Neurosci. Abstr.* 45: 164-102.
- Rubin A, Geva N, Sheintuch L and Ziv Y. (2015) Hippocampal ensemble dynamics timestamp events in long-term memory. *Elife* 4: e12247.
- Rule ME, O'Leary T and Harvey CD. (2019) Causes and consequences of representational drift. *Curr. Opin. Neurobiol.* 58: 141-147.
- Salz DM, Tiganj Z, Khasnabish S, Kohley A, Sheehan D, Howard MW and Eichenbaum H. (2016) Time Cells in Hippocampal Area CA3. *J Neurosci* 36(28): 7476-7484.
- Sargolini F, Fyhn M, Hafting T, McNaughton BL, Witter MP, Moser MB and Moser EI. (2006) Conjunctive representation of position, direction, and velocity in entorhinal cortex. *Science* 312(5774): 758-762.
- Savelli F, Yoganarasimha D and Knierim JJ. (2008) Influence of boundary removal on the spatial representations of the medial entorhinal cortex. *Hippocampus* 18(12): 1270-1282.
- Schmidt R, Diba K, Leibold C, Schmitz D, Buzsaki G and Kempter R. (2009) Single-trial phase precession in the hippocampus. *J Neurosci* 29(42): 13232-13241.
- Sederberg TW. (2012) Computer Aided Geometric Design. <https://scholarsarchive.byu.edu/facpub/1>: BYU ScholarsArchive.
- Sherrill KR, Chrastil ER, Ross RS, Erdem UM, Hasselmo ME and Stern CE. (2015) Functional connections between optic flow areas and navigationally responsive brain regions during goal-directed navigation. *Neuroimage* 118: 386-396.
- Sherrill KR, Erdem UM, Ross RS, Brown TI, Hasselmo ME and Stern CE. (2013) Hippocampus and retrosplenial cortex combine path integration signals for successful navigation. *J Neurosci* 33(49): 19304-19313.
- Sheynikhovich D, Chavarriaga R, Strosslin T, Arleo A and Gerstner W. (2009) Is there a geometric module for spatial orientation? Insights from a rodent navigation model. *Psychol Rev* 116(3): 540-566.
- Skaggs WE, McNaughton BL, Wilson MA and Barnes CA. (1996) Theta phase precession in hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus* 6(2): 149-172.
- Solstad T, Boccara CN, Kropff E, Moser MB and Moser EI. (2008) Representation of geometric borders in the entorhinal cortex. *Science* 322(5909): 1865-1868.
- Steffenach HA, Witter M, Moser MB and Moser EI. (2005) Spatial memory in the rat requires the dorsolateral band of the entorhinal cortex. *Neuron* 45(2): 301-313.

- Stensola H, Stensola T, Solstad T, Froland K, Moser MB and Moser EI. (2012) The entorhinal grid map is discretized. *Nature* 492(7427): 72-78.
- Sutton NM and Ascoli GA. (2021) Spiking Neural Networks and Hippocampal Function: A Web-Accessible Survey of Simulations, Modeling Methods, and Underlying Theories. *Cogn Syst Res* 70: 80-92.
- Taube JS. (1995) Head direction cells recorded in the anterior thalamic nuclei of freely moving rats. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 15(1): 70-86.
- Taube JS, Kesslak JP and Cotman CW. (1992) Lesions of the rat postsubiculum impair performance on spatial tasks. *Behav Neural Biol* 57(2): 131-143.
- Taube JS, Muller RU and Ranck JB. (1990) Head-direction cells recorded from the postsubiculum in freely moving rats. I. Description and quantitative analysis. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 10(2): 420-435.
- Terada S, Sakurai Y, Nakahara H and Fujisawa S. (2017) Temporal and Rate Coding for Discrete Event Sequences in the Hippocampus. *Neuron* 94(6): 1248-1262 e1244.
- Touretzky DS and Redish AD. (1996) Theory of rodent navigation based on interacting representations of space. *Hippocampus* 6(3): 247-270.
- Traub R, Miles R and Buzsaki G. (1992) Computer simulation of carbachol-driven rhythmic population oscillations in the CA3 region of the in vitro rat hippocampus. *J. Physiol.* 451: 653-672.
- Traub RD, Whittington MA and Cunningham MO. (2022) Simulation of oscillatory dynamics induced by an approximation of grid cell output. *Rev Neurosci*.
- Treves A and Rolls ET. (1994) Computational analysis of the role of the hippocampus in memory. *Hippocampus* 4(3): 374-391.
- Tsao A, Sugar J, Lu L, Wang C, Knierim JJ, Moser MB and Moser EI. (2018) Integrating time from experience in the lateral entorhinal cortex. *Nature* 561(7721): 57-62.
- Tulving E. (1984) Precis of Elements of episodic memory. *Behav Brain Sci* 7: 223-268.
- Vaidya SP and Johnston D. (2013) Temporal synchrony and gamma-to-theta power conversion in the dendrites of CA1 pyramidal neurons. *Nature Neuroscience* 16: 1812-1820.
- van Wijngaarden JB, Babl SS and Ito HT. (2020) Entorhinal-retrosplenial circuits for allocentric-egocentric transformation of boundary coding. *Elife* 9: e59816.
- Wallenstein GV and Hasselmo ME. (1997) GABAergic modulation of hippocampal population activity: sequence learning, place field development, and the phase precession effect. *J Neurophysiol* 78(1): 393-408.
- Wang C, Chen X and Knierim JJ. (2020) Egocentric and allocentric representations of space in the rodent brain. *Curr. Opin. Neurobiol.* 60: 12-20.
- Wang C, Chen X, Lee H, Deshmukh SS, Yoganarasimha D, Savelli F and Knierim JJ. (2018) Egocentric coding of external items in the lateral entorhinal cortex. *Science* 362(6417): 945-949.
- Wang Y, Romani S, Lustig B, Leonardo A and Pastalkova E. (2015) Theta sequences are essential for internally generated hippocampal firing fields. *Nat Neurosci* 18(2): 282-288.
- Welday AC, Shlifer IG, Bloom ML, Zhang K and Blair HT. (2011) Cosine directional tuning of theta cell burst frequencies: evidence for spatial coding by oscillatory interference. *J Neurosci* 31(45): 16157-16176.

- 1102 Wells CE, Amos DP, Jeewajee A, Douchamps V, Rodgers J, O'Keefe J, Burgess N, et al. (2013)
 1103 Novelty and anxiolytic drugs dissociate two components of hippocampal theta in
 1104 behaving rats. *J Neurosci* 33(20): 8650-8667.
- 1105 Wikenheiser AM and Redish AD. (2015) Hippocampal theta sequences reflect current goals. *Nat*
 1106 *Neurosci* 18(2): 289-294.
- 1107 Wills TJ, Barry C and Cacucci F. (2012) The abrupt development of adult-like grid cell firing in
 1108 the medial entorhinal cortex. *Front Neural Circuits* 6: 21.
- 1109 Wilmerding LK, Kondratyev I, Ramirez S and Hasselmo ME. (2023) Route-dependent spatial
 1110 engram tagging in mouse dentate gyrus. *Neurobiol Learn Mem* 200: 107738.
- 1111 Winter SS, Clark BJ and Taube JS. (2015) Disruption of the head direction cell network impairs
 1112 the parahippocampal grid cell signal. *Science* 347(6224): 870-874.
- 1113 Wood ER, Dudchenko PA, Robitsek RJ and Eichenbaum H. (2000) Hippocampal neurons
 1114 encode information about different types of memory episodes occurring in the same
 1115 location. *Neuron* 27(3): 623-633.
- 1116 Ziv Y, Burns LD, Cocker ED, Hamel EO, Ghosh KK, Kitch LJ, Gamal AE, et al. (2013) Long-
 1117 term dynamics of CA1 hippocampal place codes. *Nat. Neurosci.* 16(3): 264-266.
- 1118 Zugaro MB, Monconduit L and Buzsaki G. (2005) Spike phase precession persists after transient
 1119 intrahippocampal perturbation. *Nat Neurosci* 8(1): 67-71.
- 1120 Zutshi I, Leutgeb JK and Leutgeb S. (2017) Theta sequences of grid cell populations can provide
 1121 a movement-direction signal. *Curr Opin Behav Sci* 17: 147-154.
- 1122