

Atiyah-Singer Index Theorem

Will Chuang

Leung Center for Cosmology and Particle Astrophysics, National Taiwan
University, Taipei, Taiwan 10617

June 5, 2012

1/10 Why we need Atiyah-Singer index theorem?

Reason 1

Original work on $(2+1)$ -dimensional gravity can be traced back to 1963, when Staruszkiewicz first described the behavior of static solutions with point sources.

1/10 Why we need Atiyah-Singer index theorem?

Reason 1

Original work on $(2+1)$ -dimensional gravity can be traced back to 1963, when Staruszkiewicz first described the behavior of static solutions with point sources.

Reason 2

Twenty years latter, Deser, Jackiw, 't Hooft, and Witten let this theory rebirth.

1/10 Why we need Atiyah-Singer index theorem?

Reason 1

Original work on $(2+1)$ -dimensional gravity can be traced back to 1963, when Staruszkiewicz first described the behavior of static solutions with point sources.

Reason 2

Twenty years latter, Deser, Jackiw, 't Hooft, and Witten let this theory rebirth.

Reason 3

Now days, there are several aspects still active:

classical point sources, closed timelike curves, the classical and quantum behavior of point particles, matter couplings, supergravity, asymptotic behavior and global charges, the radial gauge, topologically massive gravity, gauge theory, the construction of observables from topological field theories,and so on and so forth.

2/10 What is A-S index theorem?

Scientists describe the world by measuring quantities and forces that vary over time and space. The rules of nature are often expressed by formulas, called differential equations, involving their rates of change. Such formulas may have an "index," the number of solutions of the formulas minus the number of restrictions that they impose on the values of the quantities being computed. i.e. **The analytical index is equal to its topological index.** The Atiyah-Singer index theorem calculated this number in terms of the geometry of the surrounding space.

2/10 What is A-S index theorem?

Formal Description [Nakahara]

Let (E, D) be an elliptic complex over an m -dimensional compact manifold M without a boundary. The index of this complex is given by

$$\text{ind}(E, D) = (-1)^{m(m+1)/2} \int_M \text{ch}\left(\bigoplus_r (-1)^r E_r\right) \left(\frac{Td(TM^{\mathbb{C}})}{e(TM)}\right)_{\text{vol}}. \quad (1)$$

In the integrand of the RHS, only m -forms are picked up, so that the integration makes sense.

Remarks: The division by $e(TM)$ can really be carried out at the formal level. If m is an odd integer, the index vanishes identically, see below. Original references are Atiyah and Singer (1968a, b), Atiyah and Segal (1968).

2/10 How to derive it? (Example on the board!)

Basic Requirements

smooth manifolds, and algebraic topology (especially cohomology). Some familiarity with basic notions of functional analysis: Hilbert spaces, bounded linear operators, L^2 -spaces

2/10 How to derive it? (Example on the board!)

Method 1 Pseudodifferential operators

Method 2 Cobordism

Method 3 K theory

Method 4 Heat equation

Method 5 Using Supersymmetry for Dirac Operator form

Method 6 For a Spin Complex form

3/10 Example 1: the Gauss-Bonnet theorem

The standard Einstein-Hilbert action in three spacetime dimensions:

3/10 Example 1: the Gauss-Bonnet theorem

The standard Einstein-Hilbert action in three spacetime dimensions:

$$I_{GR} = \frac{1}{16\pi G} \int_M d^3x \sqrt{-g} (R - 2\Lambda) + I_{matter} \quad (2)$$

G is the gravitational coupling constant. **The standard field equation in three spacetime dimensions:**

3/10 Example 1: the Gauss-Bonnet theorem

The standard Einstein-Hilbert action in three spacetime dimensions:

$$I_{GR} = \frac{1}{16\pi G} \int_M d^3x \sqrt{-g} (R - 2\Lambda) + I_{matter} \quad (2)$$

G is the gravitational coupling constant. **The standard field equation in three spacetime dimensions:**

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \Lambda g_{\mu\nu} = 8\pi G T_{\mu\nu} \quad (3)$$

3/10 Example 1: the Gauss-Bonnet theorem

The standard Einstein-Hilbert action in three spacetime dimensions:

$$I_{GR} = \frac{1}{16\pi G} \int_M d^3x \sqrt{-g} (R - 2\Lambda) + I_{matter} \quad (2)$$

G is the gravitational coupling constant. **The standard field equation in three spacetime dimensions:**

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \Lambda g_{\mu\nu} = 8\pi G T_{\mu\nu} \quad (3)$$

Property: The field equations are generally covariant; that is, they're invariant under the action of the group of diffeomorphisms of the spacetime M , which can be view as a gauge group.

7/10 Example 2: the Hirzebruch signature theorem

Properties:

Since the vacuum state now is locally flat,

- i. there are no gravitational waves in the classical theory (Weyl tensor is also vanishes.)
- ii. upon quantization, there are no quantum graviton (c.f. topologically massive gravity by adding CS-term)
- iii. Source produce curvature, but only locally at the location in space-time

10/10 Acknowledgment

First of all, I have to thank Prof. Pisin Chen who give me the opportunity to write this thesis and join his course of Cosmological Physics which is offer in National Taiwan University. My classmates Yen-Chin Wang, Shun-Hsiang Chan, Yu-Hsiang Zach Lin, Ai-Lei Sun, Yu-Chien Huang and Yi-Hsien Liu, Prof. W-Y. Pauchy Hwang, Hsien-chung Kao, my advisor Je-An Gu and my good friend Hsuan Li that thanks for interesting and inspiring discussions.