# From Parisi full support to deterministic QAOA circuits

## A long reasoning record

Draft, 26 October 2026. Mathematical sources: OpenAI's full-support manuscript of 27 September 2026 and QAOA manuscript of 25 September 2026, at the pinned public revision.

The full-support manuscript proves that an admissible integrable minimizer of the zero-temperature Parisi functional for the pure zero-field Sherrington–Kirkpatrick model has Stieltjes support equal to every overlap scale in [0,1). The proof turns a hypothetical support gap into a constant interval for the order parameter. Variational contact at the ends of that interval conflicts with a strict curvature inequality inside it. Along the auxiliary diffusion, the spatial gradient of the value function is a bounded martingale. Full support makes this martingale's second moment equal to time; its terminal correlation with Brownian motion gives the ground-state value. The manuscript's further argument gives a smooth Stieltjes density, strictly positive on $(0,1)$, and no atom at zero. Sections 9.1–9.2 derive this stronger ordinary conclusion separately from the selected formal support statement.

This record reconstructs those arguments and follows the finite Gaussian approximation into the separate QAOA manuscript. The latter claims that finite deterministic circuits approach the SK optimum when system size tends to infinity before circuit depth increases. Its circuit construction needs additional approximation, synthesis, seed-removal and transfer arguments. The two formal statements concern full support and scalar value consequences; neither states QAOA circuit convergence. Their verification scope is described at the end. See OpenAI (2026a), §§1–2 and OpenAI (2026b), Theorem 1.1.

The support and scalar arguments assume real analysis, stochastic calculus and variational minimization. Section 8 additionally uses Hilbert-space tensor products, symmetric Fock space, unbounded operators and quantum control. Its weighted-domain and control estimates are additional analytic inputs from the QAOA manuscript; the text gives their precise uses and proof destinations. The companion short paper concentrates on full support and the strictly positive interior density, while the exposition introduces the support argument through examples. References to OpenAI (2026a) and OpenAI (2026b) distinguish the two manuscripts throughout.

Sections 1–8 give the continuous proof: full support, the terminal value, finite Gaussian correlations and the construction of one ordinary QAOA word for each accuracy. Section 9 develops the density and finite-disorder consequences. Section 10 gives an independent scalar comparison and a heat-family lower bound; Section 11 treats alternative circuit constructions and further limits. The exact formal statements and model comparison are in Section 12, followed by the source and checking record in Section 13.

## 1. The SK model and the Parisi variational problem

For independent standard Gaussian couplings $J_{ij}$, put

$$H_n(\sigma) = \frac{1}{\sqrt{n}} \sum_{i<j} J_{ij}\sigma_i\sigma_j, \qquad \sigma \in \{-1,1\}^n, \qquad P_* = \lim_{n\to\infty} \frac{1}{n} \mathbb{E} \max_{\sigma} H_n(\sigma).$$

For overlap $R_{12} = n^{-1}\sum_i \sigma_i^1\sigma_i^2$, the covariance is $nR_{12}^2/2 - 1/2$. Thus the convention is $\xi(t) = t^2/2$, with $\xi'' = 1$. A centered Gaussian of variance $1/2$, common to all configurations, removes the final

constant without changing the expected maximum. Every normalization below uses this convention.

Let $\mathcal{U}$ consist of nonnegative, nondecreasing, right-continuous functions $\gamma : [0,1) \to [0,\infty)$ with $\int_0^1 \gamma(t)\,dt < \infty$. Their Stieltjes measures satisfy $\mu_\gamma([0,t]) = \gamma(t)$, including $\mu_\gamma(\{0\}) = \gamma(0)$. They are locally finite, but need not be probability measures or have finite total mass. Support is relative to [0,1).

Let $B$ be standard Brownian motion. The control value and Parisi functional are

$$\Phi_\gamma(t,x) = \sup_{|\alpha|\le 1} \mathbb{E}\left[\left|x + B_1 - B_t + \int_t^1 \gamma(s)\alpha_s\,ds\right| - \frac{1}{2}\int_t^1 \gamma(s)\alpha_s^2\,ds\right],$$

$$\mathcal{P}(\gamma) = \Phi_\gamma(0,0) - \frac{1}{2}\int_0^1 t\gamma(t)\,dt.$$

Controls are progressively measurable for Brownian increments after $t$. For a general admissible coefficient, the solution and its spatial derivatives are locally absolutely continuous in time on compact strips; the time equation holds almost everywhere. The control formula fixes the solution. The PDE is $\Phi_t = -\frac{1}{2}(\Phi_{xx} + \gamma\Phi_x^2)$, with $\Phi(1,x) = |x|$. Auffinger and Chen's zero-temperature Parisi formula and attainment identify $P_*$ with $\min_{\gamma\in\mathcal{U}} \mathcal{P}(\gamma)$. The support theorem itself assumes a minimizer and proves a property of it; it does not separately supply one. See OpenAI (2026a), §§1–2 and Auffinger and Chen (2017), Theorem 1 and proof of Lemma 3.

Write $u = \Phi_x$, $a = \Phi_{xx}$, and $dX_t = \gamma(t)u(t,X_t)\,dt + dB_t$, $X_0 = 0$. Write $q(t) = \mathbb{E}[u(t,X_t)^2]$ for the second moment of this gradient along $X$. The first result assumes an admissible minimizer. It constructs its Stieltjes measure and an associated diffusion, proves $\operatorname{supp}\mu_\gamma = [0,1)$ and strict increase of $\gamma$, and obtains $\mathbb{E}u(t,X_t)^2 = t$ and $\mathbb{E}a(t,X_t)^2 = 1$ for every associated diffusion. The selected formal statement has exactly this conditional scope; it does not assert a smooth density. Its Lean implementation at the fixed source revision implements the separately stated formal challenge. The Comparator configuration identifies the intended comparison; the evidence retained with this draft does not establish a successful research-target replay.

The second result concerns the scalar value. A bounded martingale ends at a spin-valued random variable whose Brownian correlation represents $P_*$. Finite Gaussian coefficients approximate that correlation at each fixed $T < 1$, followed by $T \uparrow 1$. Its formal statement is separate and retains the minimizer and diffusion hypotheses where needed. The QAOA theorem goes further: for each requested accuracy it selects one finite deterministic circuit, independent of size and disorder, whose limiting expected energy approaches $P_*$. Neither formal statement asserts that circuit conclusion. The scalar clauses are in a separate Lean implementation, with their own formal challenge and Comparator configuration. Those links identify the selected statements, not a completed replay receipt for either research target.

## 2. Analytic preliminaries and prior work

### 2.1 Structure of the argument

The proof begins with regularity on strips bounded away from terminal time. It makes the diffusion $X$ and the gradient moment $q$ available, so a first variation can show where the minimizer is in contact. On a constant interval, the shape inequalities and the rational certificate give the strict curvature needed to contradict that contact. The zero-interval case uses a separate heat argument,

and the possible terminal plateau needs its own exclusion. These steps lead to full support and the two moment equalities.

In this manuscript route, the value argument then uses bounded-martingale convergence and the attained Parisi formula to express $P_*$ as a terminal Brownian correlation and an integral of expected curvature. Section 6 defines the terminal martingale variable and proves these equalities. The fixed-$T$ Gaussian approximation recovers this integral through Euler coupling, orthogonal normalization and mean-preserving rounding. The QAOA theorem additionally requires rooted energy identification, classical synthesis, removal of temporary seeds and transfer to fixed angles. The main route is §§3–5, then §§6–7, then §§8.1–8.6; §12 compares the two formal interfaces. The original analytic details are in OpenAI (2026a), §§2–7, and the circuit construction is in OpenAI (2026b), §§2–6.

The density theorem in §§9.1–9.2 is a further consequence of support and the PDE, not a premise of the scalar or circuit construction. The positive-temperature route in §11.1 independently supplies the same bounded classical input used for synthesis. The probability consequence in §11.2 requires a separate fixed-word second-moment theorem. These branches explain why the two selected formal statements are not a formal proof of the complete circuit theorem.

These destinations separate premises of the support proof from its consequences. They locate the detailed arguments behind the short paper's support and density theorem.

**The diffusion and moment identity.** For any admissible $\gamma$, the compact-strip estimates justify defining $X, q$ and computing $q' = \mathbb{E}a^2$; see §2.4 and §3.1.

**Variational contact.** Minimality, admissible mass addition and compact mass removal give $G \geq 0$, its vanishing on support and the contact slopes. The steps are in §3.2, §3.3 and §3.4.

**Strict curvature.** Admissibility and a positive constant $c$ on an interval suffice; minimality is not needed. The shape comparisons, algebra and limiting argument are in §4.3, §4.4 and §4.5.

**Exclusion of gaps.** Combine contact with the appropriate positive, zero or terminal argument. The cases have separate destinations: terminal, bounded and initial gaps.

**Positive density.** Full support comes first. Differentiated moments then give smoothness in §9.1, and a local comparison gives strict positivity in §9.2.

**Value and circuits.** Retain the same minimizer and diffusion through the value identity in §6 and the finite Gaussian construction in §7. The circuit argument in §8 requires additional theorems.

### 2.2 The regularity available before the proof starts

**Proposition 2.1 (compact-strip regularity).** For $\gamma \in \mathcal{U}$ and $0 \leq T < 1$, the value has bounded, jointly continuous spatial derivatives of every positive finite order on $[0, T] \times \mathbb{R}$. If admissible coefficients converge in $L^1$ and their smooth, even, convex, one-Lipschitz terminal values converge uniformly to $|x|$, these spatial derivatives converge uniformly on each such strip. The bounds depend on the derivative order, the fixed strip and a common coefficient mass bound. Time derivatives need only exist almost everywhere across coefficient jumps. On an open constant interval, the heat formula supplies the additional time smoothness used in §4. The terminal estimate is

$$|x| \leq \Phi_\gamma(t, x) \leq |x| + \sqrt{2(1-t)/\pi} + \frac{1}{2}\int_t^1 \gamma(s)\, ds.$$

It controls approach to $|x|$ without differentiating that cusp. The strip estimates justify the interior calculations and finite meshes; the terminal bound and integrability justify the later endpoint limits. The two roles must remain separate. Their uses depend on the region and limit.

On a fixed $[0,T]$ with $T<1$, bounded spatial derivatives and locally absolutely continuous time dependence justify the Itô identities and Euler coupling. Across a jump of $\gamma$, the spatial derivative moments remain continuous and the time equations hold almost everywhere; this is enough for contact slopes and the endpoint chord comparison. Inside a constant interval, classical time differentiation permits the strict-curvature calculation.

As $T\uparrow 1$, the controls change: the displayed value bound, integrable coefficient tails and martingale convergence justify first variation and terminal value. No differentiated estimate at the terminal cusp is used.

These are OpenAI (2026a), Lemma 2.1 and its diffusion consequences; §2.4 reconstructs their uniformity.

The hypotheses also have distinct uses. Zero field supplies evenness, $u(0,0)=0$ and reflection in the shape argument. The convention $\xi''=1$ fixes the curvature coefficients. Monotonicity gives bounded coefficients before time one and nonnegative coefficient jumps; it also turns a support gap into a constant interval. Integrability controls terminal tails and Fubini interchanges, while local finiteness of the Stieltjes measure is enough for compact mass removal. Minimality first enters through the variational inequality in §3; the constant-interval curvature estimate itself applies to an admissible coefficient without assuming it is a minimiser. OpenAI (2026a), §§1–2; OpenAI (2026a), §2; OpenAI (2026a), §6.

### 2.3 Which prior results play which role

The imported ground-state formula supplies two facts with different uses. Auffinger and Chen's Theorem 1 identifies the ground-state limit with an infimum over $\mathcal{U}$. To choose a coefficient attaining that infimum, one must also use their construction in the proof of Lemma 3. In the cited arXiv v2, the theorem is on p. 3, the compactness construction is on pp. 7–9, and the lower-bound proof is on pp. 11–12.

The support proof is conditional on a minimizer. From that hypothesis, continue directly with the compact-strip estimates in §2.4. The construction below explains how attainment supplies the minimizer later used in the value and circuit arguments.

Here is the part of that construction which supplies the witness. Let $\alpha_\beta$ be a positive-temperature Parisi distribution function. The source bounds $\beta\alpha_\beta(t)$ uniformly on each interval ending before one. Helly selection and a diagonal subsequence produce a nonnegative, nondecreasing right-continuous limit $\gamma_0$. In the SK normalization, the source's weighted integral bound also controls $\int_0^1 \beta\alpha_\beta(t)\,dt$. For every nonnegative nondecreasing $h$, splitting at $1/2$ gives $\int_0^1 h(t)\,dt \le 4\int_0^1 th(t)\,dt$: the first half is bounded by $h(1/2)/2$, and on the second half $t\ge 1/2$. Fatou's lemma then gives $\gamma_0\in\mathcal{U}$.

Mass can concentrate at time one during this limit, so compact convergence alone does not identify the limiting control problem. Write the limiting measure of the time coefficients as

$$\nu_0(dt)=\gamma_0(t)\,dt+m\delta_1(dt),\qquad m\ge 0.$$

For an arbitrary admissible control, let $S$ denote its terminal state with coefficient $\gamma_0$. To see why the terminal choice is admissible, let the original control be $\alpha$ and put $S_s=B_s+\int_0^s \gamma_0(r)\alpha_r\,dr$. For

a cutoff $\tau < 1$ use $\alpha_s$ before $\tau$ and $g_k(S_s)$ afterwards, where $g_k$ is a continuous function bounded by one and converging pointwise to the sign with $\operatorname{sgn}(0) = 1$. This control uses only information available at time $s$. At fixed $\tau, k$, first take the temperature limit; then let $\tau \uparrow 1$ and $k \to \infty$. The continuous terminal evaluation has thereby approached $\operatorname{sgn}(S)$ without anticipating a future state. The endpoint atom then contributes

$$|S + m\operatorname{sgn}(S)| - |S| = m$$

to the reward and

$$\tfrac{1}{2}m(\operatorname{sgn}(S)^2 + 1) = m$$

to the combined control and Parisi penalties. These contributions cancel. Taking the supremum over the remaining controls proves $P_* \geq \mathcal{P}(\gamma_0)$. The upper bound proved earlier in that paper gives $P_* \leq \inf_{\gamma \in \mathcal{U}} \mathcal{P}(\gamma)$. Therefore

$$P_* \leq \inf_{\gamma \in \mathcal{U}} \mathcal{P}(\gamma) \leq \mathcal{P}(\gamma_0) \leq P_*.$$

Thus $\gamma_0$ attains the infimum. The cancellation is a step in the imported existence proof; it is not an additional assumption about the Stieltjes measure $d\gamma_0$. In particular, $\nu_0$ here is the limit of coefficient-times-Lebesgue measures, not the order parameter's Stieltjes measure. The two measures must not be identified. Auffinger and Chen, arXiv:1606.05335v2, pp. 7–12.

The cited theorem is stated for the almost-sure normalized maximum. Its expected-value formulation used here follows from Gaussian concentration. For $L_n = n^{-1}\max_\sigma H_n(\sigma)$, the Lipschitz constant in the Gaussian couplings is $\sqrt{(n-1)/(2n^2)}$. Hence $L_n - \mathbb{E}L_n \to 0$ in probability. Since the almost-sure limit is the deterministic Parisi value, the deterministic sequence $\mathbb{E}L_n$ has the same limit. The full-support theorem below remains conditional on a minimizer; neither of the selected formal interfaces constructs one. OpenAI full-support manuscript, Introduction, ground-state formula and its footnote.

Auffinger–Chen–Zeng's infinite support does not exclude an interior gap. Their terminal-plateau observation anticipates one endpoint obstruction; the argument below supplies the quantitative boundary-layer proof used here. The full-support manuscript develops a rational coercivity route and credits Chen's 2026 theorem for the same full-support conclusion and smooth absolute continuity. Chen's Proposition 4.2 has the conditional crossing form $q''(t) = 0 \Rightarrow q'''(t) > 0$; the cited rational certificate gives $q'''(t) > 0$ throughout a positive constant interval. These are different sufficient mechanisms, and this record makes no priority claim for full support. Auffinger, Chen and Zeng (2020), Theorem 1 and Remark 7; Chen (2026), Theorem 1.3 and Proposition 4.2; OpenAI (2026a), §§1–2. The transformed density comparisons also have antecedents in Lopatto's positive-temperature analysis and Chen's zero-temperature analysis; the rational coercive identity is the different step in the full-support proof. Zhou's positive-temperature interval-support result applies near the critical temperature, whereas Lopatto's cited theorem covers the low-temperature zero-field phase. Neither transfers full Stieltjes support to zero temperature on its own: for the minimizers, $\beta\mu_\beta$ converges vaguely on $[0, 1)$ to the zero-temperature Stieltjes measure, while the unscaled Parisi probability measures converge weakly on $[0, 1]$ to $\delta_1$. These are different measures and topologies. Convergence without an interval-specific lower bound can lose support information. The direct zero-temperature gap proof below is therefore essential. Chen (2026), Remarks 1.4–1.5; OpenAI (2026a), §§1–2.

The coefficient classes in this comparison can be translated explicitly. In the positive-temperature convention, write

$$U_t = -\frac{\beta^2}{2}(U_{yy} + \alpha_\nu U_y^2), \qquad U(1, y) = \log\cosh y.$$

Set $\Phi_\beta(t,x) = \beta^{-1}U(t,\beta x)$ and $\gamma_\beta = \beta\alpha_\nu$. Direct differentiation gives

$$(\Phi_\beta)_t = -\tfrac{1}{2}\big((\Phi_\beta)_{xx} + \gamma_\beta(\Phi_\beta)_x^2\big), \qquad \Phi_\beta(1,x) = \beta^{-1}\log\cosh(\beta x).$$

Dividing the associated diffusion by $\beta$ likewise gives unit Brownian noise and drift $\gamma_\beta(\Phi_\beta)_x$. Conversely, every bounded nonnegative, nondecreasing, right-continuous coefficient on $[0,1)$ has this form: choose $\beta > 0$ at least its supremum, take distribution function $\alpha_\nu = \gamma/\beta$ below one, and put the remaining probability mass at one. This identifies coefficient classes, not minimizers at different temperatures. The terminal values differ from $|x|$ by at most $(\log 2)/\beta$, but passing a strict inequality to the cusp and to coefficients unbounded near one still needs uniform estimates. A qualitative crossing statement alone supplies no positive limiting fourth-moment margin. Lopatto, arXiv:2607.11756v4, equation (1.2), §2 and Proposition 5.1.

Two related works sharpen the comparison without entering the pinned September proof. Lopatto, arXiv:2607.11756v4, Proposition 5.1 and (5.8)–Corollary 5.6, proves a conditional curvature crossing on a positive constant interval. Its integrating factor makes a transformed curvature derivative strictly increasing, so the associated contact derivative decreases and then increases; it does not assert an unconditional positive third derivative. Lopatto, arXiv:2610.00127v1, Theorem 2.3 and §8, works at positive temperature and positive external field. Its comparison requires one-sided bounds on the coefficient before and after the chosen time, a moment inequality and a stationarity equality. Several of these conditions are automatic in its density application: monotonicity supplies the one-sided bounds at a continuity point, contact on a full-support interval makes the scalar inequality an equality, and the quotient formula for the coefficient supplies stationarity. Its moment bootstrap then gives smooth, strictly positive density on a nontrivial support interval; a singleton support is a separate case. Appendix B, Theorem B.1 and equation (B.1), also prove strict positivity in the zero-field positive-temperature setting, where the extra noncentered-law condition is unnecessary. These results are therefore close analytic antecedents. The distinction used below is the unconditional quantitative fourth-moment margin for the comparison coefficient and its controlled passage to the zero-temperature cusp.

Here is an explicit example of the information lost in such a limit. It concerns admissible measures, not Parisi minimizers. Put $g = \mathbf{1}_{[0,1/3]\cup[2/3,1)}$ and $q_n = 1 - n^{-2}$, and for $n \geq 2$ define

$$\mu_n(dt) = \frac{g(t) + n^{-1}}{n}\mathbf{1}_{[0,q_n]}(t)\,dt + \left(1 - \frac{1}{n}\int_0^{q_n}(g(t) + n^{-1})\,dt\right)\delta_{q_n}(dt).$$

The continuous part has mass at most $(2/3 + 1/n)/n < 1$, so these are probability measures with a positive endpoint atom. Their density is positive throughout $[0, q_n]$, giving $\operatorname{supp}\mu_n = [0, q_n]$. Their continuous mass tends to zero and $q_n \to 1$, hence $\mu_n \Rightarrow \delta_1$ weakly on $[0,1]$. For a compactly supported continuous test function on $[0,1)$, the endpoint atom eventually lies outside its support. Consequently $n\mu_n \to g(t)\,dt$ vaguely there, although the limit vanishes on $(1/3, 2/3)$. Even the cumulative coefficients converge in the topology used for the PDE: with $\gamma_n(t) = n\mu_n([0,t])$ and $\gamma(t) = \int_0^t g(s)\,ds$, their difference is $t/n$ below $q_n$, and $\gamma_n = n$ on the terminal strip of length $n^{-2}$. Thus

$$\|\gamma_n - \gamma\|_{L^1(0,1)} \leq \frac{1}{2n} + \frac{1}{n} \longrightarrow 0.$$

Every $\gamma_n$ is admissible, but $d\gamma$ has an interior gap. Minimality and the gap argument supply information that these convergences alone do not. An atom added at time one is likewise invisible to the coefficient on $[0,1)$ and hence to its PDE and variational functional; the zero-temperature problem does not select an endpoint mass.

Control and gradient-martingale methods have antecedents in Auffinger–Chen and Jagannath–Tobasco. A finite-temperature theorem with smooth terminal value $\log\cosh$ does not alone establish the result with terminal cusp $|x|$. The required compact-strip and terminal estimates are supplied in OpenAI (2026a), §2; Chen–Handschy–Lerman supplies earlier zero-temperature regularity, control and uniqueness results (Theorem 4 for uniqueness). This is distinct from the earlier finite-temperature uniqueness result of Auffinger and Chen. Montanari and El Alaoui–Montanari–Sellke inform normalised discretisation and optimisation. The particular shifted Gaussian coefficient identity used here is proved in OpenAI (2026a), §7.

### 2.4 Recovering the compact-strip estimates

The curvature argument needs estimates uniform over the approximating coefficients. Here is the route from the control formula to those estimates; all bounds are taken on a fixed strip ending before time one.

For finite-step coefficients and a smooth, even, convex, one-Lipschitz terminal function $\psi$, the evolution through a step of height $c$ is

$$\Phi(t,x) = \begin{cases} c^{-1}\log P_{b-t}(e^{c\Phi(b,\cdot)})(x), & c>0,\\ P_{b-t}\Phi(b,\cdot)(x), & c=0, \end{cases}$$

where $P_h$ denotes convolution with a centred Gaussian of variance $h$. The heat formula preserves evenness, convexity and the spatial Lipschitz bound. Along a controlled path $dY_s = \gamma(s)\alpha_s\,ds + dB_s$, the drift of the value minus the accumulated control cost is $-\gamma(s)(\alpha_s - \Phi_x(s,Y_s))^2/2$. It is nonpositive, and is zero for the feedback control. This proves the stated control formula for the smooth step problem, including attainability by its feedback diffusion.

Using the same control in two such problems and then taking suprema gives

$$\|\Phi_{\gamma,\psi} - \Phi_{\tilde\gamma,\tilde\psi}\|_\infty \le \|\psi - \widetilde{\psi}\|_\infty + \frac{3}{2}\|\gamma - \tilde\gamma\|_{L^1(0,1)}.$$

The drift part costs at most the coefficient difference in $L^1$, by the terminal Lipschitz bound; the quadratic cost contributes another half of that difference. The supremum norm is of the difference of the values, which is bounded even though each value grows linearly in $x$. With $\psi_M(x) = M^{-1}\log(2\cosh Mx)$, $0 \le \psi_M - |x| \le (\log 2)/M$. Choose simultaneously $\gamma_m \to \gamma$ in $L^1$ and $M_m \to \infty$. These are two distinct requirements: $M_m \ge \|\gamma_m\|_\infty$ alone does not force terminal convergence. This compatible approximation determines a unique uniform limiting value. Its terminal bound, given in §2.2, also follows directly: choose the zero control for the lower bound, and use $|\alpha| - \alpha^2/2 \le 1/2$ for the upper bound.

To obtain spatial derivatives, choose $T < b < 1$. Monotonicity gives the uniform coefficient estimate

$$\gamma(t) \le \frac{\|\gamma\|_1}{1-b} \quad (0 \le t \le b).$$

An $L^1$-convergent family thus has a common coefficient bound $\Gamma$ on this larger strip. For its smooth step approximants, put $U = \Phi_x$. The differentiated mild equation is

$$U(t) = P_{b-t}U(b) + \int_t^b \gamma(s)P_{s-t}[U(s)U_x(s)]\,ds.$$

Since $\|U\|_\infty \le 1$ and $\|\partial_x P_h F\|_\infty \le Ch^{-1/2}\|F\|_\infty$,

$$\|U_x(t)\|_\infty \le \frac{C}{\sqrt{b-t}} + C\Gamma \int_t^b \frac{\|U_x(s)\|_\infty}{\sqrt{s-t}}\, ds.$$

Substituting this inequality once into its own integral uses $\int_t^s ((r-t)(s-r))^{-1/2}\, dr = \pi$. It reduces the estimate to an ordinary backward Gronwall inequality with integrable forcing. This bounds $U_x$ uniformly on $[0,T]$ without any terminal derivative bound. Repeating on successively smaller strips bounds each higher spatial derivative: put one derivative on the heat kernel and the rest on $UU_x$; only the term $U\partial_x^m U$ contains the new highest derivative. The remaining products have already been bounded.

Uniform convergence of the values then yields convergence of every positive spatial derivative. For a bounded twice differentiable difference $F$ with bounded second derivative, Taylor's formula gives

$$\|F'\|_\infty \le \frac{2\|F\|_\infty}{h} + \frac{h}{2}\|F''\|_\infty \qquad (h > 0).$$

First choose $h$ small using the common second-derivative bound, then make $\|F\|_\infty$ small. Iterate on derivative differences. Passing to the integral form of the differentiated PDE proves local absolute continuity in time and its almost-everywhere equations. No classical time derivative at a coefficient jump, and no derivative of $|x|$ at its cusp, is being assumed.

Finally, couple all feedback diffusions with the same Brownian motion. On $[0,T]$, let $L$ be a common spatial Lipschitz bound for their drifts. The difference of the two integral equations gives, for example,

$$\sup_{s\le T} |X_s^n - X_s| \le e^{LT}\left(\|\gamma_n - \gamma\|_1 + \|\gamma_n\|_1 \|u_n - u\|_{\infty,[0,T]\times\mathbb{R}}\right) \longrightarrow 0.$$

This is a deterministic bound under the coupling. The derivative bounds one order higher ensure uniform continuity when derivatives are evaluated at these moving points. Every fixed polynomial in the bounded derivatives therefore converges in expectation. This is precisely the convergence needed for the integrated curvature expression in §4; none of the forward density scores or their quotients need converge. OpenAI (2026a), Lemmas 2.1–2.3.

### 2.5 Order of approximation

The same symbol for a small error must not conceal changing background objects. The following account fixes those objects before the indicated limit. A single admissible minimiser and its associated diffusion are retained throughout the scalar construction.

**Compact-strip regularity.** Fix a derivative order $k \ge 1$, a cutoff $T < 1$ and a common coefficient $L^1$ bound. Coefficients converge in $L^1$ within that bound; spatial derivatives converge uniformly on the strip.

**Shape inequalities on a plateau.** Fix the height $c > 0$ and a closed interior neighbourhood. Choose step coefficients equal to $c$ there and smooth terminal values with $M_m \ge \|\gamma_m\|_\infty$ and $M_m \to \infty$. The coefficient and terminal approximations may be chosen as one compatible sequence.

**First variation.** Fix $\gamma$, a feasible $\eta \in L^1$ and initially $\epsilon > 0$. Remove the terminal cutoff before letting $\epsilon \downarrow 0$. The terminal remainder divided by $\epsilon$ is bounded by an $L^1$ tail of $\eta$.

**Terminal value.** Keep the chosen minimiser and diffusion fixed. Bounded martingale convergence and integrability of $\gamma$ justify $t \uparrow 1$; no terminal derivative bound is needed.

**Finite Gaussian coefficients.** Fix $\gamma, X$ and $T < 1$. Refine $N$ using the fixed-strip bounds, then control the remaining value tail by $1 - T$.

For the plateau transfer, tail estimates for a fixed approximant may depend on that approximant. The passage to the limit uses the integrated polynomial in the scaled second, third and fourth spatial derivatives of the value function and the common strip bounds; it does not require uniform bounds on the auxiliary density scores. Sections 3 and 4 explain the variation and plateau-transfer arguments, and §7 gives a complete scalar error budget. The further choices used to produce a circuit appear in §8.6. OpenAI (2026a), Lemma 2.1, Lemma 6.1, Proposition 5.1 and Proposition 7.2.

## 3. From minimality to contact

**Proposition 3.1 (variational contact).** For an admissible minimizer, the potential $G(s) = \int_s^1 (q(t) - t)\, dt$ is nonnegative on $[0, 1)$ and vanishes on $S = \operatorname{supp} \mu_\gamma$. At every $s \in S$, $q(s) = s$ and $q'(s) \le 1$, with a right derivative at zero.

The minimisation takes place over nonnegative Stieltjes measures with an integrable cumulative function. Their total mass is free. We first derive the variation of the functional, then use changes of measure which preserve this admissible class. The result will be contact on the *existing* support; the exclusion of gaps comes afterwards.

### 3.1 The moment function before minimisation

Fix any $\gamma \in \mathcal{U}$, without yet assuming that it minimises $\mathcal{P}$. Compact-strip regularity and Itô's formula give

$$u(t, X_t) = \int_0^t a(s, X_s)\, dB_s, \qquad q(t) := \mathbb{E}u(t, X_t)^2 = \int_0^t \mathbb{E}a(s, X_s)^2\, ds.$$

The initial term vanishes because zero field makes $\Phi$ even and $u(0, 0) = 0$. Thus $q(0) = 0$, $0 \le q(t) \le 1$, and

$$q'(t) = \mathbb{E}a(t, X_t)^2 \qquad (0 \le t < 1),$$

with a right derivative at zero. The integrand is continuous on each compact time strip, so $q \in C^1[0, 1)$. These facts hold for every admissible coefficient. They do not yet say that $q(t) = t$. The source proves them by smooth approximation on $[0, T]$, $T < 1$; it never differentiates the terminal cusp $|x|$ at time one. OpenAI (2026a), Lemmas 2.1–2.2.

### 3.2 The feasible right derivative

Let $\eta \in L^1(0, 1)$ and suppose that $\gamma + \varepsilon\eta \in \mathcal{U}$ for $0 \le \varepsilon \le \varepsilon_0$, where $\varepsilon_0 > 0$. This is a feasibility hypothesis: an arbitrary signed bump in $\gamma$ need not preserve monotonicity. Since $\mathcal{U}$ is convex, it is enough to check that the endpoint $\gamma + \varepsilon_0\eta$ is admissible. The formula below is the zero-field, $\xi'' = 1$ specialization of Chen, Handschy and Lerman, arXiv:1609.04368v4, Proposition 4. Their chord $\gamma_\theta = (1 - \theta)\gamma + \theta\gamma_1$ gives this direction by taking $\gamma_1 = \gamma + \varepsilon_0\eta$ and $\theta = \varepsilon/\varepsilon_0$. Their class $\mathcal{U}$ permits unnormalized cumulative measures with finite cumulative integral, so this change of parameter also covers the scaling direction $-\gamma$. We reconstruct the detailed PDE proof from OpenAI's Lemma 6.1.

For $0 < \varepsilon \le \varepsilon_0$, put $\beta = \gamma + \varepsilon\eta$, $\tilde{u} = \Phi_{\beta,x}$ and $D = \Phi_\beta - \Phi_\gamma$. Subtracting the two PDEs gives

$$D_t = -\frac{1}{2}D_{xx} - \frac{1}{2}\beta(\tilde{u} + u)D_x - \frac{\varepsilon}{2}\eta u^2.$$

The choice of diffusion

$$dY_t^\varepsilon = \frac{1}{2}\beta(t)(\tilde{u} + u)(t, Y_t^\varepsilon)\,dt + dB_t, \qquad Y_0^\varepsilon = 0,$$

cancels the first-order spatial term when Itô's formula is applied to $D(t, Y_t^\varepsilon)$. Its drift has absolute value at most $\beta(t)$, which is integrable. On $[0, T]$ the resulting identity is

$$D(0,0) = \mathbb{E}D(T, Y_T^\varepsilon) + \frac{\varepsilon}{2}\int_0^T \eta(t)\mathbb{E}[u(t, Y_t^\varepsilon)^2]\,dt.$$

At coefficient jumps the time derivatives need only exist almost everywhere. The source obtains this identity by approximating both coefficients by monotone step functions with the same smooth terminal datum, and then passing to the limit using the compact-strip estimates.

The control representation, applied with the same control on each side, gives the time-local difference bound

$$\sup_x |D(T, x)| \le \frac{3\varepsilon}{2}\int_T^1 |\eta(t)|\,dt.$$

Hence $T \uparrow 1$, at fixed $\varepsilon$, removes the terminal term even after division by $\varepsilon$. We obtain the exact finite-change identity

$$\frac{\Phi_{\gamma+\varepsilon\eta}(0,0) - \Phi_\gamma(0,0)}{\varepsilon} = \frac{1}{2}\int_0^1 \eta(t)\mathbb{E}[u(t, Y_t^\varepsilon)^2]\,dt.$$

As $\varepsilon \downarrow 0$, the coefficients converge in $L^1$ and have uniformly bounded $L^1$ norms. On each fixed strip, their gradients converge uniformly and have common spatial Lipschitz bounds. Coupling with the same Brownian motion and applying Gronwall's inequality therefore gives $Y^\varepsilon \to X$ uniformly on that strip. The integrand converges to $\eta(t)q(t)$ for each $t < 1$, and its absolute value is at most $|\eta(t)|$. Dominated convergence, followed by subtraction of the linear Parisi penalty, now yields

$$D_+\mathcal{P}(\gamma)[\eta] := \lim_{\varepsilon\downarrow 0}\frac{\mathcal{P}(\gamma + \varepsilon\eta) - \mathcal{P}(\gamma)}{\varepsilon} = \frac{1}{2}\int_0^1 \eta(t)(q(t) - t)\,dt.$$

If $\gamma$ is a minimiser, this derivative is nonnegative for every feasible direction. It need not vanish: the minimisation has one-sided constraints. OpenAI (2026a), Lemma 6.1, equations (6.1)–(6.2).

### 3.3 Which changes of measure are allowed?

Write $\mu = \mu_\gamma$ and $S = \operatorname{supp}\mu$, with support relative to $[0, 1)$. Tonelli's theorem gives the useful identity

$$\int_0^1 \gamma(t)\,dt = \int_{[0,1)}\int_s^1 dt\,\mu(ds) = \int_{[0,1)}(1 - s)\,\mu(ds) < \infty.$$

This is weighted finiteness, not a bound on $\mu([0, 1))$. In particular, there is no normalisation constraint requiring mass added at one point to be removed elsewhere.

Define the tail potential

$$G(s) = \int_s^1 (q(t) - t)\, dt \qquad (0 \le s < 1).$$

It is continuous, satisfies $|G(s)| \le 1 - s$, and has $G'(s) = s - q(s)$ and $G''(s) = 1 - q'(s)$ on compact strips, with one-sided interpretations at zero. The following two competitors suffice.

| Change to the Stieltjes measure | Direction in the cumulative function | Allowed parameter | Consequence at a minimiser |
|---|---|---|---|
| Add $\varepsilon\delta_s$, $0 \le s < 1$ | $\eta(t) = \mathbf{1}_{[s,1)}(t)$ | $\varepsilon \ge 0$ | $G(s) \ge 0$ |
| Remove $\varepsilon\nu$, where $\nu = \mu\vert_K$ and $K \subset [0,1)$ is compact | $\eta_K(t) = -\nu([0,t])$ | $0 \le \varepsilon \le 1$ | $\int_K G(s)\, \mu(ds) \le 0$ |

For atom addition, the new cumulative function is nonnegative, nondecreasing and right-continuous. Its integral increases by $\varepsilon(1 - s)$, so it remains in $\mathcal{U}$. This includes $s = 0$: an atom there adds a constant to $\gamma$ on all of $[0, 1)$. The point 1 is outside the measure's domain and is not an additional competitor. The derivative formula gives $D_+\mathcal{P}(\gamma)[\mathbf{1}_{[s,1)}] = G(s)/2$, proving $G \ge 0$ everywhere below one.

For removal, choose $T < 1$ with $K \subset [0, T]$. Then $\nu(K) \le \gamma(T) < \infty$. The measure $\mu - \varepsilon\nu$ is nonnegative for $0 \le \varepsilon \le 1$, so its cumulative function is nonnegative, nondecreasing and right-continuous. It lies between zero and $\gamma$, hence is integrable; also $|\eta_K| \le \nu(K)$. Fubini is justified quantitatively by

$$\int_K \int_s^1 |q(t) - t|\, dt\, \mu(ds) \le \int_K (1 - s)\, \mu(ds) \le \nu(K) < \infty.$$

It follows that

$$0 \le D_+\mathcal{P}(\gamma)[\eta_K] = -\frac{1}{2} \int_0^1 \nu([0,t])(q(t) - t)\, dt = -\frac{1}{2} \int_K G(s)\, \mu(ds).$$

Since $G \ge 0$, its integral on every compact $K$ is zero. The compact intervals $[0, T]$ exhaust $[0, 1)$, so $G = 0$ $\mu$-almost everywhere. If a support point $s$ had $G(s) > 0$, continuity would give a relative neighbourhood on which $G$ stays positive. After restricting to a compact interval containing that neighbourhood, the defining property of support would give it positive $\mu$-mass, contradicting the zero integral. Therefore

$$G(s) = 0 \qquad (s \in S).$$

This step applies to atoms, diffuse mass and singular continuous mass. Removing individual atoms alone would not establish it. OpenAI (2026a), Lemma 6.2.

### 3.4 Contact values and endpoint slopes

At $s \in S \cap (0, 1)$, the function $G$ has an interior minimum of value zero. Thus $G'(s) = 0$ and $G''(s) \ge 0$, giving

$$q(s) = s, \qquad q'(s) \le 1.$$

If $0 \in S$, the argument needs the zero-field initial condition in addition to the one-sided minimum. We have $G(0) = 0$ and $q(0) = 0$, so

$$G(h) = G(0) + \int_0^h (t - q(t))\, dt = \frac{1}{2}(1 - q'(0))h^2 + o(h^2).$$

Nonnegativity for $h > 0$ implies $q'(0) \le 1$. This works whether or not zero is an atom. The equality $q(0) = 0$ comes from symmetry, rather than from differentiating a one-sided minimum. If $0 \notin S$, no condition $G(0) = 0$ has been obtained or used. OpenAI (2026a), Lemma 6.2.

We have now proved, without any no-gap assumption,

$$G \ge 0 \text{ on } [0,1), \qquad G = 0 \text{ on } S, \qquad q(s) = s \text{ and } q'(s) \le 1 \text{ for } s \in S.$$

These are precisely the contact data used in §5. They make no assertion that every zero of $G$ belongs to $S$, or that $q(t) = t$ at a point outside $S$. Terminal, interior and initial gaps remain to be excluded.

### 3.5 What the first-order criterion does, and does not, establish

The contact data needed for the gap proof are already available. Readers following that proof can continue to the curvature argument in §4. This subsection gives an alternative contact proof and the converse optimality criterion.

There is also a global alternative to compact mass removal. Atom addition still gives $G \ge 0$. Now replace $\mu$ by $(1 - \varepsilon)\mu$, or equivalently $\gamma$ by $(1 - \varepsilon)\gamma$, for $0 \le \varepsilon \le 1$. This preserves admissibility, and its direction $-\gamma$ lies in $L^1$, although it may be unbounded. The first variation in §3.2 therefore applies. The signed interchange is justified by

$$\int_{[0,1)} \int_s^1 |q(t) - t|\, dt\, \mu(ds) = \int_0^1 \gamma(t) |q(t) - t|\, dt \le \int_0^1 \gamma(t)\, dt < \infty.$$

Consequently

$$0 \le D_+ \mathcal{P}(\gamma)[-\gamma] = -\frac{1}{2} \int_{[0,1)} G(s)\, \mu(ds).$$

Together with $G \ge 0$, this gives $\int G\, d\mu = 0$. Continuity then gives $G = 0$ on relative support, including zero, as in §3.3. Finite weighted mass is sufficient throughout; no finite total mass or density of $\mu$ has been assumed. This ordinary argument uses the same integrable-direction first variation as the local removal proof.

For the actual moment function associated with $\gamma$, the two potential conditions $G \ge 0$ and $G = 0$ on $S$ are also sufficient for minimality. This ordinary deduction explains exactly how much variational information has been extracted. For another admissible coefficient $\beta$, let $\nu = \mu_\beta$. The weighted identity above and $|G(s)| \le 1 - s$ make both measure integrals absolutely convergent, even when either measure has infinite total mass. Fubini gives

$$2D_+ \mathcal{P}(\gamma)[\beta - \gamma] = \int_{[0,1)} G(s)\, \nu(ds) - \int_{[0,1)} G(s)\, \mu(ds) \ge 0.$$

Here the second integral is zero because $\mu$ is concentrated on its relative support. The control formula also shows that $\mathcal{P}$ is convex in its coefficient: for a fixed control, the terminal absolute

value is convex in the coefficient and both penalties are linear; taking the supremum over the same control class preserves convexity. Therefore

$$\mathcal{P}(\beta) - \mathcal{P}(\gamma) \geq D_+\mathcal{P}(\gamma)[\beta - \gamma] \geq 0.$$

This proves the converse. It agrees with the chord optimality criterion of Chen, Handschy and Lerman, arXiv:1609.04368v4, Proposition 4, printed p. 14. Chen's zero-temperature treatment obtains the same contact integral by testing 0 and $2\gamma$ and then using weighted Fubini Chen (2026), Proposition 3.2, equations (3.11)–(3.12). The compact-removal proof above keeps the local mass operation explicit.

The converse concerns $G$ built from the PDE and diffusion of the same $\gamma$. It does not say that an arbitrary curve with these contact properties comes from a Parisi solution. Nor does it prove full support: the missing input is the strict curvature on an interval where the coefficient is constant, established next. The formal chord statement and the ordinary derivative bridge are distinguished in §12.1.

## 4. Strict curvature on a constant interval

The gap argument needs a property of the derivative $q'$, rather than convexity of the spatial function $\Phi$. We establish the following analytic input before using minimality. Let $\gamma \in \mathcal{U}$ be constant with value $c$ on an open interval $I \subset (0,1)$, and let $X$ be its diffusion started at zero. Then

$$\begin{cases} q'''(t) \geq \dfrac{1}{1000c^2}\mathbb{E}\big[(c\Phi_{xx}(t,X_t))^4\big] > 0, & c > 0, \\ q''(t) = \mathbb{E}[\Phi_{xxx}(t,X_t)^2] > 0, & c = 0, \end{cases} \qquad t \in I.$$

Thus $q'$ is strictly convex for $c > 0$ and strictly increasing for $c = 0$. The coefficient may jump outside $I$ and may be unbounded near one, provided it is nonnegative, nondecreasing, right-continuous and integrable. No support property or minimising assumption enters this statement. The positive-time restriction matters when a diffusion density is used; interval endpoints are handled later by continuity. OpenAI (2026a), Proposition 5.1.

The proof has two stages. In §§4.1–4.4 the coefficient is a positive monotone finite-step function with smooth terminal data. Shape comparisons and boundary estimates justify the polynomial correction and its zero expectation. In §4.5, approximants $\gamma_n, \Phi_n, X^n$ approach an arbitrary admissible coefficient. Only the integrated expression in spatial derivatives passes to the limit; the auxiliary forward scores need not converge. Bounds depend on a fixed compact strip and a common coefficient mass bound. The strictly positive right side is then justified for the limiting solution itself.

### 4.1 Why the uncorrected expression has no pointwise sign

On a positive constant interval put $r = c\Phi_x$, $v = r_x$, $w = r_{xx}$ and $z = r_{xxx}$. In forward time, $r_t = -\frac{1}{2}r_{xx} - rr_x$ and $dX_t = r(t,X_t)\,dt + dB_t$. Differentiating the PDE and applying Itô's formula gives

$$dr(t,X_t) = v(t,X_t)\,dB_t,$$

$$dv(t,X_t) = -v(t,X_t)^2\,dt + w(t,X_t)\,dB_t, \qquad dw(t,X_t) = -3v(t,X_t)w(t,X_t)\,dt + z(t,X_t)\,dB_t.$$

Since $q' = c^{-2}\mathbb{E}v^2$, the drift of $v^2$ yields $c^2 q'' = \mathbb{E}(w^2 - 2v^3)$. The drifts of $w^2$ and $v^3$ are, respectively, $z^2 - 6vw^2$ and $3vw^2 - 3v^4$. Therefore

$$c^2 q''' = \mathbb{E}\big[z^2 - 12vw^2 + 6v^4\big].$$

These are interior-time identities. On a smaller constant interval the heat formula gives classical time smoothness and bounded spatial derivatives of sufficiently high order, so the local martingale terms have zero mean and the expectation differentiations are justified. Nothing here assigns a sign to $q''$. OpenAI (2026a), Lemma 2.3.

There is a concrete obstruction to reading the last display pointwise. For the smooth finite-step problem $c = M = 1$ on $[0, 1)$, with terminal value $\log(2\cosh x)$, the exact solution is

$$\Phi(t, x) = \log(2\cosh x) + \frac{1 - t}{2}, \qquad r = \tanh x.$$

Indeed, $\Phi_{xx} + \Phi_x^2 = 1$. Writing $R = r^2$ gives $v = 1 - R$, $w = -2rv$ and $z = -2v^2 + 4Rv$. At $R = 1/2$, $v = 1/2$, $w^2 = 1/2$ and $z = 1/2$, so

$$z^2 - 12vw^2 + 6v^4 = -\frac{19}{8} < 0.$$

This example lies in the smooth analytic class used below. Its diffusion density is $f_t(x) = e^{-t/2}\cosh(x)p_t(x)$, obtained from the heat transition formula, and integrates to one because $\mathbb{E}\cosh(B_t) = e^{t/2}$. It illustrates the necessity of the integration argument; it uses a smooth terminal value and makes no claim to be the zero-temperature minimiser. The displayed negative value is an exact elementary calculation. The heat formula itself is OpenAI (2026a), Lemma 2.1.

The same pointwise obstruction occurs with the actual terminal datum $|x|$. For the admissible constant coefficient $\gamma_c \equiv c > 0$, write $\tau = 1 - t > 0$ and use the same heat formula:

$$\Phi_c(t, x) = c^{-1}\log \mathbb{E}e^{c|x + \sqrt{\tau}Z|}.$$

At $c = 0$ its continuous extension is the heat value, whose derivatives are

$$a_0 = \sqrt{\frac{2}{\pi\tau}}e^{-x^2/(2\tau)}, \qquad b_0 = -\frac{x}{\tau}a_0, \qquad d_0 = \left(\frac{x^2}{\tau^2} - \frac{1}{\tau}\right)a_0.$$

At $x = \sqrt{\tau}$, $d_0 = 0$ and $b_0^2 = a_0^2/\tau > 0$. For fixed $t, x$, Gaussian convolution may be differentiated in $x$ through its kernel and expanded in $c$ near zero: exponential moments dominate every resulting polynomial factor. The apparent singularity of $c^{-1}\log \mathbb{E}e^{c|x+\sqrt{\tau}Z|}$ is removable, and its spatial derivatives through order four are smooth in $c$. Hence

$$d_c^2 - 12ca_c b_c^2 + 6c^2 a_c^4 = -\frac{12a_0^3}{\tau}c + O(c^2) < 0$$

for all sufficiently small positive $c$, at that fixed point. This is an ordinary consequence of the constant-coefficient formula, not an arbitrary assignment of unrelated derivative values. It neither asserts minimality of $\gamma_c$ nor contradicts the positive expectation proved below; it shows why that expectation cannot be replaced by a pointwise inequality even in the absolute-value-terminal model.

### 4.2 The smooth finite-step problem and its boundary control

Take $0 = t_0 < \cdots < t_N = 1$, $0 < c_1 \le \cdots \le c_N \le M$, and $\gamma(t) = c_i$ on $[t_{i-1}, t_i)$. The terminal value is $M^{-1}\log(2\cosh Mx)$. On a step of height $c$, set $l = c\Phi$ and $h = e^l$. Then $h_t = -\frac{1}{2}h_{xx}$. For $a < t$ in that step the diffusion transition density is

$$p_{t-a}(x-y)\frac{h(t,x)}{h(a,y)}, \qquad p_t(x) = (2\pi t)^{-1/2}e^{-x^2/(2t)}.$$

Its mass is one by $h(a,\cdot) = P_{t-a}h(t,\cdot)$. Factoring the positive-time density as $f = e^l g$ gives $g_t = \frac{1}{2}g_{xx}$ on each step. Put $s = -\partial_x \log g$. On the first step, $g(t,x) = p_t(x)e^{-l(0,0)}$ and $s = x/t$. At a forward jump $\delta = c_+ - c_- \ge 0$, continuity of $f$ gives $g_+ = g_- e^{-\delta\Phi}$. The functions $r$ and $s$ are odd. OpenAI (2026a), §3, before Lemma 3.1.

The comparison arguments take place on unbounded spatial domains, so the boundary estimates must precede the signs. For these fixed smooth data, Gaussian convolution and the finite jump rules give, as $x \to +\infty$,

$$l(t,x) = cx + C(t) + O(e^{-\kappa x}),$$
$$g(t,x) = \exp\left(-\frac{x^2}{2t} + \beta(t)x + C_0(t)\right)(1 + O(e^{-\kappa x})).$$

The errors may be differentiated any fixed finite number of times; the estimates are uniform on each closed step, with positive time bounded away from zero for the forward estimate. Their constants may depend on the step data. For example, after completing the square in a Gaussian convolution, the error is an expectation at $Y_x = \lambda x + b + \sqrt{V}Z$ with $\lambda$ bounded below by a positive constant. Split at $Y_x = \mathbb{E}Y_x/2$. The first part inherits the exponential tail; on the second, the Gaussian probability dominates the permitted exponential growth of the error and all its specified derivatives. This proves the differentiated estimates by finite induction through heat evolution and jumps. In particular, $s = x/t - \beta(t) + O(e^{-\kappa x})$, $s_x = 1/t + O(e^{-\kappa x})$ and $s_{xx} = O(e^{-\kappa x})$. OpenAI (2026a), Lemma 3.1.

The comparison principle used below is also explicit. If $U_\tau - \frac{1}{2}U_{xx} - bU_x - dU \ge 0$ on a compact cylinder, $d \le C$, the initial values are nonnegative and the lateral values are at least $-\varepsilon$, then $U \ge -\varepsilon e^{(\max(C,0)+1)(\tau-\tau_0)}$. Multiply by the corresponding decreasing exponential and apply the minimum principle. Exhaustion is legitimate when the upper bound for $d$ is independent of the cylinder and the lateral errors tend to zero. The first-order coefficient need only be bounded on each cylinder. This observation permits the $1/x$ coefficients which appear below. OpenAI (2026a), comparison preceding Lemma 3.2.

### 4.3 Backward and forward comparisons, including jumps

Use backward time $\tau$ on a single step. The equation is $r_\tau = \frac{1}{2}r_{xx} + rr_x$. The logarithmic heat formula preserves strict positivity of $v$: its second spatial derivative is the tilted mean of the old second derivative plus the tilted variance of the old first derivative. Initially $v = cM\operatorname{sech}^2(Mx) > 0$, and positive rescaling at a backward jump preserves this property. Consequently $r(x) > 0$ for $x > 0$.

Twice differentiating gives $w_\tau = \frac{1}{2}w_{xx} + rw_x + 3vw$. At the terminal endpoint $w = -2cM^2\operatorname{sech}^2(Mx)\tanh(Mx) \le 0$ for $x > 0$. The boundary value at zero is zero, and the

differentiated tails make the far boundary tend uniformly to zero. Comparison applied to $-w$ gives $w \le 0$ on the positive half-line.

Define

$$H = \frac{r_\tau}{r} = \frac{w}{2r} + v.$$

Oddness and $r_x(0) = v(0) > 0$ make $w/r$, hence $H$, smooth and even at zero. Direct differentiation gives

$$\begin{aligned} H_\tau &= \tfrac{1}{2}H_{xx} + (v/r + r)H_x + vH, \\ (H_x)_\tau &= \tfrac{1}{2}(H_x)_{xx} + (v/r + r)(H_x)_x + ((v/r)_x + 2v)H_x + wH. \end{aligned}$$

The terminal signs are $H = (cM - M^2)\operatorname{sech}^2(Mx) \le 0$ and $H_x \ge 0$ on $x > 0$, using $M \ge c$. For the first comparison write $v/r + r = 1/x + \eta(x)$, where $\eta$ is smooth and odd near zero. The radial extension $H(|y|)$ to $\mathbb{R}^3$ turns $\frac{1}{2}H_{xx} + x^{-1}H_x$ into $\frac{1}{2}\Delta H$. It is smooth at the origin, as is the extra drift, and $H$ tends uniformly to zero at infinity. Comparison on expanding balls proves $H \le 0$.

For the second comparison, $wH \ge 0$ and $(v/r)_x = w/r - v^2/r^2 < 0$, so the zeroth-order coefficient has upper bound $2 \sup v$. On $[\varepsilon, A]$, the two lateral errors are $H_x(\varepsilon) = O(\varepsilon)$ and $H_x(A) = o_A(1)$, uniformly on the closed step. Taking $\varepsilon \downarrow 0$ and $A \to \infty$ gives $H_x \ge 0$. At a backward jump $r_{\text{new}} = \alpha r_{\text{old}}$ with $0 < \alpha \le 1$,

$$H_{\text{new}} = H_{\text{old}} - (1 - \alpha)v_{\text{old}}, \qquad (H_x)_{\text{new}} = (H_x)_{\text{old}} - (1 - \alpha)w_{\text{old}}.$$

Thus both signs survive every jump. OpenAI (2026a), Lemma 3.2.

For the forward score use $a_s = s_x$, temporarily distinguishing it from the Hessian $a = \Phi_{xx}$. The forward heat equation gives

$$\begin{aligned} s_t = \tfrac{1}{2}s_{xx} - ss_x, \qquad (a_s)_t &= \tfrac{1}{2}(a_s)_{xx} - s(a_s)_x - a_s^2, \\ (s_{xx})_t &= \tfrac{1}{2}(s_{xx})_{xx} - s(s_{xx})_x - 3a_s s_{xx}. \end{aligned}$$

The first step has $s = x/t$. Thereafter $a_s - 1/t$ satisfies the linear equation with drift $-s$ and zeroth-order coefficient $-(a_s + 1/t)$. For fixed smooth data this coefficient is bounded above on a closed step, independently of spatial truncation; this uses the tail bound and continuity, before asserting its sign. Comparison gives $a_s \ge 1/t$. The last displayed equation then gives $s_{xx} \le 0$ on $x > 0$. At a forward jump,

$$s_+ = s_- + \delta\Phi_x, \quad (a_s)_+ = (a_s)_- + \delta\Phi_{xx}, \quad (s_{xx})_+ = (s_{xx})_- + \delta\Phi_{xxx}.$$

The backward signs and $\delta \ge 0$ preserve the forward signs.

A further comparison, not concavity alone, is needed. On $x > 0$ set $d_s = s_x/s$, $d_r = r_x/r$ and $\rho = d_s - d_r$. The heat equation gives $s_t = \frac{1}{2}s_{xx} - ss_x$, whereas $r_t = -rH$. Taking logarithmic derivatives yields

$$(d_s)_t = \tfrac{1}{2}(d_s)_{xx} + d_s(d_s)_x - s_{xx}, \qquad (d_r)_t = -H_x.$$

Since $\frac{1}{2}(d_r)_{xx} + d_r(d_r)_x = \frac{1}{2}(w/r)_x = H_x - w$, substitution of $d_s = d_r + \rho$ gives its equation

$$\rho_t = \tfrac{1}{2}\rho_{xx} + d_s\rho_x + (d_r)_x\rho + 2H_x - w - s_{xx}.$$

All three terms in the source are nonnegative by the comparisons already proved, and $(d_r)_x < 0$. On the first step, concavity gives $r(x) = \int_0^x v(y)\,dy \geq xv(x)$, so $\rho = 1/x - v/r \geq 0$. Odd Taylor expansions give $d_s = 1/x + O(x)$, $d_r = 1/x + O(x)$ and $\rho = O(x)$ near zero; the tail estimates give $\rho \to 0$ at infinity. Comparison on $[\varepsilon, A]$ therefore proves $\rho \geq 0$ on each subsequent step provided its initial sign is preserved. Define

$$j = s\rho = s_x - (s/r)v.$$

At a forward jump, with $\alpha = c_+/c_-$ and $\lambda = (c_+ - c_-)/c_-$, $r_+ = \alpha r_-$, $v_+ = \alpha v_-$, $s_+ = s_- + \lambda r_-$ and $(s_x)_+ = (s_x)_- + \lambda v_-$. Substitution cancels the added terms and gives $j_+ = j_-$. Since $s_+ > 0$ on $x > 0$, the sign of $\rho$ is preserved. The quotient $s/r$ extends smoothly and evenly across zero, with $j(0) = 0$; reflection gives $j \geq 0$ everywhere. OpenAI (2026a), Lemma 3.3.

### 4.4 How the signs make a correction with zero expectation useful

Here the coefficient is a positive constant $c$ on the step containing the chosen time. The variables are the rescaled spatial derivatives

$$r = c\Phi_x, \qquad v = c\Phi_{xx}, \qquad w = c\Phi_{xxx}, \qquad z = c\Phi_{xxxx}.$$

Write $h = e^{c\Phi}$ and factor the forward diffusion density as $f = hg$; its auxiliary score is $s = -\partial_x \log g$. These are the backward and forward quantities compared in §4.3. On the positive half-line, where $r, s > 0$, set

$$H = \frac{w}{2r} + v, \qquad j = s_x - \frac{s}{r}v, \qquad \rho = \frac{s_x}{s} - \frac{v}{r}, \qquad j = s\rho.$$

In this certificate only, $b = -w/r$ is a quotient; it is not the derivative $\Phi_{xxx}$ denoted by $b$ in the unscaled curvature formula. At a fixed positive time inside the step define

$$R = r^2, \quad K = rs, \quad b = -w/r, \quad B = b - 2v, \quad L = -rs_{xx}, \quad S_1 = z + bv - 2Rb.$$

The subscript on $S_1$ distinguishes it from the support set. The previous comparisons give

$$R, K, v, B, j, L, S_1 \geq 0, \qquad b = B + 2v \geq 0, \qquad Rb^2 = w^2,$$

because $B = -2H$ and $S_1 = 2rH_x$. Here $s = -\partial_x \log g$ belongs to the forward heat factor, while $\partial_x \log f = r - s$ is the logarithmic derivative of the diffusion density. The signs have separate sources: $B \geq 0$ comes from $H \leq 0$, $S_1 \geq 0$ from $H_x \geq 0$ on the positive half-line, $j = s\rho \geq 0$ from the relative-slope comparison, and $L = -rs_{xx} \geq 0$ from forward concavity. Reflection extends these signs. The quotients were defined for $x > 0$, where $r > 0$ and $v > 0$ by the tilted mean-and-variance argument in §4.3. At zero, $r$ and $w$ are odd, $r_x(0) = v(0) > 0$, and division by $r$ is therefore division by a function with a simple zero. This gives the smooth even extension $b(0) = -z(0)/v(0)$; the other comparison identities give $j(0) = 0$ and $S_1(0) = 0$. Thus zero is not an omitted boundary of the integration.

For $\mathcal{D} = r\partial_x$, direct differentiation gives

$$\mathcal{D}(R, K, v, b, j) = (2Rv,\ 2vK + Rj,\ -Rb,\ -z - bv,\ -jv + Kb - L).$$

Also $f_x/f = r - s$. Hence $\mathcal{W}(F) = \mathcal{D}F + (v + R - K)F$ satisfies

$$f\mathcal{W}(F) = \partial_x(rfF).$$

This identity does not itself make the integral vanish. The tails give $r = O(1)$; exponential decay of $v, w, z, b$ and their derivatives; $s = x/t + O(1)$; $j = 1/t + O((1+x)e^{-\kappa x})$; and $K = O(1+x)$. Every polynomial $F$ in $R, K, v, b, j$ and its differentiated expression therefore has at most polynomial growth. Meanwhile $f(t, x) \le C\exp(-x^2/(2t) + C|x|)$. Thus every term used below is absolutely integrable and $rfF \to 0$ at both infinities. Smooth extension across zero prevents an interior boundary term. Integrating on $[-A, A]$ and taking $A \to \infty$ now gives $\mathbb{E}_f\mathcal{W}(F) = 0$. OpenAI (2026a), Lemma 3.4.

The proof has four separate checks: the comparisons supply these signs; the tail estimates make the correction integrate to zero; the rational identity below gives a positive remainder; and §4.5 passes the resulting expectation inequality to the original coefficient. The polynomial check verifies the third step. Set

$$(A, C, k, T_0, g_0, h_0) = \left(\frac{1}{13}, \frac{53}{24}, \frac{19}{16}, \frac{73}{48}, \frac{2331}{10000}, \frac{87}{2500}\right),$$
$$(x_0, y, m, n_0) = \left(\frac{4663}{5000}, \frac{6853}{10000}, -\frac{1051}{400}, -\frac{1073}{500}\right).$$

Define

$$\begin{aligned} Q &= Avb + Cv^2 + kKv - 2kRb + T_0Rv, \qquad T_1 = v(g_0R + h_0j), \\ F_2 &= (Av - 2kR)b^2 + b(2Cv^2 + 2kKv + (g_0 + 2T_0)Rv + h_0jv) \\ &\quad + x_0v^2j + yv^3 + mKv^2 + n_0Rv^2. \end{aligned}$$

The choice is motivated by the terms containing $z$. In fact,

$$\begin{aligned} \partial_bF_2 &= 2(Av - 2kR)b + 2Cv^2 + 2kKv + (g_0 + 2T_0)Rv + h_0jv \\ &= 2Q + T_1, \\ [z]\mathcal{W}(F_2) &= -\partial_bF_2 = -(2Q + T_1). \end{aligned}$$

Here $[z]$ means the coefficient of $z$; only $\mathcal{D}b = -z - bv$ contributes it. The square $(z+Q)^2$ and $S_1T_1$ contribute $2Q + T_1$, so these terms cancel exactly. Therefore subtracting $(z+Q)^2 + S_1T_1 + \mathcal{W}(F_2)$ from the curvature expression cancels every term involving $z$. The other favourable derivative is $\partial_jF_2 = x_0v^2 + h_0bv \ge 0$: the $-L$ in $\mathcal{D}j$ leaves a nonnegative correction.

Put

$$P_0 = Rj(b-7v)^2, \qquad P_1 = RB(b-\tfrac{7}{3}v)^2,$$
$$P_2 = Kv(b-\tfrac{9}{2}v)^2, \qquad P_3 = vB(K+\tfrac{1}{4}R-\tfrac{1}{7}v)^2,$$
$$P_4 = (vb-\tfrac{14}{5}v^2-\tfrac{13}{4}Kv+\tfrac{4}{3}Rv)^2,$$
$$(p_0,p_1,p_2,p_3,p_4) = \left(\frac{337}{10000}, \frac{183}{2500}, \frac{411}{10000}, \frac{11461}{5000}, \frac{647}{10000}\right).$$

Exact expansion, with $b = B + 2v$, gives

$$z^2 - 12Rvb^2 + 6v^4 = (z+Q)^2 + S_1T_1 + \sum_{i=0}^{4} p_iP_i + \mathcal{W}(F_2) + L(x_0v^2 + h_0bv) + P_{\mathrm{rem}}.$$

For completeness, the entire remainder is specified by this table; every unlisted coefficient is zero.

| Monomial | Coefficient | Monomial | Coefficient |
|---|---|---|---|
| $R^2v^2$ | 223/160000 | $R^2vB$ | 4007/240000 |
| $R^2B^2$ | 1097/120000 | $RKv^2$ | 83/80000 |
| $RKvB$ | 3343/24000 | $Rv^3$ | 16259/23400000 |
| $Rv^2B$ | 30179/5460000 | $Rv^2j$ | 26/625 |
| $RvB^2$ | 107/15000 | $RvBj$ | 3/2500 |
| $RB^3$ | 121/32500 | $RB^2j$ | 11/10000 |
| $K^2v^2$ | 579/20000 | $K^2vB$ | 207/2500 |
| $Kv^3$ | 911/975000 | $Kv^2B$ | 349/420000 |
| $Kv^2j$ | 5011/5000 | $KvB^2$ | 133/130000 |
| $KvBj$ | 87/2500 | $v^4$ | 1661507/1521000000 |
| $v^3B$ | 633799/621075000 | $v^2B^2$ | 10657/1690000 |

All square prefactors $Rj, RB, Kv, vB, 1$ are nonnegative, as are $S_1T_1$, $L(x_0v^2 + h_0bv)$ and every remainder monomial. Integration removes only $\mathcal{W}(F_2)$, giving

$$\mathbb{E}_f[z^2 - 12vw^2 + 6v^4] \geq \frac{1661507}{1521000000}\mathbb{E}_f v^4 \geq \frac{1}{1000}\mathbb{E}_f v^4.$$

The surplus of the first coefficient over $1/1000$ is exactly $140507/1521000000 > 0$. The rational identity and positive table can be checked by finite arithmetic. Their application to this diffusion also uses the signs, quotient behaviour, integrability and vanishing flux proved above. Dividing the integrated inequality by $c^2$ returns

$$\mathbb{E}[\Phi_{xxxx}^2 - 12c\Phi_{xx}\Phi_{xxx}^2 + 6c^2\Phi_{xx}^4] \geq \frac{c^2}{1000}\mathbb{E}\Phi_{xx}^4.$$

This is the unscaled estimate needed by the gap argument. Section 4.5 passes it to the original coefficient and supplies strictness for the limiting solution. OpenAI (2026a), Lemma 4.1 and Proposition 4.2.

### 4.5 A fixed-time inequality and its plateau application

**Proposition 4.1 (coercivity at a fixed time).** Let $\gamma \in \mathcal{U}$ and let $X$ be its associated diffusion, started at zero. Suppose that $0 < t < 1$, $\gamma$ is continuous at $t$, and $c = \gamma(t) > 0$. With $a = \Phi_{xx}$, $b = \Phi_{xxx}$ and $d = \Phi_{xxxx}$, evaluated at $(t, X_t)$, one has

$$\mathcal{C}_{\gamma}(t) := \mathbb{E}[d^2 - 12cab^2 + 6c^2a^4] \geq \frac{c^2}{1000}\mathbb{E}a^4. \tag{4.1}$$

Neither minimality nor a constant neighbourhood of $t$ is assumed. This is the common estimate behind plateau curvature and positive density. When $\gamma$ is smooth near $t$, Itô's formula gives $q'' = \mathbb{E}b^2 - 2\gamma\mathbb{E}a^3$ and

$$(\mathbb{E}b^2)' = \mathbb{E}d^2 - 6\gamma\mathbb{E}(ab^2), \qquad (\mathbb{E}a^3)' = 3\mathbb{E}(ab^2) - 3\gamma\mathbb{E}a^4.$$

Differentiating the first identity therefore gives

$$q'''(t) + 2\gamma'(t)\mathbb{E}a^3 = \mathcal{C}_{\gamma}(t). \tag{4.2}$$

On a plateau the coefficient derivative vanishes. At a smooth minimizer, $q = t$ makes the third derivative of $q$ vanish instead.

**Proof.** Fix $t < T < T' < 1$ and a shrinking interval $[t - \delta_n, t + \delta_n) \subset (0, T)$. Approximate $\gamma$ by

$$h_n = \max\{\epsilon_n, \min\{K_n, \eta_n\lfloor\gamma/\eta_n\rfloor\}\},$$

where $0 < \epsilon_n < c < K_n$, $\epsilon_n, \eta_n \downarrow 0$ and $K_n \uparrow \infty$. Use $\min(c, h_n)$ to the left of the interval, $c$ on it, and $\max(c, h_n)$ to its right. Monotonicity makes these positive, nondecreasing, right-continuous finite-step coefficients $\gamma_n$. Clipping toward $c$ outside the interval cannot increase the error, and

$$\|\gamma_n - \gamma\|_1 \leq \epsilon_n + \eta_n + \int_0^1 (\gamma - K_n)_+ \, ds + 2\delta_n \sup_{|s-t|\leq\delta_n} |\gamma(s) - c| \longrightarrow 0. \tag{4.3}$$

Integrability controls the capped tail and continuity at $t$ the last term. Choose smooth terminal parameters $M_n \geq \max(n, K_n)$. The terminal values then converge uniformly to $|x|$, and the coefficient masses have a common bound. The finite-step inequality of §4.4 applies at $t$, where every $\gamma_n$ equals the same positive $c$.

On a fixed strip containing $t$ and ending before one, compact-strip regularity gives uniform spatial convergence and bounds through order four. The next derivative bounds give a common spatial Lipschitz constant for those derivatives. Coupling with the same Brownian motion and applying Gronwall gives a deterministic bound $\sup_{s\leq T} |X_s^n - X_s| \to 0$. If $D_k^n = \partial_x^k \Phi_n$ and $D_k = \partial_x^k \Phi$, then, for $1 \leq k \leq 4$,

$$|D_k^n(t, X_t^n) - D_k(t, X_t)| \leq \|D_k^n - D_k\|_{\infty,[0,T]\times\mathbb{R}} + C_{k,T}|X_t^n - X_t| \longrightarrow 0.$$

After the nonnegative terms are discarded, the two surviving integrands $z_n^2 - 12v_nw_n^2 + 6v_n^4$ and $v_n^4$ are uniformly bounded, with $v_n = cD_2^n$, $w_n = cD_3^n$ and $z_n = cD_4^n$. Dominated convergence therefore passes their expectations to the limit. For completeness, let $V, W, Z$ bound both the limiting and approximating $v, w, z$, and let $e_v, e_w, e_z$ bound their coupled evaluation errors. Uniform spatial

convergence and the path bound above give these errors with $e_v, e_w, e_z \to 0$. Difference-of-powers identities give

$$|\mathbb{E}\mathcal{C}(v_n, w_n, z_n) - \mathbb{E}\mathcal{C}(v, w, z)| \le (12W^2 + 24V^3)e_v + 24VWe_w + 2Ze_z,$$
$$|\mathbb{E}v_n^4 - \mathbb{E}v^4| \le 4V^3 e_v,$$

where $\mathcal{C}(v, w, z) = z^2 - 12vw^2 + 6v^4$. These constants depend on the common strip and coefficient mass bound, not on plateau width. After division by $c^2$, this is (4.1). Only spatial moments have been passed to the limit. This transfer requires neither convergence of $s_n$ or $j_n$ nor a uniform bound on their tail constants: those auxiliary quantities have already disappeared after integration. OpenAI (2026a), Lemmas 2.1–2.3; Proposition 5.1.

**The spatial inequality at a coefficient jump.** Continuity at $t$ is a sufficient hypothesis in Proposition 4.1 and in the retained sampled formal interface. The ordinary spatial-moment argument permits a little more. Fix $c = \gamma(t) > 0$ and replace $\gamma$ by $c$ on $[t - \delta, t + \delta)$, leaving it unchanged elsewhere. Monotonicity preserves both joins and right continuity. For fixed $0 < \delta_0 < \min(t, 1 - t)$ and $0 < \delta \le \delta_0$,

$$\|\gamma_\delta - \gamma\|_1 \le 2\delta\, \gamma(t + \delta_0) \longrightarrow 0.$$

Thus local boundedness suffices for the same strip stability and coupled expectation limit. Each replacement has a positive plateau at $t$, so (4.1) holds at a jump too, with the stated right-continuous value $c$. This extension concerns spatial moments only: it supplies no time derivative $q'''(t)$ or $\gamma'(t)$ at a jump and does not remove a hypothesis from the retained Lean declaration. Neither application below requires this extension.

**Corollary 4.2 (positive plateaus).** Suppose now that $\gamma = c > 0$ on an open interval $I \subset (0, 1)$ and $t \in I$. Then (4.2) gives $q'''(t) = \mathcal{C}_\gamma(t)$. To establish strictness for this limiting solution, choose $b \in I$ later than $t$ and put $\psi = \Phi_\gamma(b, \cdot)$. This is smooth, convex, even and a bounded distance from $|x|$, so $\psi'$ cannot be constant. The Cole–Hopf formula gives

$$\Phi_{xx}(t, x) = \mathbb{E}^{\mathrm{tilt}}_{t,x}\psi'' + c\,\mathrm{Var}^{\mathrm{tilt}}_{t,x}(\psi'),$$
$$\nu_{t,x}(dy) = \frac{e^{c\psi(y)} p_{b-t}(y - x)}{\int_{\mathbb{R}} e^{c\psi(z)} p_{b-t}(z - x)\, dz}\, dy.$$

The tilted density is positive everywhere. Its variance of the nonconstant continuous function $\psi'$ is strictly positive; its mean of $\psi''$ is nonnegative. Thus $v(t, x) > 0$ for every $x$, and $\mathbb{E}v(t, X_t)^4 > 0$. Combining the limiting inequality with the differentiated identity proves $q'''(t) \ge (1000c^2)^{-1}\mathbb{E}v(t, X_t)^4 > 0$. This proves strict convexity of $q'$ on $I$. OpenAI (2026a), Lemma 2.3; Proposition 5.1.

### 4.6 Why the zero interval remains a separate case

When $c = 0$, the heat equation gives $q''(t) = \mathbb{E}\Phi_{xxx}(t, X_t)^2$ without division by $c$. Start the drift-free evolution at an earlier point of $I$; its Gaussian transition kernel makes the law of $X_t$ have a positive density everywhere. If the expectation were zero, continuity would force $\Phi_{xxx}(t, \cdot) \equiv 0$. Then $\Phi(t, \cdot)$ would be an even quadratic polynomial. Its one-Lipschitz bound forces that polynomial to be constant, contradicting its bounded distance from $|x|$. Therefore $q''(t) > 0$, giving strict increase of $q'$. OpenAI (2026a), Lemma 2.3.

The analytic certificate is thus a proved input to the gap argument in this source, with the stated model and regularity assumptions. An abstract polynomial or flux lemma assumes those analytic hypotheses; the finite-step comparisons and the transfer argument discharge them. Inspection of the corresponding Lean sources records that construction, while compilation, scoped dependency extraction and Comparator replay remain distinct checks, with their observed outcomes reported in §13.

## 5. Excluding terminal, interior and initial gaps

**Theorem 5.1 (full support).** In the pure zero-field SK model with $\xi(t) = t^2/2$, let $\gamma \in \mathcal{U}$ minimize $\mathcal{P}$. Then $\operatorname{supp}\mu_\gamma = [0,1)$ and $\gamma$ is strictly increasing. For every associated diffusion started at zero, $\mathbb{E}u(t,X_t)^2 = t$ and $\mathbb{E}a(t,X_t)^2 = 1$ for $0 \le t < 1$.

Keep the same minimiser, diffusion and moment function $q$ throughout. Section 3 has established $G \ge 0$, $G = 0$ on the support $S = \operatorname{supp}\mu_\gamma$, and the contact conditions $q(s) = s$, $q'(s) \le 1$ at support points. At zero, the value identity comes from symmetry, and the slope bound requires $0 \in S$. Section 4 supplies curvature on constant intervals for every admissible coefficient. These premises are available before any support gap is excluded. We first deal with the endpoint at one, then use the curvature estimate on the remaining intervals.

For a terminal interval $(a,1)$, including empty support $[0,1)$, the cusp estimate gives $1 - q(t) \gtrsim \sqrt{1-t}$ and hence $G < 0$ near one. On a positive plateau with two contact endpoints, strict convexity puts $q'$ below its endpoint chord, although its integral must equal the interval length. On a zero plateau the heat argument makes $q'$ strictly increasing, below the right contact slope. At an initial gap, use the one-sided contact slope if $0 \in S$; otherwise the coefficient is zero and symmetry supplies $q(0) = 0$.

**Why a terminal plateau violates the variational inequality.** Suppose that $\gamma = c$ on $(a,1)$, where $0 \le a < 1$ and $c \ge 0$ is finite. The endpoint 1 is outside the support domain, so there is no contact condition there to complete an endpoint-chord argument. Instead we will prove

$$1 - q(t) \ge C\sqrt{1-t} \tag{5.1}$$

for some $C > 0$ and every $t$ sufficiently close to one. The proof separates two facts: the diffusion puts positive probability density near zero, and the smoothed terminal slope has a definite deficit from $\pm 1$ in a shrinking window around zero.

Fix $s \in (a,1)$ and let $\rho_s$ be the law of $X_s$. Choose $R < \infty$ such that $p_0 = \rho_s([-R,R]) > 0$. For $c > 0$, set

$$h(t,y) = \mathbb{E}e^{c|y+\sqrt{1-t}Z|}, \qquad \Phi(t,y) = c^{-1}\log h(t,y),$$

where $Z$ is standard normal. The drift on this interval is $c\Phi_x = h_x/h$, and the transition density is

$$k_{s,t}(x,y) = p_{t-s}(y-x)\frac{h(t,y)}{h(s,x)}, \qquad p_v(z) = (2\pi v)^{-1/2}e^{-z^2/(2v)}. \tag{5.2}$$

The heat-semigroup identity normalises this kernel; equivalently, for Brownian motion started at $(s,x)$, $h(t,B_t)/h(s,x)$ is the positive mean-one change of measure which adds the drift $h_x/h$. For $c = 0$, the same kernel formula applies with $h = 1$, and the value function is the ordinary heat evolution of $|x|$. OpenAI (2026a), Lemma 6.3, proof of (6.6).

Put $L = 1 - s$ and $\delta = L/2$. For $s + \delta \le t < 1$, $|x| \le R$ and $|y| \le 1$,

$$p_{t-s}(y-x) \ge (2\pi L)^{-1/2}e^{-(R+1)^2/(2\delta)} =: k_* > 0,$$

while $h(t,y) \geq 1$ and $h(s,x) \leq e^{cR}\mathbb{E}e^{c\sqrt{L}|Z|} =: M < \infty$. Integrating (5.2) over the part of $\rho_s$ in $[-R,R]$ therefore gives a density $f_t$ of $X_t$ with

$$f_t(y) \geq p_0 k_*/M =: m > 0 \quad (s+\delta \leq t < 1,\ |y| \leq 1). \tag{5.3}$$

Only positive mass of a compact set was needed at time $s$; no density assumption on $\rho_s$ has entered.

Now write $\varepsilon = 1-t$ and restrict to $|y| \leq \sqrt{\varepsilon}$. Differentiating the terminal heat formula gives, with the unweighted interpretation for $c = 0$,

$$u(t,y) = \frac{\mathbb{E}[\operatorname{sgn}(y+\sqrt{\varepsilon}Z)e^{c|y+\sqrt{\varepsilon}Z|}]}{\mathbb{E}e^{c|y+\sqrt{\varepsilon}Z|}}.$$

The denominator is at most $D = e^c\mathbb{E}e^{c|Z|} < \infty$. Each sign has unnormalised mass at least $p_* := \mathbb{P}(Z \geq 1) > 0$: the threshold $|y/\sqrt{\varepsilon}|$ is at most one, and the exponential weight is at least one. Hence each sign has tilted probability at least $\theta = p_*/D \in (0, 1/2)$. It follows that

$$|u(t,y)| \leq 1-2\theta, \qquad 1-u(t,y)^2 \geq 4\theta(1-\theta) =: \kappa > 0.$$

Multiplying this uniform deficit by the density lower bound (5.3) and the window width $2\sqrt{\varepsilon}$ yields

$$1-q(t) = \int_{\mathbb{R}} (1-u(t,y)^2) f_t(y)\,dy \geq 2m\kappa\sqrt{1-t}.$$

This proves (5.1) with $C = 2m\kappa$. OpenAI (2026a), Lemma 6.3, density and terminal-window estimates.

The sign of the contradiction is now explicit. For $s'$ sufficiently close to one, the estimate holds at every $t \in [s',1)$, so

$$\begin{aligned} G(s') &= \int_{s'}^1 (q(t)-t)\,dt \\ &\leq \frac{(1-s')^2}{2} - \frac{2C}{3}(1-s')^{3/2} < 0. \end{aligned}$$

The final inequality holds once $\sqrt{1-s'} < 4C/3$. It contradicts the nonnegative cost of adding an atom at $s'$. This proof covers $c = 0$ as well as $c > 0$. In particular, empty support is impossible: then $\mu_\gamma = 0$ and $\gamma = 0$ on all of $[0,1)$. The terminal-plateau conclusion was already recorded in Auffinger, Chen and Zeng (2020), Remark 7; the estimates above supply the quantitative argument needed here.

The terminal gap is now excluded. The support proof resumes with the remaining finite components; the next argument gives a shorter route to the same exclusion, without its quantitative bound.

**A shorter qualitative exclusion.** The existing curvature input also excludes a terminal support gap without estimating its boundary layer. Here is the elementary calculus step. Suppose $q \in C^1([r,1))$, $q(r) = r$, $q'(r) \leq 1$, $q \leq 1$, and $q'$ is strictly convex on $(r,1)$. If $q(t) \geq t$ at some $r < t < 1$, strict integration below the finite endpoint chord gives

$$t - r \leq \int_r^t q'(v)\,dv < \frac{t-r}{2}(q'(r)+q'(t)),$$

so $q'(t) > 1$. For $t < s < 1$, convexity then gives

$$q'(s) \geq q'(t) + \frac{s-t}{t-r}(q'(t)-q'(r)) \geq q'(t) > 1.$$

Consequently $q(s) \geq t+(s-t)q'(t) > 1$ for $s$ sufficiently close to one, a contradiction. Thus $q(t) < t$ throughout $(r, 1)$. All chord endpoints lie below one; no derivative at one is used.

To close the empty-support case using the same preceding inputs, observe that empty Stieltjes support forces $\gamma = 0$ everywhere. The zero-interval argument makes $q$ strictly convex on $[0, 1)$, whereas the positive-plateau argument above makes $q'$ strictly convex. Here $q(0) = 0$ and $0 \leq q \leq 1$. For $0 < t < s < 1$, convexity gives

$$q(t) \leq \frac{t}{s} q(s) \leq \frac{t}{s}.$$

Letting $s \uparrow 1$ in this bound gives $q(t) \leq t$. Now fix $t < u < 1$; strict convexity on the finite interval $[0, u]$ gives

$$q(t) < \frac{t}{u} q(u) \leq t.$$

Thus the bounded continuous integrand $q(t) - t$ is strictly negative on $(0, 1)$, so $G(0) < 0$, contrary to the allowed addition of mass at zero. Neither a value nor a derivative at one has been assumed. The explicit zero-coefficient calculation additionally gives $D_+\mathcal{P}(0)[1] = 1/4 - 1/\pi$; the qualitative argument needs only its sign.

A nonempty support with a terminal complementary component has a last point $r < 1$. Its plateau has positive height, since a neighbourhood of $r$ has positive measure. The contact values and slope at $r$, including the one-sided case $r = 0$, and positive-plateau curvature meet the calculus hypotheses. Since $0 \leq q \leq 1$, integration gives $G(r) < 0$, contradicting atom addition. This is an ordinary consequence of the previously stated inputs. The quantitative proof above is retained for its stronger estimate (5.1).

**The remaining gaps have a finite right contact point.**

The support is relatively closed in $[0, 1)$. Its complement is a union of relative open intervals, and the Stieltjes measure vanishes on each such interval. For $x < y$ in one interval,

$$\gamma(y) - \gamma(x) = \mu_\gamma((x, y]) = 0,$$

so $\gamma$ is constant there. The terminal argument excludes a component ending at one. Every remaining component therefore ends at a point $b < 1$ of $S$.

First consider a gap $(a, b)$ with both endpoints in $S$, allowing $a = 0$. Compact-strip regularity gives continuity of $q'$ on $[a, b]$, even if $\gamma$ jumps at an endpoint. Contact gives

$$q(a) = a, \qquad q(b) = b, \qquad q'(a), q'(b) \leq 1. \tag{5.4}$$

If the constant coefficient is $c > 0$, Section 4 makes $q'$ strictly convex in the gap. Continuity extends the chord comparison to its endpoints, giving

$$q'(t) < \frac{b-t}{b-a} q'(a) + \frac{t-a}{b-a} q'(b) \leq 1, \qquad a < t < b. \tag{5.5}$$

Integrating contradicts $q(b) - q(a) = b - a$. Equivalently, the mean value theorem applied to (5.4) requires an interior point with $q' = 1$, whereas (5.5) excludes every such point. The endpoint derivatives need only be at most one; equality has not been assumed.

With $h = q'$ and $C_{a,b}$ its endpoint chord, the contradiction is the exact area identity

$$(b-a) - \int_a^b h = (b-a)\left(1 - \frac{h(a)+h(b)}{2}\right) + \int_a^b (C_{a,b} - h) > 0.$$

The first term is nonnegative; strict convexity makes the second positive. Only continuity of $q'$ is needed at the ends. For example, a jump $\Delta\gamma$ at $s$ gives the one-sided identity

$$q''(s+) - q''(s-) = -2\Delta\gamma\, \mathbb{E}a(s, X_s)^3.$$

It follows by applying Itô's formula to $a^2$ and using continuity of its derivative moments on a compact strip. Taking $\gamma = 0$ before $s = 1/2$ and $\gamma = 1/2$ afterward makes this jump strictly negative; the terminal Cole–Hopf solution has positive Hessian. Thus continuity of $q'$ does not license classical higher time derivatives across coefficient jumps. OpenAI (2026a), proof of Corollary 6.4.

If $c = 0$, the separate heat argument in Section 4 makes $q'$ strictly increasing. Then $q'(t) < q'(b) \le 1$ for $a < t < b$, and the same contradiction follows. The left slope bound is unnecessary in this case. OpenAI (2026a), proof of Theorem 1.1 after Lemma 6.3.

There are two distinct possibilities at the left endpoint. If $(0, b)$ is a gap and $0 \in S$, the preceding proof applies using the one-sided contact slope bound at zero. Because $(0, b)$ contains no mass, the support condition at zero forces an atom there and hence $c = \gamma(0) > 0$. Its absence has not been proved at this stage. If instead the complementary component is $[0, b)$ with $0 \notin S$, there is no mass anywhere in $[0, b)$. Thus $\gamma = 0$ there. The identity $q(0) = 0$ comes from zero-field symmetry, and the right endpoint gives $q(b) = b$, $q'(b) \le 1$. Here $q''(t) > 0$ for $0 < t < b$; at zero evenness gives $\Phi_{xxx}(0, 0) = 0$ and hence $q''(0) = 0$. Strict increase on the open interval still forces $q' < 1$ throughout $(0, b)$, an impossibility.

**Full support and the moment identities.**

The cases are exhaustive, so $S = [0, 1)$. Only now can contact be applied at every $0 < t < 1$, giving $q(t) = t$ throughout the domain; the value at zero was already known. Differentiating and using the gradient-martingale identity gives

$$\mathbb{E}a(t, X_t)^2 = q'(t) = 1, \qquad 0 \le t < 1,$$

with the right derivative at zero.

Finally, every nonempty open subinterval $(a, b)$ receives positive mass. Therefore

$$\gamma(b) - \gamma(a) = \mu_\gamma((a, b]) > 0 \qquad (0 \le a < b < 1).$$

The proof establishes full support, strict increase and the two moment identities. Smooth absolute continuity, including the later conclusion $\gamma(0) = 0$, uses the source's further argument from saturated consistency; it has not been assumed to obtain contact or to exclude an initial gap. OpenAI (2026a), conclusion of the proof of Theorem 1.1. A weaker analytic condition suffices for the bounded-gap contradiction. If $h \in C([a, b]) \cap C^2((a, b))$ has the displayed endpoint and integral data, then $h' = 0 \Rightarrow h'' > 0$ is already impossible: either $h \le 1$ everywhere, forcing $h \equiv 1$ by its integral, or it attains an interior maximum above one, where $h' = 0$ and $h'' \le 0$. This is an elementary deduction from these premises. Chen's conditional crossing theorem, with its own hypotheses, is of this form; §4 proves the stronger $q''' > 0$ throughout a positive plateau, without a minimizer premise. See Chen, arXiv:2607.18032v1, Proposition 4.2 and the proof of Proposition 4.3. Neither statement supplies terminal contact at the excluded endpoint one.

## 6. Recovering the value from a martingale

**Proposition 6.1 (terminal value).** For the minimizer and diffusion of Theorem 5.1 there is a spin-valued terminal variable $U_1$ such that $u(t, X_t) = \mathbb{E}[U_1 \mid \mathcal{F}_t]$ and

$$P_* = \mathbb{E}[U_1 B_1] = \int_0^1 \mathbb{E}a(t, X_t)\, dt. \tag{6.1}$$

Its finite-cutoff tail lies in $[0, 1-T]$. This proposition uses the attained Parisi formula in addition to the support theorem.

The bounded martingale $u(t, X_t)$ converges almost surely and in $L^2$ to $U_1$. Its second moment tends to one because $q(t) = t$. Since $|U_1| \le 1$, $\mathbb{E}U_1^2 = 1$ forces $U_1^2 = 1$ almost surely. Uniform integrability identifies $u(t, X_t) = \mathbb{E}[U_1 \mid \mathcal{F}_t]$, which is the conditional-expectation identity used below.

The drift bound $|\gamma u| \le \gamma \in L^1$ gives convergence of $X_t$ to $X_1$. Uniform terminal convergence of $\Phi$ to $|x|$, together with convexity, gives $u(t, x_t) \to \operatorname{sgn}(x)$ when $x_t \to x \neq 0$. Thus $U_1 X_1 = |X_1|$, including when $X_1 = 0$; no proof that this event has probability zero is needed.

Itô's formula gives

$$\mathbb{E}\Phi(t, X_t) = \Phi(0,0) + \frac{1}{2}\int_0^t \gamma(s) s\, ds.$$

The terminal limit first gives $\Phi(0,0) = \mathbb{E}|X_1| - \frac{1}{2}\int_0^1 s\gamma(s) ds$. Subtracting the second half from the definition of the functional yields $\mathcal{P}(\gamma) = \mathbb{E}|X_1| - \int_0^1 s\gamma(s)\, ds$. The ground-state Parisi formula identifies this with $P_*$. The diffusion equation, conditional expectation and Fubini give

$$\mathbb{E}[U_1 X_1] = \mathbb{E}[U_1 B_1] + \int_0^1 \gamma(s)\mathbb{E}[u(s, X_s)^2]\, ds.$$

Bounded $u, U_1$ and integrable $\gamma$ justify these terms. Combining the identities and using stochastic integral covariance gives

$$P_* = \mathbb{E}|X_1| - \int_0^1 t\gamma(t)\, dt = \mathbb{E}[U_1 B_1] = \int_0^1 \mathbb{E}a(t, X_t)\, dt.$$

The last integral is absolutely bounded by one because $\mathbb{E}|a| \le (\mathbb{E}a^2)^{1/2} = 1$. Full support supplies $\mathbb{E}a^2 = 1$, whereas the value uses $\int \mathbb{E}a$. Substituting the second moment into the value identity would change the assertion. OpenAI (2026a), §7. The prediction error and the energy tail have different sizes:

$$\|U_1 - u(T, X_T)\|_2^2 = 1 - T,$$

$$P_* - \int_0^T \mathbb{E}a(t, X_t)\, dt = \mathbb{E}[(U_1 - u(T, X_T))(B_1 - B_T)] \in [0, 1-T].$$

The first identity is orthogonal projection onto $\mathcal{F}_T$. Martingale orthogonality removes the past Brownian component in the second; Cauchy–Schwarz bounds the product by $1-T$, and convexity makes it nonnegative. No uniform terminal curvature bound is needed. The stochastic value calculation has a direct antecedent in Sellke, arXiv:2105.03506v4, Definition 1.3 and Lemma 3.11. There, "optimizable" means the contact identity $\mathbb{E}u(t, X_t)^2 = t$ on $[q, 1)$, with $q$ the infimum of the support of the coefficient function. Lemma 3.11 derives the finite-cutoff value identity by applying

Itô's formula to $\Phi$ and $xu$, before taking the terminal limit. This is the same value-recovery mechanism used here, specialized to zero field and $\xi'' = 1$.

A cutoff must retain the full Parisi penalty. Set $\Psi(T,x) = \Phi(T,x) - xu(T,x)$ and $I(T) = \int_0^T \mathbb{E}a(t,X_t)\,dt$. Subtracting the complete penalty from the finite-time Itô identity gives

$$\mathcal{P}(\gamma) - I(T) = \mathbb{E}\Psi(T,X_T) + \frac{1}{2}\int_0^T \gamma(t)(q(t)-t)\,dt - \frac{1}{2}\int_T^1 t\gamma(t)\,dt.$$

Under contact the middle term vanishes. The final term is absent from the displayed equality in El Alaoui, Montanari and Sellke, arXiv:2001.00904v1, §6.2, pp. 32–33. Its sign is nonpositive, so the upper bound by $\mathbb{E}\Psi(T,X_T)$ used there remains valid; its magnitude tends to zero by integrability. This finite-cutoff correction is consistent with the exact terminal identity above.

**A coefficient budget from the first curvature moment.** Full support also bounds the minimizing coefficient. Put $m(t) = \mathbb{E}a(t,X_t)$. The compact-strip identity

$$da(t,X_t) = -\gamma(t)a(t,X_t)^2\,dt + b(t,X_t)\,dB_t$$

holds before any smoothness conclusion for $\gamma$; the bounded strip derivatives make its stochastic integral a martingale. OpenAI (2026a), proof of Corollary 6.4. At zero, $X_0 = 0$, $a(0,0) \geq 0$ and $\mathbb{E}a(0,X_0)^2 = 1$ give $a(0,0) = 1$. Taking expectations and using $\mathbb{E}a(t,X_t)^2 = 1$ therefore yields

$$m(t) = 1 - \int_0^t \gamma(s)\,ds, \qquad 0 \leq t < 1.$$

Convexity gives $m(t) \geq 0$. Letting $t \uparrow 1$ and using monotonicity of $\gamma$ on every tail gives

$$\int_0^1 \gamma(s)\,ds \leq 1, \qquad 0 \leq \gamma(t) \leq \frac{m(t)}{1-t} \leq \frac{1}{1-t}.$$

Indeed, $(s-t)\gamma(t) \leq \int_t^s \gamma \leq m(t)$ for $t < s < 1$. Integrating the first-moment identity also gives

$$I(T) = T - \int_0^T (T-s)\gamma(s)\,ds, \qquad P_* = 1 - \int_0^1 (1-s)\gamma(s)\,ds.$$

The second equality uses (6.1), so it retains that proposition's ground-state identification. These are ordinary consequences in the normalization $\xi'' = 1$. The budget alone does not identify the terminal first moment. The following endpoint argument supplies that additional information.

**Proposition 6.2 (terminal normalization in zero field).** For the minimizer and diffusion of Theorem 5.1,

$$\lim_{t\uparrow 1} m(t) = 0, \qquad \int_0^1 \gamma(s)\,ds = 1, \qquad m(t) = \int_t^1 \gamma(s)\,ds,$$

and consequently

$$P_* = \int_0^1 s\gamma(s)\,ds.$$

Thus $\gamma(s)\,ds$ has total mass one; this does not normalize the Stieltjes measure $d\gamma$. This is an ordinary analytic consequence, separate from the selected formal endpoints.

**Proof.** The backward comparison in §4.3 gives $\Phi_{xxx}(t,x) \le 0$ for $x > 0$ in every positive finite-step approximation. Compact-strip convergence of the third spatial derivative passes this inequality to $\gamma$. Together with evenness and convexity, it says that $a(t,x)$ is nonnegative, even and nonincreasing in $|x|$. These inputs are OpenAI (2026a), Lemma 3.2 and Lemma 2.1; they do not require differentiability of $\gamma$. Uniform terminal convergence to $|x|$ and convex difference quotients give $u(t,x) \to 1$ at every fixed $x > 0$. For each $\delta > 0$,

$$0 \le a(t,\delta) \le \frac{2}{\delta} \int_{\delta/2}^{\delta} a(t,x)\,dx = \frac{2}{\delta}(u(t,\delta) - u(t,\delta/2)) \longrightarrow 0.$$

The spatial monotonicity is essential here: convergence of convex functions alone would not imply convergence of their second derivatives.

It remains to control the probability of the shrinking region where curvature can be large. Oddness and monotonicity give $u(t,x) \ge 0$ for $x \ge 0$, so the drift points away from zero. Tanaka's formula for $R_t = |X_t|$ gives

$$R_t = W_t + V_t + L_t, \qquad W_t = \int_0^t \operatorname{sgn}(X_s)\,dB_s, \qquad V_t = \int_0^t \gamma(s)u(s,R_s)\,ds.$$

Here $L$ is the local time of $X$ at zero. The occupation-time formula and $d\langle X\rangle_s = ds$ imply $\int_0^t \mathbf{1}_{\{X_s=0\}}\,ds = 0$. Thus $\langle W\rangle_t = t$, and the continuous martingale $W$ is Brownian motion. The process $V$ is nondecreasing and starts at zero; $L$ increases only when $R = 0$. The reflection formula therefore gives, pathwise,

$$R_t = W_t + V_t - \min\left(0, \inf_{s\le t}(W_s + V_s)\right) \ge W_t - \min\left(0, \inf_{s\le t} W_s\right) =: R^0_t.$$

Indeed $0 \le V_s \le V_t$ bounds the subtracted minimum by $\min(0, \inf_{s\le t} W_s) + V_t$. No independence of $V$ and $W$ is needed. Reflected Brownian motion $R^0_t$ has the one-time law $|B_t|$, so for $t > 0$,

$$\Pr(|X_t| \le \delta) \le \Pr(|B_t| \le \delta) \le \sqrt{\frac{2}{\pi}}\frac{\delta}{\sqrt{t}}.$$

Splitting at $|X_t| = \delta$, using spatial monotonicity outside and Cauchy–Schwarz with $\mathbb{E}a(t,X_t)^2 = 1$ inside, yields

$$0 \le m(t) \le a(t,\delta) + \sqrt{\Pr(|X_t| \le \delta)}.$$

First let $t \uparrow 1$ and then $\delta \downarrow 0$. This proves $m(1-) = 0$. The compact-strip identity for $m$ now gives total mass one and the tail formula. Tonelli and (6.1) give the stated expression for $P_*$. In particular,

$$0 \le P_* - I(T) = \int_T^1 (s-T)\gamma(s)\,ds \le (1-T)m(T) = o(1-T).$$

No numerical rate for $m(T) \to 0$ or for the required spatial derivative bounds follows from this proof.

The endpoint distinction also matters when comparing editions. Sellke, arXiv:2105.03506v4, Lemma 2.9, equation (2.7) states the expected-Hessian tail equality. In version 6, Lemma 2.9, equation (2.8), and Remark 2.10, the general statement is an inequality; the remark obtains equality indirectly for even mixtures. The specialized zero-field proof above supplies its own endpoint argument from the shape and contact identities, rather than substituting $t = 1$ into a compact-strip formula or relying on the older equality in its full stated generality.

## 7. A finite Gaussian spin and its adjacent correlations

**Proposition 7.1 (finite Gaussian approximation).** For a fixed minimizer and its diffusion, the construction below has positive active normalizers for all sufficiently fine meshes at every fixed $0 < T < 1$. Its adjacent correlation sum converges to $\int_0^T \mathbb{E}a(t, X_t)\, dt$, and this integral tends to $P_*$ as $T \uparrow 1$. The limits are taken in that order.

Keep the minimizer and its diffusion from §6. The object to construct is a finite Gaussian sign $F_N$ with two correlation vectors whose *adjacent* pairing

$$S_{T,N} = \sum_{j=0}^{N} A_j B_{j+1} \longrightarrow \int_0^T \mathbb{E}a(t, X_t)\, dt.$$

The shift by one is essential for the rooted energy identity in §8.1; §10.4 computes a case where the unshifted pairing has a different value. We first fix $0 < T < 1$, then a mesh $N \geq 1$, $\Delta = T/N$, $t_j = j\Delta$. On independent standard normals $Z_1, \dots, Z_{N+1}$, set

$$Y_0 = 0, \quad Y_j = Y_{j-1} + \sqrt{\Delta} Z_j + \Delta\gamma(t_{j-1})u(t_{j-1}, Y_{j-1}),$$

$$c_j = (\mathbb{E}a(t_{j-1}, Y_{j-1})^2)^{-1/2}, \qquad g_j = c_j a(t_{j-1}, Y_{j-1}) Z_j.$$

For sufficiently large $N$, all $c_j$, $1 \leq j \leq N$, are finite and positive. Couple the increments to Brownian motion. On the fixed strip, the gradient and curvature are bounded and Lipschitz. Monotonicity gives, for the left-grid projection $\pi_N$,

$$\int_0^T |\gamma(s) - \gamma(\pi_N(s))|\, ds \leq \Delta(\gamma(T) - \gamma(0)).$$

Drift error estimates and discrete Gronwall yield $\mathbb{E} \max_{j \leq N} |Y_j - X_{t_j}|^2 \to 0$, hence $\max_j |c_j - 1| \to 0$. The $g_j$ are orthonormal in $L^2$: they have unit variance and conditional mean zero given their past. Their predictable multipliers may depend on earlier coordinates, so orthonormality does not assert independence.

With $F_{\mathrm{G}}$ the standard Gaussian distribution function, define

$$F_N = \operatorname{sgn}(u(T, Y_N) + 2F_{\mathrm{G}}(Z_{N+1}) - 1).$$

$$A_j = \mathbb{E}[F_N g_j], \quad B_i = \mathbb{E}[F_N Z_i], \quad A_0 = 0.$$

The extra coordinate is independent and its distribution function is uniform. For any $v \in [-1, 1]$, the random sign of $v + 2U - 1$ has mean $v$. Therefore $\mathbb{E}[F_N \mid Z_1, \dots, Z_N] = u(T, Y_N)$ exactly. A deterministic sign of the gradient would not preserve this conditional mean. Rounding preserves correlations with functions of the first $N$ coordinates, which is exactly what the two coefficient vectors need. It does not make the whole rounded spin close in $L^2$ to that conditional mean. Indeed, with $m_N = u(T, Y_N)$,

$$\mathbb{E}[(F_N - m_N)^2] = 1 - \mathbb{E}m_N^2 \longrightarrow 1 - T.$$

Here $m_N \to u(T, X_T)$ in $L^2$ at fixed $T$ and full support gives $\mathbb{E}u(T, X_T)^2 = T$. The independent rounding variance disappears only in the later endpoint limit.

The approximation

$$M_N = \sum_{j=1}^{N} \sqrt{\Delta}\, a(t_{j-1}, Y_{j-1}) Z_j = u(T, Y_N) + o_{L^2}(1)$$

gives a common residual for both coefficient families. In $\mathbb{E}[M_N Z_i]$, a summand with $j < i$ is independent of centered $Z_i$; one with $j > i$ vanishes on conditioning on the first $j-1$ coordinates; and the $j = i$ term is $\sqrt{\Delta}\,\mathbb{E}a(t_{i-1}, Y_{i-1})$. Also $\mathbb{E}[M_N g_j] = \sqrt{\Delta}/c_j$ by orthonormality. Bessel's inequality applied separately to the orthonormal family $(g_j)$ and to the family $(Z_i)$ gives the vector estimates. Their union need not be orthogonal:

$$(A_j)_{j=1}^{N} = (\sqrt{\Delta}/c_j)_{j=1}^{N} + o_{\ell^2}(1),$$

$$(B_i)_{i=1}^{N} = (\sqrt{\Delta}\,\mathbb{E}a(t_{i-1}, Y_{i-1}))_{i=1}^{N} + o_{\ell^2}(1).$$

Coordinatewise errors would not control a sum of growing length. Put

$$e_N = \|u(T, Y_N) - M_N\|_2, \quad \alpha_j = \sqrt{\Delta}/c_j, \quad \beta_j = \sqrt{\Delta}\,\mathbb{E}a(t_{j-1}, Y_{j-1}) \quad (1 \le j \le N).$$

The fixed-$T$ Euler argument gives $e_N \to 0$. Bessel's inequality gives $\|A_{1:N} - \alpha\|_2, \|B_{1:N} - \beta\|_2 \le e_N$. Also $\|B_{1:N}\|_2 \le 1$ and $\|\alpha\|_2 \le C_T$ for sufficiently large $N$, with $C_T$ independent of the mesh. Cauchy–Schwarz gives

$$\begin{aligned}\left|\sum_{j=1}^{N-1} (A_j B_{j+1} - \alpha_j \beta_{j+1})\right| &\le \|A - \alpha\|_2 \|B_{2:N}\|_2 + \|\alpha_{1:N-1}\|_2 \|B - \beta\|_2 \\ &\le (1 + C_T) e_N.\end{aligned}$$

The unindexed differences in this inequality use coordinates $1, \dots, N$; restricting to the indicated subvector only decreases their norms. The rounding coordinate contributes at most $|A_N B_{N+1}| \le e_N + \sqrt{\Delta}/c_N$. Thus the dimension-free estimate is

$$\left|\sum_{j=0}^{N} A_j B_{j+1} - \Delta \sum_{j=1}^{N-1} \frac{\mathbb{E}a(t_j, Y_j)}{c_j}\right| \le (2 + C_T) e_N + \frac{\sqrt{\Delta}}{c_N} \longrightarrow 0.$$

The index shift selects the Hessian at $t_j$; $Z_{N+1}$ serves rounding rather than another Euler increment. This estimate is a fixed-strip scalar correspondence, not a circuit energy identity. Hence

$$\sum_{j=0}^{N} A_j B_{j+1} = \Delta \sum_{j=1}^{N-1} \frac{\mathbb{E}a(t_j, Y_j)}{c_j} + o(1) \longrightarrow \int_0^T \mathbb{E}a(t, X_t)\, dt.$$

The same vector estimates also give an unshifted sum with this limit:

$$\left|\sum_{j=1}^{N} A_j B_j - \sum_{j=1}^{N} \alpha_j \beta_j\right| \le (1 + C_T) e_N \longrightarrow 0,$$

$$\sum_{j=1}^{N} A_j B_j = \Delta \sum_{j=1}^{N} \frac{\mathbb{E}a(t_{j-1}, Y_{j-1})}{c_j} + o(1) \longrightarrow \int_0^T \mathbb{E}a(t, X_t)\, dt.$$

The latter is a left Riemann sum: $c_j \to 1$ uniformly and the Euler coupling replaces $Y_{j-1}$ by $X_{t_{j-1}}$. Equal continuum limits therefore do not identify the finite expression needed by the circuit. The

rooted energy calculation in §8.1 selects the adjacent pairing; the two finite pairings can differ, as §10.4 shows.

Only afterward let $T \uparrow 1$, obtaining $P_*$. The mesh count $N$ is unrelated to the spin-system size $n$. As in OpenAI (2026a), Proposition 7.2, these coefficients describe a finite Gaussian function, not yet a circuit on finite SK disorder.

The QAOA scalar input also uses the tail estimate $0 \le P_* - \int_0^T \mathbb{E}a(t, X_t)\, dt \le 1 - T$, from nonnegative curvature and $\mathbb{E}a^2 = 1$. This estimate and the fixed-strip regularity facts must be retained beside the coefficient limit; they are not all clauses of the selected finite-Gaussian statement. They enter the construction in OpenAI (2026b), §3.

Thus, for every accuracy, the same chosen minimizer supplies a cutoff $T$ and then one finite function $F_N$ with $|S_{T,N} - P_*|$ below that accuracy. This is the finite scalar input. Turning it into spins for a finite SK matrix uses the separate AMP theorem in §9.4; turning it into a common QAOA word uses §§8.1–8.6. Neither operation is part of the Gaussian limit.

### 7.1 Error estimates at a fixed cutoff

Fix the minimizer first and then $T < 1$. The bounds below have constants depending on that minimizer and strip; they give an $N^{-1/2}$ rate there, not a rate uniform as $T \uparrow 1$. Those derivative bounds are finite but have not been supplied as numerical inputs to an algorithm for choosing a mesh. The calculation is an ordinary deduction from the source's strip estimates, beyond the qualitative selected formal statement. Write $S_{T,N} = \sum_{j=0}^{N} A_j B_{j+1}$ and $f(t) = \mathbb{E}a(t, X_t)$, so $0 \le f(t) \le 1$. On the fixed strip let $K_T = \sup_{t \le T, x} |a(t, x)|$ and choose a spatial Lipschitz constant $L_T$ for $a$. Define

$$\delta_N = \max_{0 \le j \le N} \|Y_j - X_{t_j}\|_2, \qquad d_N = \max_{1 \le j \le N} \left|\mathbb{E}a(t_{j-1}, Y_{j-1})^2 - 1\right|.$$

These errors have a rate at each fixed strip, even if $\gamma$ has jumps. Put $\Gamma = \gamma(T)$ and $V = \gamma(T) - \gamma(0)$. Bounded spatial derivatives and the PDE give temporal Lipschitz constants

$$J = \tfrac{1}{2}\|u_{xx}\|_\infty + \Gamma K_T, \qquad H = \tfrac{1}{2}\|a_{xx}\|_\infty + \Gamma(K_T^2 + L_T)$$

for $u$ and $a$, respectively, on this strip. No derivative of $\gamma$ is used. Brownian coupling gives $\|X_s - X_{\pi_N(s)}\|_2 \le \sqrt{\Delta} + \Gamma\Delta$. Subtract the Euler and diffusion equations and split the drift difference on $[t_j, t_{j+1}]$ as

$$\gamma(s)(u(s, X_s) - u(t_j, Y_j)) + (\gamma(s) - \gamma(t_j))u(t_j, Y_j).$$

Put $A_T = \int_0^T \gamma(s)\, ds \le 1$ and $w_j = \int_{t_j}^{t_{j+1}} \gamma(s)\, ds$. Taking a pathwise maximum over grid times, Minkowski and discrete Gronwall give, with

$$R_N = A_T K_T \sqrt{\Delta} + [A_T(J + \Gamma K_T) + V]\Delta,$$

the bound

$$\delta_N \le \left\| \max_{j \le N} |Y_j - X_{t_j}| \right\|_2 \le e^{K_T A_T} R_N \le e^{K_T} R_N.$$

The recurrence weights the preceding maximal norm on interval $j$ by $K_T w_j$; their sum is $K_T A_T$. The term $V\Delta$ is the monotone quadrature bound for the coefficient. For the martingale residual, Itô isometry and the same increment estimate give

$$e_N \le K_T \delta_N + \sqrt{T}\{L_T \delta_N + L_T\sqrt{\Delta} + (H + \Gamma L_T)\Delta\}.$$

Thus $\delta_N, e_N = O_{\gamma,T}(\sqrt{\Delta})$. These are deductions from the compact-strip bounds and stochastic identities, not quantitative clauses of the selected Lean statement.

Since $a$ is bounded and Lipschitz, $d_N \le 2K_T L_T \delta_N \to 0$. Whenever $d_N \le 1/2$,

$$\sqrt{2/3} \le c_j \le \sqrt{2}, \qquad |1/c_j - 1| \le d_N, \qquad \|\alpha\|_2 \le \sqrt{T(1+d_N)}.$$

These estimates give an explicit sufficient condition for all the required normalisers to be positive. They do not assume positivity for every coarse mesh.

The first-moment identity in §6 makes $f = m$ nonincreasing, with $f(0) = 1$. Its right Riemann sum therefore satisfies

$$0 \le I(T) - \Delta \sum_{j=1}^{N} f(t_j) \le \Delta(f(0) - f(T)).$$

Removing the last term adds $\Delta f(T)$, so $0 \le I(T) - \Delta \sum_{j=1}^{N-1} f(t_j) \le \Delta$. The difference between the shifted interior sum and the integral is bounded by

$$\left| \Delta \sum_{j=1}^{N-1} \frac{\mathbb{E}a(t_j, Y_j)}{c_j} - \int_0^T f(t)\, dt \right| \le TK_T d_N + TL_T \delta_N + \Delta.$$

The first two terms replace the normalisers and Euler positions; the last is the preceding monotone quadrature estimate. Combining this with the vector and rounding-endpoint estimate gives

$$\begin{aligned} |S_{T,N} - P_*| \le (1-T)m(T) + (2 + \sqrt{T(1+d_N)})e_N + \sqrt{\Delta(1+d_N)} \\ + TK_T d_N + TL_T \delta_N + \Delta. \end{aligned}$$

In particular, for each fixed minimizer and $T < 1$ there are finite $C_{\gamma,T}$ and $N_0(\gamma, T)$ such that

$$|S_{T,N} - P_*| \le (1-T)m(T) + C_{\gamma,T}\sqrt{T/N} \qquad (N \ge N_0(\gamma, T)).$$

The constants are not uniform as $T \uparrow 1$. The coefficient budget bounds $\int \gamma$ and $\gamma$ on each fixed strip, but numerical bounds for all strip derivative constants and an effective mesh threshold are not supplied. The scalar tail bound $(1-T)m(T) \le 1 - T$ uses only monotonicity of the first moment; Proposition 6.2 further makes it $o(1-T)$. All terms after this scalar tail tend to zero at each fixed $T < 1$. Given a scalar tolerance $\zeta > 0$, first choose $T$ with $1 - T < \zeta/2$, then choose $N_0(\gamma, T, \zeta)$ so that their sum is below $\zeta/2$ for every $N \ge N_0$. This is the required finite choice of a Gaussian spin. The formula exposes the errors; it does not supply numerical values of $K_T, L_T, \delta_N, e_N$ or an effective mesh threshold. OpenAI (2026a), Lemma 2.1; OpenAI (2026a), Proposition 7.2.

A diagonal can be chosen using these thresholds. For prescribed $T_m \uparrow 1$, choose increasing $N_m$ large enough to make the mesh-dependent part of the fixed-$T_m$ bound less than $1/m$. An arbitrary increasing mesh schedule does not follow from the iterated limit alone. For example, the scalar array

$$h(T, N) = \min\{1, [N(1-T)]^{-1}\}$$

tends to zero in $N$ at every fixed $T < 1$, but $h(1 - 1/N, N) = 1$ for every $N \ge 2$. This is an illustration of the quantifier issue, not an asserted error law for the SK construction.

## 8. From the scalar formula to one ordinary QAOA word

On a single spin, let $Z = \mathrm{diag}(1,-1)$ and $X = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$; $Z_i$ and $X_i$ act on spin $i$ and as the identity on the other spins. Also set $Y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ and $|\pm\rangle = (|0\rangle \pm |1\rangle)/\sqrt{2}$. Then $X|\pm\rangle = \pm|\pm\rangle$ and $Z|+\rangle = |-\rangle$. The cost and mixer are

$$C_{n,J} = n^{-1/2} \sum_{i<j} J_{ij} Z_i Z_j, \qquad B_n = \sum_i X_i.$$

The two evolutions act differently:

$$e^{-i\theta C_{n,J}}|\sigma\rangle = e^{-i\theta H_n(\sigma)}|\sigma\rangle, \qquad e^{-i\beta B_n} = \prod_{i=1}^{n} (\cos\beta\, I - i \sin\beta\, X_i).$$

The cost evolution changes phases while preserving spin-basis probabilities; the mixer combines amplitudes of different spin configurations. Cost evolution alone leaves the initial uniform probabilities unchanged, and a mixer applied first only multiplies $|+\rangle^{\otimes n}$ by $e^{-in\beta}$. Alternating them lets the cost-dependent phases change subsequent measurement probabilities. Farhi, Goldstone, Gutmann and Zhou (2022), §2, equations (1)–(6).

A depth-$p$ state, with $p$ counting cost–mixer pairs, is

$$|\psi_{n,p,J}(\gamma,\beta)\rangle = e^{-i\beta_p B_n} e^{-i\gamma_p C_{n,J}} \cdots e^{-i\beta_1 B_n} e^{-i\gamma_1 C_{n,J}} |+\rangle^{\otimes n}.$$

Here the real angle vectors $\gamma, \beta$ are deterministic; $\gamma$ is separate from the Parisi coefficient $\gamma(t)$. A *word* means a finite ordered product of cost and mixer exponentials. Its pattern and angles are fixed independently of $n, J$; its cost operator depends on $n, J$. Define

$$v_p(\gamma,\beta) = \lim_{n\to\infty} \frac{1}{n} \mathbb{E}_J \langle \psi_{n,p,J} | C_{n,J} | \psi_{n,p,J} \rangle, \qquad Q_p = \sup_{\gamma,\beta \in \mathbb{R}^p} v_p.$$

The fixed-depth, fixed-angle infinite-size limit is the input from Farhi, Goldstone, Gutmann and Zhou; it does not contain the later depth conclusion; see Farhi, Goldstone, Gutmann and Zhou (2022). The source target is $\lim_{p\to\infty} Q_p = P_*$. Its needed lower bound has the literal quantifiers

$$\forall \varepsilon > 0 \quad \exists p \in \mathbb{N}, \gamma, \beta \in \mathbb{R}^p : \quad \lim_{n\to\infty} \frac{1}{n} \mathbb{E}_J \langle \psi_{n,p,J} | C_{n,J} | \psi_{n,p,J} \rangle \geq P_* - \varepsilon.$$

The same finite list of gates and angles is used for every $n, J$; this is a limiting expected-energy conclusion. A cost expectation is bounded above by its maximum eigenvalue, and zero-angle padding makes $Q_p$ nondecreasing. It therefore suffices to construct one finite word for every accuracy. Fixed-instance convergence at increasing depth is a different quantifier order. Farhi, Goldstone and Gutmann's original QAOA argument obtains that convergence by adiabatic evolution and its Trotter approximation, with depth allowed to depend on the instance; it does not supply the size-independent word required here. Farhi–Goldstone–Gutmann (2014), equation (10) and §VI, pp. 10–11; OpenAI (2026b), Theorem 1.1.

The construction first describes local operations on a large tree, where neighbouring subtrees supply Gaussian inputs. Temporary random labels let those operations implement the finite classical function of §7. The hard steps are to implement its aggregations using cost and mixer gates, and

then to replace every use of the labels by ordinary gates while preserving the energy. Only the resulting ordinary word is transferred to SK.

The choices are nested: fix $\varepsilon$, choose $T < 1$, choose a finite mesh, fix the bounded classical formula and helper probabilities, then choose the finite gate approximations and remove the seeds. The constants in a replacement may depend on the surrounding word and on those positive helper probabilities. §8.6 collects the error allocation before the final $n \to \infty$ limit. The principal circuit inputs can be read as four local lemmas. Their proofs remain in the cited manuscript; §§8.2–8.5 reconstruct the mechanisms and show where each hypothesis enters.

**Smooth replacement.** Fix the height, a finite expression and channel list, and the required derivatives. Uniform recursive weighted bounds, convergence of operators and adjoints, and strongly commuting signals with the stated joint exponential bounds permit substitution inside a fixed surrounding sequence in the exclusion-gate class defined in §8.2. Constants depend on that fixed family. This is the input used in §8.3 to replace classical aggregates by finite gates; see the foundations, smooth topology and gate calculus.

**Helper contraction and synthesis.** Fix a bounded, finite, type-excluded odd formula, zero on helpers, and positive helper masses. The contraction converges smoothly at every cut and for the specified responses. For each $\eta > 0$, finite seeded synthesis gives

$$\mathcal{E}(v) \geq \mathcal{E}(m) - 2\eta - 2\sqrt{p_H}.$$

Section 8.4 receives this one fixed seeded word. The source inputs are the synthesis statements and helper theorem.

**Seed removal.** Fix that finite seeded word and an energy tolerance. Reuse the same Gaussian at every occurrence on a vertex. Packet control and the one-particle comparison then produce one finite ordinary word with the required energy error. The seed-removal theorem supplies an energy conclusion, not merely convergence of seed marginals. It makes the ordinary-word transfer in §8.5 applicable.

**Fixed-word transfer.** Fix one ordinary word and all its angles before either size or degree varies. Its positive tree scaled-edge limit equals its SK expected-energy limit, with the convention conversion in §8.5. This fixed-word transfer closes the thermodynamic-first theorem in §8.6.

**One cost–mixer pair.** A single cost–mixer pair fixes the energy normalization before we introduce the rooted representation. Use the positive cost $C_{n,J}$ of §8 and $U = e^{-i\beta B_n} e^{-i\theta C_{n,J}}$; the symbol $\theta$ here is a cost angle, not the Parisi coefficient. Put $c = \cos(2\beta)$ and $s = \sin(2\beta)$. The Pauli identities are

$$e^{i\beta X} Z e^{-i\beta X} = cZ + sY, \qquad e^{i\alpha Z_i Z_j} Y_j e^{-i\alpha Z_i Z_j} = Y_j \cos(2\alpha) + Z_i X_j \sin(2\alpha).$$

The mixed term $Z_i Y_j$ therefore has cost-state expectation

$$\sin(2\theta J_{ij}/\sqrt{n}) \prod_{k \neq i,j} \cos(2\theta J_{jk}/\sqrt{n}).$$

The other mixed term exchanges $i$ and $j$. The $Z_i Z_j$ term has zero expectation. The $Y_i Y_j$ term can be nonzero on an individual complete graph, but its expectation after the cost gate is independent of $J_{ij}$: $Y_i Y_j$ commutes with that edge cost. Multiplying by $J_{ij}$ and averaging therefore removes it, even in the presence of triangles. For a standard Gaussian $J$,

$$\mathbb{E}\cos(aJ) = e^{-a^2/2}, \qquad \mathbb{E}[J \sin(aJ)] = a e^{-a^2/2}.$$

Averaging the mixed terms and summing the $n(n-1)/2$ unordered edges gives

$$\frac{1}{n}\mathbb{E}_J\langle C_{n,J}\rangle = \frac{n-1}{n}\,\theta\sin(4\beta)e^{-2\theta^2(n-1)/n}.$$

This agrees with Farhi et al., equation (62). With energy maximization, $\theta = 1/2$, $\beta = \pi/8$ gives the limiting value $1/(2\sqrt{e})$; their minimization example uses $\beta = -\pi/8$ and the negative value.

For the positive cost $D^{-1/2}\sum_{\{i,j\}} Z_iZ_j$ on the $(D+1)$-regular tree, each endpoint has $D$ other neighbours. The same calculation now has no common-neighbour contribution and yields

$$\langle Z_oZ_v\rangle = \sin(4\beta)\sin(2\theta/\sqrt{D})\cos^D(2\theta/\sqrt{D}).$$

Multiplying by $\sqrt{D}/2$ and letting $D\to\infty$ gives $\theta e^{-2\theta^2}\sin(4\beta)$, the same limit as finite SK. This checks the factor $1/2$ and the degree convention in §8.5.

The value just obtained will be recovered once more from the rooted insertion in §8.1.1. That calculation explains which part of the limiting state carries the energy.

### 8.1 Rooted realisation and the observable it preserves

The fixed-depth spin–boson representation has an antecedent in Boulebnane et al., Theorem 2 (informal), equations (10)–(11), p. 3: as the degree grows with depth fixed, the symmetric tree state converges to a spin coupled to finitely many bosonic modes. The manuscript credits that representation, then proves the occupation, insertion and surrounding-circuit estimates needed below. Its additional task is to control the energy observable and substitutions inside a circuit, beyond identifying a limiting state. OpenAI (2026b), §2.

The immediate task is to realise the finite Gaussian spin from §7 in a space where its shifted correlation sum becomes an energy observable. We will identify that sum with a rooted quadratic form, then relate the same form to the scaled edge energy of a fixed circuit; §8.5 supplies the final transfer to SK.

For an ordinary depth-$p$ word, this tree calculation uses only finitely many spins. Conjugate $Z_iZ_j$ backwards through the circuit. A mixer rotates the spins already in its support without enlarging that support. In a cost layer all edge factors commute, so factors disjoint from the current support commute past the other factors and cancel. The remaining factors enlarge support by at most one graph step. After $p$ pairs, the edge expectation therefore depends only on its radius-$p$ backward light cone. On the $(D+1)$-regular tree this contains at most $2\sum_{r=0}^{p} D^r$ vertices; the spins outside factor out of the product-state expectation. Thus fixed depth gives fixed tree height, although the number of children grows with $D$. This locality explains the finite tree calculation; the equality with dense SK is the separate transfer theorem in §8.5. Farhi, Goldstone and Gutmann (2014), §II, equations (14)–(26).

On a rooted tree, factor a circuit as $U = (\bigotimes_j U_j)u$, where each $U_j$ contains operations wholly in child subtree $j$. Conjugating a root observable $P$ reduces to $u^*Pu$; its insertion vector is $u^*Pu\,\mathrm{vac}$, centered when its vacuum mean is zero. This factorization is why child operations enter the recursion below. The first change of setting preserves a specified quadratic form. At height zero the pointed space is $\mathsf{k} = \mathbb{C}^2\otimes L^2(\text{seed})$, with distinguished vector $\omega = |+\rangle\otimes 1$. Recursively,

$$\mathcal{H}_0 = \mathsf{k}, \qquad \mathcal{H}_{h+1} = \mathsf{k}\otimes\Gamma_s(\mathcal{H}_h^\circ),$$

where $\mathcal{H}_h^\circ$ is the orthogonal complement of the vacuum and $\Gamma_s$ is symmetric Fock space. A particle here is one child subtree whose state differs from its vacuum. It is not one physical spin. The occupation embedding identifies a symmetric state with $m$ nonvacuum children among $D$ available children with its $m$-particle vector, using the factor $\binom{D}{m}^{-1/2}$. Adding empty descendants gives compatible embeddings between heights. OpenAI (2026b), §2, pointed spaces and occupation embeddings.

For a child direction $f$, write $q(f) = a^*(f) + a(f)$ and $W(f) = e^{iq(f)}$. Explicitly, $\Gamma_s(H) = \bigoplus_{m\geq 0} H^{\otimes_s m}$, with vacuum $\Omega$ in sector zero. On sector $m$, $a^*(f)\psi = \sqrt{m+1}\,\mathrm{Sym}(f\otimes\psi)$, and $a(f)$ is its adjoint. The number operator is multiplication by $m$; more generally $d\Gamma(A)$ sums $A$ over those $m$ child factors. The vacuum characteristic function $\langle\Omega, e^{itq(f)}\Omega\rangle = e^{-t^2\|f\|^2/2}$ fixes the Gaussian normalization. For any finite list of fixed centred real child vectors, their parent positions $\mathsf{G}(f) = q(f)$ are jointly centred Gaussian; for $f, g$,

$$\mathbb{E}[\mathsf{G}(f)\mathsf{G}(g)] = \langle f, g\rangle, \qquad \|[\mathsf{G}(f) - \mathsf{G}(g)]\mathrm{vac}\| = \|f - g\|.$$

The directions are fixed elements of the child space, even when their values depend on the child's Gaussian coordinates. The receiving root's own seed is a separate, independent tensor factor. The root shift

$$\mathsf{L}f = \omega \otimes a^*(f)\Omega$$

is an isometry on centred vectors, and is extended by zero on the vacuum. Its adjoint extracts the root-vacuum, one-child component, including a constant receiving-root seed. For a classical function of orthonormal Gaussian positions, this is its first Gaussian chaos. These covariance, independence and projection properties have distinct roles in the next calculation. OpenAI (2026b), §2, root shift and classical first chaos OpenAI (2026b), §2, classical Gaussian positions.

Start with the standard root seed $g_0 = S$ and recursively set $Z_{j+1} = \mathsf{G}(g_j)$. Suppose $g_0, \dots, g_{j-1}$ are centred orthonormal vectors. The joint Gaussian law of their lifts and the covariance identity give independent standard normals $Z_1, \dots, Z_j$. In the scalar construction of §7, the multiplier of $Z_j$ in $g_j$ depends only on its predecessors. Conditional centring therefore makes $g_j$ orthogonal to $g_1, \dots, g_{j-1}$, and the chosen normaliser $c_j$ gives unit norm. Independence of the receiving-root seed gives orthogonality to $g_0$. This closes the induction. Compatible height embeddings ensure that the earlier variables are the same variables when the formula is embedded at greater height.

Oddness of $u$ and evenness of $a$ make $Y_j$ and $g_j$ odd under simultaneous seed reversal. The extra centred rounding variable $2F_{\mathrm{G}}(Z_{N+1}) - 1$ is odd as well, so $F_N$ is centred. It is independent of the receiving-root seed, hence $\langle F_N, g_0\rangle = 0 = A_0$. With $A_j, B_{j+1}$ as in §7, first-chaos projection gives

$$\mathsf{L}^* F_N = \sum_{j=0}^{N} B_{j+1} g_j, \qquad \mathcal{E}(F_N) := \mathrm{Re}\langle F_N, \mathsf{L}F_N\rangle = \sum_{j=0}^{N} A_j B_{j+1}.$$

Thus the construction identifies both the scalar law and the required rooted energy. The Gaussian covariance rule does not declare distinct calculations on shared descendant subtrees independent. OpenAI (2026b), §3, "A finite Gaussian formula and its rooted energy".

The connection to an edge observable is another statement. Let $f_D$ be centred, invariant under rooted permutations of complete child subtrees, and supported at a fixed finite radius. Suppose its occupation embedding converges strongly to $v$, with a uniformly bounded expected top-generation occupation $\langle f_D, \mathsf{N} f_D\rangle$. Put $M = D + 1$ and $R_M = \sqrt{1 - \mathsf{N}/M}$. To see the decomposition, delete the distinguished edge and project onto the vacuum on either resulting half-tree. There are four

pairings. The vacuum–vacuum pairing is zero by centring. If only one side is excited, the other side forces the root and all remaining branches into their vacuum; root symmetry gives the term $M^{-1/2}\langle \mathsf{L}^* f_D, R_M f_D\rangle$ or its conjugate. In the pairing where both sides are excited, each projected vector has squared norm $M^{-1}\langle f_D, \mathsf{N} f_D\rangle$ by symmetry among the $M$ branches. Cauchy–Schwarz bounds their pairing by that same quantity. Finally, deleting one vacuum child multiplies the $m$-particle sector by $\sqrt{\binom{M-1}{m}/\binom{M}{m}} = \sqrt{1-m/M}$, which identifies $R_M$. Thus the exact edge decomposition is

$$\langle f_{D,i}, f_{D,j}\rangle = \frac{2}{\sqrt{M}}\,\mathrm{Re}\langle \mathsf{L}^* f_D, R_M f_D\rangle + \varepsilon_D, \qquad |\varepsilon_D| \le \frac{\langle f_D, \mathsf{N} f_D\rangle}{M}.$$

Vacuum deletion also gives $\|R_M f_D - f_D\|^2 \le M^{-1}\langle f_D, \mathsf{N} f_D\rangle$. Consequently,

$$\lim_{D\to\infty} \frac{\sqrt{D}}{2}\langle f_{D,i}, f_{D,j}\rangle = \mathcal{E}(v).$$

Strong convergence alone does not control the scaled remainder. The occupation estimate is supplied later for the relevant finite circuits. The occupation assumption cannot be dropped merely because vectors converge in norm. For orthogonal unit vectors $e_m$ in sector $m$, $\psi_k = \sqrt{1-1/k}\,\Omega + k^{-1/2} e_{k^2}$ tends to $\Omega$ in norm, whereas $\langle \psi_k, \mathsf{N}\psi_k\rangle = k$.

The elementary continuity bound

$$|\mathcal{E}(v) - \mathcal{E}(w)| \le (\|v\| + \|w\|)\|v - w\|$$

will control every subsequent approximation after the observable has been identified. OpenAI (2026b), §2, "Energy as a root shift," including its exact finite-degree decomposition.

The three identifications use different information:

First, passing from the finite Gaussian formula to the rooted classical sector preserves the joint Gaussian law and the same $F_N$. This uses covariance, conditional centring and independence of the receiving-root seed.

Second, the root-vacuum and first-chaos projection identify the energy itself: $\mathcal{E}(F_N) = S_{T,N}$.

Third, for a finite-degree circuit insertion tending to $v$, root invariance, fixed radius, strong convergence and uniform occupation control imply that the scaled edge expectation tends to $\mathcal{E}(v)$.

The first two calculations were given above. The third uses the source's edge decomposition; the requisite circuit estimates enter in §8.2.

**8.1.1 The same energy in the rooted representation** We return to the value $\theta e^{-2\theta^2}\sin(4\beta)$ obtained in the finite-SK and tree calculation. Retain its cost angle $\theta$, mixer angle $\beta$, $c = \cos(2\beta)$ and $s = \sin(2\beta)$. Here $\Omega$ is the Fock vacuum, $\mathrm{vac} = |+\rangle \otimes \Omega$, and $q(e) = a^*(e) + a(e)$. The root shift $\mathsf{L}$ defined in §8.1 creates one child excitation and puts the receiving root in its vacuum. Let $e = Z\mathrm{vac} = |-\rangle \otimes \Omega$ and $Q = q(e)$ in the rooted description. Its vacuum law is standard normal. The depth-one root operation is $u = e^{-i\beta X} e^{-i\theta Z Q}$. Using $Y|+\rangle = -i|-\rangle$, its insertion is

$$v = u^* Z u\,\mathrm{vac} = |-\rangle \otimes [c - is\cos(2\theta Q)]\Omega + |+\rangle \otimes s\sin(2\theta Q)\Omega.$$

The root shift retains only the plus-root, one-particle component. The first-chaos coefficient of the sine is $\mathbb{E}[G\sin(2\theta G)] = 2\theta e^{-2\theta^2}$, so

$$\mathsf{L}^* v = 2\theta s e^{-2\theta^2} e, \qquad \mathrm{Re}\langle v, \mathsf{L} v\rangle = 2\theta s c e^{-2\theta^2} = \theta e^{-2\theta^2}\sin(4\beta).$$

Here $\mathrm{Re}\langle e, v\rangle = c$ supplies the final factor. Thus the observable, not just the Gaussian law, agrees in all three representations. The example uses the ordinary root updates and root shift. It calibrates one depth; the higher-depth occupation and replacement hypotheses remain necessary.

### 8.2 Finite regularisation and the calculus needed for compilation

Fix $T, N$ first. The independent continuous rounding coordinate makes the event at the sign discontinuity null. Bounded continuous odd approximations to that sign therefore converge in $L^2$ by dominated convergence. If the bounded output is $m^*$, shrink it to $(1-\kappa)m^*$, with fixed $0 < \kappa < 1$, before taking $\theta_\kappa = \frac{1}{2}\arcsin((1-\kappa)m^*)$. This margin keeps the derivative of inverse sine bounded on its used range; the output loss is at most $\kappa$ in $L^2$. Then truncate the transmitted functions and approximate the resulting bounded maps by trigonometric polynomials. The source makes these choices from output toward leaves: each outer Lipschitz function is fixed before its inputs are approximated. The Gaussian isometry above converts an error in a transmitted child vector into the same $L^2$ error in its receiving position. Energy continuity then bounds the accumulated loss. The bounded output has the form $m = \sin(2\theta)$; it is the transmitted columns that are bounded, while the Gaussian fields themselves remain unbounded. OpenAI (2026b), §3, bounded trigonometric approximation.

**The formulas to be synthesized.** A mask is an event in the local seed space invariant under seed reversal. Fix a finite partition into masks, and write $p_R$ for the probability of mask $R$; probabilities used as denominators are positive. For disjoint masks $R, S$, aggregation on the finite regular tree means

$$(A_{RS}f)_i = \mathbf{1}_R(i)D^{-1/2}\sum_{j\sim i}\mathbf{1}_S(j)f_j.$$

A formula has a finite expression tree, with bounded measurable functions of the local seed as leaves. Internal operations are such aggregations or pointwise functions of previously formed quantities. Pointwise functions are smooth in their aggregate arguments, with polynomial bounds on all derivatives, uniformly in the seed. Seed dependence itself need not be smooth. Transmitted quantities are odd under simultaneous seed reversal; sent nonlinear functions and output angles are bounded, while aggregate Gaussian fields may be unbounded.

An aggregation from $S$ to $R$ excludes $R$ if its transmitted formula uses no vertex of type $R$ below that call. All ancestor exclusions are retained, so a mask cannot recur on one active expression branch. It may recur on a different branch, and distinct branches may share descendants; exclusion asserts no independence between them. Gate targets are likewise excluded from the cross-edge dependencies of their parameters. This makes the physical all-neighbour calculation agree with the intended descendant calculation. For synthesis, reserve disjoint helper masks of positive mass, one per positive aggregation depth, absent from the output dependencies; the bounded odd output angle vanishes on their union. OpenAI (2026b), Definition 3.3; Theorem 4.1. The types can be encoded together with a Gaussian seed in one raw Gaussian variable: its sign supplies the odd symmetry, while disjoint binary digits of a uniform transform of its absolute value supply independent coordinates. The resulting finite type label is even and the Gaussian seed is odd under reversal. The associated pullback preserves vacuum, parity and $\mathsf{L}$. Thus adding types preserves the energy identification, rather than assuming an additional circuit resource. OpenAI (2026b), §3, colours, exclusions and helpers.

For the already fixed formula, the exclusion estimate is

$$\|m^{\mathrm{col}} - m\|^2 \le C(\alpha + p_H),$$

where $\alpha$ is the largest nonhelper type mass and $p_H$ is the total helper mass. Its constant is independent of the number of types. The proof uses orthogonality of the disjoint masks before applying the Lipschitz estimate in the original, fixed list of formula coordinates. Treating the refined types as an arbitrary growing list of independent arguments would not provide this bound. The resulting classical near-optimality proposition supplies a finite bounded odd formula, zero on helpers, with energy as close to $P_*$ as required. OpenAI (2026b), §3, "Type exclusion at small cost" and "Classical near-optimality in synthesis form".

**Which convergence permits the later substitutions?** The gate calculus uses recursive weights $\mathsf{W}_{h+1} = \mathsf{W}_0 + d\Gamma(\Lambda_h)$, where $\Lambda_h$ is the previous weight restricted to the centred child space. The local weight includes the unbounded raw seed when needed. Its smooth domain is the intersection of all weighted domains. Smooth bounds mean polynomial bounds, with a finite loss of weight powers, for the maps, their adjoints and every specified parameter derivative. The finite height, fixed expression and fixed number of channels matter in these estimates. Strong convergence together with uniform stronger weighted bounds gives convergence in each weaker weighted norm by a spectral cutoff. This is the topology in which one replaces a gate inside a fixed surrounding circuit. OpenAI (2026b), §2, weights and the topology of convergence.

More precisely, the imported smooth-convergence contract is: for each output power $m$ and parameter multi-index $\alpha$, there are an input power $r$, exponent $k$ and constant $C$ such that

$$\|\mathsf{W}_D^m \partial_x^\alpha F_D(x)\xi\| \le C(1+|x|)^k \|\mathsf{W}_D^r \xi\|, \qquad \xi \in \bigcap_{q\ge 0} \mathrm{Dom}(\mathsf{W}_D^q).$$

The bound is uniform in degree $D$ and parameter $x$. The same bound and strong convergence on smooth inputs are required for the adjoints and all specified parameter derivatives, with compatible limiting adjoints. Constants may depend on the fixed expression, height, channel list and derivative order. Bounded adjoint derivatives alone do not supply their convergence. At bounded local seeds $\mathsf{W}_0 = 1$; for polynomial Gaussian growth the source uses the vacuum-preserving weight $P_0 + Q_0(1+|s|)Q_0$, where $P_0$ projects to the local vacuum and $Q_0 = 1 - P_0$. These recursive weights control every generation. They differ from the top-generation number weight in §8.4's packet argument. No uniformity in growing word length, vanishing helper mass or the later seed-removal parameter is implicit in this contract. OpenAI (2026b), §2, smooth-family bound.

One of the essential inputs retains a term beyond the Gaussian field. Let $K_D(x)$ be a smooth unitary family with $K_D(0) = 1$, uniformly controlled derivatives and adjoint derivatives that converge smoothly in the stated weighted topology to those of a limiting family $K$, and centred first vacuum derivatives. Write $K_\alpha = \partial_\alpha K(0) = iT_\alpha$ and $K_{\alpha\lambda} = \partial_\alpha \partial_\lambda K(0)$. For a fixed finite parameter list $a$, the source proves

$$K_D(a/\sqrt{D})^{\otimes D} \longrightarrow \exp\left\{ i\sum_\alpha a_\alpha q(T_\alpha \mathrm{vac}) + \frac{1}{2}\sum_{\alpha,\lambda} a_\alpha a_\lambda \left\langle K_{\alpha\lambda} + \tfrac{1}{2}\{T_\alpha, T_\lambda\}\right\rangle_0 \right\}.$$

The second term is imaginary by unitarity. It survives because a second-order change of size $D^{-1}$ occurs in each of $D$ children. Its scalar coefficients become a possibly nonconstant controlled operator when commuting signals replace $a$. For example, $K(x) = e^{ix^2}$ has zero first derivative at zero but $K(a/\sqrt{D})^D = e^{ia^2}$ exactly. Keeping only the first derivative would miss even this elementary contribution. The example explains why the term is necessary; the source's weighted product lemma supplies the general convergence and domain bounds. OpenAI (2026b), §2, "Products of small unitaries".

The smooth-substitution lemma makes the required joint control explicit. For a fixed finite list of strongly commuting self-adjoint signals $S_D = (S_{D,1}, \dots, S_{D,b})$, their polynomials must converge smoothly, and their joint exponentials must converge strongly on smooth inputs and satisfy, for every output weight $m$,

$$\|\mathsf{W}_D^m e^{it\cdot S_D}\xi\| \le C_m(1+|t|)^{k_m}\|\mathsf{W}_D^{r_m}\xi\|,$$

uniformly in $D$ and $t \in \mathbb{R}^b$. The coefficient family $F_D(x)$ and its adjoint obey the smooth bounds above and converge with all their jets on compact parameter sets. Each coefficient must commute with the signals' spectral projections; for unbounded coefficients this means commutation in the closed-operator sense. Seed-dependent substitution also requires strong commutation with the receiving seed multiplication algebra. These hypotheses permit the joint spectral and Fourier calculation of $F_D(S_D)$, including its derivatives and adjoints. A formal commutator identity on a single domain would not supply them. No nonsingular covariance matrix or inverse Gram matrix is required. OpenAI (2026b), Lemma 2.10, "Smooth commuting substitution".

For the formula fixed here, its finite expression tree supplies the finite channel list. Exclusion supplies the commuting physical control algebra; normalized-sum limits and the preceding product estimates supply the signal limits and weighted bounds. Conjugation by the already fixed preceding circuit preserves strong commutation, including when signals share descendants. The gate-calculus theorem applies these facts inductively to the formula and its adjoints, at each cut and inside the fixed surrounding sequence. It is this theorem, with these constructed inputs, that licenses the later replacements. Its conditional root-seed bounds also give the masked-column estimates used in §8.3. OpenAI (2026b), Theorem 2.12.

The exact cut recursion records the effect of such shared lower calculations. Let $y$ be the incoming boundary data, $V(y)$ the root matrix, and $m(y)$ its outgoing signals in the current frame. Factoring out the operations wholly inside each child gives

$$E^{[d]}(y) = V(y)\prod_{j\ \mathrm{child}} K_j^{[d-1]}(D^{-1/2}m(y)), \qquad K^{[d]}(y) = u^*E^{[d]}(0)^*E^{[d]}(y)u.$$

Every ratio $K^{[d]}$ equals one at zero. Nevertheless $m(0)$ generally does not vanish, because removing the parent input leaves the root's own signals. The actual zero-boundary root update is $u_{\mathrm{new}} = E^{[d]}(0)u$. It is not recovered by setting the entire ratio to one and discarding the baseline operation. Applying the unitary product lemma to this recursion, and then commuting substitution, is the source's smooth gate calculus. Its replacement statement covers a fixed finite surrounding sequence and its required parameter derivatives; it does not assert uniformity for arbitrary sequences whose length or channel count grows. OpenAI (2026b), §2, smooth gate calculus and its cut recursion.

### 8.3 Seeded synthesis: helper contraction and nested responses

Temporarily add longitudinal seed gates $\exp(-it\sum_i s_i Z_i)$, using an independent Gaussian $s_i$ at each vertex and reusing that same variable whenever a later seed gate acts there. The synthesis theorem turns the finite classical formula into a finite word of these gates, costs and mixers. Its ideal local $Y$ rotations have $Z$ insertion

$$v_{\mathrm{cl}} = |+\rangle m + |-\rangle n, \qquad m = \sin(2\theta).$$

Here $n$ is the remaining seed-dependent coefficient of the $|-\rangle$ component, not the number of spins. Both coefficients may depend on descendant seeds. The second term does not contribute to $\mathcal{E}$: $\mathsf{L}$ puts the root spin in $|+\rangle$, and shifting the $|-\rangle$ term puts a spin excitation into a child, orthogonal to

the classical sector. Thus $\mathcal{E}(v_{\mathrm{cl}}) = \mathcal{E}(m)$. OpenAI (2026b), §2, classical rotations OpenAI (2026b), §4, "Classical synthesis".

There are two different approximation operations. First, seed conjugations of $X$ generate $\cos(2ts)X$ and $\sin(2ts)Y$. Trigonometric approximation and matrix commutators give the required bounded measurable masked spin controls. Sign averaging of masked $X$ flips isolates an interaction between chosen types; the cost terms commute, so this isolation is exact. This sign averaging is the refocusing method of Jones and Knill (1999), §II, p. 2: orthogonal sign rows cancel unwanted Ising couplings, while identical rows retain a selected coupling. The source applies it to the finite partition of seed masks. Finite commutator products then approximate the desired pointwise matrices with smooth bounds. They can be substituted after the surrounding circuit has been fixed. The local masks are measurable functions of the seed; smoothness is required in the continuous aggregate inputs, not in the mask labels. OpenAI (2026b), §4, elementary masked controls and matrix products.

The second operation implements an aggregation. Let $I$ be the receiver region, $J$ the sender region, and $K$ an unused helper region of fixed positive probability $p_K$. They are disjoint. The lower seed-only formula $f$ excludes these regions along its cross-edge paths. With $A_{ij} = D^{-1/2}1_{i\sim j}$, the desired propagation has generator

$$H = \sum_{i\in I, j\in J} P_i A_{ij} f_j.$$

Partition $J$ into parts $J_l$ of probabilities $p_l$. Two levels of spin commutators generate the commuting Hamiltonian

$$\widetilde{H}_{\mathcal{P}} = \frac{1}{p_K} \sum_l \sum_{i\in I,\, j,j'\in J_l,\, k\in K} P_i A_{ij} X_j A_{jk} X_k A_{kj'} X_{j'} f_{j'}.$$

For a fixed receiver–sender edge $i \sim j$, the terms with $j = j'$ contain $X_j^2 = 1$; $f_j$ is seed-only and commutes with these spin factors. On helper-pinned states $X_k = 1$, so their sum is exactly

$$\frac{1}{p_K} P_i A_{ij} f_j \sum_{k\in K} A_{jk} A_{kj} = P_i A_{ij} f_j \frac{N_K(j)}{D p_K},$$

where $N_K(j)$ counts helper neighbours of $j$. Conditional on the receiver edge, its endpoint $i \in I$ is not a helper and the other $D$ neighbours have mean helper count $Dp_K$. Its binomial variance is $Dp_K(1-p_K)$, so the ratio tends to one in $L^2$ at fixed $p_K > 0$. Terms with $j \neq j'$ are simple paths and must disappear under partition refinement. The factor $1/p_K$ is harmless only after a positive helper mass has been fixed. OpenAI (2026b), §4, helper generators and contraction.

Helper pinning is a statement about every helper spin in the light cone. Its finite-degree projection is

$$\Pi_{K,V} = \prod_{v\in V} \left[1 - 1_K(s_v) \frac{1 - X_v}{2}\right].$$

It retains the seed variables and helper-labelled descendant spaces while fixing the helper spins in $|+\rangle$. The ideal propagation, the exact Hamiltonian $\widetilde{H}_{\mathcal{P}}$, and the surrounding gates that avoid $K$ preserve this compressed space. Multiplying an insertion by $1_{K^c}(s_o)$ only masks the root label and does not impose this descendant condition. Individual finite commutator factors may leave the compressed space. Accordingly, the proof first establishes the exact contraction there, then chooses finite commutator products on the full space to approximate it. OpenAI (2026b), §4, "Simultaneous helper contraction".

The estimate controlling a growing partition is

$$\rho^2 = \sum_l p_l^2 \le p_J \max_l p_l \longrightarrow 0.$$

At a helper root, each part supplies two commuting fields with columns $d_l, e_l$ on disjoint sender masks, of weighted norm at most $C\sqrt{p_l}$. The remaining correction is a sum of centred quadratics

$$\sum_l [q(d_l)q(e_l) - \langle d_l, e_l\rangle].$$

Normal ordering and disjoint-mask orthogonality bound its quadratic kernels by $C\rho$ in Hilbert–Schmidt norm. The source separately bounds the exponentials in the weighted topology, uniformly in the number of parts. Smallness of the generator on smooth vectors alone would not justify inserting its exponential into the surrounding circuit. OpenAI (2026b), §4, "Quadratics with many orthogonal columns".

The helper estimate alone does not control all cuts. Put $\epsilon_{\mathcal{P}} = \max_l \sqrt{p_l}$. The receiver and helper corrections have weighted size $O(\rho)$; a sender correction is $O(\sqrt{p_l})$ conditional on its label. The remaining lower-formula responses have size $O(\epsilon_{\mathcal{P}})$. Both quantities tend to zero with $f$, the number of lower channels and $p_K > 0$ fixed.

There remains a response inside the lower formula $f$. A cut can pass through a descendant used to compute $f$, rather than through one of the displayed three-edge paths. The source first writes the exact difference of the affected Hamiltonian terms before applying the preceding subtree circuit. That circuit is fixed with respect to the new cut input. Exclusions ensure that the downward coefficients are independent of that active boundary argument. The resulting additive cut-response rule retains both first derivatives and mixed second derivatives:

$$\sum_\alpha a_\alpha q(H_\alpha \mathrm{vac}) + \frac{1}{2}\sum_{\alpha,\lambda} a_\alpha a_\lambda \langle H_{\alpha\lambda}\rangle_0.$$

Here $H_\alpha$ is centred by parity; no limit of the extensive zero-input Hamiltonian is required. The self-adjoint generator is obtained from the limiting time group of cut ratios. OpenAI (2026b), §2, "Additive cut responses".

After substituting the signals of a lower formula, a boundary difference contains, in particular,

$$\begin{aligned}&\sum_\alpha [m_\alpha(y) - m_\alpha(0)]q(r_\alpha)\\ &\quad + \frac{1}{2}\sum_{\alpha,\lambda}[m_\alpha(y)m_\lambda(y) - m_\alpha(0)m_\lambda(0)]c_{\alpha\lambda}.\end{aligned}$$

The mixed coefficients $c_{\alpha\lambda}$ must remain even when two channels share descendants. A bounded scalar example makes the issue visible: for a shared standard normal $Q$, take $H(y_1, y_2) = \cos(Q + y_1 + y_2) - \cos Q$. At zero both first derivatives are $-\sin Q$ and are centred, but $\mathbb{E}H_{12} = -e^{-1/2}$. The two cross terms in the half-sum therefore contribute $-e^{-1/2}a_1a_2$. Centred first responses do not remove a mixed second response. This example checks the algebraic reason for retaining the term; the cut-response theorem supplies the operator and domain justification. At the actual root there is no parent input to subtract: the corresponding unsubtracted $m(0)$ terms enter the root update. The source propagates small terminal responses through the fixed depth and fixed list of lower channels. It refines the calling sender mask while keeping those lower formulas fixed. This

is what controls all cuts in a simultaneous replacement of several calls. A root-seed scalar phase may remain on helper labels; it cancels in insertions and child conjugations. OpenAI (2026b), §4, lower-formula responses and completion of the contraction proof.

Compilation now proceeds from outer aggregations to inner ones. Reserve a separate unused helper type for each aggregation rank. At a given rank, establish the simultaneous exact contraction, fix the sender partitions, and only then choose finite commutator approximations. Subsequent inner calls use their own helpers, which remain unused. Each inner accuracy is chosen after its surrounding finite sequence is fixed. This order avoids assuming uniformity over all helper masses, ranks or circuit lengths. OpenAI (2026b), §4, order of compilation.

If the resulting insertion $v$ obeys $\|1_{H^c}(v - v_{\mathrm{cl}})\| < \eta$, seed commutation and unitarity give $\|1_H v\| = \sqrt{p_H}$. First omit the helper part of $v$, at energy cost at most $2\sqrt{p_H}$. Compare the two insertions projected off helpers; their norm difference is below $\eta$, so their energies differ by at most $2\eta$. The projected classical insertion still has energy $\mathcal{E}(m)$ because $m = 0$ on helpers. Thus

$$|\mathcal{E}(v) - \mathcal{E}(m)| \le 2\eta + 2\sqrt{p_H}.$$

All projected insertion vectors have norm at most one. This completes the interface from a bounded classical formula to a finite seeded word. It uses the source's smooth gate and helper-contraction theorems; it is not a consequence of the scalar theorem or of a finite-degree central limit theorem alone. OpenAI (2026b), §4, synthesis statement and its proof.

### 8.4 Removing the seed controls without changing their joint law

Seed removal must simulate one Gaussian variable per vertex, reused at all seed gates. Replacing each occurrence with a fresh independent Gaussian would change the circuit. The source proves a theorem for any already fixed finite seeded word and any energy tolerance, producing one finite ordinary word. Its proof works in the fixed-word tree limit; it does not take a degree limit simultaneously with its small parameter. OpenAI (2026b), §5, "Removal of seed controls".

**A small component supplies a large response.** Fix $b > 0$ and $0 < r < 1$, and put

$$R_* = e^{-i\pi X/4}, \qquad \delta_r = \arcsin r, \qquad \gamma_r = b/r, \qquad e = Z\mathrm{vac}, \qquad \ell_r = Y e^{-2i\gamma_r Z q(e)}\mathrm{vac}.$$

Apply, in chronological order, cost $\gamma_r$, mixer $\delta_r$, cost $-\gamma_r$ and mixer $\pi/4$. Just before the inverse cost the insertion is $v_{2,r} = \cos(2\delta_r)e + \sin(2\delta_r)\ell_r$. Define

$$d_{0,r} = \gamma_r(v_{2,r} - e) = -2br\, e + 2b\sqrt{1 - r^2}\, \ell_r, \qquad W_Z(f) = e^{iZq(f)}.$$

Expanding the first mixer and using the Weyl identity gives the exact split

$$u_r = a_{0,r}U_{O,r} + a_{1,r}U_{S,r}, \qquad U_{O,r} = R_* W_Z(d_{0,r}), \quad U_{S,r} = R_* X W_Z(-2\gamma_r e - d_{0,r}),$$

where

$$a_{0,r} = \cos\delta_r\, e^{i\vartheta_r}, \quad a_{1,r} = -ire^{-i\vartheta_r}, \quad \vartheta_r = \gamma_r^2\, \mathrm{Im}\langle v_{2,r}, e\rangle.$$

Both component operators are unitary. The special component $\psi_{S,r} = a_{1,r}U_{S,r}\mathrm{vac}$ therefore has norm exactly $r$. Its normalized contribution to the insertion is

$$w_r = u_r^* Z \psi_{S,r}/r, \qquad \|w_r\| = 1. \tag{8.1}$$

This is the direction amplified in the echo below. The two components need not be orthogonal. The ordinary component has a bounded displacement $d_{0,r} = 2b\ell_r + o(1)$, whereas the special component contains a displacement of order $1/r$. Subsequent unitaries preserve each component's norm. Only the undisplaced packet tails have bounded occupation moments; norm remainders are carried by unitarity without assigning such bounds.

The source's high-direction lemma gives that $\ell_r, w_r$ are centred unit vectors, become mutually orthogonal, and escape every fixed occupation sector. Splitting a uniformly bounded family of occupation moments at a fixed sector cutoff shows that its pairings with these directions vanish: first let $r \downarrow 0$, then send the cutoff to infinity. The special component must nevertheless be retained because the later echo magnifies its insertion contribution by $1/r$. OpenAI (2026b), §5.2, exact initialization and high directions.

A packet has the form $W(d_r/r)\eta_r$, where the scaled centre $d_r$ is bounded and the tail $\eta_r$ has uniformly bounded occupation moments. Packets with centres separated by a fixed positive amount have overlap smaller than every fixed power of $r$. A short probe along a Pauli axis $P$ sees a finite rotation determined by $-2\,\mathrm{Im}\langle v_P, d_r\rangle$, as well as a small Weyl displacement. The direction $v_P = u^*Pu\,\mathrm{vac}$ is the actual current insertion. Toggling a cost rotates both its root Pauli axis and this direction; they cannot be selected independently. OpenAI (2026b), §5, packet estimates.

For the finite list of nonzero special centres $h_j \in \mathbb{R}^2$, the source constructs the same rotation $e^{-i\theta X}$ at every centre, with *zero clocks*: the signed pulse durations on each of two perpendicular axes in the $ZY$ plane sum exactly to zero. Choose the axes so that $c_{j,a} = h_j \cdot P_a \neq 0$, and the square pairs $(c_{j,1}^2, c_{j,2}^2)$ are distinct except for equal or opposite centres. Brackets have zero clock components and generate $iX$ coefficients $c_{j,1}c_{j,2}F(c_{j,1}^2, c_{j,2}^2)$. Polynomial interpolation makes all these coefficients one. Equal or opposite centres have the same product, so they cause no obstruction to this common rotation.

This does not give arbitrary independent rotations: opposite centres obey $U_{-h} = XU_hX$. Exact finite reachability of the required common rotation follows by taking finitely many reachable conjugates spanning the generated Lie algebra; their product map contains an identity neighbourhood, and subdivision reaches the whole one-parameter path. Closure alone would not give exact cancellation. The Lie-bracket and polynomial method follows Li and Khaneja (2005), §II, Examples 1–2, pp. 2–3; the finite-family, equal/opposite-centre and exact-clock assertions are proved in the QAOA source itself. The distinction is exact: zero net probe clocks and the common target rotation are assertions about the finite ensemble of limiting centres. A finite pulse list realizing those assertions is fixed first. Its action on the actual finite-$r$ packets is approximate. The separate accuracy lemma then follows the actual insertion drift through every probe prefix. Its ordinary-component error is $o(r)$; the error on the normalised special component is $o(1)$ and hence also contributes $o(r)$ to the state. These bounds are uniform over the bounded-tail families needed for insertions, and their constants refer to the chosen finite pulse list. OpenAI (2026b), §5, "Selective rotations with zero clocks" and "Accuracy of selective probes".

The shared seed family is essential even for a single spin. Two gates $e^{-itSZ}$ applied to $|+\rangle$, with the same standard Gaussian $S$, give mean $X$ expectation $\mathbb{E}\cos(4tS) = e^{-8t^2}$. Replacing the second seed by an independent copy gives $e^{-4t^2}$. Agreement of marginal seed laws does not preserve a circuit's value.

The error scale needed below follows from the source's quantitative probe estimate. For a fixed finite pulse list, separated packet centres and bounded designated tail moments, write $\varepsilon_r = o(r)$ for

the discarded vacuum remainder. At each probe prefix $j$, with current insertion $v_{P,r}^{(j)}$ and huge-cost direction $v_r$, the proof gives

$$\max_j |\operatorname{Im}\langle v_{P,r}^{(j)}, v_r\rangle| \le C_M(r^2 + \varepsilon_r + r^M).$$

Choose a fixed $M > 1$. The right side is $o(r)$: within each ordinary packet the relevant Hermitian expectation is real, distinct packet centres have overlap $O(r^M)$, and terms containing only special packets have size $O(r^2)$. The norm remainder is carried by unitarity; it receives no unproved occupation bound. The exact zero clocks then cancel the first-order displacement on ordinary packets, leaving $o(r)$ error there. The prescribed action on a normalized special packet has $o(1)$ error, multiplied by its $O(r)$ amplitude. These are the unequal accuracy requirements that the next echo uses. OpenAI (2026b), §5, accuracy of selective probes.

**The estimates must hold on the next input.** The repeated echo uses more than the initial special norm in (8.1). Let $\mathsf{N}$ count particles in the top child Fock space. Call a family $(\eta_r)$ bounded smooth when $\sup_r \|(1+\mathsf{N})^k \eta_r\| < \infty$ for every fixed $k$. At each stage between simulated gates, the source propagates a linear ordinary/special decomposition on every such family, with

$$\begin{aligned} u_r\eta_r &= \Psi_{O,r}(\eta_r) + \Psi_{S,r}(\eta_r), \\ \|\Psi_{S,r}(\eta_r)\| &= r\|\eta_r\|, \\ \Psi_{O,r}(\eta_r) &= \zeta_r A_r \eta_r + o(r). \end{aligned}$$

Here $|\zeta_r| = 1$, and the low unitary $A_r$ and its adjoint preserve all these seminorms uniformly. A separate part of the induction transports the special insertion:

$$u_r^* Z \psi_{S,r}/r = w_r + o(1),$$

with the negative sign between a selective flip and its undoing. The norm remainders remain separate from the designated smooth tails; no occupation estimate is imposed on them.

This forward estimate gives the insertion approximation used at the next gate. For $P = Y, Z$, the vector $\hat{v}_{P,r} = A_r^* P A_r \mathrm{vac}$ is itself a bounded smooth family. Since $A_r \hat{v}_{P,r} = P A_r \mathrm{vac}$, unitarity gives

$$\begin{aligned} \|u_r^* P u_r \mathrm{vac} - \hat{v}_{P,r}\| &= \|P u_r \mathrm{vac} - u_r \hat{v}_{P,r}\| \\ &\le \|P(u_r - \zeta_r A_r)\mathrm{vac}\| \\ &\quad + \|(u_r - \zeta_r A_r)\hat{v}_{P,r}\| = O(r). \end{aligned}$$

The second term explains why an estimate on the vacuum alone would not suffice: its input changes with $r$. The scalar phase cancels in this comparison. This $O(r)$ estimate supplies smooth approximants for actual insertions; it is not divided by $r$. The singular echo instead uses the more accurate difference derived next. OpenAI (2026b), §5, packet invariant and insertion-approximation lemma.

**The echo amplifies a controlled difference.** Consider a forward cost $t/r$, a selective flip of the special component, the reverse cost $-t/r$, and a selective undoing of that flip. The packet invariant transports the special component so that its contribution to the insertion changes sign. For the actual insertion directions before and after the first flip, it gives

$$v_r' - v_r = -2r\, w_r + o(r).$$

The Weyl product of the two costs is formed with these actual directions before making any approximation. Their large centres cancel, leaving a bounded displacement

$$\frac{t}{r}(v'_r - v_r) = -2t\, w_r + o(1).$$

Approximating each direction by a low component with merely $O(r)$ error before this subtraction would leave an uncontrolled order-one error. The remaining scalar Weyl phase is independent of the root spin and cancels in insertions; it need not itself converge. The second flip restores the transported special component so that the same construction can be used again. Mixers receive analogous selective compensations. OpenAI (2026b), §5, packet invariant, insertion approximation and "One echo".

In the limit, $\ell_r$ and $w_r$ become two orthogonal external real Gaussian modes at each root. The initialization uses $q(\ell)$ and every simulated seed gate uses the same $q(w)$: choosing $t = \sigma/2$ implements $e^{-i\sigma Z q(w)}$. Different vertices have independent local copies. This is the required joint law of the seeded word. A finite induction uses the Weyl identity

$$\langle W(f)\Omega, W(g)\Omega\rangle = \exp\bigl(-\tfrac{1}{2}\|g-f\|^2 + i\,\mathrm{Im}\langle f, g\rangle\bigr)$$

to match the relevant finite Gram matrices. Escaping high directions supply precisely the vanishing pairings between the new external modes and endogenous child directions. OpenAI (2026b), §5, external-mode limit.

**The energy pairing after seed removal.** Matching the seed law and its finite Gram matrices is not yet enough. For a centred low insertion $h = \sum_j s_j \otimes W(f_j)\Omega$, the one-particle coefficient of $W(f_j)\Omega$ is $ie^{-\|f_j\|^2/2} f_j$. Hence

$$\langle h, \mathsf{L}h\rangle = -i \sum_j \langle s_j, +\rangle e^{-\|f_j\|^2/2} \langle f_j, h\rangle.$$

In the seeded representation, $\mathsf{L}$ places the root-seed factor in its vacuum and creates $h$ only in the endogenous child space. External components therefore contribute zero to the last pairing. The high-direction estimate yields the same zeros in the actual limit. The actual insertion differs from its low insertion by $O(r)$, so energy continuity completes the comparison. Matching Gram matrices without this one-particle calculation would not identify $\mathsf{L}$ or its energy. OpenAI (2026b), §5, "Finite Gram induction," including the energy calculation.

**Why the selective rotations are available at every required stage.** The limiting projection of each special centre must be nonzero. The source imposes finitely many visibility conditions on the angles of the formal seeded word. At each relevant standard stage they have form

$$-2\,\mathrm{Im}\langle v_Z^{\mathrm{form}}, -2bz\, e + yD^{\mathrm{form}}\rangle \neq 0, \qquad z, y \in \{\pm 1\},$$

where $e = Z\mathrm{vac}$ and $D^{\mathrm{form}}$ accumulates echoes whose reverse costs have already occurred. These functions are real analytic by the finite Weyl formulas. Setting all target angles to zero, while retaining initialization, gives $4bze^{-8b^2} \neq 0$, so none is identically zero. An arbitrarily small angle perturbation avoids their finite union of zero sets and the zero echo-time hyperplanes. The huge-cost stages need a separate transverse-projection estimate; their limiting projection pairs then also have a nonzero coordinate. Finite pulse lists can now be chosen successively from limits already established at earlier stages. No list is chosen from a limit that assumes its own successful implementation. OpenAI (2026b), §5, huge-stage transverse projection, visibility and construction order.

The choices are therefore nested: make $b$ small to control initialization; perturb the fixed target angles to ensure visibility; choose all finite probe lists; finally choose $r$ small. A value of $r$ fixes every large angle and every finite product. The theorem allows those angles to be large or negative. It makes no claim of uniform bounds as the target word length grows or as the tolerance tends to zero. OpenAI (2026b), §5, proof of seed removal.

### 8.5 Transfer of the resulting ordinary word

After all these choices there is one finite ordinary word $W$, fixed independently of degree, size and disorder. On the $(D+1)$-regular tree use $C_D = D^{-1/2}\sum_{\{i,j\}\in E} Z_iZ_j$, counting each edge once. The edge expectation is defined on the finite light cone described in §8.1. The source applies the fixed-word transfer theorem

$$v_W = \lim_{D\to\infty} \frac{\sqrt{D}}{2}\langle +|W_D^* Z_i Z_j W_D|+\rangle.$$

The full preprint of Basso et al., arXiv:2110.14206v3, Theorem 1, equates the limiting tree recursion with the infinite-size SK recursion for every fixed depth and fixed angle vector. Its proof is in §6, especially (6.6)–(6.31), with the symmetry statement proved in Appendix B and the iteration stabilisation in Appendix A.4. The shorter proceedings edition states the theorem but refers to this full preprint for the proof.

Basso et al. use a negative tree cost in (2.3). Let $V = \prod_{i\in A} X_i$ on one bipartition of the finite light cone. Every edge has exactly one endpoint in $A$, hence

$$VZ_iZ_jV = -Z_iZ_j, \qquad VB_DV = B_D, \qquad V|+\rangle = |+\rangle.$$

Consequently, if $C_D^- = -C_D$, then $VW(C_D^-, B_D)V = W(C_D, B_D)$ for the same fixed word and angles, while its measured edge expectation changes sign. This converts the negative cut convention in their (2.6) to the positive energy above. Their SK cost (6.1) already has the positive convention used here. The result applies after seed removal; it does not directly transfer an arbitrary seeded or controlled word. It also supplies no joint limit in which depth or angles change with $D$ or $n$. OpenAI (2026b), §2, "Fixed-word transfer".

For an ordinary cost or mixer the root updates are, respectively,

$$u \longmapsto e^{-itZq(v_Z)}u, \qquad u \longmapsto e^{-i\beta X}u, \qquad v_Z = u^*Zu\,\mathrm{vac}.$$

The source proves smooth convergence for these updates, their adjoints and fixed angle derivatives. The resulting occupation bounds supply the hypothesis of the root-shift energy proposition in §8.1. Thus its energy is the scaled edge limit required for transfer. Finally merge consecutive gates of the same type and insert zero-angle gates where needed to express $W$ in alternating QAOA form. OpenAI (2026b), §2, "Ordinary root updates".

### 8.6 Fixing the word before the thermodynamic limit

**Approximating a specified finite observable.** First fix $T, N$ and the positive finite normalizers in §7. The rooted construction in §8.1 has energy exactly $S_{T,N}$. Let $z$ be a smoothed version of its rounded spin $F_N$, let $m = \sin(2\theta)$ be the bounded formula's spin output, and let $m^{\mathrm{col}}$ be its color-excluded approximation. Write

$$\delta_s = \|z - F_N\|_2, \qquad \delta_b = \|m - z\|_2, \qquad \delta_c = \|m^{\mathrm{col}} - m\|_2.$$

These measure errors of the spin outputs, not of the angles. The rooted shift is a contraction, so

$$|\mathcal{E}(f) - \mathcal{E}(g)| \le (\|f\|_2 + \|g\|_2)\|f - g\|_2.$$

All these outputs have norm at most one. Apply synthesis with tolerance $\eta_s$ and total helper probability $p_H$, then remove the seeds with energy tolerance $\eta_r$. The source's absolute synthesis bound and seed-removal theorem, followed by fixed-word transfer, give one ordinary word $W$ with

$$|v(W) - S_{T,N}| \le 2(\delta_s + \delta_b + \delta_c) + 2\eta_s + 2\sqrt{p_H} + \eta_r. \tag{8.2}$$

The bounded formula is chosen before the colors and helper law. The excluded receiver and ancestor colors prevent a dependency from returning through a traversed branch; reserved helper colors are absent from every dependency and the output is zero on helpers. Color refinement and small positive helper probabilities jointly control $\delta_c$ and $p_H$. Only then is synthesis performed with that law fixed, using the same sitewise Gaussian seed at every seed gate. Each term in (8.2) can thus be made smaller than its allotted tolerance in this order. OpenAI (2026b), Lemmas 3.1–3.4; synthesis energy bound; seed removal.

This establishes an absolute approximation of each specified finite $S_{T,N}$ in the construction, without assuming it is already near $P_*$. It preserves the energy; it does not assert reproduction of the entire classical output law. Near-optimality now adds the cutoff and mesh choices.

Allocate errors after choosing the desired $\varepsilon > 0$. Apply the classical near-optimality proposition with the two tolerances $\varepsilon/4$ and $\eta_H = (\varepsilon/8)^2$. It supplies one formula and its helper law together, with classical loss below $\varepsilon/4$ and $0 < p_H < \eta_H$. In its proof the uncoloured formula is fixed first, then the small positive helper probabilities and the excluded-colour approximation are selected. Thus changing a helper law does not silently preserve an already fixed formula. With $\eta = \varepsilon/8$, synthesis gives

$$\mathcal{E}(v) \ge \mathcal{E}(m) - 2\eta - 2\sqrt{p_H} > P_* - 3\varepsilon/4.$$

Seed removal uses less than $\varepsilon/4$. The resulting fixed ordinary word therefore satisfies $v_W > P_* - \varepsilon$ after the fixed-word transfer. Its thermodynamic limit is taken only after all approximation choices have been made. Since $Q_p \le P_*$ and zero padding makes $Q_p$ nondecreasing, this proves the source's thermodynamic-first conclusion from its stated analytic inputs. OpenAI (2026b), §6.

The order is: accuracy; minimizer and fixed $T < 1$; finite scalar mesh; smoothing and bounded approximations; finite types and positive helper masses; exact helper contractions and finite commutator products; one fixed seeded word; initialization size; generic angle perturbation; finite selective pulse lists; sufficiently small $r$; one ordinary word; its $n \to \infty$ limit. The degree limits identify the rooted calculus and the transferred value for each already fixed word; they do not license changing that word with the finite instance.

The reconstruction makes the source's circuit mechanisms explicit, but the evidence classes remain separate. The scalar construction and its formal interfaces in §§6–7 and 12 do not by themselves establish the weighted unitary calculus, simultaneous helper contraction or seed removal. Those analytic theorems are additional inputs from the QAOA manuscript, with the domain, commutation and uniformity hypotheses specified above. No quantitative depth bound, finite-size rate, efficient angle-selection procedure, or independent formal verification of this entire circuit construction follows from the selected formal statements.

This completes the main support-to-circuit proof. The later comparisons have separate purposes: §11.1 gives another classical input; §11.2 adds a second-moment sampling theorem; and §§11.3–11.5

compare quantifiers, other models and later finite-size estimates. They are not further premises of the proof just completed.

## 9. Density and finite-disorder consequences

### 9.1 Smoothness of the minimizing coefficient

Full support does not by itself prove that the Stieltjes measure has a density. In the source manuscript, that conclusion uses the PDE again after the gap contradiction has established $\mathbb{E}a(t, X_t)^2 = 1$. We reconstruct the additional argument here because it explains a further conclusion of the original paper. It is not a clause of the selected full-support formal statement.

Put $f_k = \partial_x^k \Phi_\gamma$, so $f_1 = u$ and $f_2 = a$. All formulas in this subsection are on a fixed compact time strip, where every finite collection of spatial derivatives is bounded. Differentiating the PDE and applying Itô's formula along the associated diffusion gives

$$df_k(t, X_t) = -\frac{\gamma(t)}{2} \sum_{j=1}^{k-1} \binom{k}{j} f_{j+1}(t, X_t) f_{k-j+1}(t, X_t)\, dt + f_{k+1}(t, X_t)\, dB_t.$$

The sum is empty when $k = 1$. This identity needs no derivative of $\gamma$. For an initially nonsmooth coefficient, apply it to finite-step approximations and pass to the integrated identity: the coefficient converges in $L^1$, the spatial factors converge uniformly, and the stochastic integrals converge by the Itô isometry. OpenAI full-support manuscript, smooth-density proof, lines 35–50.

Set $b = f_3$. The case $k = 2$ gives $da = -\gamma a^2 dt + b\, dB_t$, and consequently

$$0 = \frac{d}{dt}\mathbb{E}a^2 = \mathbb{E}b^2 - 2\gamma(t)\mathbb{E}a^3 \quad \text{for almost every } t < 1.$$

Convexity gives $a \geq 0$; the already established second moment gives $\mathbb{E}a^3 \geq (\mathbb{E}a^2)^{3/2} = 1$. Thus the quotient

$$\tilde{\gamma}(t) = \frac{\mathbb{E}b(t, X_t)^2}{2\mathbb{E}a(t, X_t)^3}$$

is well defined and continuous, before any continuity of $\gamma$ is known. It equals $\gamma$ almost everywhere. Approaching an arbitrary $t < 1$ from the right through the full-measure equality set, continuity of $\tilde{\gamma}$ and right continuity of $\gamma$ give equality at that $t$ as well. This proves continuity of $\gamma$. At zero, $X_0 = 0$, and evenness of $\Phi(0, \cdot)$ gives $b(0, 0) = 0$; hence $\gamma(0) = 0$. OpenAI full-support manuscript, lines 52–66; Chen, Proposition 4.6, pp. 25–26.

The bootstrap must treat all polynomial moments simultaneously. For a polynomial $P$ in finitely many variables, write

$$M_P(t) = \mathbb{E}P(f_1(t, X_t), \dots, f_m(t, X_t)).$$

The preceding stochastic identities and Itô's product rule provide polynomials $Q_P, R_P$, involving only finitely many additional derivatives, such that

$$M_P(t) - M_P(0) = \int_0^t \left(M_{Q_P}(s) + \gamma(s) M_{R_P}(s)\right) ds.$$

Explicitly, if

$$D_k(y) = -\tfrac{1}{2} \sum_{j=1}^{k-1} \binom{k}{j} y_{j+1} y_{k-j+1},$$

then

$$Q_P = \tfrac{1}{2}\sum_{k,l} \partial_{kl} P\, y_{k+1} y_{l+1}, \qquad R_P = \sum_k \partial_k P\, D_k.$$

All these moments start continuous. Once $\gamma$ is continuous, the integral formula makes every moment $C^1$, and the quotient formula makes $\gamma$ itself $C^1$. If all moments and $\gamma$ are $C^r$, the same argument makes them $C^{r+1}$. Induction proves $\gamma \in C^\infty([0,1))$, with derivatives at zero taken from the right. Since $\gamma(0) = 0$, there is no atom at zero; its Stieltjes measure is $\gamma'(t)\,dt$, and monotonicity gives $\gamma' \geq 0$. This is the polynomial-moment method credited by the source to Auffinger–Chen and to Chen's zero-temperature treatment. OpenAI full-support manuscript, lines 68–101; Auffinger and Chen, proof of Theorem 2(ii), p. 16; Chen, Appendix B.6, p. 38.

The exact arXiv edition of Auffinger–Chen's polynomial-moment proof has a local display discrepancy: p. 16 of arXiv:1303.3573v1 prints a square in $F_3$ below (40), whereas Proposition 3, p. 9, has the cube. The identity $d(a^2) = (-2\gamma a^3 + b^2)dt + 2ab\,dB$ requires the cube used above; the moment induction itself is unchanged.

### 9.2 Why the density is strictly positive away from zero

This is a further consequence of full support, not an input to it or to §6. A smooth increasing function can have zero derivative at an interior point. For example, $g(t) = (t - \frac{1}{2})^3 + \frac{1}{8}$ on $[0,1)$ is nonnegative and strictly increasing, and $dg$ has full support, although $g'(\frac{1}{2}) = 0$. This example is not a minimizing coefficient; it disproves only the inference from strict increase to a positive derivative. The minimizing density needs a further estimate.

The smoothness just proved permits (4.2), now with $q(t) = t$. Writing $b = \Phi_{xxx}$ and $d = \Phi_{xxxx}$, it becomes

$$2\gamma'(t)\mathbb{E}a^3 = \mathcal{C}_\gamma(t).$$

Fix $t \in (0,1)$ and set $c = \gamma(t) > 0$, using full support and $\gamma(0) = 0$. The coefficient is now smooth, so Proposition 4.1 applies at this time to the same diffusion. Its proof uses nonminimizing local constant replacements, with compact-strip bounds independent of the widths of the inserted plateaus. Combining (4.1) with the preceding identity gives

$$2\gamma'(t)\mathbb{E}a^3 \geq \frac{\gamma(t)^2}{1000}\mathbb{E}a^4 > 0, \qquad 0 < t < 1.$$

The final strict inequality uses $\mathbb{E}a^4 \geq (\mathbb{E}a^2)^2 = 1$. Cauchy–Schwarz also gives $\mathbb{E}a^3 \leq \sqrt{\mathbb{E}a^2\mathbb{E}a^4} = \sqrt{\mathbb{E}a^4} \leq \mathbb{E}a^4$. Therefore the same estimate proves the quantitative ordinary consequence

$$\gamma'(t) \geq \frac{\gamma(t)^2}{2000} > 0 \quad (0 < t < 1), \qquad \mu_\gamma((r,s]) \geq \frac{s-r}{2000}\gamma(r)^2 \quad (0 < r < s < 1).$$

The second inequality follows by integration and monotonicity. Its lower bound depends on this minimizer and the left endpoint; it is not a uniform mass estimate for a family of finite-temperature measures. This argument proves only $\gamma'(0) \geq 0$ at the left endpoint. It does not claim that the minimizing coefficient itself is constant on a neighbourhood, and it does not pass the approximants' third time derivatives to the limit. Only a spatial-derivative expectation is transferred. OpenAI full-support manuscript, lines 102–152. The density result is additional to the scalar approximation in §7. The next consequence instead concerns the geometry of near-ground spin configurations.

### 9.3 Near-ground configurations at a prescribed overlap

Choose one minimizer supplied by the attained Parisi formula, and apply the full-support theorem to that same coefficient. The external landscape theorem below then applies at every point of its Stieltjes support. For every fixed $r \in [0,1)$ and $\varepsilon, \eta > 0$, there is a constant $K = K(r, \varepsilon, \eta) > 0$, independent of $n$, such that

$$\mathbb{P}\left(\exists \sigma, \tau \in \{-1,1\}^n : \left|\frac{1}{n}\sum_{i=1}^{n} \sigma_i \tau_i - r\right| < \varepsilon, \quad \frac{H_n(\sigma)}{n}, \frac{H_n(\tau)}{n} \geq P_* - \eta\right) \geq 1 - Ke^{-n/K}, \qquad n \geq 1.$$

The probability is over the SK couplings; the configurations may depend on them. This is the specialization of Auffinger–Chen–Zeng, Proposition 1, equation (5), p. 5 to the full-support minimizer above. Their theorem applies to each point of its Stieltjes support. Its proof, on pp. 15–16, uses the coupled free energy constrained to a neighbourhood of that overlap, passes to the coupled maximum and applies Gaussian concentration. The coupled maximum approaches $2P_*$, while the unrestricted maximum approaches $P_*$; these two facts force both members of a maximizing pair to be near ground energy.

The covariance normalization needs one small adjustment. Writing $R = n^{-1}\sum_i \sigma_i \tau_i$, our Hamiltonian has covariance $nR^2/2 - 1/2$. Adjoin a common independent Gaussian $C$ of variance $1/2$. Then $\widetilde{H}_n(\sigma) = H_n(\sigma) + C$ has covariance $n\xi(R)$ with $\xi(t) = t^2/2$, as in the cited mixed-spin theorem. Its expected maximum equals that of $H_n$, so its ground-state limit is the same $P_*$. Apply the theorem with energy tolerance $\eta/2$. On $|C| \leq \eta n/2$, the resulting pair also satisfies the displayed energy inequalities for $H_n$. The omitted event has probability at most $2e^{-\eta^2 n^2/4}$, which can be absorbed into $Ke^{-n/K}$ by enlarging $K$. Since the conclusion for $H_n$ depends only on its couplings, this also removes the auxiliary Gaussian from the probability statement.

The pointwise result also gives a simultaneous statement at any fixed positive resolution. Given $\varepsilon, \eta > 0$, choose finitely many centres $r_j \in [0,1)$ whose open $\varepsilon/2$-neighbourhoods cover $[0,1]$. Apply the preceding result at each centre with overlap tolerance $\varepsilon/2$ and energy tolerance $\eta$. A finite union bound gives constants $C, c > 0$, depending only on these tolerances, such that

$$\mathbb{P}(\forall r \in [-1,1]\ \exists \sigma, \tau \in \{-1,1\}^n : |R(\sigma,\tau) - r| < \varepsilon, \quad H_n(\sigma)/n, H_n(\tau)/n \geq P_* - \eta) \geq 1 - Ce^{-cn}. \tag{9.1}$$

Indeed, on the event where every centre has a suitable pair, choose a nearest centre and use the triangle inequality. For negative targets, reverse one spin configuration: the pure two-spin Hamiltonian preserves its energy and changes the overlap's sign. The finite net covers the endpoints as well, with centres strictly below one.

There is a useful contrast at a fixed finite size. If $\sigma, \tau$ differ at $k$ sites, precisely the $k(n-k)$ edges crossing that set change their spin product. Their independent Gaussian contributions give

$$\operatorname{Var}(H_n(\sigma) - H_n(\tau)) = \frac{4k(n-k)}{n}.$$

For $0 < k < n$ this is positive, so the two energies tie with probability zero. A finite union over pairs, together with $H_n(-\sigma) = H_n(\sigma)$, shows that almost surely the exact maximum has one maximizing pair $\{\sigma_*, -\sigma_*\}$. The overlap conclusion above concerns nearly optimal pairs at intermediate separations. Its threshold $P_* - \eta$ is deterministic; setting $\eta = 0$ would not turn it into exact maximization of a finite sample. This elementary deduction uses only the displayed Gaussian Hamiltonian and is separate from the external asymptotic landscape theorem.

The overlap deduction uses fixed tolerances and target-dependent pairs. It gives neither exact arbitrary real overlaps at finite $n$, a typical-pair law, a search procedure, nor a guarantee for tolerances shrinking with $n$. It uses the external landscape theorem, full support and attainment, and is outside the two selected formal statements in §12. The separate algorithmic consequence is in §9.4.

### 9.4 A classical optimization algorithm

The scalar Gaussian construction is not itself an algorithm on the matrix of SK couplings. A finite-disorder algorithm is obtained by importing the theorem of El Alaoui, Montanari and Sellke. In their cited preprint, Assumption 2 asks for a strictly increasing minimizer of the zero-temperature Parisi functional on $\mathcal{U}$. The support theorem gives this property, and the imported attainment argument supplies the minimizer. Their Corollary 2.2 then returns, for each fixed $\varepsilon > 0$, a randomized spin configuration within $\varepsilon n$ of the finite-instance maximum with probability tending to one. Its operation count is the one in the displayed Theorem 3 on p. 7; the preprint's running references call that result "Theorem 3.2." El Alaoui, Montanari and Sellke, arXiv:2001.00904v1, pp. 7–8.

To match that theorem's Gaussian input exactly, form a symmetric matrix with

$$A_{ij} = J_{ij}/\sqrt{n} \quad (i < j), \qquad A_{ii} = \sqrt{2/n}\, G_i,$$

where the $G_i$ are additional independent standard Gaussians. For $\widehat{H}_n(x) = \frac{1}{2} x^{\mathsf{T}} A x$,

$$\mathbb{E}\widehat{H}_n(\sigma)\widehat{H}_n(\tau) = \frac{(\sigma^{\mathsf{T}}\tau)^2}{2n}, \qquad \widehat{H}_n(\sigma) = H_n(\sigma) + \frac{1}{\sqrt{2n}} \sum_i G_i.$$

The final term is the same for all spin configurations. Therefore the maximizing spin and every optimization gap are unchanged. The extra diagonal randomness is included in the algorithm's randomness. This checks the input convention without changing the target optimization problem. OpenAI full-support manuscript, algorithm application, lines 30–47.

The imported operation count is $C(\varepsilon)(\chi + n) + n\chi_1$, where $\chi$ is the cost of evaluating the full gradient and $\chi_1$ is the cost of one coordinate of the distinct-index gradient. Here $\nabla\widehat{H}_n(x) = Ax$ costs $O(n^2)$, and one coordinate of $\nabla H_n(x)$ costs $O(n)$. The resulting count is $C(\varepsilon)n^2$, with $C(\varepsilon)$ independent of $n$. Real sums and multiplications have unit cost in the cited model; the scalar functions and normalizers prescribed for the chosen accuracy are part of that construction. This statement supplies neither a bound for their numerical preprocessing nor a finite-precision bit-complexity bound. El Alaoui, Montanari and Sellke, Remark 2.1 and Theorem 3, p. 7; OpenAI full-support manuscript, lines 1–11 and 49–59.

A distinct classical construction is Jekel, Sandhu and Shi, *Potential Hessian Ascent*, version 3, Assumption 1.4 and Theorem 1.5. Its hypothesis is finite-temperature interval support $\operatorname{supp}\mu_\beta = [0, q_\beta^*]$, with $q_\beta^* \to 1$. Its inputs include oracles for the Parisi PDE, its derivatives and the minimizer's distribution function. For each fixed accuracy the stated runtime is polynomial in $n$, with an exponential dependence on inverse accuracy; it does not include implementing those oracles. Its Ginibre Hamiltonian has covariance $nR^2$: on spins it has the law $\sqrt{2}H_n$ plus a Gaussian term independent of the configuration. Thus its energy and accuracy must be divided by $\sqrt{2}$ to compare with this paper's normalization. This is a separate algorithm theorem with separate analytic inputs, rather than another interpretation of the finite Gaussian spin in §7.

The consequence is classical. It does not supply an efficient construction of QAOA angles, a quantum depth bound, or a clause of either selected Lean interface. The proof of the separate QAOA statement still requires its rooted energy, synthesis, seed-removal and transfer arguments.

## 10. An independent scalar comparison

### 10.1 Contact from convexity on constant intervals

There is a second way to obtain $q = t$ from the variational inequalities. It uses convexity on intervals where the tail potential is positive, rather than first proving full Stieltjes support. This is the argument used by the scalar implementation; it does not depend on the formal full-support theorem.

In the scalar model put $q(t) = \mathbb{E}[u(t, X_t)^2]$, $f(t) = q(t) - t$ and $G(t) = \int_t^1 f(s)\, ds$. The source first obtains $G \geq 0$, $G(1) = 0$, $q(0) = 0$ and $G' = -f$. Compact-mass variations also show that on any interval where $G > 0$ the coefficient is constant: if its cumulative mass increased between two points there, integration of the strictly positive $G$ against that positive increment would contradict the mass-removal variational inequality. This is the content of the scalar contact-integral argument.

Suppose now that $G(x) > 0$ for some $0 < x < 1$. Choose the component $(a, b)$ of this positivity set containing $x$, allowing $a = 0$ or $b = 1$. Its coefficient is constant, say $c \geq 0$, and $G(b) = 0$. At the left endpoint, $f(a) = 0$: this is the contact derivative when $a > 0$, and $q(0) = 0$ when $a = 0$.

If $c > 0$, one also has $G(a) = 0$. The only extra case is $a = 0$: right continuity gives $\gamma(0) = c > 0$, so the mass-removal condition at that initial atom forces $G(0) = 0$. The mean-value theorem on $[a, x]$ and $[x, b]$ now provides points $r < s$ with $f(r) < 0 < f(s)$. There is a later point $z \leq b$, with $z < 1$, for which $f(z) \leq f(s)$. For $b < 1$ take $z = b$, since contact gives $f(b) = 0$. For $b = 1$, choose $z$ sufficiently close to one and use $f(z) = q(z) - z \leq 1 - z < f(s)$.

The three intervals $[a, r]$, $[r, s]$ and $[s, z]$ therefore have secant slopes for $f$ of signs respectively negative, positive and nonpositive. Applying the mean-value theorem once more gives three ordered points where $f'$ has those signs. This is impossible when $f' = q' - 1$ is convex: its middle value cannot exceed the larger of its two exterior values. The scalar source supplies precisely this nonstrict convexity on a constant interval. See the positive-hump argument and its endpoint condition and the curvature-moment convexity lemma.

If $c = 0$, the source instead gives convexity of $f = q - t$. A mean-value argument between $x$ and $b$ gives $f(s) > 0$; the same right-endpoint argument produces $z > s$ with $f(z) \leq f(s)$. Since $f(a) = 0$,

$$\frac{f(s) - f(a)}{s - a} > 0 \geq \frac{f(z) - f(s)}{z - s},$$

contradicting monotonicity of adjacent secant slopes of a convex function. This case does not need $G(a) = 0$; see the zero-coefficient hump lemma.

Thus $G$ cannot have a positive component, and $G \equiv 0$ on $(0, 1)$. Differentiation gives $q(t) = t$ there, with the origin supplied separately. The scalar assembly then obtains the unit curvature moment, finite Gaussian approximation and terminal value identities from this contact equality. Nonstrict convexity suffices for this scalar route. The strict curvature estimate in Sections 4–5 has the further job of excluding gaps in Stieltjes support. These are source-level proof reconstructions, not a fresh Lean acceptance or a claim that the two formal models have been composed.

### 10.2 A universal bound for predictable correlations

The scalar implementation obtains the value equality by two inequalities. Its finite-spin limit follows from superadditivity and a uniform upper bound. A finite-profile Gaussian comparison supplies $P_* \le \mathcal{P}(\gamma)$ for every admissible coefficient. The reverse inequality at a minimizer comes from a more general finite-dimensional fact: predictable orthonormal Gaussian functions paired with any spin have a shifted correlation sum at most $P_*$. We reconstruct that fact below. It supplies the inequality that §7's mesh limit consumes, without invoking the literature's Parisi formula used in §6.

Write $e_n = n^{-1}\mathbb{E}\max_\sigma H_n(\sigma)$, with the Hamiltonian of §1. The finite-spin convergence and upper comparison are separate inputs to this subsection; their precise suppliers are the superadditive limit and the upper comparison. The proof below concerns the lower comparison, not a finite-disorder algorithm or a circuit.

**Proposition 10.1 (predictable Gaussian comparison).** Fix $K < \infty$, independent standard normals $z_0, \dots, z_K$, and real functions $f_j = f_j(z_0, \dots, z_j)$ for $0 \le j < K$. Assume measurability, square integrability and $\mathbb{E}[f_i f_j] = \delta_{ij}$. For any measurable $S = S(z_0, \dots, z_K) \in \{-1, 1\}$, the conclusion is

$$\sum_{j=0}^{K-1} \mathbb{E}[f_j S]\,\mathbb{E}[S z_{j+1}] \le P_*.$$

These assumptions allow nonconstant, dependent $f_j$; even $\mathbb{E}f_j = 0$ is unnecessary. Their order of dependence ensures $\mathbb{E}[f_i z_{j+1}] = 0$ whenever $i \le j$. The next fresh coordinate, rather than centring of $f_i$, is what will remove the projection terms. Scalar implementation, predictable-coefficient theorem.

**10.2.1 Revealing one Gaussian column** For $w \in \mathbb{R}^n$, put $Q(w) = w \otimes w/\sqrt{2}$. An independent standard Gaussian matrix $G$ gives a centered field $\langle Q(w), G\rangle$ with covariance $\langle w, w'\rangle^2/2$. Given a unit vector $v$, define

$$R_v w = w - \langle v, w\rangle v, \qquad C_v(w) = \langle v, w\rangle R_v w + \frac{\langle v, w\rangle^2}{\sqrt{2}} v.$$

Write $\alpha = \langle v, w\rangle$, $\beta = \langle v, w'\rangle$ and $r = \langle R_v w, R_v w'\rangle$. For an independent standard Gaussian column $Z$ and matrix $G'$, the covariance of $\langle C_v(w), Z\rangle + \langle Q(R_v w), G'\rangle$ is

$$\alpha\beta r + \frac{1}{2}\alpha^2\beta^2 + \frac{1}{2}r^2 = \frac{1}{2}(r + \alpha\beta)^2 = \frac{1}{2}\langle w, w'\rangle^2.$$

Thus the new field has the old joint Gaussian law. The same statement holds for $v = 0$: then $R_v$ is the identity and $C_v = 0$. Allowing that case will avoid assuming empirical linear independence.

For a finite nonempty family of candidates $w_s$ and offsets $d_s$, define

$$\mathcal{V}(w, d) = \mathbb{E}_G \max_s \{\langle Q(w_s), G\rangle + d_s\}.$$

The covariance identity and Fubini give the exact recursion

$$\mathcal{V}(w, d) = \mathbb{E}_Z \mathcal{V}((R_v w_s)_s, (d_s + \langle C_v(w_s), Z\rangle)_s).$$

Also $d_s \le \mathcal{V}(w, d)$ for each $s$, since the residual Gaussian term has mean zero. These are the two facts needed for adaptive choices. The source derives the equality by Gaussian comparison in both directions and proves the required integrability in the one-column decomposition.

**10.2.2 Adaptive directions and a finite-spin inequality** Choose each zero-or-unit direction measurably from columns already revealed. Conditional on that history, the current residual candidate vectors and offsets are fixed, so the preceding recursion applies. More precisely, conditional on the revealed history, the expected next continuation value, including its accumulated offsets, equals the current one. Iteration therefore preserves the initial expected maximum. For any measurable choice of a candidate from the final history, its accumulated offset is bounded by the remaining conditional expected maximum. Hence

$$\mathbb{E}\, d^{\text{final}}_{s(\text{history})} \le \mathcal{V}(w, 0).$$

Contraction of each residual, the finite candidate set and Gaussian column moments justify the expectations. Adaptivity does not turn a future column into part of the conditioning history. Scalar implementation, adaptive recursion.

Take all spins $w_s \in \{-1, 1\}^n$ as candidates. The full quadratic matrix has off-diagonal coefficients $(G_{ij} + G_{ji})/\sqrt{2}$, which are independent standard normals, and a centered diagonal contribution common to every spin. Thus

$$\mathcal{V}(\text{spins}, 0) = \sqrt{n}\, \mathbb{E} \max_{\sigma} H_n(\sigma).$$

Dividing the offset inequality by $n\sqrt{n}$ gives a bound by $e_n$. The implementation proves this normalization through equality of Gaussian increment distances in the finite adaptive SK comparison.

To obtain the desired coefficients, take independent rows $z^{(1)}, z^{(2)}, ...$, each with the law of $(z_0, \dots, z_K)$, and put

$$\hat{g}_n = n^{-1/2}(g(z^{(1)}), \dots, g(z^{(n)})).$$

Let $v_{n,j}$ be the normalized Gram–Schmidt vector obtained from $\hat{f}_{0,n}, \dots, \hat{f}_{j,n}$; a zero remainder is normalized to zero. Start the adaptive procedure with a zero direction to reveal column $z_0$. After seeing columns through $z_j$, $v_{n,j}$ is available, and the next column is $z_{j+1}$. This explains both the extra column and the shift.

Let $R_{n,j}$ project away $v_{n,0}, \dots, v_{n,j-1}$ and set

$$E_{n,j} = \langle C_{v_{n,j}}(R_{n,j}\widehat{S}_n), \hat{z}_{j+1,n} \rangle.$$

The raw spin and fresh column are each $\sqrt{n}$ times their empirical versions; $C_v$ is quadratic in its input. Consequently their revealed energy is exactly $n\sqrt{n}E_{n,j}$. Transposing the array between independent rows and independent columns preserves its Gaussian law. The finite comparison therefore gives, for every $n > 0$,

$$\mathbb{E} \sum_{j<K} E_{n,j} \le e_n.$$

This is an actual choice of the adaptive directions and final rowwise spin, not an uninstantiated conditional inequality. Empirical directions and normalization; row–column transposition and finite bound.

**10.2.3 Why orthogonalization preserves the limiting correlations** Keep $K$ fixed while $n \to \infty$. The strong law applies to the finitely many needed products, which are integrable by Cauchy–Schwarz. The empirical Gram matrix tends to the identity. Inductively in Gram–Schmidt, every projection coefficient tends to zero and every remaining norm tends to one; hence $\|v_{n,j} - \hat{f}_{j,n}\| \to 0$ almost surely. In particular,

$$\langle v_{n,j}, \widehat{S}_n \rangle \to a_j := \mathbb{E}[f_j S], \qquad \langle \widehat{S}_n, \hat{z}_{j+1,n} \rangle \to b_{j+1} := \mathbb{E}[S z_{j+1}],$$

while $\langle v_{n,i}, \hat{z}_{j+1,n}\rangle \to 0$ for $i \le j$. The last limit uses precisely the fresh-coordinate cancellation stated at the start of §10.2. Gram–Schmidt limits.

Expand the revealed energy before taking its limit:

$$E_{n,j} = \langle v_{n,j}, \widehat{S}_n\rangle\langle R_{n,j+1}\widehat{S}_n, \hat{z}_{j+1,n}\rangle + \frac{\langle v_{n,j}, \widehat{S}_n\rangle^2}{\sqrt{2}}\langle v_{n,j}, \hat{z}_{j+1,n}\rangle.$$

All terms introduced by projecting away earlier directions vanish, as does the second line. The remaining product tends to $a_j b_{j+1}$. Thus $\sum_{j<K} E_{n,j} \to \sum_{j<K} a_j b_{j+1}$ almost surely. This establishes the random-variable limit, but an expectation bound still requires the next step. Scalar implementation, empirical energy limit.

**10.2.4 Passing the expectation bound and returning to the value** Since $S^2 = 1$, $\|\widehat{S}_n\| = 1$ for $n > 0$, and every residual projection contracts. The column formula gives $|E_{n,j}| \le 1 + \|\hat{z}_{j+1,n}\|^2$. Define

$$c_n = \sum_{j<K}(1 + \|\hat{z}_{j+1,n}\|^2).$$

Then $\sum_j E_{n,j} + c_n \ge 0$, $\mathbb{E}c_n = 2K$ exactly, and $c_n \to 2K$ almost surely. Fatou's inequality, the finite bound and $e_n \to P_*$ imply

$$\sum_{j<K} a_j b_{j+1} + 2K \le \liminf_n \mathbb{E}[\sum_{j<K} E_{n,j} + c_n] \le P_* + 2K.$$

Cancelling $2K$ proves the claimed predictable-coefficient inequality. The source's constant-limit helper establishes the same nonnegative comparison by clipping at the limiting constant and applying dominated convergence. It shifts to dimensions $n + 1$ to avoid dimension zero. It does not apply Fatou directly to the signed energies or presume uniform integrability. Column bound and correction; expectation limit and final theorem.

For the Euler construction in §7, take $K = N$, relabel its $Z_{j+1}$ as $z_j$, and set $f_j = g_{j+1}$. Each normalized increment depends only on $z_0, \dots, z_j$ and the increments are orthonormal when their raw second moments are positive. The rounded sign is spin-valued almost surely: its tie event is null by the independent last coordinate. The scalar implementation uses a measurable Bool spin equal to that sign almost surely. It follows that

$$S_{T,N} = \sum_{j=1}^{N} A_j B_{j+1} \le P_*.$$

This finite inequality assumes admissibility, $0 \le T < 1$, $N > 0$ and positive active second moments, but no minimizer or diffusion. Literal scalar specialization.

At a minimizer, the separate contact and diffusion arguments establish the fixed-strip inputs and eventual positivity. First take the mesh limit to obtain $I(T) \le P_*$, then let $T \uparrow 1$. The scalar terminal assembly identifies that last limit with $\mathcal{P}(\gamma)$; together with $P_* \le \mathcal{P}(\gamma)$ this proves equality. The third scalar endpoint constructs its own diffusion before making this application. These sources therefore supply an independent route to the equality, alongside the imported Parisi formula in §6. Contact assembly. The inspection here does not change the research replay outcome in §13.

### 10.3 The zero coefficient

The admissible coefficient $\gamma_0 = 0$ makes the use of minimality in the normalization step explicit. Its diffusion is $X_t = B_t$, and the zero-coefficient heat formula gives

$$u_0(t,x) = 2F_{\mathrm{G}}\left(\frac{x}{\sqrt{1-t}}\right) - 1, \qquad a_0(t,x) = \sqrt{\frac{2}{\pi}}\frac{e^{-x^2/(2(1-t))}}{\sqrt{1-t}}.$$

Since $\mathbb{E}e^{-\lambda B_t^2} = (1+2\lambda t)^{-1/2}$,

$$\mathbb{E}a_0(t,B_t) = \sqrt{\frac{2}{\pi}}, \qquad \mathbb{E}a_0(t,B_t)^2 = \frac{2}{\pi\sqrt{1-t^2}}, \qquad q_0(t) = \frac{2}{\pi}\arcsin t.$$

The last identity follows by integrating $q_0' = \mathbb{E}a_0^2$ from $q_0(0) = 0$. This heat and late-step variation argument is the zero-field specialization of Chen, Handschy and Lerman, Theorem 6 and its proof, arXiv:1609.04368v4, pp. 13–16. Strict convexity of arcsin gives $q_0(t) < t$ for $0 < t < 1$. Thus $G_0(s) = \int_s^1 (q_0(t) - t)\,dt < 0$ for every $s < 1$. Inserting an atom at $s$ is admissible, and its first variation is $G_0(s)/2 < 0$. In particular, the derivative for inserting mass at zero is $1/4 - 1/\pi < 0$. Hence this coefficient does not minimize.

Here the Euler scheme is exact: $Y_j = B_{t_j}$ under the Brownian coupling. Nevertheless its normalizers are

$$c_j = \sqrt{\frac{\pi}{2}}\,(1 - t_{j-1}^2)^{1/4},$$

so they approximate this nonconstant function of time, rather than converging uniformly to one. Already $c_1 = \sqrt{\pi/2}$ for every mesh. The directions $g_j$ remain orthonormal by construction and the rounding still preserves conditional means; the missing input to the value limit is the minimizing identity $\mathbb{E}a(t,X_t)^2 = 1$.

The terminal spin also survives: $U_1 = \operatorname{sgn}(B_1)$ almost surely. But its correlation is

$$\mathbb{E}[U_1 B_1] = \sqrt{\frac{2}{\pi}} = \mathcal{P}(\gamma_0),$$

the value of this nonminimizing candidate. The negative first variation proves that this value is strictly above $P_*$. Admissibility, exact Euler sampling, and a spin-valued terminal martingale therefore do not supply the minimizing normalization or identify the ground-state value.

### 10.4 An exact one-step calculation

For a finite check of the definitions, use the admissible heat coefficient $\gamma_0 \equiv 0$, whose solution has

$$u_0(t,x) = 2F_{\mathrm{G}}(x/\sqrt{1-t}) - 1, \qquad a_0(0,0) = \sqrt{2/\pi}.$$

This coefficient is not a minimiser; the calculation tests the construction and its indices, not the minimiser-dependent convergence theorem. Take $T = 4/5$ and $N = 1$. Then

$$Y_1 = \frac{2}{\sqrt{5}}Z_1, \qquad c_1 = \sqrt{\pi/2}, \qquad g_1 = Z_1, \qquad m_1 = u_0(T,Y_1) = 2F_{\mathrm{G}}(2Z_1) - 1.$$

Strict monotonicity and symmetry of the Gaussian distribution function give

$$F_1 = \operatorname{sgn}(F_{\mathrm{G}}(2Z_1) + F_{\mathrm{G}}(Z_2) - 1) = \operatorname{sgn}(2Z_1 + Z_2)$$

almost surely. Thus the additional Gaussian is visibly the rounding coordinate. Set $V = (2Z_1 + Z_2)/\sqrt{5}$, a standard normal. Gaussian regression gives $\mathbb{E}[Z_1 \mid V] = 2V/\sqrt{5}$ and $\mathbb{E}[Z_2 \mid V] = V/\sqrt{5}$. Since $\mathbb{E}|V| = \sqrt{2/\pi}$,

$$A_0 = 0, \qquad A_1 = B_1 = 2\sqrt{\frac{2}{5\pi}}, \qquad B_2 = \sqrt{\frac{2}{5\pi}}.$$

Consequently

$$\sum_{j=0}^{1} A_j B_{j+1} = \frac{4}{5\pi} \quad \text{whereas} \quad A_1 B_1 = \frac{8}{5\pi}.$$

Dropping the index shift changes the answer by a factor of two. At $N = 1$ the interior Riemann sum is empty and the entire product comes from the rounding endpoint. Its later disappearance is an asymptotic statement, not an exact finite-mesh identity. The displayed coefficients follow from exact Gaussian regression; no numerical quadrature or finite computation is needed for this example.

**10.4.1 The same test at every mesh size** For the nonminimizing coefficient $\gamma = 0$, Euler sampling is exact: $Y_j = B_{t_j}$. The curvature moment in §10.3 gives

$$c_j = \sqrt{\pi/2}(1 - t_{j-1}^2)^{1/4}.$$

Monotonicity and symmetry of the Gaussian CDF turn the rounded spin into

$$F_N = \operatorname{sgn}(B_T + \sqrt{1-T}\, Z_{N+1}) \quad \text{almost surely.}$$

The argument of this sign is standard normal. Gaussian regression gives $B_i = \sqrt{2/\pi}\sqrt{\Delta}$ for $1 \le i \le N$ and $B_{N+1} = \sqrt{2/\pi}\sqrt{1-T}$. To calculate $A_j$, condition on Brownian motion through $t_j$ and use the heat martingale. Gaussian integration by parts then gives

$$\mathbb{E}[Z_j u_0(t_j, x + \sqrt{\Delta} Z_j)] = \sqrt{\Delta}\, a_0(t_{j-1}, x).$$

Multiplication by $c_j a_0(t_{j-1}, B_{t_{j-1}})$ and expectation yield $A_j = c_j\sqrt{\Delta}\, \mathbb{E} a_0(t_{j-1}, B_{t_{j-1}})^2 = \sqrt{\Delta}/c_j$. Consequently, for every $N \ge 1$,

$$S_{T,N} = \sqrt{\frac{2}{\pi}} \left( \Delta \sum_{j=1}^{N-1} \frac{1}{c_j} + \frac{\sqrt{\Delta(1-T)}}{c_N} \right).$$

The one-step value above is its $N = 1$ case. At fixed $T < 1$, its limit is

$$\frac{2}{\pi} \int_0^T (1 - t^2)^{-1/4}\, dt,$$

whereas the unnormalized candidate covariance is $I_0(T) = T\sqrt{2/\pi}$. The subsequent $T \uparrow 1$ limit is $4\Gamma(3/4)/(\sqrt{\pi}\,\Gamma(1/4)) \approx 0.76275976$, distinct from $\mathcal{P}(0) = \sqrt{2/\pi} \approx 0.79788456$. The general predictable comparison of §10.2 turns this limit into a lower bound for $P_*$, as shown next. It is not a numerical determination of $P_*$ or the value of the zero coefficient. Exact Euler simulation,

orthonormal predictable increments and correct rounding all hold here. The missing identity is $\mathbb{E}a_0(t, B_t)^2 = 1$; consequently the normalizers do not approach one. This calculation extends the heat and rounding derivations of §§10.3–10.4 and isolates the role of minimality in the convergence theorem.

**Corollary 10.2 (the heat test).** In this normalization,

$$P_* \geq \frac{4\Gamma(3/4)}{\sqrt{\pi}\,\Gamma(1/4)} \approx 0.7627597635. \tag{10.2}$$

For each fixed $T < 1$ and $N \geq 1$, the heat normalizers $c_j$ are positive. The increments $g_j$ have unit second moments and are orthogonal, since their multipliers depend only on preceding Gaussians. The rounded $F_N$ is a measurable spin after an arbitrary choice on its null tie event. Thus Proposition 10.1 applies with $K = N$, $z_j = Z_{j+1}$ and $f_j = g_{j+1}$, giving $S_{T,N} \leq P_*$. First let $N \to \infty$ at fixed $T$, using the displayed exact sum, and then let $T \uparrow 1$. The integrand is nonnegative and its endpoint singularity has exponent $1/4 < 1$. Substitution $s = t^2$ yields

$$P_* \geq \frac{2}{\pi}\int_0^1 (1-t^2)^{-1/4}\,dt = \frac{1}{\pi}B\Big(\frac{1}{2},\frac{3}{4}\Big) = \frac{4\Gamma(3/4)}{\sqrt{\pi}\,\Gamma(1/4)}.$$

This is a consequence of the heat calculation and the general Gaussian comparison. Minimality is unnecessary for the inequality; it is what made the different, optimizing construction converge to $P_*$ itself.

## 11. Alternative constructions and circuit limits

### 11.1 The separate positive-temperature route

The positive-temperature appendix supplies a separate classical input to the same synthesis and seed-removal theorems. Fix $\beta_0 > 1$, let $\mu_{\beta_0}$ be its Parisi measure, and write $q = q_{\beta_0} = \max\operatorname{supp}\mu_{\beta_0}$. Here $\gamma(t) = \beta_0\mu_{\beta_0}([0,t])$, $m = \phi_{\beta_0,x}$, $a = \phi_{\beta_0,xx}$ and $X$ belong to the positive-temperature PDE, whose terminal value is $\beta_0^{-1}\log(2\cosh(\beta_0 x))$. They are distinct from the zero-temperature objects in §§1–8. The input credited to Lopatto, arXiv:2607.11756v3, Theorem 1.1 gives $\operatorname{supp}\mu_{\beta_0} = [0,q]$ for $\beta_0 > 1$; its smooth density below $q$ gives the regularity used on a fixed $[0,T]$ with $0 < T < q$.

The normalization can be checked directly against Lopatto's PDE. Writing $\widehat{\Phi}_{\beta_0}(t,y)$ for his solution with terminal value $\log\cosh y$ and covariance factor $\beta_0^2$, set

$$\phi_{\beta_0}(t,x) = \frac{\widehat{\Phi}_{\beta_0}(t,\beta_0 x) + \log 2}{\beta_0}, \quad m = \widehat{\Phi}_y(t,\beta_0 x), \quad a = \beta_0\widehat{\Phi}_{yy}(t,\beta_0 x).$$

His diffusion $\widehat{X}$ has noise coefficient $\beta_0$; hence $X = \widehat{X}/\beta_0$ has unit noise and drift $\beta_0\mu_{\beta_0}([0,t])m$. His identity $\beta_0^2\mathbb{E}\widehat{\Phi}_{yy}(t,\widehat{X}_t)^2 = 1$ is exactly $\mathbb{E}a(t,X_t)^2 = 1$. This is a fixed-temperature change of variables, not convergence of either Parisi measure as $\beta_0 \to \infty$. Lopatto, arXiv:2607.11756v3, equations (1.3), (2.1)–(2.2) and Lemma 2.2.

The source derives $\mathbb{E}a(t,X_t)^2 = 1$ on $[0,q]$ and $1 - q \leq \beta_0^{-1}$. The value estimate is

$$\int_0^T \mathbb{E}a(t,X_t)\,dt \geq P_* - \frac{1/4 + \log 2}{\beta_0} - (q - T).$$

To recover its constants, martingale contact first gives $\int_0^t \mathbb{E}a(s,X_s)^2 ds = t$ on $[0,q]$; continuity yields unit second moment including the endpoints. On $[q,1]$, $\gamma = \beta_0$, so Cole–Hopf gives

$$\phi(q,x) = \beta_0^{-1}\log(2\cosh\beta_0 x) + \beta_0(1-q)/2, \quad m(q,x) = \tanh(\beta_0 x), \quad a(q,x) = \beta_0 \operatorname{sech}^2(\beta_0 x).$$

The endpoint identity $\mathbb{E}a(q,X_q)^2 = 1$ is also needed. Cauchy–Schwarz gives

$$1-q = \mathbb{E}\operatorname{sech}^2(\beta_0 X_q) \le \left(\mathbb{E}\operatorname{sech}^4(\beta_0 X_q)\right)^{1/2} = \beta_0^{-1}.$$

This is the endpoint argument in OpenAI (2026b), Appendix A, Lemma "A positive-temperature value bound". Itô's product rule and the value evolution give

$$\int_0^q \mathbb{E}a = \mathbb{E}[X_q m(q,X_q)] - \int_0^q t\gamma(t)dt, \quad \mathbb{E}\phi(q,X_q) = \phi(0,0) + \tfrac{1}{2}\int_0^q t\gamma(t)dt.$$

To compare this integral with the spin optimum, define the expected free energy in energy units by

$$F_{n,\beta_0} = (n\beta_0)^{-1}\mathbb{E}\log\sum_\sigma e^{\beta_0 H_n(\sigma)}, \qquad F_{\beta_0} = \lim_{n\to\infty} F_{n,\beta_0}.$$

The fixed-temperature limit is supplied by the positive-temperature Parisi formula cited below. The finite-size soft-maximum inequality gives $F_{\beta_0} \ge P_*$. Using $x\tanh(\beta_0 x) \ge \beta_0^{-1}\log(2\cosh\beta_0 x) - \beta_0^{-1}\log 2$ and retaining the penalty on $[q,1]$ yields

$$\int_0^q \mathbb{E}a \ge F_{\beta_0} - \frac{\beta_0(1-q)^2}{4} - \frac{\log 2}{\beta_0}.$$

The coefficient $1/4$ comes from the two tail contributions, since $\gamma = \beta_0$ above $q$:

$$\frac{\beta_0}{2}\int_q^1 t\,dt - \frac{\beta_0}{2}(1-q) = -\frac{\beta_0}{4}(1-q)^2.$$

The cutoff removes at most $q-T$, because $0 \le \mathbb{E}a \le 1$. This proves the displayed estimate. Choose first a sufficiently large finite $\beta_0$, then $T < q$ close enough to $q$. This route obtains the existence and value of the ground-state limit from the positive-temperature Parisi formula of Talagrand (2006), Theorem 1.1, pp. 222–223. At each fixed $\beta_0$, his covariance hypotheses (1.3)–(1.5) hold with $\xi(t) = \beta_0^2 t^2/2$ and error $c(n) = \beta_0^2/(2n)$. His partition sum (1.14) is unnormalized; dividing its logarithmic free energy by $\beta_0$ gives the convention here, including the $\log 2/\beta_0$ entropy term. The finite-size inequality used above is

$$F_{n,\beta_0} - (\log 2)/\beta_0 \le n^{-1}\mathbb{E}\max H_n \le F_{n,\beta_0}.$$

It does not invoke the zero-temperature companion to justify that step. Write $M_n = n^{-1}\mathbb{E}\max H_n$. The same uniform entropy bound gives

$$|F_{n,\beta_0} - P_*| \le |M_n - P_*| + \frac{\log 2}{\beta_0}.$$

Thus arbitrary joint $n,\beta_0 \to \infty$ are permitted for these scalar free-energy values. This says nothing about joint convergence of measures, PDE derivatives, message calculations or circuit words. OpenAI (2026b), Appendix A.

The remaining limits have distinct tasks. At fixed $\beta_0, T, \delta, L$, with $\delta > 0$, integer $L \geq 1$ and $L\delta \leq T$, the degree limit $D \to \infty$ identifies the finite message law and its edge value. Let $U_0, U_1, \ldots$ be independent standard normals. Put

$$X_0^\delta = \sqrt{\delta}\, U_0, \qquad X_{r+1}^\delta = X_r^\delta + \delta\gamma(r\delta)m(r\delta, X_r^\delta) + \sqrt{\delta}\, U_{r+1}, \qquad c_{\delta,r} = (\mathbb{E}a(r\delta, X_r^\delta)^2)^{-1/2}.$$

The finite-degree output is a message calculation on the $(D+1)$-regular tree, where $\partial i$ is the set of neighbours of vertex $i$. Write $t_r = r\delta$ and $b(t,x) = \gamma(t)m(t,x)$. Initialise independent standard normal labels $g_i$, with $u_{i\to j}^0 = u_i^0 = g_i$, $A_{i\to j}^{-1} = 1$ and $x_{i\to j}^0 = x_i^0 = \sqrt{\delta}\, g_i$. For $0 \leq r < L$ set

$$u_{i\to j}^{r+1} = D^{-1/2} \sum_{\substack{v \in \partial i \\ v \neq j}} A_{v\to i}^{r-1} u_{v\to i}^r, \qquad u_i^{r+1} = (D+1)^{-1/2} \sum_{v \in \partial i} A_{v\to i}^{r-1} u_{v\to i}^r.$$

Each $x$ then takes the Euler step $x^{r+1} = x^r + \delta b(t_r, x^r) + \sqrt{\delta}\, u^{r+1}$ with its corresponding directed-edge or vertex message. The directed-edge law fixes the common normaliser:

$$c_{D,r} = (\mathbb{E}a(t_r, x_{i\to j}^r)^2)^{-1/2}, \quad A_{i\to j}^r = c_{D,r} a(t_r, x_{i\to j}^r), \quad B_i^r = c_{D,r} a(t_r, x_i^r), \quad z_i^L = \sqrt{\delta} \sum_{r=1}^{L} B_i^{r-1} u_i^r.$$

The one-step lag in $A^{r-1}u^r$ gives conditional centring from the new outer Gaussian layer; successive finite-degree messages need not be independent. At fixed $\delta, L$, the joint vertex messages converge to the independent $U_r$ above, and $z_i^L$ converges to $\sqrt{\delta}\sum_{r=1}^{L} c_{\delta,r-1} a(t_{r-1}, X_{r-1}^\delta) U_r$. The source's separate edge calculation gives

$$\lim_{D\to\infty} \frac{D+1}{2\sqrt{D}} \mathbb{E}[z_o^L z_v^L] = \delta \sum_{r=1}^{L-1} c_{\delta,r} \mathbb{E}a(r\delta, X_r^\delta), \qquad o \sim v.$$

Only afterwards take $\delta \downarrow 0$ with $L = \lfloor T/\delta \rfloor$. The source bounds the discrepancy from $\int_0^T \mathbb{E}a(t, X_t)\, dt$ by $C(\beta_0, T)\sqrt{\delta}$. A separate norm-convergence argument with a uniform first occupation moment identifies this edge limit with the rooted energy. A finite-dimensional central limit theorem alone does not identify that observable. OpenAI (2026b), Appendix A.

**Source comparison for the finite weighted-tree formula.** This is an auxiliary check of the positive-temperature route. Readers continuing the circuit argument can go directly to §11.1.1 or §11.2. The comparison here is with El Alaoui–Montanari–Sellke, arXiv:2111.06813v2, 3 February 2023. Proposition 2.1 (p. 7) sums the adjacent-time term over $\ell = 2, \ldots, L$. Applying Lemma 4.3, equation (4.6) (p. 14), replaces the expectation in that summand by the coefficient at $r = \ell - 1$, so the resulting range is $r = 1, \ldots, L-1$. The displayed ranges in (4.5) (p. 13) and (4.18) (p. 16) are instead $2, \ldots, L$. Separately, the same-time display (2.21) in Lemma 2.8 (p. 9) omits a factor $1/k$ that is present in its proof and in Proposition 2.1. The QAOA manuscript's Appendix A already uses the latter proposition, the reindexed sum and spatial Lipschitz bounds for both $a$ and $b$. The two following calculations establish the finite discrepancies and their precise scope. They do not compare formulae in the journal edition.

**The test weights expose the missing factor.** Work on the $k$-regular tree, $k \geq 2$, and let all initial labels $g_i$ be independent standard Gaussians. In particular, the adjacent vertices $o, v$ have distinct labels $g_o, g_v$. Set $u_i^0 = u_{i\to j}^0 = g_i$, $A_{i\to j}^{-1} = 1$, and use constant algorithmic update functions $F_r = H_r = 1$. These satisfy EAMS Assumption 2 and its later boundedness Assumption 3. For the auxiliary tests in Lemma 2.8 choose $Y_o^r = g_o$ and $\widetilde{Y}_v^r = g_v$ for every $r \geq 0$. Each test

is measurable with respect to its required initial-label sigma-field, and all moments are finite. The tests are separate from the constant update functions.

At $t = 1$, the recursion gives

$$u_o^1 = k^{-1/2} \sum_{x \sim o} g_x, \qquad u_v^1 = k^{-1/2} \sum_{y \sim v} g_y.$$

Hence

$$\mathbb{E}[u_o^1 u_v^1 g_o g_v] = \frac{1}{k} \sum_{x \sim o,\, y \sim v} \mathbb{E}[g_x g_y g_o g_v] = \frac{1}{k}.$$

Only $x = v, y = o$ survives. The printed right side of (2.21) is $\mathbb{E}[g_v^2 g_o^2] = 1$; multiplying it by $1/k$ agrees with the calculation. Both initial labels matter. With both test weights equal to one, both sides are zero; with just the neighbouring test omitted, the left side is an odd Gaussian moment and is again zero.

A bounded version uses $Y_o^r = \operatorname{sgn}(g_o)$ and $\widetilde{Y}_v^r = \operatorname{sgn}(g_v)$. Only the same edge pair survives, so

$$\mathbb{E}[u_o^1 u_v^1 \operatorname{sgn}(g_o) \operatorname{sgn}(g_v)] = \frac{1}{k} (\mathbb{E}|g|)^2 = \frac{2}{\pi k},$$

whereas the printed right side is $2/\pi$. These tests are bounded and initial-label measurable. Both witnesses concern the auxiliary weighted identity; neither substitutes arbitrary test weights into a Parisi output theorem.

**The shifted endpoint violates the printed uniform finite bound.** For a separate test retain the initial labels and set

$$\delta = \varepsilon = \frac{1}{4}, \qquad L = 2, \qquad b(t, x) = 0, \qquad a(t, x) = \begin{cases} 1, & 0 \le t < 3/8, \\ -1, & 3/8 \le t < 1. \end{cases}$$

Use the source's actual initial states $x_i^0 = x_{i \to j}^0 = \sqrt{\delta}\, g_i$ and its definitions $A_{i \to j}^r = a(r\delta, x_{i \to j}^r)$, $B_i^r = a(r\delta, x_i^r)$. The state initialization has not been set to zero; it drops out of these coefficients because $a$ is independent of $x$. This fixed function is bounded by $K = 1$, has spatial Lipschitz constant zero, and satisfies $a^2 = 1$, so all directed-edge normalizers are exactly one. The drift has Lipschitz constant $M = 0$, and $L\delta = 1/2 \le 1 - \varepsilon$. It therefore meets the printed hypotheses of Theorem 4.1 and Assumptions 2–3. That finite theorem does not require temporal continuity or positivity of $a$.

At the mesh points, $a_0 = a_1 = 1$ and $a_2 = -1$. Thus

$$u_i^2 = \frac{1}{\sqrt{k(k-1)}} \sum_{w \sim i} \sum_{y \sim w,\, y \ne i} g_y, \qquad z_i^2 = \sqrt{\delta}\, (u_i^1 + u_i^2).$$

The equal-time covariances across the edge are zero. Each adjacent-time covariance is $\sqrt{k-1}/k$: exactly $k - 1$ labels occur in both sums, and their product normalization is $k\sqrt{k-1}$. Therefore

$$k\mathbb{E}[z_o^2 z_v^2] = 2\delta\sqrt{k-1}, \qquad \frac{k}{2\sqrt{k-1}} \mathbb{E}[z_o^2 z_v^2] = \delta.$$

The target printed in (4.5) uses $a(2\delta, x_o^2) = -1$, so it is $-2\delta\sqrt{k-1}$. Its absolute error is $4\delta\sqrt{k-1}$, which is unbounded as $k \to \infty$ while all arguments of the claimed constant $C(M, K, \delta, \varepsilon)$ remain

fixed. This refutes auxiliary Theorem 4.1 as printed. The example is not a Parisi curvature: it changes sign. Consequently it is not by itself a counterexample to the specialized assertion (4.18) in Proposition 4.7; the endpoint correction there comes from the exact proposition and reindexing explained above. Neither test refutes the main EAMS Theorem 1.7.

**A finite estimate with the required hypotheses.** The source argument does provide a usable replacement. Fix $\delta > 0$, $L \geq 1$ and $L\delta < 1$. Allow the coefficients $a_k, b_k$ to depend on degree, but require bounds independent of $k$ at the finitely many mesh times: $|a_k| \leq K$ with $K \geq 1$, and spatial Lipschitz constants $L_a, L_b$ for $a_k, b_k$. Use the original initialization and the same normalized function for node and directed-edge coefficients, with $\mathbb{E}(A^r_{i\to j})^2 = 1$. Then

$$\left| k\mathbb{E}[z_o^L z_v^L] - 2\sqrt{k-1}\,\delta \sum_{r=1}^{L-1} \mathbb{E}B_o^r \right| \leq C(K, L_a, L_b, \delta, L).$$

The constant is independent of $k$, and the sum is empty when $L = 1$. The coefficient expectation on the right uses the vertex law; its normalization was defined using the directed-edge law. The following proof keeps the two laws distinct. It reconstructs the finite estimate from EAMS Proposition 2.1 and Lemmas 3.2–3.4 and 4.2–4.3; it is not a claim that changing an index alone repairs every hypothesis of the printed theorem.

Put $D = k-1$. Conditional on the earlier cavity labels, the new message is centered Gaussian with variance $(\tau^r)^2$. Its recursion is

$$(\tau^{r+1}_{i\to j})^2 = \frac{1}{D} \sum_{\substack{w\in\partial i \\ w\neq j}} (A^{r-1}_{w\to i})^2 (\tau^r_{w\to i})^2, \qquad \tau^0 = 1.$$

This conditional Gaussianity gives $\Delta_r = \mathbb{E}[((\tau^r)^2 - 1)^2] \leq V_r/D$, where

$$V_0 = 0, \qquad V_{r+1} = 2K^4 V_r + 2(K^4 + 1).$$

Indeed, split the conditional-variance recursion into the average of $(A^{r-1})^2((\tau^r)^2 - 1)$ and the average of $(A^{r-1})^2 - 1$. Jensen bounds the first square, while independence of child cavities and exact coefficient normalization bound the second by a constant over $D$. The moment recursion also gives $\mathbb{E}(u^r_{i\to j})^2 \leq K^{2r}$.

Splitting off the missing neighbour as in the source gives $\mathbb{E}|u^r_i - u^r_{i\to j}|^2 \leq 4K^{2r}/k$. Subtract the node and directed-edge Euler recursions. Their equal initial states and the drift Lipschitz bound give

$$\mathbb{E}|x^r_i - x^r_{i\to j}|^2 \leq d_r/k, \quad d_0 = 0, \qquad d_{r+1} = 2(1 + \delta L_b)^2 d_r + 8\delta K^{2(r+1)}.$$

The bound $\|B^r_i - A^r_{i\to j}\|_2 \leq L_a\sqrt{d_r/k}$ now uses spatial Lipschitzness of $a$. This property is invoked in the proof of EAMS Lemma 4.2 but omitted from the printed hypotheses of Theorem 4.1; it is available in the QAOA manuscript's compact-strip application.

For $s = \ell - 1$, the adjacent-time summand of Proposition 2.1 is

$$Q_s = \mathbb{E}[A^{s-1}_{v\to o}(u^s_{v\to o})^2 B^{s-1}_v B^s_o].$$

Let $\mathcal{G}^{r,+}_{i\to j}$ be the labels within distance $r$ of $i$ in its cavity after removing edge $ij$. Condition on $\mathcal{G}^{s-1,+}_{v\to o} \vee \mathcal{G}^{s,+}_{o\to v}$. The three coefficient factors are measurable there: their vertex balls split into these two disjoint cavity balls. The newest layer driving $u^s_{v\to o}$ is outside this field; adding the initial label

at $v$ to the source's minus field does not change the conditional law, since that message and its variance exclude that label. Conditioning therefore replaces $(u^s_{v\to o})^2-1$ by $(\tau^s_{v\to o})^2-1$. Its contribution is bounded by $K^3\sqrt{V_s/D}$. Replacing the two node coefficients by directed-edge coefficients then costs at most $K^2L_a(\sqrt{d_{s-1}}+\sqrt{d_s})/\sqrt{k}$. The two directed-edge trees are independent; exact normalization reduces their product expectation to $\mathbb{E}A^s_{o\to v}$. One final comparison with $\mathbb{E}B^s_o$ proves

$$|Q_s-\mathbb{E}B^s_o|\le\frac{C_s}{\sqrt{D}},\quad C_s=K^3\sqrt{V_s}+K^2L_a\sqrt{d_{s-1}}+(K^2+1)L_a\sqrt{d_s}.$$

For the same-time remainder in that proposition, $|R_\ell|\le K^{2\ell+2}$ by boundedness and Cauchy–Schwarz. Hence the asserted finite bound holds, for example, with

$$C=2\delta\sum_{s=1}^{L-1}C_s+\delta\sum_{\ell=1}^{L}K^{2\ell+2}.$$

No constant uniform in increasing horizon, temperature or terminal cutoff has been asserted.

**Why the continuum value survives the correction.** The QAOA manuscript already checks the convergence and uniform boundedness of its finitely many degree-dependent normalizers, and supplies both spatial Lipschitz bounds, before taking the degree limit. Its limiting coefficient $m_r=c_{\delta,r}\mathbb{E}a(t_r,X^\delta_r)$ satisfies $|m_r|\le1$ by Cauchy–Schwarz and the definition of $c_{\delta,r}$. The shifted and corrected sums differ exactly by

$$\delta\sum_{r=2}^{L}m_r-\delta\sum_{r=1}^{L-1}m_r=\delta(m_L-m_1),\qquad|\delta(m_L-m_1)|\le2\delta.$$

Thus the error in the printed unnormalized finite theorem can grow like $\sqrt{k}$ at fixed mesh even though the correction disappears in the subsequent normalized continuum limit. The fixed-temperature coupling and separate strong realization in the QAOA manuscript justify that limiting application. These checks neither reject nor independently certify the whole EAMS or QAOA theorem. OpenAI (2026b), Appendix A.

After clipping and bounded approximation, the result enters the same synthesis and seed-removal route as the primary construction. Its source dependencies and fixed-temperature choice of parameters remain separate from the retained zero-temperature formal interfaces. OpenAI (2026b), Appendix A.

### 11.1.1 Where the temperature routes meet

Both routes produce a bounded finite classical calculation whose rooted energy is close to $P_*$. They reach that common input by different theorems and different finite approximations.

**Variational input and support.** The zero-temperature route fixes an attained minimizer $\gamma\in\mathcal{U}$ with terminal value $|x|$ and uses full relative Stieltjes support on $[0,1)$. The positive-temperature route fixes a finite $\beta_0>1$ and its Parisi probability measure, with terminal value $\beta_0^{-1}\log(2\cosh\beta_0x)$; it uses Lopatto's interval-support result on $[0,q_{\beta_0}]$.

**Scalar error.** Before discretization, the zero-temperature loss is at most $1-T$ for $T<1$. The positive-temperature loss is at most

$$(1/4+\log2)/\beta_0+(q_{\beta_0}-T),\qquad T<q_{\beta_0}.$$

**Finite energy.** In the zero-temperature route, a first-chaos projection identifies the explicit Gaussian coefficient sum with $\mathrm{Re}\langle F_N, \mathsf{L} F_N\rangle$. The alternative route first calculates finite-degree messages and edge energy, then uses a rooted norm and occupation argument.

**Order of approximation.** At zero temperature, fix $T$ before refining the scalar mesh, and choose a finite mesh before regularization and synthesis. At positive temperature, fix $\beta_0, T, \delta, L$ for the message degree limit, then refine $\delta$ at fixed $\beta_0, T$.

Both routes then use the same bounded approximation, exclusions, seeded synthesis, seed removal and fixed-word transfer. The quantum construction is shared; its classical input has been supplied independently.

Here the displayed positive-temperature estimate is exactly the one derived above, rather than an assertion of convergence of zero-temperature supports. One can choose $\beta_0$ and $T$ finitely for the required error and then stop. The later circuit argument needs that finite choice; it never asks for a joint limit of the temperature, message mesh, graph degree and circuit depth. OpenAI (2026b), Appendix A, Proposition "The independent positive-temperature construction"; §6.

### 11.1.2 Finite-temperature values and exact Gibbs samples

This comparison samples the exact Gibbs distribution. The QAOA Born law in §11.2 is a separate distribution; none of the following bounds identifies the two. For one fixed instance, let $L_n = \max_\sigma H_{n,J}(\sigma)/n$, $\mathcal{F}_{n,\beta} = (n\beta)^{-1} \log \sum_\sigma e^{\beta H_{n,J}(\sigma)}$, and let $\mathcal{U}_{n,\beta}$ be the mean energy per spin under the Gibbs law proportional to $e^{\beta H_{n,J}(\sigma)}$. Then

$$0 \le \mathcal{F}_{n,\beta} - L_n \le \frac{\log 2}{\beta}, \qquad 0 \le L_n - \mathcal{U}_{n,\beta} \le \frac{\log 2}{\beta}.$$

The first is the log-sum bound. The second uses $\mathcal{F} = \mathcal{U} + S/(n\beta)$, where the Gibbs entropy satisfies $0 \le S \le n \log 2$, together with $\mathcal{U} \le L_n \le \mathcal{F}$. Thus both disorder averages approach $P_*$ along every joint $n \to \infty$, $\beta \to \infty$, by the already established convergence of $\mathbb{E} L_n$.

There is also a finite probability estimate. Every configuration has Gaussian coefficient-vector norm $\sqrt{(n-1)/(2n^2)}$ after division by $n$, and their maximum has the same Lipschitz bound. Gaussian concentration gives, for $n \ge 2$,

$$\Pr_J(|L_n - \mathbb{E} L_n| > s) \le 2 \exp\{-n^2 s^2/(n-1)\}.$$

For an exact Gibbs draw $\Sigma^{\mathrm{G}}$, count the at most $2^n$ configurations with deficit at least $s$ and bound the partition sum below by its maximal term. This gives, for every instance,

$$\Pr\{H_{n,J}(\Sigma^{\mathrm{G}})/n \le L_n - s \mid J\} \le \min\{1, e^{n(\log 2 - \beta s)}\}.$$

The lower tail below $P_* - s$ is bounded by $\Pr_J(L_n < P_* - s/2) + \min\{1, e^{n(\log 2 - \beta s/2)}\}$; the upper tail is at most the upper tail of $L_n$. Hence an exact Gibbs sample also approaches $P_*$ in joint probability. These are elementary scalar consequences of the Gibbs law, entropy and Gaussian concentration. They supply neither an efficient sampler nor QAOA angles.

## 11.2 What is averaged, and a fixed-word probability consequence

The probability conclusion here uses an additional fixed-word second-moment theorem, beyond the expected-value construction of §§8.1–8.6. The theorem in §8 is an existence statement about limiting expected energy. There are two random choices behind that expectation. First the couplings

$J$ are sampled; then a computational-basis measurement of the state produces a spin string $\Sigma$. For the single word $W$ already chosen, put

$$E_n(J,\Sigma) = H_{n,J}(\Sigma)/n, \qquad \overline{E}_n(J) = \mathbb{E}[E_n(J,\Sigma) \mid J].$$

The quantity defining $v_W$ is $\mathbb{E}_J \overline{E}_n(J)$. It is neither the energy of every measured string nor a worst-disorder guarantee. The temporary Gaussian labels used during synthesis are not an extra source of randomness in the final angle vectors.

Farhi, Goldstone, Gutmann and Zhou supply that additional input. They prove, with the same Hamiltonian and gate-sign convention, that for every fixed word

$$\mathbb{E}_{J,\Sigma} E_n \longrightarrow v_W, \qquad \mathbb{E}_{J,\Sigma} E_n^2 \longrightarrow v_W^2.$$

Their second expression is the quantum expectation of $(C_{n,J}/n)^2$ averaged over disorder, not merely the square of the conditional mean. Consequently

$$\mathbb{E}_{J,\Sigma}(E_n - v_W)^2 \longrightarrow 0, \qquad \Pr_{J,\Sigma}(|E_n - v_W| > s) \longrightarrow 0 \quad (s > 0).$$

The law of total variance separates what vanishes:

$$\mathbb{E}(E_n - v_W)^2 = \mathbb{E}_J \operatorname{Var}(E_n \mid J) + \mathbb{E}_J(\overline{E}_n(J) - v_W)^2.$$

Both terms are nonnegative and hence tend to zero. Concentration of the conditional mean alone would not control a measured string. This is the elementary second-moment deduction from Farhi et al., arXiv:1910.08187v4, §4, equations (13)–(22).

Now run the circuit construction with accuracy $\varepsilon/2$ and freeze its resulting word. Since $v_W \geq P_* - \varepsilon/2$,

$$\Pr_{J,\Sigma}\{E_n < P_* - \varepsilon\} \leq \frac{4}{\varepsilon^2}\mathbb{E}_{J,\Sigma}(E_n - v_W)^2 \longrightarrow 0.$$

This is a consequence of the expected-value construction **together with** the external second-moment theorem. It is not a clause of either selected formal interface. If $b_n(J) = \Pr(E_n < P_* - \varepsilon \mid J)$, then for every fixed $\tau > 0$,

$$\Pr_J\{b_n(J) > \tau\} \leq \mathbb{E}_J b_n(J)/\tau \longrightarrow 0.$$

Thus the same word also has small conditional measurement-failure probability for most disorder instances, in this precise sense. No claim is made for every instance, for arbitrarily varying depths, or with a quantitative size threshold.

For $K$ fresh state preparations at the same disorder instance, the conditional failures are independent, so their joint failure probability is $\mathbb{E}_J b_n(J)^K$, not $(\mathbb{E}_J b_n(J))^K$. A fraction $\alpha$ of instances that always fail gives $\alpha$ for the former at every $K$. A bound $b_n(J) \leq \tau$ on a particular instance does give $\tau^K$ there. Repeatedly measuring an already collapsed state is a different experiment.

**Comparison with the same finite instance.** The concentration of the finite maximum in §11.1.2 strengthens this to a near-optimal sampled configuration. For every $\varepsilon > 0$, there is one finite deterministic word $W$, independent of $n$ and $J$, such that

$$\Pr_{J,\Sigma}\Big\{H_{n,J}(\Sigma) \geq \max_\sigma H_{n,J}(\sigma) - \varepsilon n\Big\} \longrightarrow 1.$$

To see this, choose the word with $v_W \geq P_* - \varepsilon/4$ and write $r_n = \mathbb{E}(E_n - v_W)^2 \to 0$ and $M_n = \max_\sigma H_{n,J}(\sigma)/n$. The convergence $\mathbb{E}M_n \to P_*$ makes its bias at most $\varepsilon/4$ for all sufficiently large $n$. The Gaussian coefficient-vector bound from §11.1.2 and Chebyshev's inequality for $E_n$ then give, for $n \geq 2$,

$$\Pr_J\{M_n > P_* + \varepsilon/2\} \leq 2\exp\left\{-\frac{n^2\varepsilon^2}{16(n-1)}\right\}, \qquad \Pr_{J,\Sigma}\{E_n < v_W - \varepsilon/4\} \leq \frac{16r_n}{\varepsilon^2}.$$

Outside these two events, $M_n - E_n \leq \varepsilon$. Their union therefore has probability tending to zero. The maximum and measured energy use the same disorder; the union bound needs no independence. One fresh preparation and measurement suffices for this asymptotic statement. Its inputs are the ordinary-word theorem, the external fixed-word second-moment theorem and Gaussian concentration; it is not a clause of the selected formal endpoints. The unspecified sequence $r_n$ and mean-maximum bias still give no effective size threshold or efficient angle-selection procedure.

### 11.3 Depth, angle choice and the order of quantifiers

The complete witness is a finite ordered word, including every real angle. Its existence for each error tolerance asserts uniformity in system size and disorder that fixed-instance convergence does not provide. The quantifiers should be read literally.

The source theorem says that for every $\varepsilon > 0$, there is one finite word $W_\varepsilon$ such that

$$\lim_{n\to\infty} \mathbb{E}E_n(W_\varepsilon) \geq P_* - \varepsilon.$$

The cost operator uses $J, n$, but this word and its angles do not.

In fixed-instance QAOA convergence, by contrast, the chosen depth and angles may depend on the fixed finite cost matrix and the requested accuracy. The performance of a prescribed family of nonzero angles with $p = p(n) \to \infty$ needs further estimates. A depth allowance alone can be met by the padded existence construction below.

The distinction already occurs for the elementary array $a_{n,p} = p/(n+p)$. For each fixed $n$, its limit as $p \to \infty$ is one; for each fixed $p$, its limit as $n \to \infty$ is zero. This is a quantifier example, not a model of QAOA. The finite-dimensional convergence theorem of Binkowski, Koßmann, Ziegler and Schwonnek, arXiv:2302.04968v1, Theorem 12 fixes the optimization problem before selecting a sufficiently long circuit. It does not replace the construction of a size-independent word here.

**An existential schedule under a prescribed depth allowance.**

There is a narrower diagonal consequence for sampled output. Choose $W_k$ with $P_* - 1/k \leq v_{W_k} \leq P_*$. Fixed-word second-moment convergence supplies $N_k$ so that, for every $n \geq N_k$, $\mathbb{E}(E_n(W_k) - v_{W_k})^2 \leq k^{-2}$. Given any prescribed integer depth allowance $d(n) \to \infty$, enlarge these strictly increasing thresholds until $d(n) \geq \mathrm{depth}(W_k)$ for every $n \geq N_k$. On $N_k \leq n < N_{k+1}$, use $W_k$ padded with zero-angle layers to depth $d(n)$. Padding changes no state, and

$$\mathbb{E}(E_n(W_k) - P_*)^2 \leq 2\mathbb{E}(E_n(W_k) - v_{W_k})^2 + 2(v_{W_k} - P_*)^2 \leq 4/k^2.$$

Thus a suitable schedule approaches the optimum in joint $L^2$, even under an arbitrarily slowly diverging allowance. The word family and thresholds are selected nonconstructively. This says nothing about an independently prescribed family of nonzero angles, and is not an interchange of limits or an effective depth bound.

**Computational meaning of a finite word.**

Nor does a supremum over angles assert that an optimizing angle vector exists at any particular depth. The proof constructs a sufficiently good word directly; monotonicity from identity padding then gives the limit of $Q_p$. Each QAOA round contains a full cost evolution and a mixer evolution. In the dense SK model a cost evolution involves all $\binom{n}{2}$ pair couplings, so the round count is not itself a bound on two-qubit gate depth, connectivity cost, precision or physical runtime.

Three computational questions remain distinct. Evaluating the limiting energy at supplied angles is one task; finding suitable angles is another; implementing and sampling the resulting quantum circuit is a third. The Basso et al. iteration evaluates the fixed-angle limit in $O(p^2 4^p)$ arithmetic time. That is not a bound on the search needed to find the existence theorem's angles. Spin–boson representations provide another route to evaluation. The earlier Boulebnane et al. mapping, arXiv:2505.07929v2, and the later mixed-spin decoupling paper, arXiv:2609.40309v1, §§II–III and Supplement S2.1, must be distinguished: the latter is a separate work, not the reference called BKL2026 in the original QAOA manuscript. Its fixed-depth mapping and numerical tensor-network performance do not supply an efficient angle-selection theorem for the present construction. Its discussion also gives fixed-word finite-size estimates; the absence of a finite-size rate **in the construction here** is not a claim that the wider literature contains none.

**Why the two-spin hypothesis matters.**

The statement is specific to the quadratic, zero-field SK model. The pure even higher-spin model has a different thermodynamic-first conclusion. For every even interaction order $q \geq 4$, Basso, Gamarnik, Mei and Zhou, arXiv:2204.10306v2, Theorem 4, (3.15), and its proof in §4.2 show

$$v_p^{(q)}(\theta, \beta) \leq \eta_{\mathrm{OGP}}(G_q) < \eta_{\mathrm{OPT}}(G_q)$$

for every finite $p$ and every fixed angle vector, in that source's normalization. The same ceiling bounds their supremum over all finite $p$ and angles. The proof combines its dense–sparse value identity with the coupled-overlap-gap obstruction of Chou, Love, Sandhu and Shi, arXiv:2108.06049v3, Parameter 4.1 and Corollary 4.4. That local-algorithm obstruction applies to sparse random signed Max-$q$-XOR for even $q \geq 4$, at sufficiently large fixed degree and under the source's logarithmic-depth bound. The Basso–Gamarnik–Mei–Zhou transfer above only needs the resulting obstruction at each fixed depth. It requires no interchange of depth and system size. The exclusion $q \geq 4$ is essential; this is not an obstruction for the SK two-spin model treated here.

For comparison, Goh, arXiv:2404.06087v6, Theorem 4.2 and Corollary 4.2.1, treats even interaction order at least four. Its logarithmic-depth obstruction concerns sparse random-regular Max-q-XORSAT; the dense spin-glass corollary is explicitly conditional on swapping limits. The source states on p. 12 that its sparse/dense equivalence is only established at fixed depth, so it does not prove a dense logarithmic-depth obstruction. These hypotheses do not include quadratic SK. Neither a conclusion for all mixed spin glasses nor a general quantum advantage follows from §8.

### 11.4 The regular-MaxCut consequence has its own limits

The original manuscript also gives a consequence for MaxCut. For each $\eta > 0$ it selects one finite depth and normalized angles, and then a degree threshold, such that every finite simple unweighted $(D+1)$-regular graph of girth greater than $2p+1$ has expected cut fraction at least

$$\frac{1}{2} + \frac{P_* - \eta}{\sqrt{D}}$$

for all sufficiently large $D$. The usual MaxCut cost is $H_{\mathrm{MC}} = \frac{1}{2}\sum_{\{i,j\}}(I - Z_iZ_j)$, so its angles are $2\gamma/\sqrt{D}$: the normalized angles are fixed, while these physical cost angles depend on degree. The sign conversion uses a bipartition of the relevant tree neighbourhood of each edge; the whole graph need not be bipartite.

The further optimality comparison is for a uniform simple labelled $(D+1)$-regular graph $G_{n,D+1}$, along sizes with $n(D+1)$ even. Write $\mathrm{OPT}(G)$ for the maximum cut size divided by $|E(G)|$, $q_{p,G}$ for the circuit's expected cut fraction, and $\Delta_D(G) = \mathrm{OPT}(G) - q_{p,G}(2\gamma/\sqrt{D}, \beta)$. The same word can be chosen so that

$$\lim_{D\to\infty} \limsup_{\substack{n\to\infty \\ n(D+1)\ \text{even}}} \Pr_G\{\sqrt{D}\,\Delta_D(G_{n,D+1}) > \eta\} = 0.$$

Size goes to infinity first at fixed degree and word. The probability is over the random graph; the cut fraction has already averaged the quantum measurement. For fixed $D, p$, McKay, Wormald and Wysocka (2004), Theorem 1, gives tight counts of cycles of lengths at most $2p+3$. Every edge with a nontree neighbourhood is within bounded distance of one of these cycles, so their fraction tends to zero. The optimum is supplied separately by Dembo, Montanari and Sen, arXiv:1503.03923v2, Theorem 1.6(b): dividing their cut size per vertex by $(D+1)/2$ gives the leading cut fraction $1/2 + P_*/\sqrt{D+1}$, with an error $o(D^{-1/2})$ in the stated iterated regime. A high-girth lower bound alone would not prove optimality on every high-girth graph. See OpenAI (2026b), Corollary 1.2 and its complete proof. This separate consequence is recorded to preserve the source's scope; it is not a new selected formal endpoint or a joint size–degree limit.

### 11.5 A later finite-size estimate and its limited depth consequence

A separate, later source gives a quantitative comparison with the fixed-angle limit. Boulebnane, Khan, Subrahmanya, Herman, Farhi, Villalonga and Shaydulin, arXiv:2609.40309v1, Corollary S3.1, pp. 46–47, use the ordered-tuple Gaussian mixed-spin cost with covariance polynomial $\xi(t) = \sum_q c_q^2 t^q$ of finite degree. Let $\Gamma > 0$ bound every cost angle in a depth-$p$ word. Choose $x > 0$ bounding $\xi(2), \xi'(1), \xi''(33)$, with $\Gamma^2 p^2 x \geq 1$. No bound on the mixer angles is needed. Set

$$B_p = 256p^4(2e\Gamma^2p^2x)^p, \qquad E_p = 70\Gamma^2p^2x + p\log 2, \qquad R_p = (2E_p/\log 2)^{2p}.$$

For

$$0 < \varepsilon \leq 4\Gamma px \min\{e^{-1/2}, 2^{-(8p^2+3)/100}\},$$

the source asserts $|\nu_n - \nu_\infty| \leq \varepsilon$ whenever

$$n \geq T_p := \max\left\{2^{8p^2+3}B_p\left(\frac{2^{\deg\xi+4}\Gamma^2p^2x^2}{\varepsilon^2}\right)^{R_p}\left[2\log_2\frac{4\Gamma px}{\varepsilon} + \deg\xi\right]^2, \frac{16\Gamma px}{\varepsilon}\right\}.$$

Here $\nu_n$ is the disorder-averaged energy per spin. The estimate applies to each near-optimal word from §8.6 separately. Given $\zeta > 0$, choose a finite word $W_\zeta$ with $\nu_\infty(W_\zeta) \geq P_* - \zeta/2$. Its depth $p_\zeta$ and cost angles are finite; pad by a zero layer if needed and take $\Gamma_\zeta = \max\{1, \max_j|\theta_j(W_\zeta)|\}$. Choose a positive $\varepsilon_\zeta \leq \zeta/2$ satisfying the displayed small-error condition. With these fixed parameters, the threshold gives

$$\nu_n(W_\zeta) \geq P_* - \zeta \qquad (n \geq T_{p_\zeta}).$$

The threshold is explicit in the chosen word and error. The existential construction supplies no quantitative bounds for $p_\zeta$ or $\Gamma_\zeta$ as $\zeta \downarrow 0$, so it does not yield a bound in accuracy alone. A common

angle bound is needed for the uniform growing-depth consequence below, not for this fixed-word application.

For SK take $c_2 = 1/\sqrt{2}$. Combining the two ordered off-diagonal couplings recovers our standard Gaussian $J_{ij}$; the diagonal adds an independent scalar Gaussian of variance $1/2$. That scalar changes the circuit by a global phase and has zero mean, so the expected-energy comparison uses the same normalization. A fixed $x \geq \max\{2, \Gamma^{-2}\}$ satisfies the above bounds for every $p \geq 1$.

For fixed mixture, $\Gamma$ and $x$, put $\varepsilon_p = e^{-p^2}$. The small-error condition holds eventually, and direct logarithms give

$$\log R_p = 4p \log p + O(p), \qquad \log \log T_p = 4p \log p + O(p + \log p).$$

Consequently, for any deterministic sequence of words $W_n$ with depths $p(n) \to \infty$, one common cost-angle bound $\Gamma$, and $p(n) \log p(n) = o(\log \log n)$, the displayed hypotheses hold and $|\nu_n(W_n) - \nu_\infty(W_n)| \leq e^{-p(n)^2}$ eventually. This is a uniform comparison along that specified bounded-angle class. It gives neither an efficient way to find near-optimal angles nor concentration of a quantum measurement in this growing-depth regime. The fixed-word second-moment theorem in §11.2 has separate quantifiers.

## 12. What the formal statements establish

### 12.1 The support statement and the derivative bridge

The first formal statement assumes a Brownian system and an admissible minimizer. It supplies a Stieltjes measure, an associated diffusion, full relative support and strict increase, and the two moment identities for every associated diffusion. Its proof assembles contact, exclusion of a terminal plateau, curvature for positive and zero constant intervals, and uniqueness to transfer the moments between associated diffusions. This describes the inspected proof's mathematical route; no successful research-target replay is claimed. Its quantifiers are $\forall W\, \forall \gamma$, followed by the minimizer hypothesis; it supplies both witnesses on the given Brownian system. The moment conclusion is universal over every diffusion satisfying the specified equation, and covers exactly $0 \leq t < 1$. The diffusion equation itself holds through time one, simultaneously for all those times outside one null set. Controls are progressive for the natural filtration of Brownian increments and bounded by one pointwise. These are not additional regularity or minimizer-existence assumptions. See the exact support statement and the companion source-interface review.

There is one useful distinction between the manuscript's first variation and its formal counterpart. The formal result initially gives a derivative along chord lengths $1/(n+1)$. For this particular control functional, that sampled limit determines the ordinary feasible right derivative. With a control fixed, the terminal absolute value is convex in the coefficient while the penalty is linear. The supremum over the same control class, and then subtraction of the linear Parisi penalty, preserve convexity. The secant slopes therefore vary monotonically as the chord shortens; each intervening chord lies between two successive sampled chords and has the same limiting slope. If $\eta$ is an integrable feasible direction, choose $\varepsilon_0 > 0$ so that $\gamma + \varepsilon_0 \eta$ is admissible and rescale the chord. This bridge is an ordinary deduction from the definitions and sampled limit, not an additional formally named theorem.

At the origin, the manuscript smoothly extends auxiliary quotient variables. The formal coercivity proof for the spatial diffusion density instead works on the positive half-line, controls the zero flux at the origin and reflects. What must agree is the resulting integrated inequality; auxiliary values

at zero need not coincide. In particular, the declared generic flux lemma has a global derivative hypothesis that is not the supplier used here: the finite-density proof uses the Gaussian half-line coercivity lemma and reflection. A totalized quotient value at zero cannot silently satisfy the generic global hypothesis. The precise source spans are in the correspondence record. The fixed-time supplier `actual_coercivity` assumes positive time and continuity of the coefficient at that time. Its scaled polynomial inequality is (4.1) multiplied by $c^2$; division requires $c > 0$. This gives a direct source destination for Proposition 4.1 without adding a minimizing or plateau premise.

### 12.2 The separate scalar statement

The second formal statement has three clauses. Given a minimizer and an associated diffusion, it establishes convergence of the finite-spin ground-state values, a bounded martingale with a spin-valued terminal limit, convergence of the gradient and the value identities. Under the same hypotheses and fixed $0 < T < 1$, it gives eventual positivity of the Gaussian normalizers and convergence of the finite coefficient sum. These are the declarations `scalar_value_target` and `finite_gaussian_target`. Their caller supplies the same minimizer and associated diffusion. The implementation derives compact-strip bounds from `sourceStripRegularity` and the unit curvature moment from contact for those objects before applying its fixed-strip helper. Those internal inputs are not additional hypotheses of the exported clauses. The convergence clause establishes existence of the finite-spin limit.

The index match in the Gaussian statement is exact. For $\Delta = T/N$ and $t_j = j\Delta$, the Euler path uses $Z_1, \dots, Z_N$ and produces $Y_0, \dots, Y_N$. The source normalizer indexed from zero corresponds to manuscript $c_1, \dots, c_N$; its eventual positivity covers precisely those normalizers. A separate $Z_{N+1}$ performs mean-preserving rounding: it enters $B_{N+1}$, not another Euler increment. Thus the finite sum is $\sum_{j=0}^{N} A_j B_{j+1}$ with $A_0 = 0$, using exactly the displayed index ranges. Total definitions at $N = 0$ make no extra mathematical claim about the construction for $N \geq 1$. More explicitly, the zero-based formal quantities `coordinate N j`, `rawCoefficient ... j` and `normalizer ... j` correspond to $Z_{j+1}$, $a(t_j, Y_j)$ and $c_{j+1}$ for $0 \leq j < N$. The positivity quantifiers are $\exists N_0\, \forall N \geq N_0\, \forall j < N$, simultaneously over the growing mesh, for each already fixed $T$. The extra coordinate with formal index $N$ is $Z_{N+1}$. The formal martingale stops at its terminal random variable $U$ for all later times; $U^2 = 1$ almost surely, and the gradient converges to $U$ almost surely and in $L^2$ from below one. The internal terminal-gradient lemma proves $UX_1 = |X_1|$ without requiring $X_1$ to have no atom at zero. That identity is used inside the implementation; it is not an extra conjunct of the endpoint. See the scalar statement and its complete dictionary.

The third declaration, `ground_state_parisi_target`, assumes a minimizer alone and concludes $P_* = \mathcal{P}(\gamma)$. Its proof chooses $X =$ `diffusionProcess` $W\ \gamma$ and proves the diffusion predicate for that process before applying the equality with a supplied diffusion. No diffusion is required from the caller, and none of these clauses constructs a minimizer. The exact declarations are MainConclusions, lines 13–27; their analytic suppliers and applications are in ContactAssembly. The independent contact and lower-bound arguments are reconstructed in §10.1 and §10.2.

### 12.3 Relating the two statements

The support and scalar models encode closely related control problems. A direct formal application still needs a specified map of their objects. The following contract identifies the obligations, rather than asserting an equivalence of their complete interfaces.

**Coefficients.** Restrict the scalar real-time coefficient to $[0, 1)$; conversely, choose an extension

compatible with the scalar right-continuity predicate. Prove equality almost everywhere of the coefficients used by each payoff and penalty. The scalar endpoint and exterior values are unconstrained by the shape conditions, so two extensions need not be literally equal.

**Brownian space.** Bundle or unbundle the same sample space, sigma algebra, probability law and Brownian process. Identify the natural filtration and every elapsed-increment filtration. Equality in law alone does not identify admissible controls on a chosen space.

**Controls and payoffs.** Bundle an admissible scalar control as a support control and conversely. With $s = t + r$, prove equality between the absolute integral over $[t, 1]$ using $\alpha(s - t)$ and the elapsed integral over $[0, 1 - t]$ using $\alpha(r)$. Preserve progressive measurability and the pointwise bound $|\alpha| \leq 1$.

**Value and minimizer.** Equality of the payoff sets gives equality of their suprema for every $x$ and $0 \leq t < 1$. Only then identify the first and second spatial derivatives and the Parisi penalties. Transport minimality against every admissible competitor, including restriction or extension of that competitor.

**Diffusion and moments.** Reindex the same process between nonnegative and real time, and stop it at one where required. Transport the integral equation on one full-measure event for all $t \in [0, 1]$, path continuity, initial value, gradient and moment integrals.

The diffusion comparison has a version issue. The support predicate asks for a globally progressive process; the scalar predicate asks for adaptedness of the stopped process and almost-sure continuity on $[0, 1]$. There is a concrete ordinary construction of the needed progressive version. For $t \geq 0$ and $k = 0, 1, ...,$ put

$$X_t^{(k)} = X_{\min(2^{-k}\lfloor 2^k t \rfloor, 1)}.$$

Each left-step process is progressive: on every bounded time interval, its value on a step is measurable at that step's left endpoint. Define $\bar{X}_t$ as the pointwise limit where the sequence converges, and zero elsewhere. The convergence set and this limit are progressively measurable. On the single event where $X$ is continuous on $[0, 1]$, the equality $\bar{X}_t = X_{\min(t,1)}$ holds for every $t \geq 0$. Thus stopping and reindexing preserve paths, time integrals and moments on that event; no completion of the natural filtration is needed merely to construct this version. The initial value follows from the scalar equation at zero and $B_0 = 0$ on their common full-measure event.

This settles the version construction at the ordinary mathematical level. Its implementation in the two formal predicates, and the coefficient, filtration and value identifications listed above, remain separate obligations. In particular, equality of values for every $x$ is needed before transporting their derivatives; agreement at $(0, 0)$ is insufficient.

Once those obligations are proved, an application must use the same transported minimizer and associated diffusion in all premises. No such cross-model transport theorem is supplied here. The manuscript proof in §§3–7 and the scalar implementation's independent argument remain available without it. These distinctions come from the defining support model and scalar model, rather than from their endpoint names.

### 12.4 Logical limits of the argument

The full-support theorem and scalar-value theorem use related but different formal representations. Their mathematical conclusions can be discussed together, but a direct formal implication would require the coefficient, Brownian law, control value and diffusion translations described in

§12, with common witnesses. The finite Gaussian result also needs its rooted energy identification before it says anything about a QAOA circuit. The boundedness, regularization, synthesis and transfer estimates in §8 are additional steps, with temperature-specific hypotheses where the positive-temperature route is used.

Infinite support allows gaps. An indefinite curvature identity does not imply strict convexity before expectation is controlled. An algebraic certificate does not establish its analytic hypotheses. Coordinatewise convergence does not control a growing coefficient sum. A finite-dimensional Gaussian law does not identify an energy observable without its operator argument. Separate witnesses cannot be merged silently. The order $\forall \varepsilon\ \exists$finite word $\lim_{n\to\infty}$ cannot be replaced by a fixed-instance depth limit.

The local comparison in §11.1 concerns two printed finite-step displays; it neither replaces the cited theorem's hypotheses nor supplies the remaining circuit estimates.

## 13. Source checks and finite calculation

The source collection retains the two manuscripts, formal statements, their implementations and build metadata. The derived project specifies Lean 4.34.1 and a pinned mathlib revision. The proof-assistant and library infrastructure is described by de Moura and Ullrich (2021), *The Lean 4 Theorem Prover and Programming Language* and The mathlib Community (2020), *The Lean Mathematical Library*. These references credit the software; the retained project metadata identifies the versions used for this source. The retained polynomial computation in §4 uses exact rational arithmetic. Its expansion matches all 22 positive remainder coefficients. The surplus of the $v^4$ coefficient above $1/1000$ is $140507/1521000000$. This finite check does not establish the PDE signs, integrability, boundary flux or limiting argument.

The source-byte audit verified the inspected edition; it did not prove its claims. Paper assembly checked identifiers, declared scope and document bindings, not semantic correspondence or reader understanding. Tool-compatibility checks and synthetic fixtures exercised their stated cases, not either selected SK theorem. Reading of the mathematical and formal sources supports the bounded comparisons reported here, without a complete independent proof acceptance or transitive axiom audit.

The retained observations must be kept separate. The original graph records unrun plans and an earlier full-support attempt that could not start because the Comparator executable was unavailable. A later retained attempt, observed on 23 October 2026, built and exported the challenge, then stopped while building the implementing solution. The retained partial-observation receipt and log preserve that event: the imported object for `Diffusion/AbsIncrement` was missing. The challenge is an admitted specification, so its successful build and export are not a proof of the theorem. There was no successful solution export or research-target comparison in that observation.

The scalar-value invocation in the same original two-target run reached a terminal failure observed on 24 October 2026. Its challenge built and exported, but its solution build stopped because the imported object for `Gaussian/IntegralPositiveMulGaussian` was missing. The retained scalar receipt and log establish no solution export, kernel acceptance or comparison acceptance. The companion source review locates both records. Both failures concern the original build configuration. A subsequent repair to the derived configuration was recorded. In the subsequent corrected replay, the full-support challenge built and its exporter was observed running. Reconciliation on 25 October found that the controller had stopped without a recovered terminal research result. At that date,

Comparator and research extraction outcomes were unknown, neither rejected nor accepted. The interruption receipt retains this observation separately from the original build failures.

A subsequent source-bound observation of 26 October 2026 records a successful local Lake build of the Support implementation and v2 extraction of the single conditional full-support root. Its selected traversal contains 1,674 declarations. The root reports no sorry, unsafe or partial flag, and its recorded transitive axiom names are `propext`, `Classical.choice` and `Quot.sound`. The selected in-scope traversal completed, but dependencies outside it remain unresolved boundaries; the extracted graph is partial. This does not establish minimiser existence, positive density, the scalar target or QAOA convergence.

The portable observation identifies the original receipts by their hashes and retains the source inventory, root and extraction limits. Its internal-consistency checks are not authenticated execution provenance. The curated paper graph records the build and documentary extraction observation separately; it is not the extracted declaration graph. Comparator acceptance, standalone kernel replay and the scalar-target results remain pending. The earlier failed and interrupted invocations remain dated history, rather than evidence of a later rejection or acceptance. Permitted axioms and checker settings describe the requested comparison; they cannot supply its missing result.

This editorial revision does not establish a new formal result. The source review through 26 October 2026 records the preceding paper revision's reading coverage, finite checks, unsuccessful observations and remaining obligations; viewing it requires repository access. This remains a draft after source-based agent review; independent mathematical acceptance and a human reconstruction study have not been performed.

## References


- Auffinger and Chen (2015), "On properties of Parisi measures," *Probability Theory and Related Fields* **161**, 817–850. Inspected preprint: arXiv:1303.3573v1.
- Farhi, Goldstone and Gutmann (2014), *A Quantum Approximate Optimization Algorithm*, arXiv:1411.4028v1.
- Binkowski, Koßmann, Ziegler and Schwonnek (2023), *Elementary Proof of QAOA Convergence*, arXiv:2302.04968v1.
- Basso, Gamarnik, Mei and Zhou (2022), *Performance and limitations of the QAOA at constant levels on large sparse hypergraphs and spin glass models*, arXiv:2204.10306v2.
- Boulebnane, Khan, Liu, Larson, Herman, Shaydulin and Pistoia (2026), *Spin–Boson Mapping of the Quantum Approximate Optimization Algorithm*, arXiv:2505.07929v2.
- Boulebnane, Khan, Subrahmanya, Herman, Farhi, Villalonga and Shaydulin (2026), *Decoupling of the QAOA into independent spin-boson systems and high-depth performance on pure and mixed spin glasses*, arXiv 2609.40309, version 1.
- Goh (2025), *The Overlap Gap Property limits limit swapping in the QAOA*, preprint, version 6.
- OpenAI (2026a), *Full support of the zero-temperature Sherrington–Kirkpatrick order parameter*, manuscript, 27 September 2026.

- OpenAI (2026b), *QAOA attains the SK ground-state energy in the thermodynamic-first limit*, manuscript, 25 September 2026.
- Auffinger and Chen (2017), "Parisi formula for the ground state energy in the mixed $p$-spin model," *Annals of Probability* **45**, 4617–4631.
- Auffinger and Chen (2015), "The Parisi formula has a unique minimizer," *Communications in Mathematical Physics* **335**, 1429–1444.
- Auffinger, Chen and Zeng (2020), "The SK model is infinite step replica symmetry breaking at zero temperature," *Communications on Pure and Applied Mathematics* **73**, 921–943.
- Jagannath and Tobasco (2016), "A dynamic programming approach to the Parisi functional," *Proceedings of the American Mathematical Society* **144**, 3135–3150.
- Chen, Handschy and Lerman (2018), "On the energy landscape of the mixed even $p$-spin model," *Probability Theory and Related Fields* **171**, 53–95.
- Montanari (2025), "Optimization of the Sherrington–Kirkpatrick Hamiltonian," *SIAM Journal on Computing* **54**, FOCS19-1–FOCS19-38.
- El Alaoui, Montanari and Sellke (2021), "Optimization of mean-field spin glasses," *Annals of Probability* **49**, 2922–2960.
- Farhi, Goldstone, Gutmann and Zhou (2022), "The Quantum Approximate Optimization Algorithm and the Sherrington–Kirkpatrick Model at Infinite Size," *Quantum* **6**, 759.
- Basso, Farhi, Marwaha, Villalonga and Zhou (2022), "The Quantum Approximate Optimization Algorithm at High Depth for MaxCut on Large-Girth Regular Graphs and the Sherrington–Kirkpatrick Model," *LIPIcs* **232**, 7:1–7:21.
- El Alaoui, Montanari and Sellke (2023), "Local algorithms for Maximum Cut and Minimum Bisection on locally treelike regular graphs of large degree," *Random Structures and Algorithms* **63**, 689–715.
- Chen (2026), *FRSB in the SK spin glass: convergence to full-interval support at zero temperature*, arXiv:2607.18032v1.
- Lopatto (2026), *Full replica symmetry breaking in the Sherrington–Kirkpatrick model*, arXiv:2607.11756v3.
- Zhou (2026), "Existence of full replica symmetry breaking for the Sherrington–Kirkpatrick model at low temperature," *Communications on Pure and Applied Mathematics* **79**, 1746–1770.
- Sellke (2021), *Optimizing Mean Field Spin Glasses with External Field*, arXiv:2105.03506v4.
- Sellke (2024), *Optimizing Mean Field Spin Glasses with External Field*, *Electronic Journal of Probability* **29**, paper 4. Inspected revision: arXiv:2105.03506v6, 17 December 2023. Lemma 2.9 and Remark 2.10 are distinguished from the preceding v4 entry.
- Jekel, Sandhu and Shi (2026), *Potential Hessian Ascent: The Sherrington–Kirkpatrick Model*, arXiv:2408.02360v3.
- Chou, Love, Sandhu and Shi (2022), *Limitations of Local Quantum Algorithms on Random Max-k-XOR and Beyond*, arXiv:2108.06049v3.

- Lopatto (2026), *Full replica symmetry breaking in the Sherrington–Kirkpatrick model*, arXiv:2607.11756v4. This later comparison is distinct from the v3 input cited above.
- Lopatto (2026), *Parisi measure of the Sherrington–Kirkpatrick model with external field*, arXiv:2610.00127v1.
- Jones and Knill (1999), "Efficient Refocussing of One Spin and Two Spin Interactions for NMR Quantum Computation," *Journal of Magnetic Resonance* **141**, 322–325. Inspected preprint: arXiv:quant-ph/9905008v1.
- Li and Khaneja (2005), *Noncommuting vector fields, polynomial approximations and control of inhomogeneous quantum ensembles*, arXiv:quant-ph/0510012v1.
- Talagrand (2006), "The Parisi formula," *Annals of Mathematics* **163**, 221–263. Published article.
- Dembo, Montanari and Sen (2017), "Extremal Cuts of Sparse Random Graphs," *Annals of Probability* **45**, 1190–1217. Inspected preprint: arXiv:1503.03923v2.
- McKay, Wormald and Wysocka (2004), "Short Cycles in Random Regular Graphs," *Electronic Journal of Combinatorics* **11**, R66. Published article.
- de Moura and Ullrich (2021), "The Lean 4 Theorem Prover and Programming Language," *CADE 28*, 625–635. Published article.
- The mathlib Community (2020), "The Lean Mathematical Library," *CPP 2020*, 367–381. Published article.