

Reciprocal Sums and the Sylvester Recurrence: Further Results and Proofs

Erdős Problem #243

WILL COOK*

September 2026

ABSTRACT

We study rational reciprocal sums with $a_{n+1}/a_n^2 \rightarrow 1$. Integer finite differences exclude nonintegral regular rates and reduce the cubic rate to a polynomial. A square condition in the cubic field and congruences modulo seven then prove irrationality. For general rational tails, we derive Sylvester recurrence criteria from persistent divisors, bounded numerator increases and weighted crossings of new maxima. We describe the arithmetic estimates still needed for Erdős Problem #243.

1 Introduction

Sylvester's sequence 2, 3, 7, 43, 1807, ... has reciprocal sum 1. Its recurrence $a_n = a_{n-1}^2 - a_{n-1} + 1$ gives the telescoping identity

$$\frac{1}{a_{n-1} - 1} = \frac{1}{a_{n-1}} + \frac{1}{a_n - 1}.$$

The tail beginning with $1/a_{n-1}$ therefore equals $1/(a_{n-1} - 1)$. Erdős Problem #243 asks whether every rational reciprocal sum with $a_n \sim a_{n-1}^2$ must eventually have this form:

Problem 1.1 (Erdős #243). Let $1 \leq a_1 < a_2 < \dots$ be a sequence of integers with

$$\lim_{n \rightarrow \infty} \frac{a_n}{a_{n-1}^2} = 1 \quad \text{and} \quad \sum \frac{1}{a_n} \in \mathbb{Q}.$$

Then $a_n = a_{n-1}^2 - a_{n-1} + 1$ for all sufficiently large n .

See [5, p. 64] and [6, p. 105]. Bloom's catalogue, accessed 28 July 2026, lists the problem as open [13]. Below we index the same recurrence as $a_{n+1} = a_n^2 - a_n + 1$, which is the shift the formal sources use; the two forms are the same statement. The polynomial $a^2 - a + 1$ is called the [Sylvester successor](#).

Clearing a rational reciprocal tail gives positive integers C_n, D_n with $C_{n+1} = a_n C_n - D_n$ and $D_{n+1} = a_n D_n$. The first update can also be written $C_{n+1} = C_n - E_n$, where

**Authorship and AI use.* Will Cook built and directed the research infrastructure and reviewed the claims when he could. AI agents did most of the research and drafting. Cook did not independently verify every claim.

$E_n = D_n - (a_n - 1)C_n$. Quadratic growth makes $E_n/C_n \rightarrow 0$. The eventual Sylvester recurrence follows once the integer error itself vanishes.

The two principal arguments control different features of this integer numerator. A precise regular rate controls its finite differences: although C_n grows, a sufficiently high difference tends to zero and therefore vanishes. The resulting polynomial must still satisfy the exact denominator recurrence, which imposes congruences on its values. A bound on upward steps gives a different use of integrality. After a common divisor stabilises, every earlier multiplier is coprime to every later numerator. The Chinese remainder theorem then produces a block that an unbounded numerator with bounded upward steps cannot cross. When common factors keep being cancelled, the later criteria instead estimate the prime powers that survive.

We prove the cubic and nonintegral-rate theorems first (Section 2). Their rate assumptions exclude the eventual Sylvester recurrence, so these results settle restricted cases by ruling out rationality. The general recurrence question remains open here. Section 3 compares the results with earlier criteria. Sections 4–5 then develop the integer tail numerators and zero-error identities used in the remaining arguments.

For further criteria we distinguish two normalisations. In Section 6, the least common multiple removes repeated factors among the earlier denominators. In Section 7, reduction to lowest terms also cancels factors shared with the tail numerator. Thus a prime can persist in the LCM and disappear from the reduced denominator. The constant and periodic errors are treated in Sections 9–10. The bounded-increase proof occupies Sections 11–12, followed by the scalar summability criterion in Section 13. Section 14 gives further questions and examples showing which recurrence hypotheses cannot be omitted. The appendices contain formal-source locations, the finite residue calculation and a result map.

We use $z_+ = \max(z, 0)$. Deleting a finite prefix is harmless for an eventual recurrence, but not for the coefficients in a higher-order rate. In particular, the index n in $1 + 3/n + o(n^{-3})$ is kept fixed throughout the cubic proof. The recurrence sections also use zero-based indexing, with that convention stated where the integer sequences are introduced.

Keywords. irrationality; Ahmes series; Sylvester’s sequence; unit fractions; Lean 4. **MSC 2020.** 11J72 (primary); 11B37, 11D68, 68V20 (secondary).

2 Irrationality at the cubic rate

A sufficiently precise rate constrains an integer tail more strongly than the quadratic limit alone. At rate $1 + 3/n + o(n^{-3})$, its numerator must eventually be cubic. The denominator recurrence will exclude every possible cubic. Here the increments of P_n/a_n grow like a positive multiple of n^2 , so the bounded-increase criterion later in the paper does not apply. The eventual Sylvester recurrence gives deviation $O(1/a_n)$ and is also excluded by this rate.

Theorem 2.1 (cubic-rate irrationality). *Lean · Comparator* If strictly increasing positive integers satisfy $a_n^2/a_{n+1} = 1 + 3/n + o(n^{-3})$, then $\sum_n 1/a_n$ is irrational.

The same extraction argument rules out nonintegral regular rates above one. These require no number-field calculation.

Theorem 2.2 (nonintegral regular-rate irrationality). [Lean](#) · [Comparator pending](#) Let $\lambda > 1$ be a real number that is not an integer. If strictly increasing positive integers satisfy $a_n^2/a_{n+1} = 1 + \lambda/n + o(n^{-\lambda})$, then $\sum_n 1/a_n$ is irrational.

The [zero-indexed Lean theorem](#) also derives convergence of the reciprocal series from the rate.

The rates retain their stated indices throughout. [Appendix A](#) gives the finite-prefix construction relating these one-based statements to the zero-based formal statements without changing the rate's error term.

For a concrete sequence satisfying the hypothesis, take

$$a_1 = 8, \quad a_{n+1} = \left\lceil \frac{na_n^2}{n+3} \right\rceil \quad (n \geq 1).$$

It begins 8, 16, 103, 5305, The inequality $a_{n+1} \geq a_n^2/4$ gives $a_n \geq 4 \cdot 2^{2^{n-1}} \geq 8$ by induction. Hence $a_{n+1} \geq a_n^2/4 \geq 2a_n$, so the sequence is strictly increasing. Rounding upwards gives

$$0 \leq 1 + \frac{3}{n} - \frac{a_n^2}{a_{n+1}} < \frac{16}{a_n^2} = o(n^{-3}).$$

Thus its reciprocal sum is irrational. The proof of the theorem uses the following arithmetic obstruction.

Theorem 2.3 (rising-factorial cubic exclusion). [Lean](#) · [Comparator](#) Let $a, C, D : \mathbb{N} \rightarrow \mathbb{Z}_{>0}$ satisfy

$$C_{n+1} = a_n C_n - D_n, \quad D_{n+1} = a_n D_n. \quad (1)$$

Then for every $A \in \mathbb{Q}_{>0}$ and every $B \in \mathbb{Q}$,

$$\liminf_{X \rightarrow \infty} \frac{\#\{n \leq X : C_n \neq A n(n+1)(n+2) + B\}}{X} > 0.$$

In particular no such orbit satisfies $C_n = A n(n+1)(n+2) + B$ for all large n .

The short paper needs only to exclude eventual equality with a cubic. Here we keep track of where each congruence fails and obtain positive lower density of disagreement. The bound may depend on the orbit and the cubic; no uniform numerical constant is claimed.

Suppose, then, that the disagreement set has lower density zero. It leaves arbitrarily late four-term blocks intact. On each such block, a third difference bounds $G_n = \gcd(C_n, D_n)$; the divisibility chain therefore stabilises. Dividing it out leaves the proposed numerator $mn(n+1)(n+2)/6 + c$, with $m > 0$ integral and $c = \pm 1$. If the middle of three consecutive numerators vanishes modulo a prime, the recurrence makes the negative product of its neighbours a square. Applied to the polynomial, this forces a square at every root modulo almost every prime. A square in the cubic field follows, and traces leave $m = 12$. Four-term blocks modulo seven exclude the two signs.

The density assertion comes from the same finite tests. A failed test recurs on an arithmetic progression, and each occurrence forces a disagreement in a window of bounded length. The next lemma counts those disagreements. After proving the arithmetic exclusion, we return to the rate assumption in [Lemma 2.11](#).

The integer tail numerator. Suppose $\sum_{n \geq 1} 1/a_n = p/q$ with positive integers p, q , and write $x_n = \sum_{k \geq n} 1/a_k$ for the tail. Put

$$D_n = q \prod_{1 \leq k < n} a_k, \quad C_n = D_n x_n.$$

Then D_n is a positive integer, and

$$C_n = D_n \left(\frac{p}{q} - \sum_{1 \leq k < n} \frac{1}{a_k} \right) = p \prod_{1 \leq k < n} a_k - q \sum_{1 \leq k < n} \prod_{\substack{1 \leq j < n \\ j \neq k}} a_j$$

is a positive integer as well. From $x_n = 1/a_n + x_{n+1}$,

$$C_{n+1} = D_{n+1} x_{n+1} = a_n D_n \left(x_n - \frac{1}{a_n} \right) = a_n C_n - D_n, \quad D_{n+1} = a_n D_n,$$

which is (1). These are the tail variables of Koizumi's Lemma 4 [12, pp. 11–12], up to a common positive factor. Section 4 constructs these variables together with the error $E_n = D_n - (a_n - 1)C_n$, which this section does not need. The subsequent cubic exclusion uses only these integer recurrences; it does not assume a reciprocal sum. To apply its zero-based statement without shifting the rate hypothesis, adjoin $a_0 = 1$, $D_0 = q$ and $C_0 = p + q$. The recurrences at 0 then give $D_1 = q$ and $C_1 = p$, while every original index $n \geq 1$ is unchanged. The auxiliary term a_0 need not satisfy the increasing-sequence hypothesis of the irrationality theorem: the exclusion requires only positive multipliers.

2.1 Reduction and lower density

For $S \subseteq \mathbb{N}$ write $\underline{d}(S) = \liminf_X \#(S \cap [1, X])/X$. Thus $\underline{d}(S) = 0$ means that these proportions approach zero along some sequence of cutoffs. Their limit need not exist. For a sequence F , let $\Delta F_n = F_{n+1} - F_n$; higher powers of Δ mean repeated forward differences. This operator is distinct from the later indexed quantity Δ_n , the Sylvester defect.

Lemma 2.4 (periodic obstruction principle). *Lean · Comparator* Let h, L be positive integers and let $j_1, \dots, j_L$ be fixed integer offsets. Suppose that for every sufficiently large n in one residue class modulo h at least one of $n + j_1, \dots, n + j_L$ lies in S . Then $\underline{d}(S) \geq 1/(Lh)$.

Proof. There are $X/h + O(1)$ relevant starting indices below X . Their witnesses lie below $X + O(1)$ because the offsets are fixed, and each element of S serves as a witness for at most L starting indices. Hence $L\#(S \cap [1, X]) \geq X/h - O(1)$, which gives the claimed lower density. $\square$

Fix $A \in \mathbb{Q}_{>0}$, $B \in \mathbb{Q}$, and suppose for contradiction that

$$P(n) = A n(n+1)(n+2) + B, \quad S = \{n : C_n \neq P(n)\}, \quad \underline{d}(S) = 0.$$

We retain this assumption through Section 2.4.

Lemma 2.5 (gcd stabilisation and the reduced polynomial). [Lean · Comparator](#) Write $G_n = \gcd(C_n, D_n)$. Then $6A$ is a positive integer, G_n divides $6A$ for every n , and G_n is eventually equal to a positive integer g . On the tail where $G_n = g$, put $u_n = C_n/g$, $v_n = D_n/g$ and $Q(n) = P(n)/g$. Then

$$u_{n+1} = a_n u_n - v_n, \quad v_{n+1} = a_n v_n, \quad \gcd(u_n, v_n) = 1, \quad \gcd(u_n, u_{n+1}) = 1, \quad (2)$$

and there are $m \in \mathbb{Z}_{>0}$ and $c \in \{-1, 1\}$ with

$$Q(n) = \frac{m}{6} n(n+1)(n+2) + c. \quad (3)$$

Proof. Equation (1) gives $G_n \mid G_{n+1}$ and $G_n \mid C_t$ for every $t \geq n$. Since $\underline{d}(S) = 0$, beyond every threshold there is a block of four consecutive indices disjoint from S : otherwise every block of four would meet S and $\underline{d}(S) \geq 1/4$. On such a block $\Delta^3 C_t = \Delta^3 P(t) = 6A$, and G_n divides each of the four values, hence divides $6A$. So $6A$ is a positive integer and the divisibility chain (G_n) is bounded and stabilises at some g .

On the primitive tail, $\gcd(u_n, v_n) = 1$ by construction, and a prime dividing u_n and u_{n+1} would divide $v_n = a_n u_n - u_{n+1}$, which gives $\gcd(u_n, u_{n+1}) = 1$.

Since $g \mid 6A$, the number $m := 6A/g$ is a positive integer. Put $c = B/g$, so

$$Q(n) = m \binom{n+2}{3} + c.$$

Choose any sufficiently late $n \notin S$. Then $Q(n) = u_n \in \mathbb{Z}$, and the binomial term is an integer, so $c \in \mathbb{Z}$. This proves (3) with $c \in \mathbb{Z}$. It also shows that Q is integer-valued: its values at integer arguments are integers, even though its coefficients need not all be integers.

It remains to exclude $c = 0$ and every c with a prime factor. Let ℓ be a prime dividing c , or any prime if $c = 0$. For $n \equiv -1 \pmod{6\ell}$, write $n+1 = 6\ell k$; then

$$\frac{n(n+1)(n+2)}{6} = \ell k(6\ell k - 1)(6\ell k + 1), \quad \frac{(n+1)(n+2)(n+3)}{6} = \ell k(n+2)(n+3),$$

so ℓ divides both $Q(n)$ and $Q(n+1)$. Agreement at both indices would give $\ell \mid \gcd(u_n, u_{n+1}) = 1$. Hence one of $n, n+1$ lies in S for every late $n \equiv -1 \pmod{6\ell}$, and Lemma 2.4 with $L = 2$, $h = 6\ell$ gives $\underline{d}(S) \geq 1/(12\ell)$. This contradiction leaves $c = \pm 1$. $\square$

Lemma 2.6 (coprimality and irreducibility). [Lean · Comparator](#) On the primitive tail, $\gcd(a_n, v_n) = 1$, the multipliers at distinct indices are pairwise coprime, infinitely many of them exceed 1, and the cubic $Q_{m,c}$ of (3) is irreducible over $\mathbb{Q}$.

Proof. A prime dividing a_n and v_n would divide u_{n+1} and v_{n+1} , against $\gcd(u_{n+1}, v_{n+1}) = 1$; and every earlier multiplier divides every later v_n , so distinct multipliers are coprime. If $a_n = 1$ for all large n then v_n is eventually a positive constant and $u_{n+1} = u_n - v_n$ decreases without bound, against positivity. So infinitely many distinct primes divide late multipliers.

Suppose $Q_{m,c}$ had a rational root. Choose a prime $\ell \nmid 6m$ dividing a late multiplier a_N , large enough that the root reduces modulo ℓ . Then $Q_{m,c}$ vanishes modulo ℓ on a full residue class, while $\ell \mid v_n$ and therefore $\ell \nmid u_n$ for every $n > N$. Agreement at any late index of that class is impossible, and Lemma 2.4 with $L = 1$, $h = \ell$ gives $\underline{d}(S) \geq 1/\ell$. $\square$

2.2 Square specialisation

For the proposed cubic $f(T) = T^3 - T + 6c/m$, the recurrence will force $r^2 - 1$ to be a square whenever r is a root modulo a good prime. We need the same conclusion for an algebraic root α in $\mathbb{Q}(\alpha)$. The issue is that a putative square root could lie in a quadratic extension of this field. Its nontrivial automorphism would fix α and change the sign of the square root. Chebotarev supplies a prime with exactly that behaviour, contradicting the finite-field condition. The following lemma isolates this passage; in the cubic application we take $H(T) = T^2 - 1$.

Lemma 2.7 (square specialisation). *Lean · Comparator pending* Let $f \in \mathbb{Q}[T]$ be irreducible with root α , and let $H \in \mathbb{Q}[T]$ satisfy $H(\alpha) \neq 0$. If for all but finitely many primes ℓ every root $r \in \mathbb{F}_\ell$ of the reduction of f has $H(r)$ a square in $\mathbb{F}_\ell^\times$, then $H(\alpha)$ is a square in $\mathbb{Q}(\alpha)^\times$.

Proof. Suppose that $H(\alpha)$ is a nonsquare in $K = \mathbb{Q}(\alpha)$, and choose β with $\beta^2 = H(\alpha)$. The nontrivial automorphism of the quadratic extension $K(\beta)/K$ extends to an element $\sigma \in \text{Gal}(L/K)$ for a finite Galois extension $L/\mathbb{Q}$ containing $K(\beta)$. In particular $\sigma\alpha = \alpha$ and $\sigma\beta = -\beta$.

By Chebotarev's density theorem in the form stated by Stevenhagen and Lenstra [10, §3, p. 15], infinitely many unramified rational primes have Frobenius conjugate to σ . Only infinitude is needed. We discard the exceptional primes in the hypothesis and the finitely many primes at which coefficients, α or β have denominators, together with 2 and the primes at which β reduces to zero. Above each remaining prime ℓ , choose a prime of L whose Frobenius is σ . Its residue field then satisfies

$$\bar{\alpha}^\ell = \bar{\alpha}, \quad \bar{\beta}^\ell = -\bar{\beta} \neq \bar{\beta}.$$

Thus $\bar{\alpha} \in \mathbb{F}_\ell$ is a root of the reduction of f . Neither of the two roots $\pm\bar{\beta}$ of $X^2 - H(\bar{\alpha})$ is fixed by Frobenius, so neither belongs to $\mathbb{F}_\ell$. Hence $H(\bar{\alpha})$ is a nonsquare there, contradicting the hypothesis. $\square$

An alternative proof of Lemma 2.7 uses zeta functions. A nonsquare algebraic integer b in a number field K is nonsquare modulo infinitely many degree-one primes of K . The underlying zeta-function argument first gives zero or a nonsquare. The zero case is excluded beyond the prime divisors of the nonzero constant coefficient of the minimal polynomial of b . If $H(\alpha)$ is not a square in $K = \mathbb{Q}(\alpha)$, this applies to $b = m^2H(\alpha)$, where the nonzero integer m makes b an algebraic integer, and the residue map at such a prime, of norm ℓ , sends α to a root $r \in \mathbb{F}_\ell$ of the reduction of f at which $H(r)$ is a nonsquare. The prime-reduction argument ([nonsquares modulo primes](#)) is proved from the simple pole of the Dedekind zeta function at $s = 1$, which is in Mathlib. If b were a nonzero square modulo all but finitely many primes of degree one, these primes would split in $K(\sqrt{b})$, and for real $s > 1$ near 1 the zeta function of $K(\sqrt{b})$ would be at least a positive constant times the square of the zeta function of K , which the two simple poles forbid.

Both arguments use the condition at every root for all but finitely many primes. Testing finitely many primes supplies no such condition. The zeta-function argument is the one used in the linked formal proof.

Proposition 2.8 (the square forced by the numerator recurrence). [Lean · Comparator](#) Write $\kappa = m/6$ and $\eta = 6c/m$, so that $Q_{m,c}(n) = \kappa f(n+1)$ for $f(T) = T^3 - T + \eta$. Then $\alpha^2 - 1$ is a square in $\mathbb{Q}(\alpha)^\times$ for a root α of f .

Proof. Eliminating v_n from (2) gives

$$u_{n+2} = (a_n + a_{n+1})u_{n+1} - a_n^2 u_n. \quad (4)$$

Let ℓ be a prime and k an index with $\ell \mid u_k$. Reading (4) at $n = k-1$ modulo ℓ gives $u_{k+1} \equiv -a_{k-1}^2 u_{k-1}$, hence

$$-u_{k-1}u_{k+1} \equiv (a_{k-1}u_{k-1})^2 \pmod{\ell}, \quad (5)$$

so $-u_{k-1}u_{k+1}$ is a square modulo ℓ .

Let $\ell \nmid 6m$ and let $r \in \mathbb{F}_\ell$ be a root of f . Then $r \neq 0$ and $r \neq \pm 1$, since $f(0) = \eta \neq 0$ and $f(\pm 1) = \eta$. Using $\eta = r - r^3$,

$$f(r-1) = -3r(r-1), \quad f(r+1) = 3r(r+1),$$

so at an index $k \equiv r-1 \pmod{\ell}$ we may reduce the polynomial values in $\mathbb{F}_\ell$. All equalities in the following calculation are in that field: $Q(k) = 0$, $Q(k-1) = -3\kappa r(r-1)$, $Q(k+1) = 3\kappa r(r+1)$, and

$$-Q(k-1)Q(k+1) = 9\kappa^2 r^2 (r^2 - 1). \quad (6)$$

If $r^2 - 1$ were a nonsquare modulo ℓ , then so would be the right side of (6), because $9\kappa^2 r^2$ is a nonzero square; agreement at all three of $k-1, k, k+1$ would then contradict (5). That prohibition recurs at every index of the class $r-1$ modulo ℓ , and Lemma 2.4 with $L = 3, h = \ell$ would give $\underline{d}(S) \geq 1/(3\ell)$. So $r^2 - 1$ is a square in $\mathbb{F}_\ell^\times$ at every root of every good reduction, and Lemma 2.7 applies with $H(T) = T^2 - 1$. $\square$

2.3 The square condition forces $m = 12$

Lemma 2.9 (classification of the scales). [Lean · Comparator](#) Let m be a positive integer, let $c \in \{-1, 1\}$ and $\eta = 6c/m$, and suppose $T^3 - T + \eta$ is irreducible over $\mathbb{Q}$ with a root α such that $\alpha^2 - 1$ is a square in $\mathbb{Q}(\alpha)^\times$. Then $m = 12$.

Proof. Choose $\beta \in \mathbb{Q}(\alpha)$ with $\beta^2 = \alpha^2 - 1$. The identity $(\alpha + \beta)(\alpha - \beta) = 1$ gives a reciprocal pair in the cubic field. Taking $z = \alpha + \beta$, we obtain

$$z^{-1} = \alpha - \beta, \quad \alpha = \frac{1}{2}(z + z^{-1}), \quad (7)$$

and z generates $\mathbb{Q}(\alpha)$. Write its minimal polynomial as $p(Z) = Z^3 + bZ^2 + dZ + w$ with $b, d, w \in \mathbb{Q}$ and $w \neq 0$. The minimal polynomial of z^{-1} is

$$\frac{Z^3 p(Z^{-1})}{w} = Z^3 + \frac{d}{w}Z^2 + \frac{b}{w}Z + \frac{1}{w}.$$

Thus Newton's identities apply directly to both z and its inverse. We use the first two traces of α to restrict these coefficients, then the third trace to impose $\eta = 6c/m$. All traces below are from $\mathbb{Q}(\alpha)$ to $\mathbb{Q}$. The polynomial $T^3 - T + \eta$ gives

$$\text{Tr } \alpha = 0, \quad \text{Tr } \alpha^2 = 2, \quad \text{Tr } \alpha^3 = -3\eta. \quad (8)$$

Newton's identities give $\text{Tr } z = -b$ and $\text{Tr } z^{-1} = -d/w$, so the first equation in (8) and (7) give

$$d = -bw. \quad (9)$$

For the second trace, Newton's identities and $d = -bw$ give

$$\text{Tr } z^2 = b^2 + 2bw, \quad \text{Tr } z^{-2} = b^2 - 2b/w.$$

Since the trace of 1 is 3, squaring (7) yields $4 \text{Tr } \alpha^2 = \text{Tr } z^2 + 6 + \text{Tr } z^{-2}$. Substituting $\text{Tr } \alpha^2 = 2$ gives $b^2 + b(w - w^{-1}) = 1$. Multiplication by w factors this equation:

$$(bw - 1)(b + w) = 0. \quad (10)$$

For the third trace, the same identities give

$$\begin{aligned} \text{Tr } z^3 &= -b^3 - 3b^2w - 3w, \\ \text{Tr } z^{-3} &= b^3 - 3b^2/w - 3/w. \end{aligned}$$

The cross terms in $(z + z^{-1})^3$ have trace $3(\text{Tr } z + \text{Tr } z^{-1}) = 0$ by (9). Thus $8 \text{Tr } \alpha^3 = \text{Tr } z^3 + \text{Tr } z^{-3}$. Using $\text{Tr } \alpha^3 = -3\eta$ now gives

$$\eta = \frac{(b^2 + 1)(w + w^{-1})}{8}. \quad (11)$$

By (10), either $b = -w$ or $b = w^{-1}$, and (11) becomes

$$\eta = \frac{(w^2 + 1)^2}{8w} \quad \text{or} \quad \eta = \frac{(w^2 + 1)^2}{8w^3}.$$

Write $w = r/s$ with $r \in \mathbb{Z} \setminus \{0\}$, $s \in \mathbb{Z}_{>0}$ and $\gcd(r, s) = 1$. Since $\eta = 6c/m$,

$$m = \frac{48c rs^3}{(r^2 + s^2)^2} \quad \text{or} \quad m = \frac{48c r^3 s}{(r^2 + s^2)^2}. \quad (12)$$

Now $\gcd(r^2 + s^2, rs) = 1$, so integrality of m and $|c| = 1$ force $(r^2 + s^2)^2 \mid 48$, hence $r^2 + s^2 \in \{1, 2, 4\}$. With $r \neq 0$ and $s \geq 1$ the value 1 is too small, and 4 is not a sum of two nonzero squares, so $r^2 + s^2 = 2$ and $|r| = s = 1$. Then (12) gives $m = 12cr$, and $m > 0$ gives $m = 12$. $\square$

The square condition leaves the two possible cubics

$$Q_{12, \pm 1}(n) = 2n(n+1)(n+2) \pm 1. \quad (13)$$

The square condition uses three consecutive numerators. The last step also uses the preceding denominator update, so it tests four consecutive numerators instead.

2.4 The two remaining cubics fail modulo seven

Lemma 2.10 (two incompatible four-term residue patterns). [Lean · Comparator](#) For $Q_{12,1}$, every sufficiently late four-index window beginning at $n \equiv 0 \pmod{7}$ contains an exceptional index. For $Q_{12,-1}$, the same holds for windows beginning at $n \equiv 1 \pmod{7}$. In either case, $\underline{d}(S) \geq 1/7$. The specified starting residue classes are part of the assertion; no claim is made for every block of four consecutive indices.

Proof. Reduce modulo 7. For $c = 1$ the values of $Q_{12,1}$ at $n = 0, 1, 2, 3$ are 1, 13, 49, 121, that is

$$(1, 6, 0, 2), \tag{14}$$

and for $c = -1$ the values at $n = 1, 2, 3, 4$ are 11, 47, 119, 239, that is

$$(4, 5, 0, 1). \tag{15}$$

Each four-term pattern recurs with period 7.

Index a proposed occurrence locally by 0, 1, 2, 3, and use a_i for the multiplier at local index i . Let $(c_0, c_1, 0, c_3)$ be the residues of u and (d_0, d_1, d_2, d_3) those of v . Thus $c_{i+1} = a_i c_i - d_i$ and $d_{i+1} = a_i d_i$ modulo 7. The middle zero gives $a_1 c_1 = d_1$, then $d_2 = a_1 d_1$ and $c_3 = -d_2$, so

$$d_1^2 = c_1 a_1 d_1 = c_1 d_2 = -c_1 c_3. \tag{16}$$

For (14) this is $-6 \cdot 2 = 2$, and for (15) it is $-5 \cdot 1 = 2$. The square roots of 2 modulo 7 are 3 and 4, so $d_1 \in \{3, 4\}$ in both cases.

The first step gives $a_0 = (c_1 + d_0)/c_0$ and hence $d_1 = d_0(d_0 + c_1)/c_0$, which is legitimate because $c_0 \in \{1, 4\}$ is invertible. For (14) this is $d_1 = d_0(d_0 + 6)$, with values

$$(0, 0, 2, 6, 5, 6, 2) \quad \text{at } d_0 = 0, 1, \dots, 6,$$

and for (15) it is $d_1 = 2d_0(d_0 + 5)$, with values

$$(0, 5, 0, 6, 2, 2, 6).$$

The incompatibility also follows without enumerating d_0 . Completing the square gives

$$d_1 + 2 = (d_0 + 3)^2 \quad (c = 1), \quad d_1 + 2 = 2(d_0 - 1)^2 \quad (c = -1).$$

Since 2 is a square modulo 7, either update makes $d_1 + 2$ a square. The required values $d_1 = 3, 4$ instead give the nonsquares 5, 6. The two displayed lists provide the same exhaustive check: both images are $\{0, 2, 5, 6\}$, disjoint from $\{3, 4\}$.

Hence every sufficiently late window $[7k, 7k + 3]$ for the plus profile, and every sufficiently late window $[1 + 7k, 4 + 7k]$ for the minus profile, contains an element of S . Windows within each family are pairwise disjoint. There are $X/7 + O(1)$ such windows contained in $[0, X]$, so choosing one exceptional index in each gives $\#(S \cap [0, X]) \geq X/7 - O(1)$ and therefore $\underline{d}(S) \geq 1/7$. This stronger local bound is not promoted to a uniform lower bound for arbitrary cubic profiles. $\square$

Proof of Theorem 2.3. Assume $\underline{d}(S) = 0$. Lemmas 2.5 and 2.6 put the primitive tail in the shape (3) with $c = \pm 1$ and $Q_{m,c}$ irreducible. Each excluded branch already contradicts the assumption by producing a positive, branch-dependent lower density, namely $1/(12\ell)$ or $1/\ell$ for the relevant prime ℓ . Proposition 2.8 and Lemma 2.9 force $m = 12$, and Lemma 2.10 then gives $\underline{d}(S) \geq 1/7$, the final contradiction. $\square$

2.5 Polynomial extraction at a regular rate

It remains to connect the arithmetic exclusion with an asymptotic rate. For the cubic rate, the natural comparison is $n(n+1)(n+2)$; its successive ratio is exactly $1 + 3/n$. For a real parameter $\lambda > 1$, the corresponding comparison sequence is $\Gamma(n + \lambda)/\Gamma(n)$, with ratio $1 + \lambda/n$. The argument first leaves an error $o(n)$. Subtracting the two recurrences then shows that the first difference of that error tends to zero. A sufficiently high difference of the original integer sequence is therefore a small integer and must vanish. Its eventual polynomial growth also forces λ to be integral.

Lemma 2.11 (integer extraction at a regular rate). [Lean](#) · [Comparator](#) *Let $\lambda > 1$ and let C_n be positive integers with*

$$\frac{C_{n+1}}{C_n} = 1 + \frac{\lambda}{n} + o(n^{-\lambda}). \quad (17)$$

Then λ is an integer $d \geq 2$, and there are $A \in \mathbb{Q}_{>0}$ and $B \in \mathbb{Q}$ with $C_n = A n(n+1) \cdots (n+d-1) + B$ for all large n .

Proof. Put $F_n = \Gamma(n + \lambda)/\Gamma(n)$, so that $F_{n+1}/F_n = 1 + \lambda/n$ exactly and $F_n \sim n^\lambda$ by the gamma-ratio asymptotic [20, 5.11.12]. Write $\varepsilon_n = o(n^{-\lambda})$ for the error in (17) and $z_n = C_n/F_n$, so that $z_{n+1}/z_n = 1 + \varepsilon_n/(1 + \lambda/n)$. Since $\lambda > 1$ the errors are absolutely summable, so the product converges and $z_n \rightarrow K$ for some $K > 0$, with $z_n - K = o(n^{1-\lambda})$. To see the stated error, put $\eta_n = \sup_{k \geq n} k^\lambda |\varepsilon_k| \rightarrow 0$; the tail of the logarithmic product has absolute value at most $O(\eta_n \sum_{k \geq n} k^{-\lambda}) = o(n^{1-\lambda})$. The factors are positive eventually, so the limiting product is nonzero. Hence

$$\delta_n := C_n - K F_n = F_n(z_n - K) = o(n). \quad (18)$$

We subtract the recurrences for C_n and $K F_n$ to obtain $\delta_{n+1} - \delta_n = (\lambda/n)\delta_n + \varepsilon_n C_n$. The first term is $o(1)$ by (18), and the second is $o(1)$ because $C_n = O(n^\lambda)$. Hence $\Delta \delta_n \rightarrow 0$. Every further difference is a finite linear combination of shifts of this sequence, so $\Delta^j \delta_n \rightarrow 0$ for all $j \geq 1$.

For the comparison sequence, the Gamma recurrence gives

$$\Delta F_n = \frac{\Gamma(n+1+\lambda)}{\Gamma(n+1)} - \frac{\Gamma(n+\lambda)}{\Gamma(n)} = \lambda \frac{\Gamma(n+\lambda)}{\Gamma(n+1)}.$$

Iterating this identity gives

$$\Delta^j F_n = \lambda(\lambda-1) \cdots (\lambda-j+1) \frac{\Gamma(n+\lambda)}{\Gamma(n+j)},$$

whose right side is $O(n^{\lambda-j})$. Choose an integer $j > \lambda$; then $\Delta^j F_n \rightarrow 0$, so $\Delta^j C_n \rightarrow 0$. These are integers, so $\Delta^j C_n = 0$ for all large n . Choose N beyond this threshold. The Newton forward-difference formula gives

$$C_{N+t} = \sum_{i=0}^{j-1} \binom{t}{i} \Delta^i C_N \quad (t \geq 0),$$

so C_n agrees eventually with a polynomial with rational coefficients. Its growth $C_n \asymp n^\lambda$ forces the degree to be λ , so $\lambda = d$ is an integer, and $d \geq 2$ because $\lambda > 1$. For that integer $F_n = n(n+1) \cdots (n+d-1)$ exactly, so (18) says that the difference of two polynomials of degree d is $o(n)$, hence constant. The leading coefficient of the rational polynomial is K , so $K \in \mathbb{Q}_{>0}$; the constant difference is rational as well. This gives the stated shape with $A = K$ and $B \in \mathbb{Q}$. $\square$

The little-oh error in (17) cannot in general be replaced by a big-oh error. The positive integers $C_n = n(n+1)(n+2) + (-1)^n$ satisfy $C_{n+1}/C_n = 1 + 3/n + O(n^{-3})$ and are not eventually polynomial. Likewise $C_n = \lceil n^{3/2} \rceil$ satisfies $C_{n+1}/C_n = 1 + 3/(2n) + O(n^{-3/2})$: rounding contributes $O(n^{-3/2})$, while the unrounded ratio has error $O(n^{-2})$. These two scalar examples prevent the proposed weakening of Lemma 2.11. Neither is asserted to satisfy the exact reciprocal-tail recurrences.

Lemma 2.12 (comparison of the numerator and denominator ratios). *Lean · Comparator*
 Let a_n be strictly increasing positive integers with $a_{n+1}/a_n^2 \rightarrow 1$ and $\sum_n 1/a_n$ rational, and let (C_n, D_n) be the integer tail. Write $\gamma_n = a_n^2/a_{n+1} - 1$. Then

$$\frac{C_{n+1}}{C_n} = 1 + \gamma_n + O(a_n^{-1}), \quad a_n \geq \exp(c 2^n) \text{ eventually for some } c > 0.$$

Proof. Since $a_{n+1}/a_n^2 \rightarrow 1$ and $a_n \rightarrow \infty$, there is an N with $a_{n+1} \geq a_n^2/2 \geq 2a_n$ for $n \geq N$. Iterating $a_{n+1} \geq a_n^2/2$ from an index where $a_n > 2$ gives $a_n \geq \exp(c 2^n)$ eventually. The same doubling gives, for $m > N$,

$$\frac{1}{a_m} \leq x_m = \sum_{k \geq m} \frac{1}{a_k} \leq \frac{2}{a_m},$$

so $x_m = (1 + \rho_m)/a_m$ with $0 \leq \rho_m = a_m x_{m+1} \leq 2a_m/a_{m+1} \leq 4/a_m$, using $a_{m+1} \geq a_m^2/2$. Hence

$$\frac{C_{n+1}}{C_n} = \frac{a_n D_n x_{n+1}}{D_n x_n} = a_n \cdot \frac{(1 + \rho_{n+1})/a_{n+1}}{(1 + \rho_n)/a_n} = \frac{a_n^2}{a_{n+1}} \cdot \frac{1 + \rho_{n+1}}{1 + \rho_n}.$$

The last factor is $1 + O(1/a_n)$ and a_n^2/a_{n+1} is bounded, so the product is $a_n^2/a_{n+1} + O(a_n^{-1})$. $\square$

The exact identity in Section 6 gives the same estimate: $C_{n+1}/C_n = 1 - E_n/C_n$, while $0 < \gamma_n + E_n/C_n < 3/a_n$ eventually. The identity follows from Theorem 5.1; its bound also uses vanishing relative error and near-quadratic growth. Equation (21) is the resulting weighted-sum application, not the identity itself.

Proof of Theorem 2.2. The stated rate implies $a_{n+1}/a_n^2 \rightarrow 1$ and convergence of the reciprocal series. Suppose its sum were rational, and form the positive integer tail numerators C_n . Lemma 2.12 gives $C_{n+1}/C_n = a_n^2/a_{n+1} + O(a_n^{-1})$ and an eventual bound $a_n \geq \exp(c 2^n)$ for some $c > 0$. Hence $a_n^{-1} = o(n^{-\lambda})$ and

$$\frac{C_{n+1}}{C_n} = 1 + \frac{\lambda}{n} + o(n^{-\lambda}).$$

Lemma 2.11 forces λ to be an integer, contrary to the hypothesis. $\square$

For a concrete nonintegral rate, take $a_1 = 8$ and

$$a_{n+1} = \left\lfloor \frac{2na_n^2}{2n+3} \right\rfloor \quad (n \geq 1).$$

The first terms are 8, 26, 387, 99846. Writing $r_n = 1 + 3/(2n)$ gives $a_{n+1} = \lfloor a_n^2/r_n \rfloor$ and $0 \leq r_n - a_n^2/a_{n+1} < r_n^2/a_n^2 \leq 25/(4a_n^2)$. Also $a_{n+1} \geq (2/5)a_n^2$, which from $a_1 = 8$ proves strict increase and exponential lower growth. Thus the ratio error is $o(n^{-3/2})$, and Theorem 2.2 proves this reciprocal sum irrational.

Proof of Theorem 2.1. The hypothesis $a_n^2/a_{n+1} = 1 + 3/n + o(n^{-3})$ gives $a_{n+1}/a_n^2 \rightarrow 1$. Suppose $\sum_n 1/a_n$ were rational and pass to the integer tail, so that (a, C, D) satisfies (1) with every C_n and D_n a positive integer. By Lemma 2.12 and $a_n \geq \exp(c2^n)$, the error term $O(a_n^{-1})$ is $o(n^{-3})$, so

$$\frac{C_{n+1}}{C_n} = 1 + \frac{3}{n} + o(n^{-3}).$$

Lemma 2.11 at $\lambda = 3$ gives $C_n = A n(n+1)(n+2) + B$ for all large n , with $A \in \mathbb{Q}_{>0}$ and $B \in \mathbb{Q}$. That contradicts Theorem 2.3. $\square$

No restriction on the rational normalisation was assumed: the proof derives the permitted primitive leading coefficient. The two residue patterns (14) and (15), the image $\{0, 2, 5, 6\}$ modulo 7 and the condition $(r^2 + s^2)^2 \mid 48$ have been calculated above.

The rationality supposition in this proof supplies the integer orbit. The rate implies $a_{n+1}/a_n^2 \rightarrow 1$ and excludes the eventual Sylvester recurrence, so the conclusion is irrationality on a restricted class of the sequences in Problem 1.1.

Writing $P_n = \prod_{k < n} a_k$, at the cubic rate P_n/a_n grows like n^3 and its increment like n^2 . Thus neither the bounded-increment criterion nor Koizumi's nonpositive upper-limit criterion applies. The rate also fails his $1 + o(1/n)$ condition, and Duverney's signed defect series diverges. We do not decide whether the LCM-weighted criteria apply; that requires information about the repeated factors in the prefix product. Conversely, Theorem 2.3 assumes only positive integer a_n, C_n, D_n and the recurrences (1), not the near-quadratic growth limit. The two arguments therefore retain distinct hypotheses.

In familiar approximation notation, $C_n = q(\prod_{k < n} a_k \cdot \sum_m 1/a_m - \sum_{k < n} \prod_{j < n, j \neq k} a_j)$ is a linear form in the reciprocal sum with integer coefficients. These integer remainders grow like n^3 . We use the precision $o(n^{-3})$ to determine their polynomial form by finite differences, then exclude that form by the recurrence. The integer example $n(n+1)(n+2) + (-1)^n$ above explains why an $O(n^{-3})$ error alone does not justify that polynomial conclusion.

3 Prior work and formalisation

Classical criteria and pseudo-greedy expansions. The rational-tail method predates these coordinates. Erdős and Straus [15, Theorem 2.1, pp. 85–86] use eventual integer remainders under a small-numerator hypothesis. In polynomial Cantor series, Hančl and Tijdeman split the numerator into finitely many shifted products. Rationality is then characterised by the vanishing of the resulting polynomial sum [16, Theorem 2.2

and its proof, pp. 39–40]. Their theorem provides methodological context, not a result for arbitrary arrays of signed integer coefficients: the rearrangement of the infinite sum needs boundary control, supplied there by the polynomial hypothesis. Neither result supplies the first-crossing argument for a lower error bound proved below.

Several classical results give sufficient conditions directly on the sequence (a_n) . Their hypotheses use different quantities, so the comparisons must be made separately. Koizumi's product criterion is implied by eventual nonnegativity of the integer error E_n , under the identification of coordinates given below. It therefore already covers the descent argument in Section 8 (see [12, Cor. 4(1) and Prop. 1(1), pp. 14–15]). The original Erdős–Straus criterion instead uses a least common multiple. Erdős and Straus proved that if $\lim a_n/a_{n-1}^2 = 1$, the reciprocal sum is rational, and (a_n) has no Sylvester tail, then

$$\limsup_{n \rightarrow \infty} \frac{[a_1, \dots, a_n]}{a_{n+1}} \left(\frac{a_{n+1}^2}{a_{n+2}} - 1 \right) > 0,$$

where $[a_1, \dots, a_n]$ is the least common multiple [1, Theorem 3, p. 132]. This is the indexing in the original theorem. With $A_r = \text{lcm}(a_1, \dots, a_{r-1})$ and $r = n + 1$, its expression is exactly $(A_r/a_r)(a_r^2/a_{r+1} - 1)$, the LCM quantity in the short note. The index change is not an additional difference between those two criteria. Erdős's 1988 survey, followed by the catalogue summary, instead prints $a_n^2/a_{n+1} - 1$ in the second factor while retaining the same prefix-LCM quotient; that displayed summary is off by one and is not followed here [6, p. 105][13]. Koizumi records the convenient sufficient rate $a_n^2/a_{n+1} = 1 + o(1/n)$ under which the Erdős–Straus criterion settles the problem [12, Remark 3, p. 16]. Tijdeman and Yuan give a criterion of the same kind for Ahmes series with positive integer numerators, weighted by the least common multiple of the earlier denominators [4].

Duverney's signed criterion gives a further comparison. For positive integers $a_n \rightarrow \infty$ and signs $\epsilon_n \in \{-1, 1\}$, its printed hypothesis (3.6) is the one-sided condition

$$\sum_{n \geq 0} \left(\frac{a_{n+1}}{a_n^2} - 1 \right) < \infty,$$

without absolute values. The stated conclusion is that $\sum_{n \geq 0} \epsilon_n/a_n$ is rational if and only if

$$a_{n+1} = a_n^2 - (\epsilon_{n+1}/\epsilon_n)a_n + \epsilon_{n+2}/\epsilon_{n+1}$$

for all large n [2, Corollary 3.2, p. 287]. For the comparison here we impose the stronger sufficient hypothesis $\sum_{n \geq 0} |a_{n+1}/a_n^2 - 1| < \infty$ and take all signs positive. The distinction concerns one step of the printed proof. On pp. 299–300, the reduced auxiliary fractions p'_n/q'_n satisfy $p'_{n+1} \mid q'_n$ and

$$p'_n \leq p'_N \prod_{k=N}^{n-1} \frac{q'_k}{p'_k}.$$

The required boundedness follows if $\prod_{k \geq N} (p'_k/q'_k)$ has a positive limit. Absolute convergence of the growth-defect series ensures this, using the summable error in Duverney's estimate (3.3). Signed convergence alone does not justify that product step, even for positive rational factors. For each integer $m \geq 2$, take the pair $1 + 1/m, 1 - 1/m$ successively m times. The deviations cancel after every pair, and the intervening partial sums

are $1/m \rightarrow 0$, so their series converges. But the product through the block $m = N$ is

$$\prod_{m=2}^N \left(1 - \frac{1}{m^2}\right)^m = \frac{(N+1)^N}{2N^{N+1}} \sim \frac{e}{2N} \rightarrow 0;$$

the equality follows by cancelling powers of consecutive integers. This example does not impose $p'_{n+1} \mid q'_n$. It refutes only the general product inference, not Duverney's arithmetic criterion. We therefore use only the absolute-convergence form; the interpretation with merely signed convergence is not needed in any proof here.

In the all-positive case $\epsilon_n = 1$, absolute convergence also gives a direct comparison with the classical product criterion. With the one-based notation $P_n = \prod_{1 \leq j < n} a_j$ used above,

$$\frac{P_{n+1}/a_{n+1}}{P_n/a_n} = \frac{a_n^2}{a_{n+1}}.$$

The ratios on the right have an absolutely convergent sum of deviations from 1, because the same is true of their reciprocals and those reciprocals tend to 1. Thus P_n/a_n has a positive finite limit, and its increments tend to zero. This case already satisfies the classical nonpositive-upper-limit condition; the bounded-increment result allows a larger class of sequences.

A different quantitative question is treated by Duverney, Kurosawa and Shiokawa [17, Theorem 1, author-version p. 2; proof in Section 3]. For rational $x_n > 1$, their result assumes eventual $x_{n+1} \geq x_n^2$ and control of the accumulated denominators of x_{n+1}/x_n^2 ; it computes the irrationality exponent of a signed reciprocal series. We use it only as a restricted comparison. In particular, the eventual Sylvester recurrence and the cubic rate considered here approach quadratic growth from the other side.

Badea's positive-term criterion is adjacent but different. For a convergent series $\sum_n b_n/a_n$ with a_n, b_n positive integers, eventual strict inequality

$$a_{n+1} > \frac{b_{n+1}}{b_n} a_n^2 - \frac{b_{n+1}}{b_n} a_n + 1$$

forces irrationality, while rationality under the corresponding non-strict inequality forces eventual equality [3, Theorem A, p. 313; Cor. 2.2, p. 316]. For $b_n = 1$, its hypothesis is $a_{n+1} \geq a_n^2 - a_n + 1$, an inequality between consecutive denominators, not the one-step sign condition $E_n \geq 0$ used in integer descent. Under the standing rational-tail and growth hypotheses, however, either inequality imposed eventually forces the Sylvester recurrence and hence the other. Their eventual forms are equivalent in this setting; that fact does not supply either condition for a general signed error.

Koizumi's pseudo-greedy expansion [12] chooses a_n by rounding $x_n^{-1} + 1$ to the nearest integer, with a half-integer rounded upwards. Here x_n is the remaining sum before subtracting $1/a_n$. Thus $a_n = \lfloor x_n^{-1} + 3/2 \rfloor$, and the gap $\epsilon_n = x_n^{-1} + 1 - a_n$ is the signed rounding error. Write the rational initial sum as $r = p/q$ with positive integers p, q . His Lemma 4 [12, pp. 11–12] produces integers $c_n > 0, d_n$ and e_n with $x_n = c_n/d_n$ and $\epsilon_n = e_n/c_n$, satisfying

$$a_n = \frac{d_n - e_n}{c_n} + 1, \quad c_{n+1} = c_n - e_n, \quad d_{n+1} = a_n d_n,$$

and the proof of that lemma also gives $c_{n+1} = a_n c_n - d_n$. These are the recurrences of Section 4, after matching the starting index and the initial normalization. For a sequence satisfying the hypotheses of Problem 1.1, the rounding rule is guaranteed only after a finite restart [12, Corollary 3, p. 9]. Restarting there with the tail in lowest terms gives (c_n, d_n, e_n) ; the original (C_n, D_n, E_n) on that tail is a fixed positive integer multiple of this triple, with the indices matched. Thus $E_n/C_n = \varepsilon_n$ is unchanged by the restart. The integer E_n is not necessarily a numerator in lowest terms, and the rounding range is not asserted before the restart. The first identity above gives $E_n = D_n - (a_n - 1)C_n$; the remaining identities are exactly the numerator and denominator recurrences, including Proposition 4.1. On an all-negative tail we later write $e_n = -E_n > 0$ for the magnitude. That local use of lower-case e_n has the opposite sign to Koizumi's signed integer e_n .

His Theorem 3 [12, pp. 12–13] proves the equivalence of two assertions. Conjecture 1 [12, p. 4] asks whether, for a positive rational r , the condition $\varepsilon_n \rightarrow 0$ forces $\varepsilon_n = 0$ eventually. Question 1 [12, p. 3] is the question of Erdős and Graham stated as Problem 1.1 above. Two implications used below already appear in these coordinates: his Lemma 3, that $\varepsilon_n = 0$ forces $\varepsilon_{n+1} = 0$, is the absorption of Theorem 5.8, and his Proposition 1(2), that $\varepsilon_n \geq 0$ for all large n forces $\varepsilon_n = 0$ for all large n , is the descent of Theorem 8.1. Both statements are given in [12, Lemma 3, p. 10; Prop. 1(2), p. 14]. Koizumi attributes Proposition 1(2) to Badea. Its contrapositive says that a counterexample must have negative errors infinitely often; it assumes the sign is eventually non-negative, while Theorem 12.1 below assumes only that the negative part is eventually bounded. It permits negative errors between $-B$ and 0 and imposes no independent upper bound on positive errors; the relative-error limit is still required.

The growth hypothesis in Problem 1.1 is calibrated by two facts about Sylvester's sequence. With $a_1 = 2$, it satisfies $a_n \sim c_0^{2^n}$ with $c_0 = 1.2640847 \dots$. Deleting initial terms and reindexing produces sequences with $a_n \sim C^{2^n}$ for arbitrarily large C whose reciprocals still sum to a rational number [7, arXiv v4, p. 2]. The classical sufficient condition for irrationality, $\lim_n a_n^{1/2^n} = \infty$, is therefore sharp. Kovač and Tao identify that condition as folklore [7, arXiv v4, p. 2]; they attribute the sharpness observation to Erdős (1975). A sequence with $a_n/a_{n-1}^2 \rightarrow 1$ has $a_n^{1/2^n}$ convergent, so that criterion says nothing about the sequences of Problem 1.1, and rationality is genuinely possible there. Sylvester's sequence is A000058 in the OEIS; A129871 is the variant 1, 2, 3, 7, 43, ... with an initial 1 prepended. For the recent literature on irrationality of Ahmes series we refer to Kovač and Tao [7], who resolve several problems of Erdős and Graham drawn from the same two sources cited above, and whose introduction gives a sample of the intermediate work, including Sándor (1984) and Badea (1987). They do not treat Problem #243; the rigidity conclusion asked for there is not among their results.

Crmaric and Kovač [18, Theorems 1–2] address the neighbouring Problem #270, not Problem #243. Allowing integer $f(n) \rightarrow \infty$ in $\sum_n (\prod_{j=1}^{f(n)} (n+j))^{-1}$ gives every positive real value; imposing nondecreasing f gives a measure-zero value set. The latter assertion does not rule out individual rational values. Their extension of Takeya's subsum argument explains why decay alone need not force irrationality when the summands remain freely selectable. The present exact denominator recurrence imposes additional compatibility, so neither direction is an implication between their theorem and ours.

The *Formal Conjectures* collection contains a mathematically equivalent unproved declaration, up to its zero-based indexing [14]. Its summand is $\mathbb{Q}$ -valued, so Lean's Summable hypothesis asserts the existence of a sum in $\mathbb{Q}$; the finite indexing shift changes that sum only by a rational prefix. The declaration therefore does encode the rationality premise, but its proof is sorry. It states the problem without proving it, and the development in this note is independent of it. Earlier formal work on the remainder method should also be distinguished from a solution of this problem: Koutsoukou-Argyragi and Li's Archive of Formal Proofs entry [19] records an Isabelle/HOL formalisation of Erdős–Straus (1974), Theorem 2.1, Corollary 2.10 and Theorem 3.1. That is formal prior art for those classical criteria, not an existing formal proof of Erdős #243.

Each criterion still requires an additional bound or convergence estimate under the unrestricted hypotheses. In the stable-gcd case, for example, the record-increment theorem gives a positive lower bound for the log-log coefficient, rather than proving it infinite. All conclusions about a persistently nonzero error assume failure of eventual Sylvester behaviour. We collect them in Section 14, where mixed-sign tails remain unclassified. The two recurrence proofs below are independent of the cubic argument: one uses a stable gcd and CRT, the other an infinite product and integer descent.

4 Integer numerators, denominators and errors

We retain an unreduced fraction C_n/D_n for each rational tail. This keeps the denominator update multiplicative: removing $1/a_n$ from the tail gives

$$D_{n+1} = a_n D_n, \quad C_{n+1} = a_n C_n - D_n. \quad (19)$$

We measure the difference from the Sylvester recurrence by $E_n = D_n - (a_n - 1)C_n$. These are Koizumi's recurrences [12, Lemma 4, pp. 11–12]; their formal definitions are the [denominator update](#), the [numerator update](#), and the [error](#).

We also consider integer solutions of (19) without initially assuming that C_n/D_n is a reciprocal tail. Here and in the following recurrence sections the indices may start at 0, and E_n always denotes the error just defined. We state positivity and growth assumptions as they are needed. The terms a_n are the multipliers, while C_n, D_n are the numerator and denominator. The pseudo-greedy rounding rule is an additional condition giving $-C_n/2 \leq E_n < C_n/2$, whereas the absorption argument below needs only $|E_n| < C_n$. At a negative error we write $e_n = -E_n > 0$. For an isolated step we use $a, D, C \in \mathbb{Z}$ without subscripts.

Proposition 4.1 (update law). [Lean](#) · [Comparator](#) For all $a, D, C \in \mathbb{Z}$,

$$aC - D = C - (D - (a - 1)C).$$

Consequently, every solution of (19) satisfies $C_{n+1} = C_n - E_n$.

Proof. $C - (D - (a - 1)C) = aC - D$. □

C_n decreases when $E_n > 0$, increases when $E_n < 0$, and stays unchanged when $E_n = 0$. This is why bounds on the negative error become bounds on upward increments.

Example 4.2 (the Sylvester sequence). Take $a_n = 2, 3, 7, 43, 1807, \dots$ with $a_{n+1} = a_n^2 - a_n + 1$, and start the state at $D_0 = C_0 = 1$. The two updates give

n	a_n	D_n	C_n	E_n
0	2	1	1	0
1	3	2	1	0
2	7	6	1	0
3	43	42	1	0
4	1807	1806	1	0

and $D_n = a_n - 1$ with $C_n = 1$ at every index: if $D_n = a_n - 1$ and $C_n = 1$ then $C_{n+1} = a_n - (a_n - 1) = 1$ and $D_{n+1} = a_n(a_n - 1) = a_{n+1} - 1$. Hence $E_n = D_n - (a_n - 1)C_n = 0$ throughout and the numerator never moves, which is Proposition 4.1 in the stationary case.

Construction from the reciprocal sum. Let $T_n = \sum_{k \geq n} 1/a_k$ and suppose $T_0 \in \mathbb{Q}$. Put D_0 equal to a common denominator and $D_n = D_0 a_0 \cdots a_{n-1}$, and set $C_n = D_n T_n$. Then $C_n \in \mathbb{Z}$ for every n , and from $T_n = 1/a_n + T_{n+1}$ one gets $C_{n+1} = a_n C_n - D_n$, which is the tail update. On Sylvester's sequence $T_n = 1/(a_n - 1)$ exactly, so $D_n = (a_n - 1)C_n$ and $E_n = 0$: the error measures deviation from the telescoping tail identity, and it vanishes identically on the Sylvester orbit.

In Lean, the elementary identities are stated for an abstract integer system, and their application to a rational reciprocal sum has separate statements, including [the product condition for a rational reciprocal sum](#). This passage uses the [tail realisation](#), the [denominator realisation](#) and the [signed update](#).

Conversely, the recurrences identify a reciprocal sum when the ratio C_n/D_n tends to zero. Regard that ratio as a real number, the [tail ratio](#). Assume $a_n > 0$, $D_n > 0$, $C_{n+1} + D_n = a_n C_n$ and $D_{n+1} = a_n D_n$. Dividing the numerator update by $a_n D_n$ gives $C_n/D_n = 1/a_n + C_{n+1}/D_{n+1}$, the [one-step reciprocal identity](#). Iterating gives

$$\frac{C_0}{D_0} = \sum_{n < N} \frac{1}{a_n} + \frac{C_N}{D_N},$$

as in the [finite telescoping identity](#). If $C_N/D_N \rightarrow 0$, taking the limit identifies $\sum_n 1/a_n = C_0/D_0$, the [series realisation](#). No growth, error bound or separate rationality hypothesis is used in these three identities. Positivity and the stated limit of C_n/D_n are still required. They prove the direction from recurrences to a series; the construction from a rational reciprocal sum was given above and is also formalised in separate declarations.

Under the relative-error hypothesis used below, the required limit is automatic. Suppose $a_n > 1$, $C_n > 0$, $D_n \geq 0$ and $E_n/C_n \rightarrow 0$ on an exact integer orbit. If $D_0 = 0$, then $D_n = 0$ and $E_n/C_n = 1 - a_n \leq -1$ at every index, a contradiction. Thus $D_0 \geq 1$ and $D_n \geq D_0 2^n$. For each $\varepsilon > 0$, the update gives $C_{n+1} \leq (1 + \varepsilon)C_n$ eventually. Taking $0 < \varepsilon < 1$ shows that $C_n/D_n \rightarrow 0$, so the telescoping identity realises the reciprocal sum as C_0/D_0 .

The same assumptions also imply the quadratic growth limit. Put $\theta_n = E_n/C_n$. Then $a_n = D_n/C_n + 1 - \theta_n \rightarrow \infty$, and the exact recurrences give

$$\frac{a_{n+1}}{a_n^2} = \frac{1}{1 - \theta_n} - \frac{1}{a_n} + \frac{1 - \theta_{n+1}}{a_n^2} \rightarrow 1.$$

In particular, the multipliers are strictly increasing eventually. Thus a global orbit with these positivity and relative-error assumptions already gives a sequence of the type in Problem 1.1 after deletion of a finite prefix. No separate tail-limit estimate is needed for such an orbit; without the relative-error assumption, that estimate remains a condition of the general telescoping argument.

Proposition 4.3 (scaling the numerator and denominator). *Lean · Comparator* For every $s, a, D, C \in \mathbb{Z}$,

$$\begin{aligned} a(sD) &= s(aD), & a(sC) - sD &= s(aC - D), \\ sD - (a - 1)sC &= s[D - (a - 1)C]. \end{aligned}$$

Multiplying the numerator and denominator by the same factor therefore preserves the recurrence and scales the error by that factor. When the factor divides all entries, we can divide it out instead. This is used in the finite calculation of Appendix B and in the induction in Theorem 10.1.

5 Zero errors and the Sylvester recurrence

To recover the Sylvester recurrence from a zero error, we eliminate D_n between two successive updates. Write

$$\Delta_n = a_{n+1} - (a_n^2 - a_n + 1),$$

the *Sylvester defect*. The identity below expresses $\Delta_n C_{n+1}$ through two consecutive errors. If both errors vanish, positivity lets us cancel C_{n+1} and recover the recurrence. Under the bound $|E_n| < C_n$, it will also make a single zero propagate to the next index.

Theorem 5.1 (defect identity). *Lean · Comparator* For all $a, a', D, C \in \mathbb{Z}$,

$$\begin{aligned} [a' - (a^2 - a + 1)](aC - D) &= a^2[D - (a - 1)C] \\ &\quad - [aD - (a' - 1)(aC - D)], \end{aligned}$$

that is, $\Delta_n C_{n+1} = a_n^2 E_n - E_{n+1}$.

Proof. A direct expansion: both sides equal $a'aC - a'D - a^3C + a^2D + a^2C - aD - aC + D$. $\square$

No sign or growth hypothesis is needed for the identity itself. Such hypotheses enter only in its consequences below.

Example 5.2 (one defect and the error it creates). Continue Example 4.2 but replace $a_3 = 43$ by $a_3 = 44$. The states $D_3 = 42$ and $C_3 = 1$ are unchanged, since they depend

only on a_0, a_1, a_2 , and $E_2 = 0$ still. The defect is $\Delta_2 = a_3 - (a_2^2 - a_2 + 1) = 44 - 43 = 1$, and $E_3 = D_3 - (a_3 - 1)C_3 = 42 - 43 = -1$, so the identity reads

$$\Delta_2 C_3 = 1 \cdot 1 = 1 = 49 \cdot 0 - (-1) = a_2^2 E_2 - E_3.$$

By Proposition 4.1 the numerator then rises, $C_4 = C_3 - E_3 = 2$. A single unit of defect at one index has produced a negative error of magnitude 1 at the next, increasing the numerator from 1 to 2.

Theorem 5.3 (two vanishing errors force the step). [Lean](#) · [Comparator](#) *Let $a, a', D, C \in \mathbb{Z}$ with $aC - D \neq 0$. If*

$$D - (a - 1)C = 0, \quad aD - (a' - 1)(aC - D) = 0,$$

then $a' = a^2 - a + 1$.

Proof. Theorem 5.1 gives $[a' - (a^2 - a + 1)](aC - D) = 0$. The second factor is nonzero, so the first factor must vanish. $\square$

The hypothesis $C_{n+1} \neq 0$ is not removable: at $C_{n+1} = 0$ the identity gives no information about a' .

5.1 Relations among three consecutive numerators

Eliminating the denominator from two successive steps gives a relation among three reduced numerators. Write u, u_1, u_2 for the reduced numerator coordinates, v, v_1 for the corresponding denominator coordinates, and h, h_1 for the two common factors removed in reduction. If the two steps have multipliers a, a_1 , then the exact hypotheses are

$$hu_1 + v = au, \quad hv_1 = av, \quad h_1u_2 + v_1 = a_1u_1.$$

Theorem 5.4 (eliminating the denominator from two steps). [Lean](#) · [Comparator](#) *Let $a, a_1, u, u_1, u_2, v, v_1, h, h_1$ be integers satisfying*

$$hu_1 + v = au, \quad hv_1 = av, \quad h_1u_2 + v_1 = a_1u_1.$$

Then $a^2u + hh_1u_2 = h(a + a_1)u_1$.

Proof. Multiply the first equation by a , replace av using the second equation, and then replace $h_1u_2 + v_1$ using the third. The remaining terms factor as the displayed right-hand side. $\square$

This is the [recurrence obtained by eliminating the denominator](#). Both cancellation factors remain in the formula. The identity adds no assumption to the two exact steps, but it still contains both multipliers and cannot by itself force $a_1 = a^2 - a + 1$; Lemma 5.6 specifies what it forgets modulo a fixed old denominator.

When neither step cancels a common factor, the same recurrence gives a square identity. Write p, p_1, p_2 for three consecutive numerators, $q = ap - p_1$ for the denominator before the first step, and a, a_1 for the two multipliers. Then

$$p_2 + a^2p = (a + a_1)p_1.$$

The resulting square identity is as follows.

Theorem 5.5 (a square identity without cancellation). [Lean · Comparator](#) *Let a, a_1, p, p_1, p_2, q be integers with $q = ap - p_1$ and $p_2 + a^2p = (a + a_1)p_1$. Then*

$$q^2 + (pp_2 - p_1^2) = (a_1 - a)pp_1.$$

Proof. Substitute $q = ap - p_1$ and $p_2 = (a + a_1)p_1 - a^2p$:

$$\begin{aligned} q^2 + pp_2 - p_1^2 &= (ap - p_1)^2 + p((a + a_1)p_1 - a^2p) - p_1^2 \\ &= (a_1 - a)pp_1. \end{aligned}$$

□

The term q^2 is nonnegative, while $pp_2 - p_1^2$ measures the difference between the product of the outer numerators and the square of the middle one. The right side records the change in multiplier, so the identity gives a necessary algebraic constraint without asserting a sign or a global monotonicity. It is checked as [the square identity when no factor is cancelled](#); no global exclusion of approximate solutions is claimed.

The reduced error update must also be compatible with the denominator recurrence. The following calculation gives the exact compatibility condition. Here e, e_1 are signed errors in reduced coordinates, not the positive magnitudes denoted by e_n in the all-negative case. Let Δ be a proposed value of the Sylvester defect. If

$$hu_1 = u - e, \quad he_1 = a^2e - \Delta(u - e),$$

then the denominator equality

$$h((a_1 - 1)u_1 + e_1) = a((a - 1)u + e)$$

holds precisely when the following difference vanishes:

$$\begin{aligned} &h((a_1 - 1)u_1 + e_1) - a((a - 1)u + e) \\ &= [a_1 - (a^2 - a + 1) - \Delta](u - e). \end{aligned}$$

Consequently, when $u - e \neq 0$,

$$h((a_1 - 1)u_1 + e_1) = a((a - 1)u + e) \iff a_1 = a^2 - a + 1 + \Delta.$$

The [equivalence with the denominator recurrence](#) is formalised using the [factored difference between the two recurrences](#). The hypothesis $u - e \neq 0$ is essential: if $u - e = 0$, the factorised mismatch cannot identify a_1 .

The identities hold under the local hypotheses displayed in their statements. It is their proposed global applications, not the identities, that would need additional information, such as control of cancellation or a sign estimate for $pp_2 - p_1^2$. No such information is deduced here for an arbitrary rational near-quadratic tail.

Lemma 5.6 (saturation modulo an old denominator). [Lean · Comparator](#) *Let $M \geq 2$. Any finite word $r_0, \dots, r_k$ of units modulo M is compatible with the cancellation-free recurrences modulo M , with all reduced denominator residues equal to zero. Consequently the eliminated two-step identity alone imposes no further restriction on such unit words when the multiplier residues are free.*

Proof. Set $v_i \equiv 0 \pmod{M}$ and choose $a_i \equiv r_{i+1}r_i^{-1} \pmod{M}$. Then $r_{i+1} + v_i \equiv a_i r_i$ and $v_{i+1} \equiv a_i v_i$. Eliminating v_i gives the two-step identity automatically. $\square$

This is finite residue compatibility, not existence of a positive integer orbit, much less a rational tail with near-quadratic growth. For $M \mid v_T$ on a tail with no further common-factor cancellation, the actual u_n are indeed units modulo M . Useful further restrictions must therefore retain information discarded here: for example multiplier size, changing moduli, or primes outside the old denominator. The square restriction used in the cubic proof concerns precisely a prime at which a middle numerator vanishes.

Theorem 5.7 (sequence form). *Lean · Comparator* Let $a, D, C : \mathbb{N} \rightarrow \mathbb{Z}$ satisfy $D_{n+1} = a_n D_n$ and $C_{n+1} = a_n C_n - D_n$. If $E_n = 0$ for all sufficiently large n and $C_{n+1} \neq 0$ for all sufficiently large n , then $a_{n+1} = a_n^2 - a_n + 1$ for all sufficiently large n .

Take the larger of the two thresholds and apply Theorem 5.3 at each later index. This is the final step of the criteria below that force E_n to vanish eventually.

The same identity shows that a zero error persists whenever $|E_n| < C_n$ at the subsequent indices.

Theorem 5.8 (zero is absorbing). *Lean · Comparator* Let $a, C, D : \mathbb{N} \rightarrow \mathbb{N}$ satisfy $C_{n+1} + D_n = a_n C_n$ and $D_{n+1} = a_n D_n$, and put $E_n = D_n - (a_n - 1)C_n$. Suppose that $|E_n| < C_n$ for every n . If $E_n = 0$ then $E_{n+1} = 0$.

Proof. Putting $E_n = 0$ in Theorem 5.1 gives $\Delta_n C_{n+1} = -E_{n+1}$, so C_{n+1} divides E_{n+1} . A multiple of C_{n+1} of absolute value smaller than C_{n+1} is zero, and $|E_{n+1}| < C_{n+1}$ by hypothesis. $\square$

Beyond an index from which $|E_n| < C_n$ always holds, either E_n reaches zero and remains zero, or it is never zero. A zero at an earlier index need not persist: in Example 5.2, the altered multiplier gives $E_3 = -1$ and $C_3 = 1$, so the required strict inequality fails. Sections 9 and 10 address the special all-negative case. The mixed-sign analysis in Sections 12 and 13 separates arbitrarily late negative errors from an eventually nonnegative tail.

6 LCM numerators and weighted records

The least common multiple clears a rational tail while removing factors repeated among earlier denominators. We prove that its integer numerator is bounded exactly when a weighted sum over new maxima converges. At each such step we may subtract a fixed amount from the increase before summing. The error may have either sign, extending the setting of Koizumi's eventual-nonnegative case in Proposition 1(2).

Applied to rational tails, this gives a Sylvester recurrence criterion. Its convergence hypothesis remains to be derived in the unrestricted problem. Proposition 7.15 later gives an auxiliary model which avoids whole moduli. That model omits both the denominator recurrence and coprimality to the individual prime factors, so its counterexamples do not apply to the theorem here.

In this section the indices start at 0. Write the full reciprocal sum as p/q , with positive integers p, q , put $x_n = \sum_{k \geq n} 1/a_k$, and take $D_0 = q$. Set

$$L_n = \text{lcm}(q, a_0, \dots, a_{n-1}), \quad M_n = D_n/L_n, \quad U_n = L_n x_n = C_n/M_n, \quad V_n = E_n/M_n.$$

The rational tail has denominator dividing L_n , so U_n is an integer. Also $V_n = L_n - (a_n - 1)U_n \in \mathbb{Z}$. This clears the denominator but need not reduce the fraction: U_n and L_n may still have common factors. With $\rho_n = \gcd(L_n, a_n)$, the exact updates are

$$M_{n+1} = M_n \rho_n, \quad \rho_n U_{n+1} = U_n - V_n, \quad V_n = L_n - (a_n - 1)U_n.$$

These are Bado's LCM coordinates, with the indexing translated explicitly. Take his denominator parameter to be q and identify his a_{n+1} with our a_n . Then his $M_n, \Delta_n, K_n, u_{n+1}$ and g_{n+1} are respectively our L_n, M_n, U_n, V_n and ρ_n . His (19) becomes the middle update above [8, Prop. 7.1 and (16)–(20), pp. 6–7].

Write $R_n = \max_{j \leq n} U_j$ and $\mathcal{R} = \{n : U_{n+1} > R_n\}$. The inequality $|V_n| < U_n$ gives $U_{n+1} < U_n$ when $\rho_n \geq 2$, so every sufficiently late strict rise has $\rho_n = 1$. The stronger eventual bound $-U_n \leq 2V_n$, supplied by $V_n/U_n = E_n/C_n \rightarrow 0$, gives the quantitative estimate $U_{n+1} \leq 3U_n/4$ when $\rho_n \geq 2$. At a sufficiently late LCM record, $\rho_n = 1$ and the actual jump is $d_n = U_{n+1} - U_n = -V_n > 0$. At a contracting step with $\rho_n \geq 2$ this identity need not hold.

For $B = 0$, divergence under unbounded growth can already be seen by comparing a record step with the interval of new heights it covers:

$$\int_{R_n}^{R_{n+1}} f(t) dt \leq (R_{n+1} - R_n)f(U_n) \leq (U_{n+1} - U_n)f(U_n).$$

The last expression is $(-V_n)f(U_n)$ at a record. The theorem retains this divergence after subtracting a fixed B at every record. That is where the congruences enter: they select a progression of heights whose first crossings necessarily have an excess jump. For the divergence direction, unboundedness of U_n suffices; no limit $U_n \rightarrow \infty$ is assumed.

Theorem 6.1 (boundedness and a weighted sum over new maxima). [Lean](#) · [Comparator](#)
Let a_n, L_n, U_n be positive integers and V_n integers satisfying

$$\begin{aligned} L_{n+1} &= \text{lcm}(L_n, a_n), & \rho_n &= \gcd(L_n, a_n), \\ \rho_n U_{n+1} &= U_n - V_n, & V_n &= L_n - (a_n - 1)U_n, & -U_n &\leq 2V_n. \end{aligned}$$

Let $f : [1, \infty) \rightarrow [0, \infty)$ be finite and nonincreasing, with $\int_1^\infty f(t) dt = \infty$. For each fixed integer $B \geq 0$,

$$\sup_n U_n < \infty \iff \sum_{n \in \mathcal{R}} (-V_n - B)_+ f(U_n) < \infty.$$

Proof. A bounded integer running maximum increases only finitely many times, so bounded U_n gives a finite sum. Suppose instead that U_n is unbounded. There are infinitely many record rises, each with $\rho_n = 1$. Its multiplier is greater than one because $d_n = (a_n - 1)U_n - L_n > 0$. If $r < s$ are record indices, then $a_r \mid L_s$ and $\gcd(a_s, L_s) = 1$,

hence $\gcd(a_r, a_s) = 1$. The record multipliers are therefore distinct and pairwise co-prime. For any fixed $B \geq 1$, choose B of them greater than B , and take T after their indices. These earlier multipliers $m_0, \dots, m_{B-1}$ all divide L_T .

Put $P = \prod_i m_i$ and choose x with $m_i \mid x+i$ by the Chinese remainder theorem. Select the heights $\tau = x + B + kP > R_T$, $k \in \mathbb{Z}$. At the first crossing of any selected height we have $U_n \leq R_n < \tau \leq U_n + d_n$, so the crossing is a record step. Here $d_n = U_{n+1} - U_n$ is the full jump from the current numerator. If $d_n \leq B$, then $U_n \in [\tau - B, \tau)$, so

$$U_n = \tau - B + i = x + kP + i \quad \text{for some } 0 \leq i < B.$$

The choice of x gives $m_i \mid U_n$. Since $m_i \mid L_n$ as well, we obtain $m_i \mid d_n = (a_n - 1)U_n - L_n$, contradicting $0 < d_n \leq B < m_i$.

Every crossing therefore has $d_n > B$. If it first crosses $h \geq 1$ selected heights, their spacing gives $(h-1)P < d_n$. Writing $r = d_n - B \geq 1$ and using $P \geq B+1$, we obtain $d_n = B + r \leq Pr$, hence $h \leq r$. Since each crossed height is above U_n , monotonicity of f gives

$$\sum_{\substack{\tau \text{ first crossed} \\ \text{at step } n}} f(\tau) \leq (d_n - B)f(U_n).$$

Each selected height has one first crossing. The first is at most $R_T + P$, and successive selected heights are P apart. For $R_N \geq R_T + P$, the intervals $[\tau, \min\{\tau + P, R_N\}]$, over selected heights $\tau \leq R_N$, cover $[R_T + P, R_N]$. Since f is nonincreasing, its integral over each interval is at most $Pf(\tau)$. Summing the crossing bounds gives

$$\sum_{\substack{T \leq n < N \\ n \in \mathcal{R}}} (-V_n - B)_+ f(U_n) \geq \frac{1}{P} \int_{R_T+P}^{R_N} f(t) dt. \quad (20)$$

Since $R_n \rightarrow \infty$, the right side diverges for every $B \geq 1$. The case $B = 0$ follows by domination. $\square$

The record multipliers themselves supplied the CRT moduli, and the bound $-V_n < U_n$ excluded records with $\rho_n \geq 2$. For reciprocal tails we also have vanishing relative error. This forces the numerator to be unbounded whenever Sylvester behaviour fails.

Theorem 6.2 (a convergent weighted sum over new maxima). [Lean · Comparator](#) Assume the growth and rationality hypotheses of Problem 1.1, and let f be as in Theorem 6.1. The sequence is eventually Sylvester if and only if, for some integer $B \geq 0$,

$$\sum_{n \in \mathcal{R}} (-V_n - B)_+ f(U_n) < \infty.$$

Proof. If the sequence is not eventually Sylvester, zero error is never reached on a sufficiently late tail. Integrality gives $1/U_n \leq |V_n|/U_n = |E_n|/C_n \rightarrow 0$, so U_n is unbounded. The same relative-error limit gives $2|V_n| < U_n$ at every sufficiently large index, and hence the bound $-U_n \leq 2V_n$ required by Theorem 6.1. Applying that theorem to this tail forces divergence for every B . Once the running maximum of the retained tail exceeds the omitted prefix maximum, deleting that prefix no longer changes which steps

set new records. Unboundedness ensures that this crossing occurs. Conversely, a sequence satisfying the eventual Sylvester recurrence telescopes to $x_n = 1/(a_n - 1)$, so $V_n = 0$ eventually. $\square$

The nonincreasing weights $1/t$ and $1/[t \log(et)]$ have divergent integrals and are admissible. The weight $1/t^2$ is excluded: its integral is finite, so the crossing lower bound would not contradict unbounded numerators. For example, $f(t) = 1/[t \log(et)]$ gives the sufficient condition

$$\sum_{n \in \mathcal{R}} \frac{(-V_n - B)_+}{U_n \log(eU_n)} < \infty.$$

Its finite lower bound in (20) is $P^{-1} \log(\log(eR_N)/\log(e(R_T + P)))$. Further fixed iterated logarithmic factors are allowed whenever the integral still diverges. These are specialisations of one crossing theorem.

The criterion also has an exact expression in the original growth defect. Put $\gamma_n = a_n^2/a_{n+1} - 1$ and $\theta_n = E_n/C_n$. The defect identity gives

$$\gamma_n + \theta_n = \frac{(1 - \theta_n)(a_n - 1 + \theta_{n+1})}{a_{n+1}}, \quad 0 < \gamma_n + \theta_n < 3/a_n$$

eventually. Thus the two nonnegative summands $U_n f(U_n)(\gamma_n - B/U_n)_+$ and $(-V_n - B)_+ f(U_n)$ differ by at most $3U_n f(U_n)/a_n$. To see summability directly, put $P_n = \prod_{j < n} a_j$. The tail estimate $x_n \sim 1/a_n$ gives $C_n/a_n \sim qP_n/a_n^2$, and

$$\frac{P_{n+1}/a_{n+1}^2}{P_n/a_n^2} = \frac{a_n^3}{a_{n+1}^2} \rightarrow 0.$$

Thus $\sum_n C_n/a_n < \infty$ by the ratio test. Since $U_n \leq C_n$ and $f(U_n) \leq f(1)$, the comparison error is summable. Consequently Theorem 6.2 is equivalent to finiteness of

$$\sum_{n \in \mathcal{R}} U_n f(U_n) \left(\frac{a_n^2}{a_{n+1}} - 1 - \frac{B}{U_n} \right)_+ \quad (21)$$

for some B . The original hypotheses do not currently supply this finiteness. In particular, termwise convergence to zero is insufficient. A Lean formulation of this comparison, with the sufficient constant 16 in place of 3 and the summand of (21) at record indices and zero elsewhere, is in the companion Lean repository [plectis-erdos-lean](#); it is not covered by the evidence record.

The crossed heights lie above R_n , but the divisibility argument uses the starting numerator U_n and the full jump $U_{n+1} - U_n$. Replacing this by $R_{n+1} - R_n$ would discard the recovery from an earlier decrease.

Integer coefficients. The first-crossing proof extends to summands with arbitrary integer numerators b_n . We record this extension separately from the unit-fraction criteria used above. In that case

$$V_n = b_n L_n - (a_n - 1)U_n, \quad \rho_n U_{n+1} = U_n - V_n.$$

Assume that $a_n \geq 2$, that L_n, U_n are positive integers, and that $L_{n+1} = \text{lcm}(L_n, a_n)$, with $\rho_n = \text{gcd}(L_n, a_n)$. Suppose $V_n \geq -B$ eventually, with an integer $B \geq 0$. Then $U_{n+1} \leq U_n + B$; if $B = 0$, boundedness follows at once. For $B \geq 1$, a record step with $R_n > B$ must have $\rho_n = 1$, since $\rho_n \geq 2$ would give

$$U_{n+1} \leq \frac{U_n + B}{2} \leq \frac{R_n + B}{2} < R_n.$$

Thus an unbounded sequence would supply infinitely many pairwise coprime record multipliers. Choose B of them larger than B , and then a CRT block of B consecutive integers above the previous maximum, each divisible by one of the chosen multipliers. The first step past the upper end of the block is a high record step, so $\rho_n = 1$. Its positive jump is at most B , hence the preceding numerator U_n lies in the block. The corresponding multiplier divides both U_n and L_n , hence divides the jump $(a_n - 1)U_n - b_n L_n$. A positive jump at most B cannot have a divisor larger than B . No growth or relative-error hypothesis was used.

With the additional limit $V_n/U_n \rightarrow 0$, choose an integer K bounding U_n . Eventually $|V_n| < U_n/K \leq 1$, so the integer V_n is zero. The recurrence becomes $\rho_n U_{n+1} = U_n$; the positive integer sequence U_n is then nonincreasing and hence eventually constant. For positive b_n , the classical comparisons are Badea's Corollary 2.2 [3, p. 316] and the criterion of Tijdeman and Yuan [4]. Further finite examples separating boundedness from stationarity are in the [coefficient proof supplement](#).

7 New maxima of reduced numerators

We next reduce each tail to lowest terms, writing u_n/v_n . This can remove a prime power which remains present in every later LCM L_j . Before applying coprimality to new maxima of u_n , we therefore need estimates for the factors that survive cancellation. Throughout this section u_n denotes the reduced numerator and U_n the LCM numerator of the preceding section.

Fractions in lowest terms. Put

$$G_n = \text{gcd}(C_n, D_n), \quad u_n = C_n/G_n, \quad v_n = D_n/G_n, \quad \tilde{e}_n = E_n/G_n.$$

Thus $\text{gcd}(u_n, v_n) = 1$ and $\tilde{e}_n$ is a signed integer. The factor $h_n = G_{n+1}/G_n$ records the common factor removed at the next step. The relation to the preceding section is exact:

$$\frac{G_n}{M_n} = \text{gcd}(U_n, L_n), \quad u_n = \frac{U_n}{\text{gcd}(U_n, L_n)}, \quad v_n = \frac{L_n}{\text{gcd}(U_n, L_n)}.$$

Indeed, $C_n = M_n U_n$ and $D_n = M_n L_n$. Clearing with an LCM and reducing to lowest terms are different operations. For example, the exact step $(C, D, a) = (3, 6, 3)$ has $E = 0$ and gives $(C', D') = (3, 18)$. With $L = 6$, its LCM numerator falls from 3 to 1, while the reduced numerator stays 1: the LCM factor is $\rho = 3$, but the reduction factor is $h = 1$. From this step onwards, the Sylvester recurrence holds.

In general, before reduction the next numerator is $w_n = a_n u_n - v_n = u_n - \tilde{e}_n$, so

$$h_n u_{n+1} = w_n, \quad h_n v_{n+1} = a_n v_n, \quad \text{gcd}(u_n, u_{n+1}) = 1.$$

The negative magnitude $e_n = -E_n$ used earlier is not $\tilde{e}_n$; neither is Koizumi's real gap ε_n .

For the maxima and their increments write

$$R_n = \max_{k \leq n} u_k, \quad H_n = \max_{j \leq n} C_j, \quad s_n = R_{n+1} - R_n.$$

At a strict rise let $d_n = u_{n+1} - u_n$. Then $s_n = (d_n - (R_n - u_n))_+$. Thus the actual jump includes recovery of the earlier decrease $R_n - u_n$, whereas the record increment does not. For example, if $R_n = 10$, $u_n = 5$ and $u_{n+1} = 12$, then $s_n = 2$ but $d_n = 7$. This illustrates the definitions, not a claimed reciprocal-tail orbit. We will use

$$m_n = (-\tilde{e}_n)_+, \quad A_n = \frac{R_n}{u_n} m_n, \quad \delta_n = \left(\frac{a_n^2}{a_{n+1}} - 1 \right)_+, \quad \ell(x) = \log_2 \log_2 \max(4, x).$$

The factor R_n/u_n in A_n measures how far the current numerator has fallen below its previous maximum. It equals one at a maximum and can be large after a decrease. The symbol A_n is distinct from the prefix product A_n used later.

Unless a statement in this section specifies an abstract recurrence, we assume the hypotheses of Problem 1.1 and use its integer tails, for which $|E_n|/C_n \rightarrow 0$ after a finite shift. We will write out the conclusion $a_{n+1} = a_n^2 - a_n + 1$ eventually rather than give it a separate symbol.

Comparison with Duverney's reduced parameters. For the positive reciprocal series, the eventually reduced parameters in Duverney's Theorem 3.1 and Section 5.2 [2, pp. 285–286, 299–300] can be taken as follows. The symbols p_n, q_n in this comparison are Duverney's auxiliary integers, not the fixed numerator and denominator of the full reciprocal sum:

$$p_n = u_n, \quad q_n = w_n = a_n u_n - v_n = h_n u_{n+1}, \quad E_n = G_n(p_n - q_n).$$

Indeed, $\gcd(u_n, w_n) = \gcd(u_n, v_n) = 1$. Since $a_n v_n = a_n^2 u_n - a_n w_n$, reduction of the next tail gives

$$h_n = \gcd(w_n, a_n v_n) = \gcd(w_n, a_n^2), \quad p_{n+1} \mid q_n.$$

For the gcd equality, subtract the multiple $a_n w_n$ and use the coprimality of u_n, w_n ; the divisibility follows from $q_n = h_n p_{n+1}$. Eliminating v_{n+1} then gives Duverney's recurrence:

$$a_{n+1} = \frac{p_n}{q_n} a_n^2 - a_n + \frac{q_{n+1}}{p_{n+1}}.$$

Thus $q_n/p_{n+1} = h_n$ is precisely the cancellation factor, and $q_n/p_n \rightarrow 1$. This identifies the reduced parameters, not every possible unreduced choice in the original theorem. An upper bound on $q_n - p_n$ controls the increase before cancellation in the reduced coordinates. To transfer that bound directly to $(-E_n)_+ = G_n(q_n - p_n)_+$ would also require control of G_n .

Lemma 7.1 (the denominator valuation transition). *Lean · Comparator* Let u, v, a be positive integers, $\gcd(u, v) = 1$, and $w = au - v > 0$. Put $h = \gcd(w, av)$ and $v' = av/h$. For a

prime p , write $r = v_p(a)$, $s = v_p(v)$ and $t = v_p(w)$. Then

$$v_p(v') = \begin{cases} \max(r, s), & r \neq s, \\ \max(0, 2s - t), & r = s. \end{cases}$$

In particular $v_p(v') \leq \max(r, s)$. A strict loss relative to s requires $r = s \geq 1$ and $t > s$.

Proof. Always $v_p(v') = r + s - \min(t, r + s)$. If $r \neq s$, primitivity implies $t = \min(r, s)$: when $s > 0$, u is a p -adic unit, and when $s = 0 < r$, v is a unit. If $r = s > 0$, both terms in $au - v$ are divisible by p^s , so $t \geq s$; substituting gives the second case. If $r = s = 0$, the same formula gives zero without needing u to be a unit. $\square$

Cancellation is not the same as a fall in the reduced denominator valuation. For example, $(u, v, a) = (2, 15, 9)$ gives $w = 3$, $h = 3$ and $(u', v') = (1, 45)$. The numerator before cancellation exceeds u by only $w - u = 1$, yet the 3-adic valuation of the reduced denominator rises from 1 to 2. This is a finite exact step, not an infinite counterexample.

Corollary 7.2 (persistence of a prime power). *Lean · Comparator* Suppose $p^k \mid v_s$. If $w_n < p^{k+1}$ at every step from s through $t - 1$, then $p^k \mid v_t$.

Proof. At a first loss of divisibility by p^k , the current exponent is some $j \geq k$ and the valuation lemma forces $v_p(w_n) > j$. This would give $w_n \geq p^{j+1} \geq p^{k+1}$, contrary to the hypothesis. $\square$

These local calculations explain the persistence threshold used below.

7.1 Growth of the running maximum

Theorem 7.3 (increments of the running maximum). *Lean · Comparator* Let $\Theta = \limsup_n (H_{n+1} - H_n) / \ell(H_n)$ on a rational-tail orbit under the standing hypotheses, and suppose E_n is not eventually zero. Then either G_n is unbounded and Θ is infinite, or G_n stabilises at a value g and $\Theta \geq g v_T / \varphi(v_T)$ for every late T , so that $\Theta > g \geq 1$. For any orbit under the standing hypotheses, therefore, $\Theta = 0$ or $\Theta > 1$, and $\Theta \leq 1$ forces the eventual Sylvester recurrence.

The constant in the next condition is measured against a double logarithm, not against C_n . This grows much more slowly than any positive power of C_n . Every fixed bound on $(-E_n)_+$ satisfies it on a nonzero tail because $C_n \rightarrow \infty$; it also permits unbounded negative parts on the double-logarithmic scale, with the coefficient specified below. In contrast, an error of size $\sqrt{C_n}$ has relative size tending to zero but violates the condition. These are comparisons of bounds, not constructions of reciprocal tails.

Corollary 7.4 (the double-logarithmic bound). *Lean · Comparator* If

$$\limsup_n \frac{(-E_n)_+}{\ell(C_n)} \leq 1,$$

then the sequence is eventually Sylvester. Every counterexample therefore satisfies

$$\limsup_n \frac{(-E_n)_+}{\ell(C_n)} > 1.$$

Proof of Theorem 7.3 and Corollary 7.4. We first prove the auxiliary lower bound $\Theta \geq 1$. This part needs only an exact positive integer orbit with $a_n > 1$, $D_0 \geq 1$, $|E_n|/C_n \rightarrow 0$ and error not eventually zero. In particular, it will also apply to the abstract orbit in Theorem 7.11. The key input is that, outside a set of indices of density zero, a_n is coprime to the entire preceding denominator D_n . This supplies almost one new coprime modulus per step. The denominator growth then places the resulting CRT block at the required double-logarithmic height.

How often a multiplier is coprime to the preceding denominator. We use the LCM factorisation from Section 6. In the present zero-based indexing, put

$$L_n = \text{lcm}(D_0, a_0, \dots, a_{n-1}), \quad M_n = D_n / L_n, \quad \rho_n = \gcd(L_n, a_n).$$

The finite telescoping identity gives

$$\frac{C_n}{M_n} = L_n \left(\frac{C_0}{D_0} - \sum_{j < n} \frac{1}{a_j} \right) \in \mathbb{Z}_{>0}.$$

Thus $M_n \mid C_n$, while $M_0 = 1$ and $M_{n+1} = \rho_n M_n$. Since L_n and D_n have the same prime divisors,

$$\#\{n < N : \gcd(a_n, D_n) > 1\} = \#\{n < N : \rho_n > 1\} \leq \log_2 M_N \leq \log_2 C_N = o(N).$$

Here the last estimate follows by summing $\log(C_{n+1}/C_n) = \log(1 - E_n/C_n) = o(1)$. This is the density-one coprimality argument of Bado [8, Theorem 9.1, p. 7], written with the exact-orbit hypotheses used here. Only the finite telescoping identity is needed; no estimate on record increments has entered this count.

Choosing the moduli and controlling their product. Absorption and vanishing relative error give $C_n \rightarrow \infty$, hence $H_n \rightarrow \infty$. The maximum is still subexponential: for every $\varepsilon > 0$, choose K_ε with $C_j \leq K_\varepsilon e^{\varepsilon j}$ for every j . Then $H_n \leq K_\varepsilon e^{\varepsilon n}$, so $\log H_n = o(n)$. Since $D_n \geq 2^n$ and $a_n = D_n/C_n + 1 - E_n/C_n$, eventually

$$a_n \geq 2^{n/2}, \quad a_n < 2D_n, \quad \ell(D_n) \leq n + O(1).$$

For the last inequality, $D_{n+1} < 2D_n^2$ gives $1 + \log_2 D_{n+1} < 2(1 + \log_2 D_n)$. Iteration from a fixed late index, followed by another logarithm, gives $\ell(D_n) \leq n + O(1)$.

Suppose $H_{n+1} - H_n \leq c\ell(H_n)$ eventually for some $0 < c < 1$. For a large integer B , take the first B indices at or after $\lceil 3 \log_2 B \rceil$ for which $\gcd(a_n, D_n) = 1$. Write the corresponding multipliers as $m_0, \dots, m_{B-1}$ and let s be the index immediately after the last choice. The density estimate just proved gives $s = B + o(B)$: deleting $O(\log B)$ initial indices and $o(s)$ exceptional indices leaves B choices. Each $m_i > 2B$ for large B , and the chosen multipliers are pairwise coprime, because each earlier one divides the denominator preceding a later one. All divide D_s . For $P = \prod_i m_i$ we therefore have

$$(2B)^B < P \leq D_s, \quad H_s < P, \quad B < P, \quad \ell(3P) \leq B + o(B).$$

The bound on H_s uses $\log H_s = o(s)$ and $s = B + o(B)$. Thus the product is large enough to place the block beyond the previous maximum, but small enough that the allowed

jump at its height is less than the block length: $c\ell(3P) < B$ for all sufficiently large B . Both comparisons are needed; existence of a distant CRT block alone would not control a height-dependent jump bound.

Crossing the block. Choose $x \in [P, 2P)$ by the Chinese remainder theorem so that $m_i \mid x + i$ for $0 \leq i < B$. At the first record $C_t \geq x$ after s , the preceding maximum is below x and its increase is at most $c\ell(2P) < B$. Thus $x \leq C_t < x + B < 3P$, so some m_i divides both C_t and D_t . The exact updates preserve this common divisor. At the next record C_r we would therefore have

$$m_i \mid C_r - C_t, \quad 0 < C_r - C_t \leq c\ell(C_t) \leq c\ell(3P) < B < m_i,$$

which is impossible. This proves the auxiliary bound $\Theta \geq 1$.

The integer normalisation matters here. Scaling (C, D, E) by a positive integer k scales the running maximum by k and its limit-superior coefficient by k , since $\ell(kx)/\ell(x) \rightarrow 1$. Thus the bound with coefficient 1 is not invariant under arbitrary clearing of denominators. The next step keeps track of the common factor rather than discarding it.

The common gcd and the sharper coefficient. For any fixed N , divide the tail from N onwards by G_N . The resulting integer orbit satisfies the same hypotheses. Its running maximum is eventually H_n/G_N , and $\ell(H_n/G_N)/\ell(H_n) \rightarrow 1$. The auxiliary bound therefore gives $\Theta \geq G_N$. If G_N is unbounded, then $\Theta = +\infty$.

Otherwise $G_n = g$ eventually. Fix a later index T with $v_T > 1$. The reduced fractions have pairwise coprime multipliers, each coprime to v_T . For a large integer L , exactly

$$k_L = \frac{\varphi(v_T)}{v_T} L + O(v_T)$$

of the offsets $0, \dots, L-1$ are coprime to v_T . Assign to those offsets the next k_L multipliers, beginning at T . Put $s = T + k_L$ and $Q = v_s = v_T \prod_{T \leq j < s} a_j$. Solve $x \equiv 0 \pmod{v_T}$ and $x \equiv -j$ modulo the multiplier assigned to offset j , and take the solution in $[Q, 2Q)$. Every integer in $[x, x + L)$ then fails to be coprime to v_s . No later reduced numerator can lie there.

The running maximum R_s of the reduced numerator is below Q for large L , by the growth estimates above. The first crossing of x after s must therefore have a record increment at least L . Its preceding maximum is below $2Q$, and $\ell(2Q) \leq s + O(1) = T + k_L + O(1)$. The corresponding record indices tend to infinity with L , so

$$\limsup_n \frac{R_{n+1} - R_n}{\ell(R_n)} \geq \lim_{L \rightarrow \infty} \frac{L}{T + k_L + O(1)} = \frac{v_T}{\varphi(v_T)}.$$

Since $H_n = gR_n$ eventually, this gives $\Theta \geq gv_T/\varphi(v_T) > g$. Under the eventual Sylvester recurrence, C_n is eventually constant and $\Theta = 0$. Finally,

$$H_{n+1} - H_n = (-E_n - (H_n - C_n))_+ \leq (-E_n)_+.$$

Together with $\ell(C_n) \leq \ell(H_n)$, this proves the corollary. $\square$

Integer solutions of the recurrences. The strict bound $\Theta > 1$ is also a Lean theorem for solutions of (19) with $a_n > 1$, $C_n > 0$, $D_0 \geq 1$ and vanishing relative error whose error is not eventually zero ([strict record bound](#)). The implication of Corollary 7.4, that $\limsup_n (-E_n)_+ / \ell(C_n) \leq 1$ forces $E_n = 0$ eventually, is a Lean theorem for solutions of (19) with $a_n > 1$, $C_n > 0$ and vanishing relative error ([double-logarithmic bound for an exact orbit](#)); there $D_0 \geq 1$ follows from the vanishing relative error. The Lean sources also contain pointwise-rise versions of the arithmetic steps: [persistence of a common divisor](#), a [CRT block starting in \$\[P, 2P\)\$](#) , the [landing of a slowly rising numerator in that block](#) and the [resulting exclusion](#). Their landing bound controls increments of C_n ; the argument above controls increments of H_n .

No upper bound on Θ is proved. If G_n is unbounded, the theorem forces $\Theta = +\infty$. If $G_n = g$ eventually, it proves only $\Theta \geq g v_T / \varphi(v_T)$ for every late T . Since $v_T \mid v_{T+1}$, the ratios $\varphi(v_T)/v_T$ are nonincreasing and have a limit $\sigma \geq 0$. If $\sigma = 0$, the lower bound again forces $\Theta = +\infty$; if $\sigma > 0$, it gives $\Theta \geq g/\sigma$ but does not determine whether Θ is finite. Replacing H_n by the running maximum of u_n divides the coefficient by g , since $\ell(gx)/\ell(x) \rightarrow 1$.

7.2 Bounds that allow for cancellation and earlier decreases

Theorem 7.5 (bounds allowing for previous decreases). [Lean · Comparator](#) *Under the standing hypotheses, the following are equivalent: eventual Sylvester behaviour; $\limsup_n A_n < \infty$; $\limsup_n R_n \delta_n < \infty$. Each of those two limits superior is 0 or $+\infty$.*

Corollary 7.6 (the critical rate). [Lean · Comparator](#) *Under the standing hypotheses and $\delta_n = O(1/n)$, with no convergence of $n\delta_n$ assumed, eventual Sylvester behaviour is equivalent to $u_n = O(n)$ and to $(-\tilde{e}_n)_+ = O(1)$. A counterexample at the critical rate therefore has $\limsup_n u_n/n = \infty$ and $\limsup_n (-\tilde{e}_n)_+ = \infty$.*

Proof of Theorem 7.5. Suppose the orbit is not eventually Sylvester and $A_n \leq K$ eventually, for an integer $K \geq 1$. Absorption and $|\tilde{e}_n|/u_n \rightarrow 0$ give $u_n \rightarrow \infty$. Negative reduced errors must occur arbitrarily late, since otherwise $h_n u_{n+1} = u_n - \tilde{e}_n$ would make u_n eventually nonincreasing.

The bound controls cancellation as well as upward motion. Fix a late index s , and let $t > s$ be the first subsequent negative-error index. At the intervening steps the reduced numerator is nonincreasing, so $u_t \leq u_{s+1}$. Since $R_t \geq u_s$ and $m_t = |\tilde{e}_t| \geq 1$,

$$A_t = \frac{R_t m_t}{u_t} \geq \frac{u_s}{u_{s+1}} = \frac{h_s}{1 - \tilde{e}_s/u_s}.$$

Thus $h_s \leq 3K/2 < 2K$ after the relative-error threshold. Also $m_n \leq A_n \leq K$, since $R_n \geq u_n$, and hence $u_{n+1} \leq u_n + K$.

We can now choose primes too large to be removed by cancellation. The density-one argument in the proof of Theorem 7.3 supplies infinitely many late multipliers a_n coprime to D_n . These multipliers are pairwise coprime. At each such step, a_n is coprime to v_n and $\gcd(u_n, v_n) = 1$, so $a_n u_n - v_n$ is coprime to both a_n and v_n . Hence $h_n = 1$ and $a_n \mid v_{n+1}$. Only finitely many of the chosen multipliers can have all their prime divisors at most $2K$, since each uses a different prime from that finite set. Choose K distinct primes

$p_0, \dots, p_{K-1} > 2K$ at these steps. Once a chosen prime divides a reduced denominator, it divides every later one: $h_n v_{n+1} = a_n v_n$ and $h_n < 2K < p_i$ prevent its removal. All the primes therefore divide v_T at a common later index T .

Consequently every u_n with $n \geq T$ is coprime to all the p_i . Choose a CRT block of K consecutive integers, one divisible by each p_i , above R_T . Divergence forces a first crossing, while $u_{n+1} \leq u_n + K$ forces it to land inside the block, a contradiction. This proves that finite $\limsup A_n$ forces the eventual Sylvester recurrence.

The comparison $|\delta_n - m_n/u_n| \leq 3/a_n$ gives $|R_n \delta_n - A_n| \leq 3R_n/a_n \rightarrow 0$. Here $R_n \leq H_n$, $\log H_n = o(n)$ and a_n grows doubly exponentially. Thus the two boundedness conditions are equivalent. Under the eventual Sylvester recurrence, $u_n = 1$ and $m_n = 0$ eventually, so $A_n \rightarrow 0$ and $R_n \delta_n \rightarrow 0$. Since both quantities are nonnegative, their limits superior can only be 0 or $+\infty$. $\square$

The estimate at the first later negative error is the lemma in the release, [negative error scaled by the previous maximum after cancellation](#): at the first later negative error, $u_t \leq u_{s+1}$ and $u_s u_t \leq R_t |\tilde{e}_t| u_{s+1}$. The exact example $15/134 = 1/10 + 1/85 + 1/5695$, with steps in reduced fractions $(15, 134) \rightarrow (4, 335) \rightarrow (1, 5695)$ and $A_1 = 15/4$, attains equality in the displayed bound.

For Corollary 7.6, the comparison $|\delta_n - m_n/u_n| \leq 3/a_n$ and the rate $\delta_n = O(1/n)$ give $m_n/u_n = O(1/n)$. If $u_n = O(n)$, then $m_n = O(1)$. Conversely, $m_n = O(1)$ gives $u_{n+1} \leq u_n + m_n$, hence $u_n = O(n)$ and $R_n = O(n)$. In either case $A_n = R_n m_n/u_n = O(1)$, so Theorem 7.5 gives eventual Sylvester behaviour. The eventual Sylvester recurrence gives $u_n = 1$ and $m_n = 0$ eventually, which proves the converse implications.

The condition bounds the negative reduced error after multiplying it by R_n/u_n . At a current maximum this is just $(-\tilde{e}_n)_+$; after a decrease the multiplier is larger. A bound on $(-\tilde{e}_n)_+$ does not directly control this additional factor. Under the critical-rate hypothesis, the preceding corollary supplies the required control. The eventual Sylvester recurrence makes the expression eventually zero. We do not derive it for every rational tail satisfying the growth hypothesis.

Lemma 7.7 (large odd prime powers in the reduced denominator). [Lean · Comparator](#)
Under the standing hypotheses, for every fixed $A > 0$ and every sufficiently large n , the reduced denominator v_n has an odd prime-power divisor $Q = p^k$ with

$$Q > (H_n + 2)^A, \quad H_n = \max_{j \leq n} C_j.$$

The prime p may depend on n ; no stable-gcd or prime-arrival assumption is imposed.

Proof. The denominator grows too fast for all its odd prime-power factors to remain small compared with H_n . Indeed, the estimate for the rational tail gives $C_{n+1}/C_n \rightarrow 1$, hence $\log H_n = o(n)$. Since $x_n = u_n/v_n \leq 2/a_n$ and $u_n \geq 1$, $v_n \geq a_n/2$. Also $v_n \mid L_n = \text{lcm}(q, a_0, \dots, a_{n-1})$, so the 2-primary part of v_n is at most $\max(q, a_{n-1}) = a_{n-1}$ for all large n . Its odd part W_n therefore satisfies

$$W_n \geq \frac{a_n}{2a_{n-1}} \geq \frac{a_{n-1}}{4}, \quad \log W_n \geq c2^{n-1} - O(1)$$

for some $c > 0$. Put $B_n = (H_n + 2)^A = \exp(o(n))$. If every exact odd prime-power factor of W_n were at most B_n , then $W_n \mid \text{lcm}(1, \dots, \lfloor B_n \rfloor)$, giving

$$\log W_n \leq B_n \log B_n = \exp(o(n)).$$

This contradicts the preceding exponential lower bound for all sufficiently large n . An offending exact prime-power factor is the required Q . $\square$

The argument compares the logarithm of the odd denominator part with the logarithm of a factorial bound. It needs neither the prime number theorem nor a lower density of new primes, and makes no such claim.

Theorem 7.8 (unit record increments). *Lean · Comparator Under the standing hypotheses, if $R_{n+1} - R_n \leq 1$ for all large n , then $a_{n+1} = a_n^2 - a_n + 1$ for all large n . Hence the sequence is eventually Sylvester if and only if $\#\{n : R_{n+1} - R_n \geq 2\}$ is finite. No hypothesis is placed on decreases in u_n , on record-setting jumps, or on the cancellation factors h_n .*

Proof. Suppose the orbit is not eventually Sylvester. Absorption gives $\tilde{e}_n \neq 0$ on a late tail, so vanishing relative error and integrality give $u_n \rightarrow \infty$. Choose s so large that the unit-increment bound and $|\tilde{e}_n| < u_n/2$ hold for every $n \geq s$, and that Lemma 7.7 applies with $A = 3$. Since $H_s \geq R_s \geq 1$, it supplies an odd $Q = p^k \mid v_s$ with $Q > (H_s + 2)^3 > 4(R_s + 2)$ and $Q \geq 16$. Let Y be the least multiple of p above R_s . Then $Y \leq R_s + p < Q/4 + p \leq 5Q/4$. At the first $t > s$ with $u_t \geq Y$, the integral unit-record increment forces $u_t = Y$. Before t , one has $w_n < 3u_n/2 < 15Q/8 < pQ = p^{k+1}$. Corollary 7.2 therefore gives $p^k \mid v_t$. But $p \mid u_t = Y$, contradicting $\gcd(u_t, v_t) = 1$. The converse follows because the eventual Sylvester recurrence gives $u_n = 1$ eventually. $\square$

The plectis-erdos-lean repository contains the [unit-increment landing lemma](#) and the [one-unit record-increment implication](#); they assume the existence of the prime power that Lemma 7.7 supplies.

The two-unit criterion discussed below bounds the actual jump at each record step, rather than the increase in the running maximum. As numerical bounds on individual steps, neither condition implies the other: the first allows $s_n = 2$, while $s_n \leq 1$ permits large jumps after an earlier decrease. This is a comparison of the bounds, not a claim that they remain logically independent under all the standing tail hypotheses; there both criteria force eventual Sylvester behaviour. The finite example $(8, 177) \rightarrow (7, 4071) \rightarrow (10, 2373393)$ with multipliers 23 and 583 has records 8 and 10, increment 2, jump 3 and $\gcd(8, 10) = 2$, which is why the parity argument for crossing an odd level does not transfer from actual jumps to increments of the running maximum.

7.3 Counting jumps before a prime power can be lost

Theorem 7.9 (counting crossings before a prime power is lost). *Lean · Comparator Let the orbit satisfy the reduced recurrences of this section, with $2|\tilde{e}_n| < u_n$ from an index s . Let $p \geq 3$ be prime, $Q = p^\ell$ divide v_s with $Q \geq 16$, put $L = pQ/2$, and assume $R_s < L/2$. Assume that $u_t \geq L$ for some $t > s$, and let τ be the first such index, let J be the set of steps in $[s, \tau)$ that first cross at least one odd multiple of p in $(L/2, L]$, and put $X = \sum_{n \in J} (d_n - 2)$. Then every $n \in J$ is a record step with $h_n = 1$ and $d_n \geq 3$, and $pQ \leq (8p + 8)|J| + 4X + 8p$.*

Proof of Theorem 7.9. For $s \leq n < \tau$, one has $u_n < L$ and $w_n < 3u_n/2 < 3pQ/4 < p^{\ell+1}$, since $Q = p^\ell$. Corollary 7.2 gives $Q \mid v_t$ throughout $[s, \tau]$, so $p \nmid u_t$ there. A first crossing of a level above R_s is a record step. If $h_n \geq 2$ then $u_{n+1} = w_n/h_n < 3u_n/4$, so every such record step has $h_n = 1$. An odd multiple of p cannot be landed on. A jump of size 1 crossing it would land on it; a jump of size 2 avoiding the landing would have two even endpoints, contrary to $\gcd(u_n, u_{n+1}) = 1$. Thus every $n \in J$ has $d_n \geq 3$. It remains to count these levels. The interval $(L/2, L]$ has length $pQ/4$, and odd multiples of p are spaced by $2p$, so it contains at least $Q/8 - 1$ of them. The levels first crossed by a step of size d_n have the same spacing, so there are at most $1 + d_n/(2p)$ of them. Summing over J gives

$$Q/8 - 1 \leq |J| + \frac{2|J| + X}{2p},$$

which rearranges to the asserted inequality. $\square$

The next series ignores record jumps of size at most two, apart from finitely many initial terms, and assigns a smaller cost to a large jump when it begins at a large numerator. The eventual Sylvester recurrence gives no late records, so both series converge. The proof shows that a non-Sylvester rational tail contributes a fixed positive amount on arbitrarily late finite intervals. Thus convergence is a genuine additional requirement, not a restatement of the pointwise limit $|E_n|/C_n \rightarrow 0$.

Theorem 7.10 (two convergence criteria for large record jumps). *Lean · Comparator Under the standing hypotheses, the sum $\mathcal{E} = \sum_{n \text{ record}} (\mathbf{1}_{d_n \geq 3} u_n^{-1/2} + (d_n - 2)_+ u_n^{-1})$ is finite if and only if the sequence is eventually Sylvester; and $\sum_{n \text{ record}} (d_n - 2)_+ u_n^{-1/2}$ is finite if and only if the sequence is eventually Sylvester.*

Proof. Assume first that the recurrence is not eventually Sylvester. Then $u_n \rightarrow \infty$. For every sufficiently late s , Lemma 7.7 with $A = 3$ gives an odd $Q = p^k \mid v_s$ with $Q > (H_s + 2)^3$, hence $Q \geq 16$ and $pQ > 4R_s$. Set $L = pQ/2$ and take its first crossing. The preceding theorem applies. Because $Q \geq p$ and $p \geq 3$,

$$\frac{8p}{L} = \frac{16}{Q} \leq 1, \quad \frac{8p + 8}{\sqrt{L}} \leq 8\sqrt{2}(1 + 1/p) < 16.$$

Dividing the preceding counting inequality by L therefore gives

$$1 \leq 16 \frac{|J|}{\sqrt{L}} + 4 \frac{X}{L} \leq 16 \sum_{n \in J} \left(\frac{1}{\sqrt{u_n}} + \frac{d_n - 2}{u_n} \right),$$

since $u_n < L$ for $n < \tau$ and $d_n \geq 3$ on J . Thus an arbitrarily late finite window contributes at least $1/16$ to $\mathcal{E}$, contradicting convergence. No disjointness of the windows is needed: every window lies in a tail of the nonnegative series. Under the eventual Sylvester recurrence, $u_n = 1$ eventually, so there are no late records and $\mathcal{E}$ is finite. Finally, term by term at a record,

$$\mathbf{1}_{d_n \geq 3} u_n^{-1/2} + (d_n - 2)_+ u_n^{-1} \leq 2(d_n - 2)_+ u_n^{-1/2}.$$

Hence finiteness of the second series implies finiteness of the first, and convergence in the converse direction again follows from eventual Sylvester behaviour. $\square$

Formalisation. The plectis-erdos-lean repository contains [the integer inequality counting crossings before a prime power is lost](#) and its counting lemmas. In Lean, [the two-unit record-rise implication](#) assumes that arbitrarily late s admit an odd $p^k \mid v_s$ with $3R_s < p^k$; Lemma 7.7 supplies such prime powers.

7.4 Bounds on the error and on the original sequence

Theorem 7.11 (slow negative part). [Lean · Comparator](#) *Let $a, C, D : \mathbb{N} \rightarrow \mathbb{N}$ satisfy (19) with $a_n > 1$, $C_n > 0$, $D_0 \geq 1$, under vanishing relative error. Suppose that for some $\delta \in (0, 1)$ and all large n with $E_n < 0$ one has $-E_n \leq (1 - \delta)\ell(C_n)$. Then $E_n = 0$ for all large n , and $a_{n+1} = a_n^2 - a_n + 1$ for all large n .*

Theorem 12.1 is the case of a constant bound, since a constant is eventually below $(1 - \delta)\ell(C_n)$ on a nonzero tail, where $C_n \rightarrow \infty$.

Proof. Suppose the error is not eventually zero. The auxiliary lower bound in the proof of Theorem 7.3 applies to this exact integer orbit: it used only $a_n > 1$, $C_n > 0$, $D_0 \geq 1$ and vanishing relative error. It gives $\limsup_n (H_{n+1} - H_n)/\ell(H_n) \geq 1$. On the other hand, the hypothesis and $C_n \leq H_n$ give

$$H_{n+1} - H_n \leq (-E_n)_+ \leq (1 - \delta)\ell(C_n) \leq (1 - \delta)\ell(H_n)$$

for all large n , a contradiction. Thus $E_n = 0$ eventually, and Theorem 5.7 gives the Sylvester recurrence. $\square$

The corresponding pointwise-rise argument also has formalised arithmetic lemmas: [persistence of a common divisor](#), [divisibility of every later error by \$\gcd\(a_n, D_n\)\$](#) , the [landing lemma](#) and the [resulting exclusion](#). The count of indices and the simultaneous choice of constants are written out in the earlier proof. Corollary 7.4 includes the boundary coefficient 1, while the preceding theorem on the running maximum also discounts rises that recover an earlier decrease. The present statement is retained for its direct bound on E_n .

To compare the bounded hypothesis with the classical criteria, number the original sequence from 1 in this subsection and put $P_n = \prod_{1 \leq j < n} a_j$. Then

$$\frac{P_n}{a_n} \left(\frac{a_n^2}{a_{n+1}} - 1 \right) = \frac{P_{n+1}}{a_{n+1}} - \frac{P_n}{a_n}.$$

Thus the bound concerns upward increments, not boundedness of P_n/a_n . The quadratic growth limit controls only the ratio of consecutive terms. Sequences satisfying the eventual Sylvester recurrence satisfy the bound, as do exact-square sequences $a_n = b^{2^{n-1}}$ with $b \geq 2$, for which the difference is zero. The latter never satisfies the Sylvester recurrence, so the bounded criterion below recovers irrationality of its reciprocal sum. The rounded examples after Corollary 7.13 show more generally how a term c/n in the ratio produces increments of order n^{c-1} . The theorem allows a finite positive upper limit where the classical product criterion requires a nonpositive one; neither bound is derived from the original problem alone.

Theorem 7.12 (bounded or slowly growing increments of the product ratio).

Lean · Comparator Let $a_1 < a_2 < \dots$ be positive integers with $a_{n+1}/a_n^2 \rightarrow 1$ and $\sum_{n \geq 1} 1/a_n = p/q$, where p, q are positive integers. Put

$$Q_n = \frac{a_1 a_2 \cdots a_{n-1}}{a_n} \left(\frac{a_n^2}{a_{n+1}} - 1 \right).$$

If $\limsup_n Q_n < \infty$, the sequence is eventually Sylvester. The same conclusion holds if, for some $\delta > 0$ and all large n ,

$$Q_n \leq \frac{1 - \delta}{q} \ell(a_1 \cdots a_{n-1}/a_n).$$

For the bounded clause, use the stated denominator q , put $P_n = \prod_{j < n} a_j$, $x_n = \sum_{k \geq n} 1/a_k$, and form the integer tail $D_n = qP_n$, $C_n = D_n x_n$, $E_n = D_n - (a_n - 1)C_n$. Exact cancellation gives

$$E_n + qQ_n = qP_n \left(\frac{1}{a_{n+1}} - (a_n - 1) \sum_{k \geq n+2} \frac{1}{a_k} \right).$$

The expression in parentheses is positive eventually, since $\sum_{k \geq n+2} 1/a_k \leq 4/a_{n+1}^2$ and $a_{n+1} > 4(a_n - 1)$ eventually. Thus $E_n \geq -qQ_n$, and $\limsup Q_n < \infty$ supplies the eventual lower bound required by Theorem 12.1. This proves the bounded clause without first estimating the absolute size of the comparison error.

For the slow-growth clause and the later summability comparisons, the same exact formula gives

$$0 < E_n + qQ_n \leq qP_n/a_{n+1} = O(P_n/a_n^2)$$

eventually. The ratio test in Section 6 shows that these errors have a convergent sum, hence tend to zero. The factor $1/q$ in the slow-growth clause cancels the factor q in $E_n + qQ_n = o(1)$. Suppose the sequence is not eventually Sylvester. Absorption and vanishing relative error give $C_n \rightarrow \infty$, and the tail estimate $C_n \sim qP_n/a_n$ implies

$$\frac{\ell(C_n)}{\ell(P_n/a_n)} \rightarrow 1.$$

For $0 < \delta < 1$, the assumed bound and $E_n + qQ_n = o(1)$ consequently give $(-E_n)_+ \leq (1 - \delta/2)\ell(C_n)$ eventually. This contradicts Theorem 7.11. If $\delta \geq 1$, the hypothesis gives $Q_n \leq 0$ eventually and the bounded clause already applies.

The same comparison permits any positive real clearing factor $F_n \leq qP_n$: multiplying $E_n + qQ_n = o(1)$ by $F_n/(qP_n)$ gives

$$F_n - (a_n - 1)F_n x_n + \frac{F_n}{a_n} \left(\frac{a_n^2}{a_{n+1}} - 1 \right) = o(1).$$

Neither $F_n x_n$ nor $F_n - (a_n - 1)F_n x_n$ need be integral. The LCM choice $F_n = L_n$ makes both integers, but its update includes the factor ρ_n of Section 6. The short note uses only this LCM specialisation; no general clearing factor is needed in its bounded-increment proof.

Attribution. Erdős and Straus require $\limsup \leq 0$ in the corresponding criterion, with the least common multiple in place of the product and the growth factor one index later, so their quantity is $[a_1, \dots, a_n] a_{n+1}^{-1} (a_{n+1}^2/a_{n+2} - 1)$ [1, Theorem 3, p. 132]; Koizumi's Corollary 4(1) uses the product form [12, Cor. 4(1), pp. 14–15]. The bounded clause follows from Theorem 12.1. For Koizumi's product expression it replaces a nonpositive upper limit by an arbitrary finite upper limit. The material distinction here is product versus least common multiple. Reindexing $r = n+1$ makes the classical LCM expression $[a_1, \dots, a_{r-1}] a_r^{-1} (a_r^2/a_{r+1} - 1)$, exactly the expression in the short note's LCM corollary.

The comparison $E_n + qQ_n = o(1)$ has its classical predecessor in Koizumi's proof of Corollary 4(1) [12, (11)–(12), p. 15].

Corollary 7.13 (the $1/n$ threshold). *Lean · Comparator* Let $a_1 < a_2 < \dots$ be positive integers with $a_{n+1}/a_n^2 \rightarrow 1$ and $\sum_n 1/a_n \in \mathbb{Q}$. If

$$\limsup_{n \rightarrow \infty} n (a_n^2/a_{n+1} - 1)_+ < 1,$$

then $a_{n+1} = a_n^2 - a_n + 1$ for all large n . The same conclusion holds if, for some $K \geq 0$ and $\varepsilon > 0$,

$$\frac{a_n^2}{a_{n+1}} - 1 \leq \frac{1}{n} + \frac{K}{n^{1+\varepsilon}} \quad \text{eventually.}$$

In particular the one-sided bound by $1/n$ is included.

Proof. Put $\gamma_n = a_n^2/a_{n+1} - 1$ and $t_n = (\prod_{j < n} a_j)/a_n$, so that $t_{n+1}/t_n = 1 + \gamma_n$. Choose $r \in (0, 1)$ and N with $\gamma_n^+ \leq r/n$ for all $n \geq N$. Then

$$t_n \leq t_N \prod_{k=N}^{n-1} \left(1 + \frac{r}{k}\right) = O(n^r),$$

so $(t_n \gamma_n)_+ = t_n \gamma_n^+ = O(n^{r-1})$ tends to zero and $\limsup_n t_n \gamma_n \leq 0$. Koizumi's Corollary 4(1) [12, pp. 14–15] gives the first conclusion. For the second, divide t_n by n to obtain

$$\frac{t_{n+1}/(n+1)}{t_n/n} = \frac{1 + \gamma_n}{1 + 1/n} \leq 1 + \frac{K}{n^\varepsilon(n+1)}.$$

The product of the upper bounds converges, so t_n/n is bounded above. Consequently $t_n(\gamma_n)_+ = O(1)$, and Theorem 7.12 applies. $\square$

The inclusive clause requires more than $\limsup n(\gamma_n)_+ \leq 1$. That weaker condition does not give the product estimate: for example, the scalar sequence $\gamma_n = (1 + 1/\log n)/n$ has $t_n \asymp n \log n$ and $t_n \gamma_n \asymp \log n$. This is a limitation of the estimate, not a counterexample to the original problem.

More generally, if $c > 0$, $\varepsilon > 0$ and

$$\frac{a_n^2}{a_{n+1}} = 1 + \frac{c}{n} + O(n^{-1-\varepsilon}),$$

then the consecutive-ratio identity gives $t_n \sim Kn^c$ for some $K > 0$ and $t_{n+1} - t_n \sim Kcn^{c-1}$. To see the constant, take logarithms and subtract $c \log(1 + 1/n)$; the remainder

is absolutely summable. Hence the increments are bounded for $c \leq 1$ and unbounded for $c > 1$. These rates are realised by increasing integer sequences: set $a_{n+1} = \lceil na_n^2/(n+c) \rceil$ and choose the seed large enough that $a_{n+1} \geq a_n^2/(1+c) \geq 2a_n$ throughout. The rounding remainder in $[0, 1)$ gives an error in the ratio bounded by $(1+c)^2/a_n^2$, smaller than every fixed inverse power of n . This calculation supplies the family comparison behind the two rounded examples retained in the short note. It concerns their growth and does not assume rationality.

The second clause has a concrete application outside Koizumi's nonpositive-upper-limit product condition. The sequence $a_1 = 4$, $a_{n+1} = \lceil na_n^2/(n+1) \rceil$ begins 4, 8, 43, 1387, ... and satisfies

$$a_n \geq 2 \cdot 2^{2^{n-1}}, \quad 0 \leq 1 + \frac{1}{n} - \frac{a_n^2}{a_{n+1}} < \frac{4}{a_n^2}.$$

The lower bound follows from $a_{n+1} \geq a_n^2/2$; the upper error bound follows by writing the rounding remainder in $[0, 1)$. Consequently $t_n \sim Kn$ and $t_n \gamma_n \rightarrow K > 0$ by the product comparison above. The corollary proves that the reciprocal sum is irrational, since the eventual Sylvester recurrence would give $\gamma_n = O(1/a_n)$ rather than $\gamma_n \sim 1/n$.

The strict clause includes Koizumi's sufficient rate $1 + o(1/n)$ [12, Remark 3, p. 16]; the inclusive clause allows a bounded positive increment rather than requiring a non-positive upper limit.

7.5 Signs of the error and of the growth ratio

We keep the preceding one-based indexing. The growth defect need not have the opposite sign to E_n : the next identity displays the correction, including its sign under the eventual Sylvester recurrence.

Proposition 7.14 (comparison of the two signs). *Lean · Comparator* The factor $M_n = D_n/L_n$ divides $G_n = \gcd(C_n, D_n)$. Thus G_n/M_n is a positive integer and $E_n/M_n = (G_n/M_n)\tilde{e}_n$ is an integer with the same sign as E_n . The sign of the growth ratio minus one also depends on a correction term. The exact recurrences give, whenever $a_{n+1}C_nC_{n+1} \neq 0$,

$$\frac{a_n^2}{a_{n+1}} - 1 = -\frac{E_n}{C_n} + \Lambda_n, \quad \Lambda_n = \frac{(1 - E_n/C_n)(a_n - 1 + E_{n+1}/C_{n+1})}{a_{n+1}}, \quad (22)$$

and, under the standing positive rational-tail hypotheses, $0 < \Lambda_n < 3/a_n$ for all sufficiently large n . The Erdős–Straus quantity of Theorem 3 is

$$Z_n^{\text{ES}} = \frac{[a_1, \dots, a_n]}{a_{n+1}} \left(\frac{a_{n+1}^2}{a_{n+2}} - 1 \right).$$

Its least common multiple includes a_n but not the clearing denominator q . It is not $Q_n = (P_n/a_n)\gamma_n$ from the preceding subsection, nor $L_n\gamma_{n+1}/a_{n+1}$ under our convention $L_n = \text{lcm}(q, a_1, \dots, a_{n-1})$. Being a positive multiple of the next growth defect, Z_n^{ES} has the sign of $\Lambda_{n+1} - E_{n+1}/C_{n+1}$. For all sufficiently large n , it is positive when $E_{n+1} \leq 0$; for $E_{n+1} > 0$, it is negative precisely when $E_{n+1}/C_{n+1} > \Lambda_{n+1}$. Under the eventual Sylvester recurrence, $E_n = 0$ and $\Lambda_n = (a_n - 1)/a_{n+1} > 0$.

Proof. Since both $C_n = D_n x_n$ and $L_n x_n$ are integers, $M_n = D_n/L_n$ divides C_n as well as D_n . It therefore divides G_n . For (22), write $\theta_n = E_n/C_n$; Proposition 4.1 gives $C_{n+1} = C_n(1 - \theta_n)$ and $D_n/C_n = a_n - 1 + \theta_n$, so $a_{n+1} - 1 + \theta_{n+1} = a_n D_n/C_{n+1} = a_n(a_n - 1 + \theta_n)/(1 - \theta_n)$. Multiplying the asserted identity by a_{n+1} and substituting reduces it to $a_{n+1} + a_n - 1 + \theta_{n+1} = a_n^2/(1 - \theta_n)$, which is the displayed relation with a_n added to both sides. The bound follows from $\theta_n \rightarrow 0$ and $a_{n+1} \geq a_n^2/2$. On Sylvester's sequence $\theta_n = 0$ and $a_{n+1} = a_n^2 - a_n + 1$, so $a_n^2/a_{n+1} - 1 = (a_n - 1)/a_{n+1} = \Lambda_n > 0$. $\square$

A nonpositive upper limit admits positive values tending to zero; it is not an eventual pointwise sign condition. For the *product* quantity $Q_n = (P_n/a_n)\gamma_n$, the identity $E_n + qQ_n = o(1)$ shows that eventual $E_n \geq 0$ implies $\limsup Q_n \leq 0$, as in Koizumi's Corollary 4(1) [12, pp. 14–15]. For the classical LCM expression Z_n^{ES} , the corresponding integer error is also taken one index later. Its published hypotheses are those of Erdős–Straus Theorem 3 [1, p. 132]. Reindexing relates it to an LCM-cleared error, not to E_n with the unchanged product factor qQ_n .

A comparison with $\mathcal{B}$ -free integers. For a fixed family $\mathcal{B} = \{m_i\}$, the integers divisible by none of the m_i are called $\mathcal{B}$ -free integers. With pairwise coprime moduli and $\sum_i 1/m_i < \infty$, this is the standard setting of [21, §1.2]. The next proposition is an elementary interval count in that setting. It tests what can follow from avoiding whole multiples alone. Unlike the reciprocal-tail argument, it imposes all the moduli from the outset and has no denominator recurrence or cancellation factors.

Proposition 7.15 (an elementary interval bound). *Lean · Comparator* Let $m_0 < m_1 < \dots$ be pairwise coprime integers at least 2 with $\theta = \sum_i 1/m_i < 1$. For all integers $x \geq 1$ and $L \geq 1$ satisfying $L > k/(1 - \theta)$, where $k = \#\{i : m_i \leq x + L\}$, the interval $[x, x + L)$ contains an integer divisible by no m_i . If also $\ell(m_i) = i + O(1)$, then for every $\epsilon > 0$ there are an index T and a strictly increasing sequence of positive integers (u_n) such that

$$m_i \nmid u_n \text{ for every } i \geq T \text{ and every } n, \quad u_{n+1} - u_n \leq (1 + \epsilon)\ell(u_n) \text{ eventually.}$$

Proof. The interval $[x, x + L)$ contains exactly L integers. Each lies in $[1, \infty)$ and is smaller than $x + L$, so a modulus exceeding $x + L$ divides none of them. Each of the k remaining moduli divides at most $L/m_i + 1$ of them, so the covered count is at most $L\theta + k < L$. This interval count itself does not use pairwise coprimality; that condition is needed for the exact CRT proportions in the later gap theorem.

For the second assertion, choose a tail of the family and reindex it so that $\theta < \epsilon/(1 + \epsilon)$. If $k(z) = \#\{i : m_i \leq z\}$, the hypothesis $\ell(m_i) = i + O(1)$ gives $k(z) \leq \ell(z) + C$ for all large z and some constant C . Choose

$$(1 - \theta)^{-1} < \rho < 1 + \epsilon, \quad L(y) = \lceil \rho \ell(y) \rceil.$$

Then $L(y) = O(\ell(y)) = o(y)$, and the definition of ℓ gives $\ell(y + 1 + L(y)) = \ell(y) + o(1)$. Consequently, for all large integers y ,

$$\frac{k(y + 1 + L(y))}{1 - \theta} \leq \frac{\ell(y) + C + o(1)}{1 - \theta} < \rho \ell(y) \leq L(y),$$

while $L(y) \leq (1 + \epsilon)\ell(y)$. The first part also supplies an admissible u_0 beyond this threshold. Apply it successively with $x = u_n + 1$ and length $L(u_n)$, and choose u_{n+1} in the resulting window. The sequence is strictly increasing, avoids every retained modulus, and has the required rise bound. $\square$

The small-increment sequence avoids only the retained moduli $m_i, i \geq T$. Discarding the prefix makes their reciprocal sum small. For the full family, Theorem 14.7 instead gives the coefficient $\prod_i (1 - 1/m_i)^{-1} > 1$ for the increasing enumeration of integers avoiding all the moduli.

Proposition 7.15 concerns avoidance of whole multiples, not coprimality to a composite modulus. The distinction matters: no integer in $[2, 5)$ is coprime to 30, although none is divisible by 30. Thus this proposition alone does not disprove a coprime-walk extension of Theorem 11.3. Proposition 14.6 below gives a separate counterexample using sparse *prime* moduli. Under the additional scale $\ell(m_j) = j + O(1)$, Theorem 14.7 determines the exact maximal-gap coefficient for these $\mathcal{B}$ -free integers. Its proof uses a finite sieve estimate and the Chinese remainder theorem, not an orbit construction. Neither static construction supplies an exact reciprocal-tail orbit.

Conditions on a possible counterexample. Together, these criteria give necessary conditions on a counterexample. For every fixed $\delta \in (0, 1)$, it must have

$$(1 - \delta)\ell(C_n) \leq -E_n = o(C_n) \quad \text{at infinitely many } n.$$

It must also have a divergent sum of relative increases, unbounded negative error scaled by the previous maximum, and infinitely many record jumps of the reduced numerator of size at least three, with $h_n = 1$. These are necessary conditions, not a construction of a counterexample. The residue-saturation lemma shows precisely one limitation of the fixed-old-modulus method; it is not an impossibility theorem about every future approach. A useful next target is interaction between new prime support, three consecutive numerators and the global bound on numerator growth.

8 Nonnegative errors

The scalar update already settles nonnegative errors. With indices starting at zero, $C_{n+1} = C_n - E_n$ makes C_n nonincreasing, and a decreasing sequence of natural numbers must stabilise.

Theorem 8.1 (descent). *Lean · Comparator* Let $C, E : \mathbb{N} \rightarrow \mathbb{N}$ satisfy $C_{n+1} + E_n = C_n$ for every n . Then $E_n = 0$ for all sufficiently large n .

Proof. The recurrence gives $C_{n+1} \leq C_n$. Choose an index at which the natural-number sequence reaches its minimum. Every later value equals that minimum, and $C_{n+1} + E_n = C_n$ then gives $E_n = 0$. $\square$

The hypothesis is exactly one-sidedness: C and E take values in $\mathbb{N}$. If $E_n < 0$ infinitely often, the numerator rises at those indices and this descent argument no longer applies. The next two sections first treat constant and periodic negative errors.

9 Constant negative errors

A constant negative error makes the numerator grow linearly. Such behaviour can persist for arbitrarily long finite intervals, even on genuine rational orbits, by the flat-transient theorem in the working report on Problem #243 [9, “Flat-transient theorem”]. We exclude infinite constant and periodic negative tails below. These exclusions supply no bound for the finite transient lengths.

Suppose the error is negative with a constant magnitude, $E_n = -m$ for a fixed $m > 0$ and every n . By Proposition 4.1 the numerator increases by m at each step, so $C_n = c + nm$ with $c = C_0$, and the definition of the error becomes the identity

$$D_n + m = (a_n - 1)(c + nm). \quad (5.1)$$

Together with $D_{n+1} = a_n D_n$ this is a closed system in (a, D) . We shall prove that it has no infinite natural-number solution with $a_n \geq 2$, after examining a finite example.

Example 9.1 (a nonintegral second step). Take $m = c = 1$, so that $C_n = 1 + n$ and (5.1) reads $D_n + 1 = (a_n - 1)(1 + n)$, and take $D_0 = 1$. Each step is now determined: at $n = 0$ the equation reads $2 = (a_0 - 1) \cdot 1$, so $a_0 = 3$, and $D_1 = a_0 D_0 = 3$; at $n = 1$ it reads $4 = (a_1 - 1) \cdot 2$, so $a_1 = 3$, and $D_2 = a_1 D_1 = 9$; at $n = 2$ it reads $10 = (a_2 - 1) \cdot 3$, which has no integer solution, and the orbit stops. Longer prefixes occur for other starting values. For $2 \leq a_0 < 5000$, the exact residue search in Appendix B gives a maximum of 17 successful updates, hence 18 multiplier values including a_0 .

Theorem 9.2 (no constant negative magnitude). *Lean · Comparator* For any $m, c \in \mathbb{N}$ with $m > 0$, there is no pair of sequences $a, D : \mathbb{N} \rightarrow \mathbb{N}$ with $a_n \geq 2$ for all n satisfying $D_{n+1} = a_n D_n$ and (5.1). The same holds if the identity is assumed only from some index onwards.

Proof when c and m are coprime. Assume first $\gcd(c, m) = 1$. Every multiplier must share a prime with m , but each such prime can occur in at most one multiplier.

Every a_j shares a prime with m . Suppose $\gcd(a_j, m) = 1$. Then m is invertible modulo a_j , so some n has $c + nm \equiv 0$, that is $a_j \mid C_n$; and $a_j \mid D_n$ for every $n > j$, since D is multiplicative with a_j among its factors. Choosing such an n beyond j and reading (5.1) modulo a_j gives $a_j \mid m$, contradicting $\gcd(a_j, m) = 1$ and $a_j \geq 2$.

A prime divisor of m occurs in at most one a_j . Suppose $p \mid m$ and $p \mid a_j$. For $n > j$ we have $p \mid D_n$, so (5.1) gives $(a_n - 1)C_n \equiv 0 \pmod{p}$. Now $C_n = c + nm \equiv c$, and $p \nmid c$ because $p \mid m$ and $\gcd(c, m) = 1$; hence $p \mid a_n - 1$, so $p \nmid a_n$. Thus p cannot divide any later multiplier.

This injects infinitely many multipliers into the finite set of prime divisors of m , a contradiction. $\square$

The case $\gcd(c, m) = 1$ is the *exclusion when c and m are coprime*. The special case $m = c = 1$, worked through in Example 9.1, where m has no prime divisors at all and the first fact is immediately contradictory, is recorded separately as the *case $c = m = 1$* .

Removing the scale. For general c , put $g = \gcd(c, m)$. Equation (5.1) shows $g \mid D_n$ for every n , so dividing D, c and m by g leaves a system of the same shape with coprime data, which the previous case excludes. $\square$

The formal statements are the [exclusion at every scale](#) and the [eventual exclusion](#). The latter follows by shifting the orbit to the first index of constant error.

Theorem 9.2 excludes an error that is eventually constant and negative, at every magnitude and every scale. Constancy fixes both the possible prime divisors and the congruence used at later indices. For varying magnitudes, even when their prime divisors belong to a fixed finite set, this argument does not supply the same later congruence. It therefore does not settle that case.

10 Periodic negative errors

Let the negative error have periodic magnitude $e_n = -E_n > 0$, so $e_{n+h} = e_n$ and $C_{n+1} = C_n + e_n$. Summing over a period gives $C_{n+h} = C_n + M$, where the positive integer $M = \sum_{j=0}^{h-1} e_j$ is the *drift*. Periodicity makes this sum independent of the starting index.

At $h = 1$ this says that the magnitude is constant and that M is its value, which is the situation of Section 9; Theorem 9.2 already excludes it without using the pointwise bound $e_n < a_n$ in the proof below. The new case is $h \geq 2$.

For a constant error, the proof used the finitely many prime divisors of its magnitude. For a periodic error we use the prime divisors of the increase M over one period. The required divisibility facts are as follows. Every multiplier a_j divides each later denominator, by [persistence of a multiplier](#). If a prime divides both D_n and a_n , the numerator update makes it divide C_{n+1} , the [one-step divisibility implication](#). Once it divides both numerator and denominator, it divides every later numerator, denominator and error, by [persistence of a common divisor](#).

Periodicity carries this divisibility back to every phase. Suppose $d \mid M$ and $d \mid a_i, a_j$ with $i < j$. Then $d \mid D_j$, and $C_{j+1} = a_j C_j - D_j$ gives $d \mid C_{j+1}$. The common divisor persists in both sequences and in every later error. For any phase r , choose k with $r + kh \geq j + 1$; periodicity gives $d \mid e_{r+kh} = e_r$. Taking $kh \geq j + 1$ also gives $d \mid C_{kh} = C_0 + kM$, hence $d \mid C_0$. This is the [repeated-divisor implication](#); it uses the [period relation](#) and the [increase over successive periods](#), each iterated through whole periods. A repeated prime can therefore be divided out of the entire system. This explains the induction on M in the proof.

Theorem 10.1 (no periodic negative magnitude). [Lean](#) · [Comparator](#) *Let $a, D, C, e : \mathbb{N} \rightarrow \mathbb{N}$ with $a_n \geq 2$, $e_n > 0$ and $e_n < a_n$ for every n , satisfying*

$$D_{n+1} = a_n D_n, \quad C_{n+1} = C_n + e_n, \quad D_n + e_n = (a_n - 1)C_n,$$

and suppose $e_{n+h} = e_n$ and $C_{n+h} = C_n + M$ for some $h > 0$ and $M > 0$. This is impossible.

Proof. We use strong induction on the drift M . A prime divisor of M that recurs among the multipliers need not give an immediate contradiction. It instead divides the whole orbit, allowing us to reduce M by that prime and apply the induction hypothesis.

Suppose first that no prime divisor of M is a common divisor of C_0 and of every magnitude. The repeated-divisor implication applies in the contrapositive: a prime divisor of M occurring in two of the multipliers would be exactly such a common divisor, so each prime divisor of M occurs in at most one multiplier. On the other hand, every multiplier shares a prime with M . Indeed, if $\gcd(a_j, M) = 1$, choose $k \geq 1$ so that

$a_j \mid C_j + kM = C_{j+kh}$. The same multiplier divides D_{j+kh} , and periodicity gives $e_{j+kh} = e_j$. The identity at $j+kh$ therefore implies $a_j \mid e_j$, contrary to $0 < e_j < a_j$. Thus infinitely many multipliers would require distinct prime divisors of the fixed integer M , a contradiction.

Otherwise some prime p divides M , divides C_0 , and divides every magnitude. Then $p \mid C_n$ for every n , since $C_{n+1} = C_n + e_n$ and both summands on the right are divisible by p , and the same identity $D_n + e_n = (a_n - 1)C_n$ then gives $p \mid D_n$ for every n . Divide D , C and e by p . The multipliers are untouched, so $a_n \geq 2$ and $e_n < a_n$ persist and the magnitudes stay positive; the three recurrences are homogeneous in (D, C, e) and so survive; the period is still h ; and the drift becomes $M/p < M$. The inductive hypothesis applies. $\square$

The first of the two cases above is the [exclusion when no prime of \$M\$ divides all phases](#), the induction is the [exclusion at every scale](#), and the eventual form is the [eventual exclusion](#).

The phase-by-phase proof uses $e_n < a_n$ to prevent a multiplier from dividing its own nonzero phase magnitude. This is not a rescaling of the orbit. Nevertheless, for an infinite solution of (19) with periodic positive magnitudes, the bound follows eventually from the other assumptions. The identity gives $C_n > 0$, and periodicity gives

$$C_n = \frac{M}{h}n + O(1), \quad \sup_n e_n < \infty,$$

and D_0 cannot be zero: otherwise all D_n vanish and the defining identity gives $e_n = (a_n - 1)C_n \geq C_n$, contradicting these bounds. Thus $D_n \geq 2^n D_0$, and

$$a_n - 1 = \frac{D_n + e_n}{C_n} \rightarrow \infty.$$

Consequently $e_n < a_n$ eventually. Applying the eventual exclusion therefore rules out periodic positive magnitudes without an independent smallness assumption. The numbered theorem retains the pointwise bound used by its checked phase-by-phase proof. The later bounded-negative theorem also excludes this case.

Remark. For the tail numerators of [12], the absolute error satisfies $|E_n| < a_n$ eventually. Indeed, under the dictionary in Section 3, Lemma 4(2) gives $-C_n/2 \leq E_n < C_n/2$, hence $|E_n| \leq C_n/2$. The proof of Lemma 4(3) gives $C_{n+1} \leq \frac{3}{2}C_n$ [12, pp. 11–12], hence $C_n \leq C_N(3/2)^{n-N}$ beyond a pseudo-greedy starting index N . On the other hand, convergence of $\sum 1/a_n$ gives $a_n \rightarrow \infty$, and $a_{n+1}/a_n^2 \rightarrow 1$ then gives $a_{n+1} \geq 2a_n$ eventually. Thus $|E_n|/a_n \rightarrow 0$. On a negative-error tail this gives $e_n < a_n$.

For integer solutions of (19), the same conclusion follows when $a_n > 1$, $C_n > 0$ and $E_n/C_n \rightarrow 0$ hold: the calculation in Section 4 already derives the reciprocal series and its growth from these hypotheses. Summability is not an independent assumption in that setting. The bound $e_n < a_n$ is still only eventual, whereas Theorem 10.1 states it at every index; its eventual form, cited above, is therefore the applicable one. For a periodic magnitude, boundedness of e_n and divergence of a_n already suffice.

11 Bounded increases and coprimality

For nonperiodic errors, the Chinese remainder theorem replaces the fixed period. It produces a block of consecutive forbidden values. An unbounded numerator with bounded upward steps must enter that block when it first crosses its lower end.

Lemma 11.1 (shifted blocks of consecutive multiples). *Lean · Comparator* Let $m_0, \dots, m_{B-1}$ be pairwise coprime and at least 2. For every bound there is a t beyond it with $m_i \mid t + i$ for each $i < B$.

Proof. Solve $t \equiv -i \pmod{m_i}$ simultaneously by the Chinese remainder theorem. Adding a sufficiently large multiple of $\prod_i m_i$ puts the solution beyond the prescribed bound. $\square$

Lemma 11.1 matches each modulus m_i to its own multiple $t + i$ inside a window of B consecutive integers. Matchings of a set of integers to distinct multiples in an interval are the subject of Erdős Problem #650, solved by van Doorn, Li and Tang: for every set S of k positive integers, every open interval of length $2 \max S$ contains distinct multiples of at least $\min(k, \lceil 2\sqrt{k} \rceil)$ elements of S , and this count is optimal [11, Theorem 2.1]. Their extremal construction uses the same Chinese remaindering for moduli that need not be pairwise coprime, with their Claim 3.2 supplying the divisibility of residue differences by greatest common divisors that the generalised theorem requires [11, pp. 4–5]. The first-crossing argument below needs only the pairwise coprime case.

Example 11.2 (a block of three consecutive multiples). Take $B = 3$ and $(m_0, m_1, m_2) = (3, 4, 5)$. The congruences $t \equiv 0 \pmod{3}$, $t \equiv 3 \pmod{4}$ and $t \equiv 3 \pmod{5}$ hold exactly when $t \equiv 3 \pmod{60}$. At $t = 3$ the block is 3, 4, 5 with $3 \mid 3$, $4 \mid 4$ and $5 \mid 5$; at $t = 63$ it is 63, 64, 65 with $3 \mid 63$, $4 \mid 64$ and $5 \mid 65$. A sequence of natural numbers that starts at 0, tends to infinity and rises by at most 3 at each step cannot step over the block $\{3, 4, 5\}$: it has a first value at least 3, that value is at most 5, and every member of $\{3, 4, 5\}$ is divisible by one of the three moduli.

Theorem 11.3 (bounded increases and coprimality to earlier moduli). *Lean · Comparator* Let $u : \mathbb{N} \rightarrow \mathbb{N}$ tend to infinity with $u_{n+1} \leq u_n + B$ for a fixed integer $B \geq 1$. Then u cannot remain coprime to infinitely many fresh pairwise coprime moduli: there is no family of pairwise coprime $m_i \geq 2$, one for each index, such that $\gcd(m_i, u_t) = 1$ whenever $i < t$.

The CRT block must lie above the initial segment where some moduli are not yet available. No monotonicity of u is assumed.

Proof. Choose $m_0, \dots, m_{B-1}$ and use Lemma 11.1 to find $t > \max(u_0, \dots, u_B)$ with $m_i \mid t + i$ for $0 \leq i < B$. Since $u_n \rightarrow \infty$, there is a first $n > B$ with $u_n \geq t$. Minimality and the rise bound give

$$t \leq u_n \leq u_{n-1} + B < t + B.$$

Hence $u_n = t + i$ for some $0 \leq i < B < n$, so $m_i \mid u_n$. The inequality $i < n$ now permits the avoidance hypothesis, which says $\gcd(m_i, u_n) = 1$; this contradicts $m_i \geq 2$. The argument uses the first crossing of the CRT block, not monotonicity of u . $\square$

Only unboundedness above is used to obtain this first crossing. Divergence is stated because it holds for the reduced rational tail in the application. This distinction also explains why the weighted proof in Section 6 works with an unbounded running maximum, without assuming that its numerator tends to infinity.

Divergence forces u to reach the forbidden block, and the uniform rise bound prevents it from jumping over the block. A bound only along a subsequence would leave the intervening steps unrestricted. Without a scale restriction on the moduli, Proposition 14.6 gives an unbounded coprime walk with rises $o(\log \log u_n)$. The same forbidden-block crossing, under a two-sided bound on the error, appears in the proof of Bado's Theorem 5.1 [8, pp. 4–5], posted in September 2026.

In the application, m_i is the i th multiplier. Its coprimality is known only for later numerators, which explains the condition $i < t$.

To apply the preceding theorem, we reduce C_n/D_n to lowest terms and require that no further common factor be cancelled. In that case the reduced numerators and denominators still satisfy the original recurrences. The relevant coprimality is as follows.

Proposition 11.4 (persistent coprimality). *Lean · Comparator* Suppose that $\gcd(u_n, v_n) = 1$ at every index and

$$u_{n+1} + v_n = a_n u_n, \quad v_{n+1} = a_n v_n.$$

Then $\gcd(a_n, v_n) = 1$, the a_n are pairwise coprime, and $\gcd(a_i, u_t) = 1$ whenever $i < t$.

Proof. A common prime divisor of a_n and v_n would divide both $u_{n+1} = a_n u_n - v_n$ and $v_{n+1} = a_n v_n$, contradicting their coprimality. Hence $\gcd(a_n, v_n) = 1$. For $i < t$, the denominator recurrence gives $a_i \mid v_t$. Combining this with $\gcd(a_t, v_t) = 1$ proves that a_i and a_t are coprime; combining it with $\gcd(u_t, v_t) = 1$ proves that a_i and u_t are coprime. $\square$

These are the [step coprimality](#), the [pairwise coprimality](#), and the [coprimality to each earlier multiplier](#). Applying Theorem 11.3 shows that a numerator satisfying these conditions cannot tend to infinity with uniformly bounded upward increments. This is the [reduced exclusion under bounded upward increments](#), together with its eventual form, the [version with an eventual increment bound](#).

Example 11.5 (coprimality in Sylvester's sequence). Put $u_n = 1$ and $v_n = D_n = 1, 2, 6, 42, 1806, \dots$ as in Example 4.2, with multipliers $a_n = v_n + 1 = 2, 3, 7, 43, 1807, \dots$. Then $\gcd(u_n, v_n) = 1$, $u_{n+1} + v_n = 1 + v_n = a_n u_n$ and $v_{n+1} = a_n v_n$, so Proposition 11.4 applies and gives the classical fact that Sylvester's numbers are pairwise coprime. Its numerator is constant, so it satisfies every hypothesis of the exclusion just stated except divergence; since the tail exists, that hypothesis cannot be dropped.

The fraction C_n/D_n in Section 4 need not be in lowest terms. To use the preceding argument, we must show that the common factor stops changing. One uniform bound on the negative magnitudes at arbitrarily late indices suffices: each gcd divides every later gcd, and at a negative index it also divides the nonzero magnitude. Thus a bound at arbitrarily late negative indices bounds the entire gcd sequence. The next proposition justifies dividing by its eventual value.

Proposition 11.6 (the tail gcd stabilises). *Lean · Comparator* Let $a, D, C : \mathbb{N} \rightarrow \mathbb{N}$ satisfy $C_{n+1} + D_n = a_n C_n$ and $D_{n+1} = a_n D_n$, and put $E_n = D_n - (a_n - 1)C_n$. Suppose some fixed integer $B \geq 1$ satisfies $-B \leq E_n < 0$ at infinitely many indices. Then $\gcd(C_n, D_n)$ is eventually constant. Dividing C_n, D_n by its eventual value gives coprime sequences satisfying the same recurrences.

Proof. First, $C_n > 0$ at every index. If $C_n = 0$, the nonnegative recurrence forces $C_{n+1} = D_n = 0$, and both sequences then vanish at all later indices. This contradicts the existence of arbitrarily late negative errors.

The two recurrences give $G_n \mid G_{n+1}$ for $G_n = \gcd(C_n, D_n)$, as in the [gcd divisibility relation](#). Also $G_n = \gcd(C_n, |E_n|)$, since $D_n = E_n + (a_n - 1)C_n$; at a negative index this is the [gcd identity with the negative magnitude](#). For any n , choose $t \geq n$ with $-B \leq E_t < 0$. Then

$$G_n \leq G_t \leq |E_t| \leq B.$$

The entire positive divisibility chain is therefore bounded and eventually constant, by [stabilisation of a bounded divisibility chain](#). In Lean this is the [gcd stabilisation result](#). Dividing both sequences by the eventual value preserves the recurrences and gives coprime numerator and denominator at every later index. $\square$

Other negative errors may exceed B in magnitude. Gcd stabilisation uses the bounded witnesses, whereas the CRT application needs a bound on *every* upward increment.

Vanishing relative error also gives sparsity of the strict gcd increases. The next proposition asserts long finite intervals of constancy, not constancy on an infinite tail. For $a, C, D : \mathbb{N} \rightarrow \mathbb{N}$ satisfying (19) with $C_n > 0$, put

$$G_n = \gcd(C_n, D_n), \quad \Gamma(N) = \#\{0 \leq j < N : G_j < G_{j+1}\}.$$

Proposition 11.7 (vanishing relative error makes strict gcd changes sparse).

Lean · Comparator Let $a, C, D : \mathbb{N} \rightarrow \mathbb{N}$ satisfy $C_{n+1} + D_n = a_n C_n$ and $D_{n+1} = a_n D_n$ with $C_n > 0$, and put $E_n = D_n - (a_n - 1)C_n$, $G_n = \gcd(C_n, D_n)$ and $\Gamma(N) = \#\{0 \leq j < N : G_j < G_{j+1}\}$. If $|E_n|/C_n \rightarrow 0$, then $\Gamma(N) = o(N)$. Moreover, for every starting bound B and block length L , some $n \geq B$ satisfies

$$G_n = G_{n+1} = \dots = G_{n+L}.$$

The first assertion is the [sublinear strict-growth theorem](#); the second is the [arbitrarily late constant-block theorem](#). The proof first converts vanishing relative error into subexponential growth of C_n . Each strict divisibility increase of G_n contributes a factor of at least 2, so $2^{\Gamma(N)} G_0 \leq G_N \leq C_N$. Taking logarithms gives $\Gamma(N) = o(N)$. For $L = 0$ the second assertion is immediate. For $L \geq 1$, if every block of L successive transitions beyond B contained a strict increase, disjoint such blocks would give $\Gamma(N) \geq (N - B)/L - O(1)$, a contradiction.

The quantifiers matter. Proposition 11.6 uses a bound on negative magnitudes at arbitrarily late indices and yields eventual constancy. Proposition 11.7 uses only vanishing relative error and yields arbitrarily late constant blocks of each prescribed finite length. This proof does not establish eventual constancy under that hypothesis, nor does it bound the negative magnitudes.

12 Bounded negative part

The lower bound $E_n \geq -B$ has two uses. At negative indices it bounds the gcd, making it stable. At every index it bounds the upward step $C_{n+1} - C_n = -E_n$. We can therefore combine the preceding coprimality and first-crossing arguments.

Without the denominator recurrence, $C_n = c + bn$, $E_n = -b$ would be a counterexample for any positive integers b, c . Theorem 9.2 excludes this exact-orbit behaviour. The separate summability criterion in the next section needs only the scalar update. For positive integer solutions of (19) with vanishing relative error, both criteria are equivalent to eventual zero error. Neither extra estimate has been derived from the unrestricted hypotheses.

Theorem 12.1 (bounded negative part). *Lean · Comparator* Let $a, C, D : \mathbb{N} \rightarrow \mathbb{N}$ satisfy $a_n > 1$, $C_n > 0$, and

$$C_{n+1} + D_n = a_n C_n, \quad D_{n+1} = a_n D_n \quad (n \geq 0).$$

Put $E_n = D_n - (a_n - 1)C_n$. If

$$\frac{E_n}{C_n} \rightarrow 0 \quad \text{and} \quad E_n \geq -B \quad \text{for all sufficiently large } n$$

for some integer $B \geq 0$, then $E_n = 0$ for all sufficiently large n .

The limit is equivalent, for positive C_n , to the integer condition $K|E_n| < C_n$ eventually for each $K \in \mathbb{N}$. This is the form used by the linked declaration. Its $K = 1$ case gives $|E_n| < C_n$ eventually.

Proof. Suppose that the error fails to vanish eventually. After the point from which $|E_n| < C_n$ always holds, Theorem 5.8 excludes every zero. Hence $|E_n| \geq 1$, and the relative limit forces $C_n \rightarrow \infty$.

There are infinitely many negative indices, since otherwise Theorem 8.1 would give eventual zero. At each sufficiently late negative index, $1 \leq -E_n \leq B$, so $B \geq 1$. Proposition 11.6 makes $G_n = \gcd(C_n, D_n)$ stabilise to an integer $g > 0$. The divided sequences $u_n = C_n/g$, $v_n = D_n/g$ satisfy the coprime recurrences, with $u_n \rightarrow \infty$ and $u_{n+1} - u_n = -E_n/g \leq B$. Their coprimality properties from Proposition 11.4 now contradict Theorem 11.3. $\square$

Corollary 12.2. *Lean · Comparator* Under the hypotheses of Theorem 12.1, the multipliers satisfy $a_{n+1} = a_n^2 - a_n + 1$ for all sufficiently large n .

Proof. Theorem 12.1 gives eventual zero error. Since $C_{n+1} > 0$, Theorem 5.7 gives the recurrence. $\square$

To apply it to Problem 1.1, delete a finite prefix and use Koizumi's Corollary 3 [12, p. 9]. The remaining sequence is the pseudo-greedy expansion of its own reciprocal sum. His Lemma 4 gives positive integer numerators c_n , denominators d_n and errors e_n with

$$c_{n+1} = a_n c_n - d_n, \quad d_{n+1} = a_n d_n, \quad e_n = d_n - (a_n - 1)c_n.$$

The terms a_n eventually exceed one, and $e_n/c_n \rightarrow 0$ by that corollary. The same construction gives $-c_n/2 \leq e_n < c_n/2$. The additional condition still required is an eventual lower bound $e_n \geq -B$. Neither Koizumi's construction nor the constant and periodic exclusions above establish this bound for every rational sum in the problem. The result therefore remains conditional.

13 Finite total relative increase

We can bound a positive integer sequence by multiplying the relative increases of its terms. If their sum converges, that product is finite. Only finitely many upward integer steps can then occur, and descent finishes the argument. No denominator recurrence is needed. For comparison, $C_n = n + 1$ and $C_n = (n + 1)^2$ have relative increases tending to zero but divergent sums. The eventual Sylvester recurrence gives an eventually constant numerator, so it satisfies the summability condition.

Theorem 13.1 (finite sum of relative increases). *Lean · Comparator* Let $C_n \in \mathbb{N}_{>0}$ and $E_n \in \mathbb{Z}$ satisfy $C_{n+1} = C_n - E_n$. If

$$\sum_{n=0}^{\infty} \frac{(-E_n)_+}{C_n} < \infty,$$

then $E_n = 0$ eventually. For an exact reciprocal-tail orbit this implies $a_{n+1} = a_n^2 - a_n + 1$ eventually. No hypothesis $|E_n|/C_n \rightarrow 0$ is required.

Proof. Put $\delta_n = (-E_n)_+/C_n$. The update gives $C_{n+1} \leq C_n(1 + \delta_n)$, so

$$C_N \leq C_0 \prod_{n < N} (1 + \delta_n) \leq C_0 \exp\left(\sum_n \delta_n\right).$$

Choose an integer upper bound K for C_n . Each strict rise of the integer sequence contributes at least $1/K$ to $\sum_n \delta_n$, so there are only finitely many rises. The remaining positive integer sequence is nonincreasing and therefore stabilises, forcing $E_n = 0$. The recurrence follows from Theorem 5.7, since $C_n > 0$. $\square$

The same proof allows any positive real C_0 , provided $E_n \in \mathbb{Z}$: after the finitely many rises, the bounded positive values lie in the finite set $(C_0 + \mathbb{Z}) \cap (0, K]$, so they stabilise. The Lean statement retains integer-valued C_n .

The two indispensable features of this argument are positivity and discrete increments. With $C_n = 1 + 1/(n + 1)$ and $E_n = 1/((n + 1)(n + 2))$, the numerator decreases forever with zero relative-increase sum, but the errors are not integers. With $C_n = -n - 1$ and $E_n = 1$, the increments are integers but positivity fails.

For the gap sequence of the pseudo-greedy expansion, the same criterion, with the same product bound and integer descent, appears in the Erdős Problem a Day working report on Problem #243, dated 12 August 2026 [9, "A global termination criterion"]. Bado's Theorem 11.1 also accounts for the factors removed in LCM clearing [8, Thm. 11.1 and Remark 11.3, p. 9]. In the notation of Section 6, the update gives

$$\log U_N \leq \log U_0 + \sum_{n < N} \frac{(-E_n)_+}{C_n} - \sum_{n < N} \log \rho_n.$$

His sufficient condition is an upper bound on the difference of these two sums. The subtractive term records the decrease caused by a repeated factor in the denominator. It may offset some relative increases; the hypothesis does not require either sum to converge separately. Unlike the scalar criterion above, this argument also uses the pseudo-greedy limit: bounded U_n and $V_n/U_n = E_n/C_n \rightarrow 0$ make the integer V_n eventually zero, after which $\rho_n U_{n+1} = U_n$ gives eventual constancy.

Corollary 3 [12, p. 9] and Lemma 4 [12, pp. 11–12] supply the same positive integer update after a finite restart. Theorem 13.1 therefore applies if the relative-increase sum converges. This is the sole additional hypothesis, asked for in Problem 14.5; the cited construction does not establish it.

Example 13.2 (the product bound at a geometric rate). Suppose $C_0 = 100$ and $(-E_n)_+/C_n \leq 2^{-n-1}$ for every n , so that the sum of relative increases is at most 1; the constraint at $n = 0$ still permits E_0 as negative as -50 . Since $1 + x \leq e^x$,

$$C_N \leq 100 \prod_{n < N} (1 + 2^{-n-1}) \leq 100e < 272$$

for every N . For all sufficiently large n , we have $2^{-n-1} < 1/272$. A strict rise would then force $(-E_n)_+/C_n \geq 1/C_n > 1/272$, a contradiction. Thus C_n is eventually nonincreasing; as a positive integer sequence it stabilises, and $E_n = 0$ thereafter. The constant 272 comes from the total mass and not from any single magnitude, which is why the hypothesis of Theorem 13.1 is summability of $(-E_n)_+/C_n$ and not a bound on it.

14 Further questions

We collect the restrictions on a possible counterexample and the estimates which would exclude it. The constant and periodic arguments concern all-negative tails. They leave open a periodic pattern seen only along the negative indices of a mixed-sign sequence. The unrestricted problem remains unresolved here.

Proposition 14.1 (necessary conditions on a counterexample). [Lean](#) · [Comparator](#) For the integer tail attached to any counterexample to Problem 1.1,

$$E_n \neq 0 \text{ eventually, } \frac{|E_n|}{C_n} \rightarrow 0,$$

and

$$\limsup_{\substack{n \rightarrow \infty \\ E_n < 0}} (-E_n) = \infty, \quad \sum_{n=0}^{\infty} \frac{(-E_n)_+}{C_n} = \infty.$$

In particular, negative indices occur infinitely often. Thus the remaining regime consists of unbounded negative errors along exact reciprocal tails for which the sum of relative increases diverges.

Proof. Koizumi's construction gives vanishing relative error after a finite shift, hence $|E_n| < C_n$ eventually. Absorption then makes $E_n \neq 0$ eventually, since eventual zero would give the Sylvester recurrence. Descent excludes an eventually nonnegative error.

Theorem 12.1 excludes a bounded negative part, and Theorem 13.1 excludes convergence of the sum of relative increases. These statements are unchanged by deleting a finite prefix. $\square$

Proposition 14.1 does not assert infinitely many prime divisors of the error magnitudes. Nor are its numerical conditions alone a reformulation of the problem: the numerator and denominator must satisfy the exact recurrences. With those recurrences, positivity and vanishing relative error, Section 4 supplies the reciprocal-series realisation. The following exact-orbit question is therefore equivalent to the original problem.

Problem 14.2 (unbounded negative errors with divergent relative sum). Exclude, or construct, natural-number sequences (a, D, C) satisfying (19), with $a_n > 1$ and $C_n > 0$, whose multipliers satisfy $\lim_n a_{n+1}/a_n^2 = 1$ and whose error satisfies vanishing relative error, and which is negative infinitely often with magnitudes unbounded along that infinite set and

$$\sum_n \frac{(-E_n)_+}{C_n} = \infty.$$

Such an orbit fails the bounded-negative-part and summable-relative-increase hypotheses. By Proposition 14.1 the integer tail of a counterexample is such an orbit after deletion of a finite prefix, so an exclusion would settle Problem #243. Conversely, a global orbit with the displayed properties has $C_n/D_n \rightarrow 0$ by Section 4. The reciprocal-series realisation then identifies its reciprocal sum as C_0/D_0 . After deletion of a finite prefix the multipliers are strictly increasing, and the infinitely many negative errors rule out a Sylvester tail. It is therefore a counterexample to Problem #243. Finite admissible prefixes alone do not provide such a construction.

Remark. The recurrences alone admit unbounded negative errors for which $\sum_n (-E_n)_+/C_n$ diverges. The following example fails both the relative-error and growth assumptions of Problem 14.2. Put

$$C_n = 2^n, \quad b_0 = 2, \quad b_{n+1} = \frac{1}{2}b_n(b_n + 2), \quad a_n = b_n + 2, \quad D_n = b_n C_n,$$

so that $(b_n) = 2, 4, 12, 84, 3612, \dots$ and $(a_n) = 4, 6, 14, 86, 3614, \dots$. Every b_n is a positive even integer, since $b_n = 2k$ gives $b_{n+1} = 2k(k + 1)$, so all four sequences are $\mathbb{N}$ -valued. They satisfy both recurrences: $C_{n+1} + D_n = 2^n(2 + b_n) = a_n C_n$ and $D_{n+1} = b_{n+1}2^{n+1} = b_n(b_n + 2)2^n = a_n D_n$. Its error is

$$E_n = D_n - (a_n - 1)C_n = b_n 2^n - (b_n + 1)2^n = -C_n,$$

so the negative magnitudes 2^n are unbounded and $\sum_n (-E_n)_+/C_n$ diverges. Here $|E_n| = C_n$ at every index, so both the strict inequality $|E_n| < C_n$ and the vanishing relative error fail. The multipliers satisfy $a_{n+1} = \frac{1}{2}(a_n^2 - 2a_n + 4)$, so $a_{n+1}/a_n^2 \rightarrow \frac{1}{2}$ and the growth hypothesis fails as well. The example therefore says nothing about Problem 1.1, and it does not isolate the error bounds: $E_n = -C_n$ at every index forces that multiplier recurrence, so growth fails with them. It is recorded only to show that the two recurrences alone do not force the required conclusion.

A scalar profile need not satisfy the recurrences. The example $C_n = n^2 + 1$, $E_n = -(2n+1)$ satisfies the numerator update, $|E_n|/C_n \rightarrow 0$, $\log C_n = o(n)$ and $\sum_n (E_n/C_n)^2 < \infty$. It is not an exact reciprocal-tail orbit. Indeed, $C_2 = 5$, $C_3 = 10$ and $C_4 = 17$. The first numerator update forces $5 \mid D_2$; the multiplicative denominator update gives $5 \mid D_3$; the next numerator update then forces $5 \mid C_4$, a contradiction. This is the failed-route example referred to at the end of the short note: subexponential size and a small relative error do not supply arithmetic compatibility.

14.1 Growth of the factors repeated in the denominator

To compare repeated prime factors in the denominator with its total size, define

$$L_0 = D_0, \quad L_{n+1} = \text{lcm}(L_n, a_n), \quad M_n = \frac{D_n}{L_n}.$$

Since $D_n = D_0 \prod_{j \leq n} a_j$, the quotient is integral and $M_n L_n = D_n$. The product counts every occurrence of a prime, whereas the least common multiple keeps only its largest exponent. The quotient M_n records the remaining factors. Problem 14.3 below asks whether failure of the Sylvester recurrence would force this quotient to grow exponentially, contradicting its known subexponential upper bound. We first record a local estimate for cancellation during a recovery interval; it does not by itself give that global lower bound.

14.2 Cancellation during recovery intervals

The LCM quotient M_n need not equal the common divisor G_n removed by reduction to lowest terms. Since M_n divides both C_n and D_n , we have $M_n \mid G_n$, but equality is not assumed. The following estimate concerns the successive reduction factors, over an interval where the reduced numerator regains its starting value. In the calculation below, h_n is the factor removed at step n , c_n is a positive integer with $c_n^2 \mid h_n$, and $\tilde{e}_n$ denotes the signed reduced error, as in Section 7. For the estimate itself, let $u, h, c : \mathbb{N} \rightarrow \mathbb{N}$ and $\tilde{e} : \mathbb{N} \rightarrow \mathbb{Z}$ satisfy

$$h_n u_{n+1} = u_n - \tilde{e}_n, \quad u_n > 0, \quad h_n > 0.$$

Fix $K, r, L \in \mathbb{N}$ with $K, L > 0$, assume

$$K|\tilde{e}_{r+i}| < u_{r+i} \quad (0 \leq i < L), \quad u_r \leq u_{r+L},$$

and let R be a finite family of moduli. Suppose every m_q , $q \in R$, divides $\prod_{r \leq n < r+L} c_n$, and that $c_n^2 \mid h_n$ throughout the interval. Then

$$K^L \text{lcm}(m_q : q \in R)^2 < (K+1)^L. \quad (23)$$

To see this, each relative-error bound gives $h_n u_{n+1} < (1 + 1/K)u_n$. Multiplication over the interval and $u_r \leq u_{r+L}$ yield

$$\prod_{r \leq n < r+L} h_n < (1 + 1/K)^L.$$

The least common multiple of the m_q divides the product of the c_n , and $c_n^2 \mid h_n$ at every step. Its square therefore divides, and is at most, the positive product on the left.

Multiplying by K^L proves the claimed bound. This is the [bound on cancellation over a recovery interval](#).

The square in (23) comes from the assumption $c_n^2 \mid h_n$. If the least common multiple of the chosen moduli is at least $2^{|R|}$, then

$$K^L 4^{|R|} < (K+1)^L.$$

The formal consequence is [the bound on the number of independent moduli](#). Taking logarithms gives the explicit bound

$$\frac{|R|}{L} < \frac{\log(1+1/K)}{\log 4} \leq \frac{1}{K \log 4}.$$

Thus the number of these moduli is small relative to the recovery length when the relative-error bound is small. This statement requires the lower bound $2^{|R|}$ on their least common multiple; it does not apply to an arbitrary family with repeated prime factors.

For a fixed number of steps, the same estimate has a simpler consequence: a sufficiently late recovery cannot include any cancellation. If $K|\tilde{e}_n| < u_n$ eventually for every K , then for each fixed $L > 0$ there is an N such that every recovery $u_r \leq u_{r+L}$ with $r \geq N$ satisfies

$$\prod_{0 \leq i < L} h_{r+i} = 1.$$

Indeed, choose K so large that $(1+1/K)^L < 2$ and apply the same product estimate beyond its error threshold. The positive integer product is then smaller than 2, so it equals 1. The same K works for every length $1 \leq j \leq L$, since $(1+1/K)^j \leq (1+1/K)^L < 2$. Thus, after a sufficiently late step with $h_n > 1$, the numerator cannot regain its starting value at any of the next L indices. This includes a recovery followed by another fall before the last endpoint. The fixed-length conclusion is formalised as [absence of late cancellation during recovery intervals of fixed length](#). The theorem does not bound a recovery length that varies with r , nor does it supply a lower bound for the least common multiple of the chosen moduli. Those are the missing inputs needed to turn (23) into a global contradiction.

Problem 14.3 (growth of repeated denominator factors). For every rational-tail orbit satisfying the hypotheses but not the conclusion of Problem 1.1, must

$$\limsup_{n \rightarrow \infty} \frac{\log M_n}{n} > 0?$$

Equivalently, must there be a $K \geq 1$ for which

$$2^n \leq M_n^K$$

at infinitely many indices?

The two displayed formulations are equivalent up to changing the positive constant; the second is not a weaker target. Section 6 already gives $1 \leq M_n \leq C_n$, hence $\log M_n/n \rightarrow 0$ on every orbit under consideration. A positive answer would therefore

be a contradiction, not a further compatible necessary condition. It would prove Problem #243. The missing step is to force exponential growth of M_n from failure of the Sylvester recurrence.

There is also a more local-looking question whose content is nevertheless the entire prefix. Put $A_n = \prod_{j < n} a_j$, so $D_n = D_0 A_n$. From

$$D_0 A_n = (a_n - 1)C_n + E_n$$

one immediately obtains

$$\gcd(A_n, a_n - 1) \mid E_n.$$

The checked [tail-height estimate](#) and vanishing relative error make $|E_n|$ subexponential in n .

Problem 14.4 (common factors of the prefix and $a_n - 1$). In every rational-tail orbit satisfying the hypotheses but not the conclusion of Problem 1.1, is

$$\limsup_{n \rightarrow \infty} \frac{\log \gcd(A_n, a_n - 1)}{n} > 0?$$

Equivalently, do there exist $\eta > 0$ and infinitely many n such that $\gcd(A_n, a_n - 1) \geq e^{\eta n}$?

By Proposition 14.1, the error is nonzero eventually. The displayed divisibility therefore bounds $\gcd(A_n, a_n - 1)$ by $|E_n|$ at every sufficiently late index. A positive answer contradicts the subexponential bound on that error. Arbitrarily long locally admissible blocks do not answer this question: the gcd uses the complete prefix.

14.3 The direct analytic question

Problem 14.5 (summability from rationality). Let (a_n) satisfy the hypotheses of Problem 1.1, and let (D_n, C_n, E_n) be its integer tail. Must

$$\sum_{n=0}^{\infty} \frac{(-E_n)_+}{C_n} < \infty?$$

An affirmative answer closes Problem #243 by Theorem 13.1. The summands are non-negative and tend to zero, so $\log(1 + t)$ is comparable to t at their values. Convergence is therefore equivalent to boundedness of the partial products $\prod_{n < N} (1 + (-E_n)_+/C_n)$, the same criterion in multiplicative form. A negative answer requires a sequence satisfying the full growth and rationality hypotheses. A locally admissible state orbit is insufficient. By Theorem 6.2, it is enough to establish the finiteness of (21) for one non-increasing weight with divergent integral and one fixed $B \geq 0$. The equivalent sum in Theorem 6.2 uses only steps where the LCM numerator exceeds its previous maximum. It subtracts B from each upward jump, takes the positive part and weights it by $f(U_n)$.

Remark. One further question is not left open here, since it is answered in [12]: whether vanishing relative error follows from $a_{n+1}/a_n^2 \rightarrow 1$ and rationality. It does. After deleting a finite prefix, Koizumi's Corollary 3 [12, p. 9] makes the sequence the pseudo-greedy expansion of its own reciprocal sum and gives $\varepsilon_n \rightarrow 0$ under exactly the hypotheses of

Problem 1.1; rationality is not needed for that step, only summability of $\sum 1/a_n$. The matched-index dictionary in Section 3 gives $\varepsilon_n = E_n/C_n$ on the restarted tail. For a rational sum, Lemma 4(2) [12, pp. 11–12] also gives $-c_n/2 \leq e_n < c_n/2$ at every index of that expansion. Thus $|E_n| < C_n$ holds beyond the restart, as required by Theorem 12.1; it need not hold at every original index. The eventual inequality $|E_n| < C_n$ also follows directly from vanishing relative error with $K = 1$. What remains unsupplied is the eventual lower bound on E_n in Theorem 12.1. That additional hypothesis is why the theorem is still conditional.

14.4 Unbounded upward increments

Proposition 14.6 (small increases when the prime moduli are sparse). [Lean · Comparator](#) *There exist strictly increasing primes p_i and a strictly increasing positive integer sequence $u_n \rightarrow \infty$ such that $\gcd(u_n, p_i) = 1$ for all i, n and*

$$u_{n+1} - u_n = O(\sqrt{\log \log(u_n + e^e)}) = o(\log \log(u_n + 3)).$$

Thus the bounded-increment hypothesis of Theorem 11.3 cannot be replaced by an $o(\log \log u_n)$ bound without a quantitative restriction on the moduli.

Proof. Choose increasing primes $p_i > \max\{\exp(\exp((i+2)^2)), 2^{i+3}\}$, for $i \geq 0$. Primes of arbitrarily large size suffice; no distribution theorem is used. Then $\theta = \sum_i 1/p_i < 1/4$ and $k(z) = \#\{i : p_i \leq z\} \leq \sqrt{\log \log z}$ whenever z is large. For integer x put $L(x) = \lceil 4\sqrt{\log \log(x + e^e)} + 8 \rceil$. Since $L(x) = o(x)$, eventually $L(x) > k(x + L(x))/(1 - \theta)$. The elementary window bound of Proposition 7.15 leaves an integer in $[x, x + L(x))$ divisible by no p_i . The set of such integers is therefore unbounded. Enumerate it increasingly as (u_n) and apply the same bound with $x = u_n + 1$ to obtain $u_{n+1} - u_n \leq L(u_n + 1)$. For prime moduli nondivisibility is exactly coprimality, proving every claim. The moduli are deliberately much sparser than the scale used in the next theorem. $\square$

We now keep the full family of moduli, rather than discarding a prefix as in Proposition 7.15. The resulting coefficient depends on $\sigma = \prod_j (1 - 1/m_j)$. For a finite prefix, the corresponding product is exactly the proportion of admissible residue classes by the Chinese remainder theorem. This explains the constant in the statement before the limiting argument is made.

Theorem 14.7 (largest gaps between integers avoiding given multiples). [Lean · Comparator](#) *Let $m_0 < m_1 < \dots$ be pairwise coprime integers at least 2 with $\ell(m_j) = j + O(1)$, where $\ell(x) = \log_2 \log_2 \max(4, x)$. Let $\sigma = \prod_j (1 - 1/m_j) > 0$ and enumerate the positive integers divisible by no m_j in increasing order as (u_n) . Then*

$$\limsup_{n \rightarrow \infty} \frac{u_{n+1} - u_n}{\ell(u_n)} = \sigma^{-1}.$$

This is a statement about avoidance of multiples of whole moduli. It is not a statement about coprimality to composite m_j , or about integer tail orbits.

Proof. Write $k(z) = \#\{j : m_j \leq z\} = \ell(z) + O(1)$. For a fixed prefix length T , set

$$M_T = \prod_{j < T} m_j, \quad \sigma_T = \prod_{j < T} (1 - 1/m_j), \quad \theta_T = \sum_{j \geq T} 1/m_j.$$

Here $\theta_T \rightarrow 0$. By the Chinese remainder theorem, avoiding the first T whole moduli selects precisely the fraction σ_T of the residues modulo M_T .

For the upper bound, an integer interval $[x, x+L)$ contains at least $\sigma_T L - M_T$ integers avoiding that prefix. The remaining moduli cover at most $L\theta_T + k(x+L)$ integers. Choose T large enough that $\sigma_T > \theta_T$, and then any $c > (\sigma_T - \theta_T)^{-1}$. For $L = \lceil c\ell(x) \rceil$, the number left is positive for all sufficiently large x , since $k(x+L) = \ell(x) + O(1)$. Thus the set being enumerated is unbounded, and every sufficiently late such interval meets it. Applying this at $x = u_n + 1$ gives

$$\limsup_n \frac{u_{n+1} - u_n}{\ell(u_n)} \leq (\sigma_T - \theta_T)^{-1}.$$

Letting $T \rightarrow \infty$ gives the upper bound σ^{-1} .

For the lower bound, fix T and let the integer length L tend to infinity. Among the offsets $0 \leq j < L$, precisely $K_L = \sigma_T L + O(M_T)$ avoid the first T moduli. Assign to these offsets distinct moduli $m_T, \dots, m_{T+K_L-1}$. Solve simultaneously $x \equiv 0 \pmod{M_T}$ and $x \equiv -j \pmod{m_j}$ modulo the modulus assigned to j . Put $Q_L = \prod_{i < T+K_L} m_i$ and choose this solution in $[Q_L, 2Q_L)$. Every integer of $[x, x+L)$ is then divisible by a modulus: either an old one or its assigned new one. Let u be the greatest admissible integer below x ; its successor is at least $x+L$, so the gap is at least L . The scale assumption gives

$$\ell(2Q_L) \leq T + K_L + O(1),$$

because $\log_2 m_i = 2^{i+O(1)}$ and these upper bounds sum geometrically. Since $u < 2Q_L$, the ratio for this gap is at least $L/(T + K_L + O(1))$, tending to σ_T^{-1} . These u tend to infinity: $x \geq Q_L \rightarrow \infty$ and admissible integers are unbounded. Hence the limsup is at least σ_T^{-1} for every T . Let $T \rightarrow \infty$ to finish. $\square$

For the Fermat moduli $m_j = 2^{2^j} + 1$, pairwise coprimality and the finite-product identity

$$\prod_{j=0}^N \left(1 - \frac{1}{2^{2^j} + 1}\right) = \frac{2^{2^{N+1}} - 1}{2^{2^{N+1}} - 1}$$

give $\sigma = 1/2$, so the coefficient is exactly 2. These results describe gaps between integers avoiding fixed moduli. They do not construct multipliers or numerator and denominator sequences satisfying the exact recurrences. In particular, the static proportion σ must not be substituted for $\varphi(v_T)/v_T$: the first excludes multiples of the whole moduli, while the second excludes every prime factor of the reduced denominator.

Formalisation targets

Erdős–Straus. Formalise Theorem 3 of [1], with its prefix-LCM quotient and correctly indexed growth factor, under the published analytic hypotheses.

Duverney. Formalise the absolute-convergence form of Corollary 3.2 of [2], including its signed numerators, and recover the all-positive specialisation used for comparison here. For the printed condition interpreted as mere signed convergence, first justify the nonvanishing-product step discussed in Section 3. That interpretation is not treated as an already verified formalisation target.

Statements and declarations

Artefact and data availability. The Lean sources, with their fixed toolchain and library manifest, are in the [public repository](#). The [evidence record](#) identifies the recorded Lean declarations, source versions and Comparator status of the results, including gaps. The external formal-conjecture entry states the problem without proving it. The ordinary proofs and their exposition require mathematical review independently of these checks.

Funding and competing interests. This work received no external funding. The author declares no competing interests.

Acknowledgements. I thank Wouter van Doorn for advice on exposition, including the explanation of restrictive hypotheses and the removal of unnecessary terminology. The problem numbering follows Bloom’s Erdős Problems catalogue [13].

A Guide to the formal sources

The margin marks and declarations below identify fixed revisions of the formal sources. The printed square-specialisation argument uses Chebotarev, while the supplied formal proof uses a Dedekind-zeta pole comparison. Agreement of their conclusions does not verify every step of the printed argument. No Lean build, Comparator replay or independent review was performed for this revision. The supplied evidence records also retain source-pin and index gaps.

The recurrence identities and arithmetic exclusions are in `ReciprocalTailRigidity.lean`, the scalar convergence theorem is in `SparseResetRecovery.lean`, and the bounds on cancelled factors are in `RepairEntropy.lean`. The Lean statement excluding a periodic error assumes $e_n < a_n$; Section 10 derives this bound eventually for a periodic positive error magnitude. The printed bounded-negative theorem uses $E_n/C_n \rightarrow 0$. The formal statement uses its division-free form, $K|E_n| < C_n$ eventually for each natural K ; the bound $|E_n| < C_n$ is its $K = 1$ case. Sections 3 and 12 give the construction from a rational reciprocal sum.

Indexing of the regular-rate theorems

Index translation for the Lean statements. Both Lean declarations use $b : \mathbb{N} \rightarrow \mathbb{N}$, including a positive b_0 , whereas the theorems above start at a_1 . A simple shift changes λ/n by order n^{-2} and may lose the stated $o(n^{-\lambda})$ precision. Instead choose a large N with $a_N > N$. Such an N exists: strict increase gives $a_n \geq n$ and the rate gives $a_n^2/a_{n+1} < 2$ eventually, hence $a_{n+1} > a_n^2/2 \geq n^2/2 > n + 1$ for large n . Define

$$b_n = \begin{cases} n + 1, & 0 \leq n < N, \\ a_n, & n \geq N. \end{cases}$$

Then b is positive and strictly increasing, with the same indexed rate for every $n \geq N$. Moreover

$$\sum_{n \geq 0} \frac{1}{b_n} - \sum_{n \geq 1} \frac{1}{a_n} = \sum_{n=0}^{N-1} \frac{1}{n+1} - \sum_{n=1}^{N-1} \frac{1}{a_n} \in \mathbb{Q}.$$

The series converge by the stated rapid-growth rate. If the original sum were rational, so would be the sum for b , contradicting the corresponding zero-indexed Lean theorem. This finite-prefix implication is proved here in ordinary mathematics; the Lean declarations check the zero-indexed theorems themselves.

The square-specialisation proofs

The Lean proofs of Lemma 2.7, Proposition 2.8, Theorem 2.3 and Theorem 2.1 assume no form of the Chebotarev theorem. The ordinary proof of Lemma 2.7 above uses the classical finite-Galois Chebotarev theorem; its Lean proof uses the simple pole of the Dedekind zeta function instead, as described after that proof, and the Lean proofs of the other three pass through the lemma. The Lean statement of Theorem 2.1 is indexed from zero; the one-based conclusion uses the finite-prefix argument above.

The nonintegral-rate formal proof uses the real-parameter extraction theorem and a quantitative canonical-tail estimate. Convergence of the reciprocal series is derived from the rate.

Recurrence declarations

Proposition 4.1. [update law](#).

Proposition 4.3. [denominator](#), [numerator](#) and [error](#).

Theorem 5.1. [defect identity](#).

Theorem 5.3. [local rigidity step](#).

Theorem 5.7. [eventual Sylvester recurrence](#).

Theorem 5.8. [absorption of a vanishing error](#), with the companion [contrapositive along a tail](#).

Theorem 8.1. [stabilisation of the nonnegative error](#), using the [eventual constancy of a nonincreasing sequence](#).

Lemma 11.1. [shifted consecutive multiples](#).

Theorem 11.3. [the Chinese-remainder first-crossing consequence](#).

Theorem 12.1. The theorem is [stated with an eventual lower bound and division-free vanishing relative error](#). The Lean proof shifts past both thresholds and applies the [version with \$|E_n| < C_n\$ and the lower bound at every index](#). It uses [divergence from vanishing relative error](#) and the [exclusion for bounded upward increments and arbitrarily late bounded negative magnitudes](#), via its formulation with [divergence as a hypothesis](#). Both exclusion statements require a uniform bound on all upward increments. Bounded negative magnitudes at arbitrarily late indices suffice for gcd stabilisation, but not for the jump bound in the CRT argument.

Theorem 13.1. The scalar conclusion is [the eventual vanishing for a positive integer sequence](#); its exact-orbit consequence is [the scalar recurrence from a convergent sum](#). A separate exact-orbit formulation retains the denominator and tail update equations, positivity, the centered-step identity and division-free normalized vanishing. Under

$\sum_n (-E_n)_+ / C_n < \infty$, the checked results give a [bound on numerator growth](#), [eventual zero error](#), and [the Sylvester recurrence](#). These are conditional hypotheses, not properties proved here for the original orbit. The [plectis-erdos-lean](#) repository contains a [formulation for the rational-tail numerators using a convergent sum](#), not covered by the evidence record. Summability remains an assumption; the elementary scalar implication needs neither growth nor vanishing relative error.

Proposition 14.1. In the Lean statement of Proposition 14.1, the divergent sum of relative increases is recorded as divergence of the partial sums.

B A factorial modulus for integral recursion

In the case $m = c = 1$ of Section 9, the identity (5.1) becomes $D_n + 1 = (a_n - 1)(n + 1)$. It determines the recursion $a_{n+1} = \text{num}(n, a_n) / (n + 2)$, where

$$\text{num}(n, a) = (n + 1)a^2 - (n + 2)a + (n + 3),$$

the [forced numerator](#). We stop at the first nonintegral quotient. The [formal survival predicate](#) requires exact division at each step. For $a_0 = 3$, as in Example 9.1, the first quotient is $\text{num}(0, 3) / 2 = 3$, but the next quotient is $\text{num}(1, 3) / 3 = 13/3$, so the recursion stops. Direct iteration can produce very large intermediate integers. To decide whether the first h divisions are exact, however, it suffices to know the initial value modulo $(h + 1)!$. Computing a pseudo-greedy orbit through residues modulo a shrinking product modulus is Koizumi's method [12, Remark 2 and Algorithm 1, pp. 13–14]; for the forced numerator that product is a factorial.

Theorem B.1 (factorial residue reduction). [Lean · Comparator](#) *Let h be a nonnegative integer and let a, b be integers with $a \equiv b \pmod{(h + 1)!}$. The first h steps of the recursion $a_{n+1} = \text{num}(n, a_n) / (n + 2)$, starting at index zero, are integral for $a_0 = a$ if and only if they are integral for $a_0 = b$.*

Proof. Define $M(0, i) = 1$ and $M(h + 1, i) = (i + 2)M(h, i + 1)$. Thus $M(h, 0) = (h + 1)!$. We prove the stronger statement that, at index i , congruent inputs modulo $M(h, i)$ give the same answer to whether the next h quotients are all integral. There is nothing to prove for $h = 0$.

For the inductive step, suppose $a \equiv b \pmod{(i + 2)M(h, i + 1)}$. Since $\text{num}(i, \cdot)$ is an integer polynomial, its values at a and b are congruent modulo that product. In particular, one is divisible by $i + 2$ if and only if the other is. If neither is divisible, both recursions stop. Otherwise their quotients are congruent modulo $M(h, i + 1)$, so the induction hypothesis applies to the remaining h updates at index $i + 1$. Taking $i = 0$ proves the claim. $\square$

The formal [factorial residue reduction](#) uses the [shrinking-modulus induction](#), [polynomial congruence](#), and [cancellation after exact division](#). Its modulus is identified by the [ascending-factorial formula](#) and its [factorial value at the initial index](#).

For $h = 1$, the modulus is $2! = 2$. The first quotient is integral if and only if $2 \mid a^2 - 2a + 3$, or equivalently if a is odd. For instance, $\text{num}(0, 3) = 6$ and $\text{num}(0, 4) = 11$. Thus the parity of a decides the first step, as Theorem B.1 asserts in this case.

Exact enumeration for $2 \leq a_0 < 5000$ gives a maximum of 17 successful updates, or 18 multiplier values including a_0 . Nine seeds attain it, the least being 1501. The seed $a_0 = 1$ is excluded: it gives $a_n = 1$ at every step and violates the hypothesis $a_n \geq 2$. This procedure tests exact division only; it does not test all the conditions required of a positive integer solution of the recurrences. The finite search does not prove the infinite exclusion; Theorem 9.2 does so under its stated hypotheses. The reduction remains useful because its shrinking-modulus argument applies to other recurrences defined by polynomial division.

C Result map and proof dependencies

The short paper proves cubic-rate irrationality in Section 3 and the bounded-increment criterion in Section 4. Their dependencies are summarised below. Details retained here are the signed-series comparison in Section 3, the integer-coefficient extension in Section 6, and the general $1 + c/n$ comparison in Section 7. Section 2 retains the full regular-rate extraction and positive-density cubic exclusion; Theorem 7.3 gives the separate double-logarithmic argument. The short note also includes the scalar failed-divisibility example from Section 14.

Bounded upward increments. Vanishing relative error gives eventual absorption. Bounded negative values at arbitrarily late indices give a stable gcd; CRT then excludes the unbounded reduced tail with bounded upward increments. The increment bound is an extra hypothesis.

A convergent sum of relative increases. The inequality $C_{n+1} \leq C_n(1 + (-E_n)_+/C_n)$ turns summability into a uniform bound on C_n . Integrality then permits only finitely many rises, after which the numerator stabilises. No vanishing relative error is needed.

New maxima and prime-power persistence. A bound on the negative reduced error scaled by the previous maximum also bounds every cancellation factor. Primes larger than that bound persist in the reduced denominator, giving the CRT contradiction in Theorem 7.5. The double-logarithmic argument uses the density-one coprimality count to obtain enough moduli, then controls the height of the CRT block. The LCM numerator gives weighted first-crossing estimates. A prime power dividing the reduced denominator persists while the unreduced next numerator stays below the threshold in Corollary 7.2. Crossings can be counted before that threshold is reached.

Cubic rate. The tail estimate and finite differences of the Gamma ratio force an eventual cubic polynomial for the integer numerator. The recurrence forces a modular square condition, which the square-specialisation lemma lifts to the cubic field. The ordinary proof uses Chebotarev; the Lean proof obtains the same implication from a Dedekind-zeta pole comparison. The trace calculation leaves $m = 12$, and both possible signs are excluded modulo seven.

Avoidance of fixed moduli. For sufficiently sparse prime moduli, a coprime integer sequence can have unbounded but very small increments. At the specified double-exponential scale, counting residue classes and applying CRT give the largest gaps be-

tween integers divisible by none of the whole moduli. These auxiliary results do not construct reciprocal-tail examples.

What remains. For the unrestricted problem, one still needs the stated summability or record bound, or a contradiction to the necessary conditions on a counterexample. The static examples and local identities supply neither.

References

- [1] P. Erdős and E. G. Straus, *On the irrationality of certain Ahmes series*, J. Indian Math. Soc. (N.S.) **27** (1964), 129–133. MR 175848.
- [2] D. Duverney, *Irrationality of fast converging series of rational numbers*, J. Math. Sci. Univ. Tokyo **8** (2001), 275–316. MR 1837165.
- [3] C. Badea, *A theorem on irrationality of infinite series and applications*, Acta Arith. **63** (1993), no. 4, 313–323, doi:10.4064/aa-63-4-313-323.
- [4] R. Tijdeman and P. Yuan, *On the rationality of Cantor and Ahmes series*, Indag. Math. (N.S.) **13** (2002), no. 3, 407–418, doi:10.1016/S0019-3577(02)80018-0.
- [5] P. Erdős and R. L. Graham, *Old and New Problems and Results in Combinatorial Number Theory*, Monogr. Enseign. Math. **28**, Geneva, 1980, p. 64.
- [6] P. Erdős, *On the irrationality of certain series: problems and results*, in A. Baker (ed.), *New Advances in Transcendence Theory*, Cambridge UP, 1988, pp. 102–109, doi:10.1017/CBO9780511897184.009.
- [7] V. Kovač and T. Tao, *On several irrationality problems for Ahmes series*, Acta Math. Hungar. **175** (2025), no. 2, 572–608, doi:10.1007/s10474-025-01528-0; arXiv:2406.17593v4. Page references are to arXiv:2406.17593v4.
- [8] I. O. Bado, *Prime-Support Rigidity and Primitive Pseudo-Greedy Dynamics: Partial Progress on Erdős Problem #243*, preprint posted September 2026, doi:10.13140/RG.2.2.36612.08325.
- [9] P. White with Claude (Anthropic), *Erdős #243: working report*, Erdős Problem a Day, page dated 12 August 2026. AI-assisted, unrefereed working report.
- [10] P. Stevenhagen and H. W. Lenstra, Jr., *Chebotařev and his density theorem*, Math. Intelligencer **18** (1996), no. 2, 26–37, doi:10.1007/BF03027290. Page references are to the linked author version dated 23 March 1995.
- [11] W. van Doorn, Y. Li and Q. Tang, *Optimal bounds for an Erdős problem on matching integers to distinct multiples*, preprint arXiv:2603.28636v1, 2026.
- [12] J. Koizumi, *Irrationality of the reciprocal sum of doubly exponential sequences*, Integers **26** (2026), Paper No. A28, 17 pp., doi:10.5281/zenodo.18714404. Locators refer to this published version.
- [13] T. F. Bloom, *Erdős Problem #243*, accessed 28 July 2026.
- [14] The Formal Conjectures Authors, *FormalConjectures.ErdosProblems.243*, Lean source at commit f776d2f, 2025. Formal problem statement, not a proof.
- [15] P. Erdős and E. G. Straus, *On the irrationality of certain series*, Pacific J. Math. **55** (1974), no. 1, 85–92.
- [16] J. Hančl and R. Tijdeman, *On the irrationality of polynomial Cantor series*, Acta Arith. **133** (2008), no. 1, 37–52, doi:10.4064/aa133-1-3. Locators refer to the published version.

- [17] D. Duverney, T. Kurosawa and I. Shiokawa, *Irrationality exponents of certain fast converging series of rational numbers*, Tsukuba J. Math. **44** (2020), no. 2, 235–250, doi:10.21099/tkbjm/20204402235. Theorem locators follow the linked 14-page author version.
- [18] T. Crmarić and V. Kovač, *On the irrationality of certain super-polynomially decaying series*, Colloq. Math. **179** (2025), 55–68, doi:10.4064/cm9628-5-2025. Theorem locators follow arXiv:2504.18712v1.
- [19] A. Koutsoukou-Argyraiki and W. Li, *Irrationality Criteria for Series by Erdős and Straus*, Archive of Formal Proofs, 12 May 2020. An Isabelle/HOL formalisation; the archive entry identifies the results formalised.
- [20] National Institute of Standards and Technology, *Digital Library of Mathematical Functions*, §5.11(iii), formula 5.11.12, accessed 16 September 2026.
- [21] E. H. el Abdalaoui, M. Lemańczyk and T. de la Rue, *A dynamical point of view on the set of $\mathcal{B}$ -free integers*, Int. Math. Res. Not. IMRN (2015), no. 16, 7258–7286, doi:10.1093/imrn/rnu164. The cited definition is also in §1.2 of arXiv:1311.3752v3.