

No Finite Separable Representation at Three Prime Generators

Kernel rank and a tail criterion for Erdős Problem #269

WILL COOK*

July 2026; exposition revised 18 September 2026

ABSTRACT

For three distinct primes, the reciprocal running-LCM kernel has nonsingular minors of every order, with the same indices in every third-coordinate layer. Row and column rescaling reduces the proof to a matrix whose entries are 1 or the reciprocal of the third prime. Thus no finite sum of products separates one exponent from the other two. Separately, for the repeated $\{2, 3, 5\}$ series we prove a tail recurrence and a residue criterion equivalent to irrationality. The unresolved step is to prove its inequalities for every positive multiplier coprime to 30 and arbitrarily late starts.

1 Introduction

For a finite prime set P , let $\mathcal{S}_P$ consist of the positive integers whose prime factors lie in P , including 1. We use $\mathbb{N} = \{0, 1, 2, \dots\}$ for exponent indices. For $x \geq 1$, put

$$L(x) = \text{lcm}\{u \in \mathcal{S}_P : u \leq x\}, \quad \mathcal{R}_P = \sum_{u \in \mathcal{S}_P} L(u)^{-1}.$$

Erdős asked whether $\mathcal{R}_P$ is irrational when $|P| \geq 2$ [1, p. 65][2, p. 106]. For $P = \{p, q, r\}$, write

$$H(x) = p^{\lfloor \log_p x \rfloor} q^{\lfloor \log_q x \rfloor} r^{\lfloor \log_r x \rfloor}, \quad K(i, j, k) = H(p^i q^j r^k)^{-1}.$$

Prime-power divisibility gives $L = H$ (Proposition 4.1). Fan used this identity to separate the two-prime kernel [7]. With three primes, even a finite sum of separated products is impossible. This is a statement about the kernel, not the arithmetic nature of its sum.

Formal proofs. A result with a kernel-checked Lean proof carries a mark in the margin. *Lean* opens the proof: the declaration itself when one declaration states the whole result, otherwise the list of declarations that together state it. *Comparator* opens the record of an independent check, in which the same statement, written again from Mathlib alone in a separate repository, was compared with our proof by Lean's Comparator tool, allowing

* *Authorship and AI use.* Will Cook built and directed the research infrastructure and reviewed the claims when he could. AI agents did most of the research and drafting. Cook did not independently verify every claim.

only the three standard axioms. A dagger on the Lean mark means that the Lean proof assumes an input named just below the result. A result without a mark has no Lean proof of its whole statement; what is checked is said below it. The [evidence record](#) gives every declaration, version and check. These checks show that the stated propositions are proved; whether they are the right propositions is a question the reader can settle by comparing them with the text.

2 Nonsingular minors of every order

Theorem 2.1 (no finite separation of the kernel). *Let p, q, r be primes with $p \neq q$, $p \neq r$ and $q \neq r$. For every $n \geq 0$ there are injective maps $I, J : \{0, \dots, n-1\} \rightarrow \mathbb{N}$ such that, for every $k \geq 0$,*

Lean
Comparator

$$\det(K(I(a), J(b), k))_{0 \leq a, b < n} \neq 0.$$

Consequently, for no finite d do there exist rational-valued functions $f_\ell(i)$ and $G_\ell(j, k)$, $0 \leq \ell < d$, satisfying

$$K(i, j, k) = \sum_{\ell < d} f_\ell(i) G_\ell(j, k) \quad \text{for all } i, j, k.$$

Proof. For fixed k , we rescale rows and columns to obtain a matrix independent of k . Put $c = r^{-1}$, $x_i = \{i \log_r p\}$ and $y_j = \{j \log_r q\}$, where braces denote fractional parts. Splitting the floor exponents gives

$$K(i, j, k) = U_i(k)^{-1} C_{ij} V_j(k)^{-1}, \quad C_{ij} = c^{\mathbf{1}_{\{x_i + y_j \geq 1\}}},$$

where

$$U_i(k) = p^i q^{\lfloor \log_q(p^i r^k) \rfloor} r^{k + \lfloor \log_r p^i \rfloor},$$

$$V_j(k) = q^j p^{\lfloor \log_p(q^j r^k) \rfloor} r^{\lfloor \log_r q^j \rfloor}.$$

The positive factors do not affect whether a minor vanishes; C records only whether $x_i + y_j \geq 1$. Distinct primes make $\log_r p$ and $\log_r q$ irrational, so each fractional-part orbit is dense in $[0, 1]$. The row and column indices are independent choices; no density of a single orbit of pairs is required. For $n \geq 1$, choose distinct rows with $0 < x_{I(0)} < \dots < x_{I(n-1)} < 1$. We choose each column so that its entries change from 1 to c at a different selected row. Writing $s_b = 1 - y_{J(b)}$, density of (y_j) lets us choose

$$s_0 \in (0, x_{I(0)}), \quad s_b \in (x_{I(b-1)}, x_{I(b)}) \quad (1 \leq b < n).$$

The intervals for the s_b are disjoint, so the column indices are distinct. The strict inequalities also avoid the threshold itself, and give $x_{I(a)} + y_{J(b)} \geq 1$ exactly when $b \leq a$. Thus multiplying row a by $U_{I(a)}(k)$ and column b by $V_{J(b)}(k)$ reduces the selected kernel minor to

$$T_n(c) = \begin{pmatrix} c & 1 & \cdots & 1 \\ c & c & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ c & c & \cdots & c \end{pmatrix}, \quad \det T_n(c) = c(c-1)^{n-1}.$$

Indeed, subtracting each preceding row from the next, working upwards from the last row, leaves diagonal entries $c, c-1, \dots, c-1$. The original determinant is therefore

$$c(c-1)^{n-1} \prod_{a < n} U_{I(a)}(k)^{-1} \prod_{b < n} V_{J(b)}(k)^{-1} \neq 0.$$

The indices were chosen from C , independently of k . For $n = 0$ the empty determinant equals 1.

For the final assertion, suppose a separation with d summands existed and fix any k . On the rows $I(0), \dots, I(d)$ and columns $J(0), \dots, J(d)$ its matrix would be the product of the $(d+1) \times d$ matrix $(f_\ell(I(a)))_{a,\ell}$ and the $d \times (d+1)$ matrix $(G_\ell(J(b), k))_{\ell,b}$. Its rank would be at most d , so its determinant would vanish, contrary to the minor just constructed. $\square$

No continuity or boundedness is assumed for the separated factors.

Corollary 2.2 (the same minors modulo integers coprime to 30). *For $(p, q, r) = (2, 3, 5)$ and every $n \geq 1$, there are injective maps $I, J : \{0, \dots, n-1\} \rightarrow \mathbb{N}$, chosen independently of B and k , such that for every $B \geq 2$ coprime to 30 and every $k \geq 0$, the selected $n \times n$ kernel matrix has unit determinant over $\mathbb{Z}/B\mathbb{Z}$, with each reciprocal prime power interpreted by its modular inverse.*

Lean
Comparator

Proof. Choose the maps from Theorem 2.1. Every row and column factor is a unit modulo B . The normalised determinant is $5^{-1}(-4/5)^{n-1}$, also a unit. $\square$

The modulus may be composite, for example 49 or 77. Coprimality with 30 makes both the reciprocal entries and the determinant units: the latter introduces only a power of 4 in its numerator.

Example 2.3. For $(p, q, r) = (2, 3, 5)$, the leading two-by-two determinant is $-1/15$: its four entries are $1, 1/6, 1/2, 1/60$, so it equals $1/60 - 1/12$.

The indices must be selected. The leading 4×4 block at $\{2, 3, 5\}$ is singular:

$$K(3, j, 0) = \frac{1}{120} K(0, j, 0) \quad (0 \leq j < 4).$$

These equalities follow by evaluating the height at 3^j and $8 \cdot 3^j$. But at $j = 4$, $K(3, 4, 0) - K(0, 4, 0)/120 = -1/19440000$. Thus the leading minors do not supply the arbitrary-order theorem.

Proposition 2.4 (rank of a matrix of threshold columns). *Let $m \geq 1$ and let c lie in a field with $c \neq 0, 1$. For $0 \leq k \leq m$ write v_k for the length- m column with a 1 in each of the first k coordinates and c thereafter. A matrix whose distinct columns are v_k for k in a nonempty set E has rank $|E| - \mathbf{1}_{\{0, m\} \subseteq E}$.*

Lean
Comparator

The correction accounts for the two constant columns: $v_0 = cv_m$. There is no other dependence among distinct threshold columns.

Proof. List $E = \{k_1 < \dots < k_t\}$. The $t - 1$ differences $v_{k_{j+1}} - v_{k_j} = (1 - c)\mathbf{1}_{\{k_j, \dots, k_{j+1}-1\}}$ have disjoint nonempty supports, hence are linearly independent. If $k_1 > 0$ or $k_t < m$, their union misses a coordinate where v_{k_1} is nonzero, so the rank is t . If $k_1 = 0$ and $k_t = m$, then $v_0 = c\mathbf{1}$ lies in the span of the differences (their sum is $(1 - c)\mathbf{1}$), so the rank is $t - 1$. $\square$

The restrictions $c \neq 0, 1$ exclude a zero column or identical columns; they hold for $c = 1/r$ over $\mathbb{Q}$. In one fixed layer k , ordering the sampled row phases puts every column in the stated form. The formula therefore determines the rank of each nonempty rectangular sample. For fixed row and column indices this rank is independent of k . Section 4 of the companion paper, *The Three-Prime Running LCM: Kernel Rank and Tail Arithmetic*, gives the exact rational comparisons.

Exact rank is not an approximation obstruction. Truncating the original kernel to $i < N$ gives a sum of N separated terms. The bound $H(x) > x^3/(pqr)$ makes the truncation error tend to zero both uniformly and in $\ell^1(\mathbb{N}^3)$. By contrast, the rescaled matrix in the rank proof has uniform distance $(1 - 1/r)/2$ from the matrices of finite rank. Rescaling preserves exact rank but removes the decay responsible for the first approximation. The companion proves both assertions, including the distinction between finite restrictions and the full infinite matrix, in Section 4.1.

3 The two-prime comparison

Fan's post of 26 June 2026 gives the repeated two-prime factorisation, its quadratic expression in one Hecke–Mahler value and its transcendence conclusion [7]. Counting the contributions at powers of each prime also gives the affine formula for the distinct-height sum below. The calculation fixes the normalisations and distinguishes the two sums.

Theorem 3.1 (both two-prime sums). *Let $p < q$ be distinct primes. Put $\theta = \log p / \log q$ and $A = \sum_{n \geq 0} p^{-n} q^{-\lfloor n\theta \rfloor}$. Let $\mathcal{R}_{p,q}$ sum the reciprocal running LCM at every positive $\{p, q\}$ -smooth integer, and let $\mathcal{D}_{p,q}$ count each distinct running LCM once. Then*

Lean†

$$\mathcal{D}_{p,q} = \frac{(q-p)A + p}{q-1}, \quad \mathcal{R}_{p,q} = \frac{(p+q-1)A - (p-1)A^2}{q-1}. \quad (1)$$

Both numbers are transcendental.

The Lean proof assumes the transcendence theorem of Bugeaud and Laurent; the two identities in (1) are proved in Lean without it.

Proof. Set $x = 1/p$, $y = 1/q$, $m_n = \lfloor n\theta \rfloor$ and $\delta_n = m_{n+1} - m_n$. Unique factorisation makes θ irrational, and $0 < \theta < 1$ gives $\delta_n \in \{0, 1\}$. The initial value and the positive powers of p contribute $A = \sum_{n \geq 0} x^n y^{m_n}$. There is one q -power strictly between p^n and p^{n+1} exactly when $\delta_n = 1$, so the positive powers of q contribute

$$B = \sum_{n \geq 0} \delta_n x^n y^{m_n+1}, \quad \mathcal{D}_{p,q} = A + B.$$

Since $0 < x, y < 1$, these series converge absolutely. The identity $y^{m_{n+1}} - y^{m_n} = \delta_n y^{m_n} (y - 1)$ and an index shift in A give $A - 1 - xA = x(y - 1)B/y$. Therefore

$$B = \frac{p - (p - 1)A}{q - 1}.$$

At $p^i q^j$ the height is $p^{i+\lfloor j/\theta \rfloor} q^{j+m_i}$, hence

$$\mathcal{R}_{p,q} = A \sum_{j \geq 0} y^j x^{\lfloor j/\theta \rfloor} = A(1 + B).$$

For the last equality, each $j \geq 1$ corresponds to $n = \lfloor j/\theta \rfloor$ with $m_n = j - 1$ and $\delta_n = 1$. These identities prove (1).

For the Hecke–Mahler series $F_\theta(x, y) = \sum_{n \geq 1} \sum_{k=1}^{m_n} x^n y^k$, geometric summation gives

$$A = \frac{1}{1-x} - \frac{1-y}{y} F_\theta(x, y). \quad (2)$$

Bugeaud and Laurent’s theorem [5, Theorem 1.1] applies with intercept $\rho = 0$, $\beta = x$ and $\alpha = y$: the slope θ is irrational and lies in $(0, 1)$, x and y are nonzero algebraic numbers, $|x| < 1$ and $|xy^\theta| = p^{-2} < 1$. Hence $F_\theta(x, y)$ is transcendental; this case $\rho = 0$ is due to Loxton and van der Poorten [6, Theorem 8, p. 40]. Thus A is transcendental. The displayed affine and quadratic polynomials are nonconstant, so an algebraic value of either would force A to be algebraic. $\square$

The same calculation applies to the monoid generated by any coprime integers $1 < p < q$, such as 4, 9, but need not describe all integers supported on their prime factors. The companion’s Section 3 gives this extension and the failure for 4, 8.

The three-prime tail argument starts again from the running LCM, retaining every multiplicity; it does not use the rank theorem. The distinct-height assertion is a separate historical statement [3].

4 The recurrence for the repeated sum

Proposition 4.1 (the running least common multiple). *Let p, q, r be pairwise distinct primes and $x \geq 1$. Then the running LCM equals the three-prime height: $L(x) = H(x)$.*

[Lean](#)
[Comparator](#)

Proof. Every smooth $n \leq x$ has prime exponents bounded by the corresponding integer logarithms, so $n \mid H(x)$. Conversely the three maximal pure powers occur among those smooth numbers; their product divides the running LCM because they are pairwise coprime. $\square$

The inequalities $x/p < p^{\lfloor \log_p x \rfloor} \leq x$ for each of the three primes also give

$$x^3/(pqr) < H(x) \leq x^3. \quad (3)$$

Proposition 4.2 (cells and jumps). *The running LCM is constant when the three integer logarithms are constant. A jump in exactly one logarithm multiplies it by the corresponding prime. The first n positive powers of each prime, together with 1, form $3n + 1$ distinct points.*

[Lean](#)
[Comparator](#)

Proof. The first two claims follow from the height formula. Positive powers of different primes cannot coincide, by unique factorisation, and none is 1. $\square$

For $(p, q, r) = (2, 3, 5)$, the smooth numbers 5 and 6 have the same running LCM, 60. They contribute 2/60 to the repeated sum, whereas the height 60 contributes only 1/60 to the distinct-height sum. The next identity retains this multiplicity when equal heights are grouped.

Proposition 4.3 (grouping terms with the same height). *For a finite exponent box $\mathcal{B}$, set $F(H) = \{(i, j, k) \in \mathcal{B} : H(p^i q^j r^k) = H\}$. Then*

Lean
Comparator

$$\sum_{(i,j,k) \in \mathcal{B}} K(i, j, k) = \sum_H \frac{\#F(H)}{H}. \quad (4)$$

Proof. Each term in $F(H)$ equals $1/H$. The multiplicities remain present in every subsequent infinite sum. $\square$

From now on let $P = \{2, 3, 5\}$ and $S = \mathcal{R}_P$. For $a \geq 0$ define

$$s_a = \sum_{\substack{x \text{ smooth} \\ 2^a \leq x < 2^{a+1}}} \frac{1}{H(x)}, \quad T_a = \sum_{j \geq 0} s_{a+j}, \quad h_a = \frac{H(2^a)}{2}, \quad X_a = h_a T_a.$$

The half-height h_a clears the finite prefix: for $a \geq 1$ and smooth $x < 2^a$, the exponent of 2 in $H(x)$ is at most $a - 1$, so $H(x) \mid H(2^a)/2$. This is why we omit the final factor 2. Here $h_0 = 1/2$; only for $a \geq 1$ is h_a an integer. Define

$$b_a = \frac{H(2^{a+1})}{H(2^a)}, \quad m_a = \sum_{\substack{x \text{ smooth} \\ 2^a \leq x < 2^{a+1}}} \frac{H(2^{a+1})}{2H(x)}. \quad (5)$$

Lemma 4.4 (integer coefficients and four possible bases). *For every $a \geq 0$, m_a is a positive integer and $b_a \in \{2, 6, 10, 30\}$. The word “numerator” does not impose the positional-digit restriction $m_a < b_a$; that restriction need not hold.*

Lean
Comparator

Proof. If $x < 2^{a+1}$, the exponent of 2 in $H(x)$ is at most a , while the other exponents are bounded by those of $H(2^{a+1})$. Hence $2H(x) \mid H(2^{a+1})$. Each summand is an integer, and the shell contains 2^a . Between consecutive powers of 2 there is at most one power of 3 and at most one power of 5: successive powers of either odd prime have ratio greater than two. These factors, when present, and the factor 2 at the right endpoint give the four possible bases. $\square$

In the shell $[2, 4)$ the terms $1/2$ and $1/6$, with $h_2 = b_1 = 6$, give $m_1 = 4$ and the recurrence $X_2 = 6X_1 - 4$. The shell $[16, 32)$ shows why these numerators are not positional digits: $m_4 = 65 > b_4 = 30$. The companion computes the carried digits separately.

Proposition 4.5 (the tail recurrence and a quadratic bound). *The series defining S, T_a converge. For every $a \geq 0$,*

Lean
Comparator

$$X_{a+1} = b_a X_a - m_a, \quad 0 < X_a \leq \frac{8640}{343} (a+1)^2 < 90(a+1)^2.$$

For every integer $B \geq 1$, either some BX_a is integral and all later states are integral, or $\text{dist}(BX_a, \mathbb{Z}) \geq 1/31$ at arbitrarily large indices.

Proof. For each pair of exponents of 3 and 5, at most one exponent of 2 places the smooth integer in $[2^a, 2^{a+1})$. The two odd-prime exponents lie between 0 and a , giving at most $(a+1)^2$ terms. By (3), $s_a \leq 30(a+1)^2/8^a$. Since $h_a \leq 8^a/2$ and $a+j+1 \leq (a+1)(j+1)$ for $a, j \geq 0$, summing the later shells gives

$$X_a \leq 15(a+1)^2 \sum_{j \geq 0} \frac{(j+1)^2}{8^j} = \frac{8640}{343}(a+1)^2.$$

This proves convergence and the bound. Splitting the first shell gives the recurrence because $m_a = h_{a+1}s_a$ and $h_{a+1} = b_a h_a$. If all sufficiently late distances are strictly below $1/31$, write $BX_a = z_a + e_a$ with $z_a \in \mathbb{Z}$ and $|e_a| < 1/31$. Since Bm_a is integral, $e_{a+1} - b_a e_a$ is an integer of absolute value less than $(1 + b_a)/31 \leq 1$, so $e_{a+1} = b_a e_a$. The lower bound $b_a \geq 2$ forces this bounded error to vanish; integrality then propagates by the recurrence. $\square$

The identity $S/2 = \sum_{a \geq 0} m_a / (b_0 \cdots b_a)$ places the recurrence in the setting of Cantor series. The classical criteria of Erdős–Straus [8, Theorem 2.1] and Hančl–Tijdeman [9, Theorem 3.1] require $m_a / (b_{a-1} b_a) \rightarrow 0$. That hypothesis fails here: the numerators grow quadratically while the bases are bounded. The companion’s Section 6 proves the growth assertion and gives the precise carry conditions. The argument below instead uses the recurrence and the proved bound $90B(a+1)^2$ directly.

Theorem 4.6 (rationality gives positive integer tails). *If $S = A/D$ in lowest terms, where $D = 2^u 3^v 5^w B$ and $\gcd(B, 30) = 1$, then for every $a \geq a_0 = u + 1 + 2v + 3w$,*

[Lean](#)
[Comparator](#)

$$d_a = BX_a \in \mathbb{Z}_{>0}, \quad d_{a+1} = b_a d_a - Bm_a, \quad d_a \leq 90B(a+1)^2.$$

Proof. For $a \geq 1$, every $H(x)$ with $x < 2^a$ divides h_a . Subtracting the finite prefix therefore gives

$$X_a = h_a S - \sum_{\substack{x \text{ smooth} \\ x < 2^a}} \frac{h_a}{H(x)}, \quad \sum_{\substack{x \text{ smooth} \\ x < 2^a}} \frac{h_a}{H(x)} \in \mathbb{Z}. \quad (6)$$

The exponent of 2 in h_a is $a - 1$. Moreover $a \geq 2v$ gives $2^a \geq 3^v$, and $a \geq 3w$ gives $2^a \geq 5^w$. Thus $2^u 3^v 5^w \mid h_a$ after the stated onset, and BX_a differs from $h_a A / (2^u 3^v 5^w)$ by an integer and is therefore integral. Positivity, the recurrence and the bound follow from Proposition 4.5. $\square$

Every positive denominator splits in the required way: B is what remains after removing all factors 2, 3 and 5. For example, a hypothetical reduced denominator $2^3 3^2 5 \cdot 7$ would give $B = 7$; the next result gives the first integral $7X_a$ at $a = 4$, whereas the sufficient bound above gives $a_0 = 11$.

Corollary 4.7 (the first index at which the denominator clears). *Under the same lowest-terms hypothesis, put $M = 2^u 3^v 5^w$ and let den denote the positive reduced denominator. For $a \geq 1$,*

[Lean](#)
[Comparator](#)

$$\text{den}(BX_a) = \frac{M}{\gcd(M, h_a)}.$$

Consequently BX_a is integral exactly when $2^a \geq \max(2^{u+1}, 3^v, 5^w)$. The first such a can be found by integer comparisons, without logarithmic rounding.

Proof. By (6), BX_a differs from $h_a A/M$ by an integer. Since $\gcd(A, M) = 1$, reducing that fraction gives the displayed denominator. The three divisibility conditions for h_a are precisely the three inequalities stated above. $\square$

5 A window test and the remaining arithmetic

Two steps of the recurrence already show the finite data we need:

$$X_{\ell+2} = b_{\ell+1}b_\ell X_\ell - (b_{\ell+1}m_\ell + m_{\ell+1}).$$

If BX_ℓ is an integer, this identity determines the residue of $BX_{\ell+2}$ modulo the product of the two bases. We will compare the least positive representative of that residue with the upper bound for the endpoint tail.

For a window starting at $\ell \geq 0$ and having $h \geq 0$ steps, define its product of bases and accumulated numerator from (5) by

$$W_{\ell,0} = 1, \quad F_{\ell,0} = 0, \quad W_{\ell,h+1} = b_{\ell+h}W_{\ell,h}, \quad F_{\ell,h+1} = b_{\ell+h}F_{\ell,h} + m_{\ell+h}.$$

Induction on h gives

$$X_{\ell+h} = W_{\ell,h}X_\ell - F_{\ell,h}. \quad (7)$$

For integers $W \geq 1$ and t , define $\text{lpr}_W(t) = 1 + ((t - 1) \bmod W)$, using a remainder in $\{0, \dots, W - 1\}$. Thus a zero residue is represented by W , not 0. Throughout this section use the upper bound

$$K(B, a) = 90B(a + 1)^2. \quad (8)$$

Lemma 5.1 (least positive residues). *If d is a positive integer with $d \leq K$ and $d \equiv -BF \pmod{W}$, where $W \geq 1$, then $\text{lpr}_W(-BF) \leq K$.*

Lean
Comparator

Proof. Every positive representative of the residue is at least its least positive representative, so $\text{lpr}_W(-BF) \leq d \leq K$. $\square$

For example, the window starting at $\ell = 1$ with length 6 has $W_{1,6} = 648000$ and $F_{1,6} = 524431$. At $B = 1$ its least positive residue is 123569, exceeding $K(1, 7) = 5760$. This rules out an integral X_1 , but not a rational denominator that would clear only at a later index. The theorem therefore asks for windows beyond every prescribed starting index, as well as for every eligible denominator.

Theorem 5.2 (a residue criterion for irrationality). *The number S is irrational if and only if*

Lean
Comparator

$$\begin{aligned} &\text{for every } B \geq 1 \text{ with } \gcd(B, 30) = 1 \text{ and every } a_0 \geq 1, \\ &\text{there are } \ell \geq a_0, h \geq 1 \text{ such that } \text{lpr}_{W_{\ell,h}}(-BF_{\ell,h}) > K(B, \ell + h). \end{aligned} \quad (9)$$

Proof. Assume (9) and suppose S is rational. Theorem 4.6 supplies a positive integral carry $d_a = BX_a$ after its onset. Choose a window in (9) beyond that onset. Equation (7) gives $d_{\ell+h} \equiv -BF_{\ell,h} \pmod{W_{\ell,h}}$, contradicting Lemma 5.1 and the upper bound.

Conversely, let S be irrational and fix $B, \ell \geq 1$. Equation (6) makes BX_ℓ nonintegral. Put $\delta = \lceil BX_\ell \rceil - BX_\ell \in (0, 1)$. Since $W_{\ell,h} \geq 2^h$ and $X_{\ell+h} = O((\ell + h + 1)^2)$, for all sufficiently large h the number $\delta + BX_{\ell+h}/W_{\ell,h}$ lies in $(0, 1)$. The window identity shows that the integer $\lceil BX_\ell \rceil W_{\ell,h} - BF_{\ell,h}$ lies strictly between 0 and $W_{\ell,h}$ and is congruent to $-BF_{\ell,h}$. Hence

$$\text{lpr}_{W_{\ell,h}}(-BF_{\ell,h}) = \delta W_{\ell,h} + BX_{\ell+h}.$$

Its exponentially growing first term eventually exceeds $K(B, \ell + h)$. This proves (9), in fact from each fixed start. $\square$

The proved bound $90B(a+1)^2$ controls every positive integral BX_a that rationality would force. Polynomial growth alone does not suffice: the zero bound would make the residue inequality automatic. The companion, Section 7, treats smaller and more general bounds. No finite search establishes the unbounded quantifiers in (9).

The remaining task has the following three equivalent formulations.

Problem 5.3 (the repeated three-prime target). Prove that $S = \mathcal{R}_{\{2,3,5\}}$ is irrational.

Problem 5.4 (no reduced tail is an integer). Prove that for every $a \geq 1$ and every integer $B \geq 1$ coprime to 30,

$$BX_a \notin \mathbb{Z}. \quad (10)$$

Problem 5.5 (residue inequalities after every starting index). Prove (9).

Indeed, one integral scaled tail makes S rational by (6), whereas denominator clearing gives such tails from a rational S . Theorem 5.2 supplies the third equivalence. These arithmetic assertions remain unproved. The companion's finite certificate (Section 8) and comparisons with other approaches (Section 9) are supplementary, not premises of this argument.

Statements and declarations

Proof sources. Theorem 3.1 uses the transcendence theorem of Bugeaud and Laurent, which its Lean proof assumes. Fan's post of 26 June 2026 on the erdosproblems.com forum has priority for the two-prime deduction [7]. Every other theorem, proposition, lemma and corollary has a Lean proof of its whole statement, marked in the margin; Appendix A describes the formal sources. Neither these proofs nor the finite calculations supply the required windows for every eligible denominator and arbitrarily late starts.

Artefact and data availability. The companion contains the detailed bounds and supplementary arguments; its source inventory explains the normalisation conventions of the Lean sources. AI agents carried out most of the research and drafting. The work has not had independent human mathematical review.

Funding and competing interests. This work received no external funding. The author declares no competing interests.

Acknowledgements. The problem numbering and its status as of 28 July 2026 are taken from the Erdős Problems catalogue maintained by Thomas Bloom [4]. Earlier mathematical inputs are credited at their points of use. We thank Wouter van Doorn for advice on exposition, especially on unexplained terminology, unnecessary notation and the need to explain restrictive hypotheses. His advice concerned a note on Problem #243; he has not reviewed the mathematics of this paper.

A Guide to the formal sources

There are two separate arguments. The rank proof uses the height formula, row and column rescaling, and a threshold determinant. The arithmetic argument uses the shell coefficients, the tail recurrence, denominator clearing and least positive residues. The Lean proof of Theorem 5.2 establishes the equivalence of the residue condition with irrationality; it does not establish the residue condition itself.

Library declarations. The margin marks link each result to its Lean proof. The main steps rest on the following library declarations, each opened at a fixed revision. Running LCM: the exact height formula; constancy on each logarithmic cell; the jumps by p , q and r ; exactly 3 count + 1 jump values; the quadratic bound on exponent shells. Dyadic blocks: at most one internal pure-power jump per channel; the radix is 2, 6, 10 or 30; the radix lies in $[2, 30]$. Kernel rank: the kernel is not of rank one; the smallest 2×2 minor equals $-1/15$; the tail-state recurrence. Residues and carries: no bounded positive state represents an escaping residue (integer form); the local-window escape condition and its exclusion of a positive reduced carry; a bounded-radix orbit is integral or cofinally far from the integers; no uniformly bounded nonzero carry lift; block-null words are channel coboundaries; the carry as a residue digit plus a coboundary; a finite-dimensional factorisation gives a finite realised span.

Scope of the supplementary literature. The companion develops the comparisons omitted from the main argument. Its Section 5 relates the bounded-ratio argument to Erdős–Taylor [10, Theorem 1, p. 600] and Fan [11, Lemma 3.1, p. 7]; Section 6 compares the exact carry restrictions and polynomial-radix assumptions of Hančl–Tijdeman [12, Theorems 2.2, 3.1 and 4.2]. The Isabelle development of Koutsoukou–Argyragi and Li [18] verifies classical criteria, not the present series.

Section 9 of the companion separates the finite-difference hypotheses of Luca–Ouaknine–Worrell [13, Definition 5, Theorems 6 and 8, Claim 10] from quadratic growth alone, and checks the obstacles to using the echoing [14], digit-complexity [19, Theorem 1] and multivariate Mahler [20] results. These comparisons do not establish the missing hypotheses for S .

For broader background, the companion distinguishes work on fixed-prime semi-groups [15, 16] from other unit-fraction irrationality problems [17]. No estimate from those papers is used in this note’s counting argument.

References

- [1] Paul Erdős and Ronald L. Graham, *Old and New Problems and Results in Combinatorial Number Theory*, Monographies de L'Enseignement Mathématique **28**, L'Enseignement Mathématique (1980), [source](#).
- [2] Paul Erdős, *On the irrationality of certain series: problems and results*, in *New Advances in Transcendence Theory*, Cambridge University Press (1988), 102–109, [doi:10.1017/CB09780511897184.009](#).
- [3] Paul Erdős, *Letter to the Editor*, Fibonacci Quarterly **12**, no. 4 (1974), 335, [source](#).
- [4] Thomas F. Bloom, *Erdős Problem #269* (2026), [source](#). Accessed 28 July 2026.
- [5] Yann Bugeaud and Michel Laurent, *Transcendence and continued fraction expansion of values of Hecke–Mahler series*, Acta Arithmetica **209** (2023), 59–90, [doi:10.4064/aa220323-18-1](#); arXiv:2203.12901.
- [6] John H. Loxton and Alfred J. van der Poorten, *Arithmetic properties of certain functions in several variables III*, Bulletin of the Australian Mathematical Society **16** (1977), 15–47, [doi:10.1017/S0004972700022978](#).
- [7] Steve Fan, *Comment on Erdős Problem #269, thread 269, post 7218* (2026), [source](#). Public forum post, 26 June 2026, thread 269, post 7218.
- [8] Paul Erdős and Ernst G. Straus, *On the irrationality of certain series*, Pacific Journal of Mathematics **55**, no. 1 (1974), 85–92, [doi:10.2140/pjm.1974.55.85](#).
- [9] Jaroslav Hančl and Robert Tijdeman, *On the irrationality of Cantor and Ahmes series*, Publicationes Mathematicae Debrecen **65**, no. 3–4 (2004), 371–380, [doi:10.5486/PMD.2004.3254](#).
- [10] Paul Erdős and S. James Taylor, *On the set of points of convergence of a lacunary trigonometric series and the equidistribution properties of related sequences*, Proceedings of the London Mathematical Society **s3-7**, no. 1 (1957), 598–615, [doi:10.1112/plms/s3-7.1.598](#).
- [11] Steve Fan, *Strongly complete sets and a conjecture of Erdős* (2026), [source](#); arXiv:2607.14071. The cited Lemma 3.1 is in arXiv v1, 15 July 2026.
- [12] Jaroslav Hančl and Robert Tijdeman, *On the irrationality of polynomial Cantor series*, Acta Arithmetica **133**, no. 1 (2008), 37–52, [doi:10.4064/aa133-1-3](#).
- [13] Florian Luca, Joël Ouaknine and James Worrell, *Transcendence of Hecke–Mahler Series*, Bulletin of the London Mathematical Society **57**, no. 5 (2025), 1360–1368, [doi:10.1112/blms.70033](#); arXiv:2412.07908. Numbered references use the published article.

- [14] Pavol Kebis, Florian Luca, Joël Ouaknine, Andrew Scoones and James Worrell, *On Transcendence of Numbers Related to Sturmian and Arnoux-Rauzy Words*, in *51st International Colloquium on Automata, Languages, and Programming (ICALP 2024)*, Leibniz International Proceedings in Informatics **297**, Schloss Dagstuhl – Leibniz-Zentrum für Informatik (2024), 144:1–144:15, doi:[10.4230/LIPIcs.ICALP.2024.144](https://doi.org/10.4230/LIPIcs.ICALP.2024.144).
- [15] Robert Tijdeman and H. G. Meijer, *On integers generated by a finite number of fixed primes*, *Compositio Mathematica* **29**, no. 3 (1974), 273–286, [source](#).
- [16] Alessandro Languasco, Florian Luca, Pieter Moree and Alain Togbé, *Sequences of integers generated by two fixed primes*, *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg* **95** (2025), 123–148, doi:[10.1007/s12188-025-00293-9](https://doi.org/10.1007/s12188-025-00293-9); arXiv:[2309.12806](https://arxiv.org/abs/2309.12806).
- [17] Vjekoslav Kovač and Terence Tao, *On several irrationality problems for Ahmes series*, *Acta Mathematica Hungarica* **175** (2025), 572–608, doi:[10.1007/s10474-025-01528-0](https://doi.org/10.1007/s10474-025-01528-0); arXiv:[2406.17593](https://arxiv.org/abs/2406.17593).
- [18] Angeliki Koutsoukou-Argyaki and Wenda Li, *Irrationality Criteria for Series by Erdős and Straus*, *Archive of Formal Proofs* (2020), [source](#). Entry dated 12 May 2020; proof-document version consulted: 6 February 2026.
- [19] Boris Adamczewski and Yann Bugeaud, *On the complexity of algebraic numbers I. Expansions in integer bases*, *Annals of Mathematics* **165**, no. 2 (2007), 547–565, doi:[10.4007/annals.2007.165.547](https://doi.org/10.4007/annals.2007.165.547).
- [20] Boris Adamczewski and Colin Faverjon, *Mahler’s method in several variables and finite automata*, *Annals of Mathematics* **204**, no. 2 (2026), 455–533, doi:[10.4007/annals.2026.204.2.1](https://doi.org/10.4007/annals.2026.204.2.1). Online 13 September 2026; locators here refer to the 68-page [author manuscript](#).