

The Binary Totient Series: a complete working record of Erdős Problem 249

Will Cook

July 25, 2026

Abstract

This document is the complete working record of one attack on Erdős Problem 249: whether $S = \sum_{n \geq 1} \varphi(n)/2^n$ is irrational. The problem is open, and nothing here decides it. What the document does is collect, in one place and in a form that can be used directly as premises, every result the project holds: the Lean-checked theorems with their exact hypotheses and source sites, the exact finite computations, the coordinates in which the constant can be written, the routes that have been closed and the mechanism that closed each one, and the exact statements that remain open. Results are sorted by the scale at which they are proved, because that is where the problem stands or falls: everything proved here holds at fixed or bounded scale, while the remaining obligation is cofinal.

How to read this document

This is a working record, not a survey. It is written to be read in full before work begins on the problem, so that every statement the project has established is available as a premise rather than as a reference to be chased. Three conventions carry that load.

Evidence bands. Every statement is tagged. [Lean] means a proof term was checked by the pinned Lean kernel. [Cert] means an exact finite computation, with no floating-point decision anywhere in it. [Math] means proved in ordinary mathematics in the sources but not formalised. [Cited] means established in the published literature. [Open] means not proved. These are never blurred, and a statement carrying one band is never described in language belonging to another.

Scale. Every parameter-indexed statement is tagged `scale:fixed` (proved at specific listed values), `scale:bounded` (proved below an explicit bound), `scale:cofinal` (proved for arbitrarily large parameters), or `scale:uniform` (proved for all parameters). This axis is the reason the problem is still open: the corpus proves at fixed and bounded scale, and the remaining obligation is cofinal. Sorting by scale puts that boundary where it can be seen.

Coordinates. Every statement records the representation it is expressed in. An obstruction is a fact about a coordinate, not about the object, and a wall measured in one representation may simply not exist in another; the atlas section gives the transport maps, so any obstruction recorded here can be re-measured elsewhere.

Erdős Problem 249 is open. No section of this document claims otherwise, and a reduction of the problem to another statement is recorded as a reduction, never as progress toward a solution.

1 The problem, and what is actually known

Let φ be Euler's totient function and put

$$S := \sum_{n \geq 1} \frac{\varphi(n)}{2^n} = 1 + \frac{1}{4} + \frac{2}{8} + \frac{2}{16} + \frac{4}{32} + \cdots = 1.6066 \dots$$

The series converges absolutely because $\varphi(n) \leq n$. Erdős Problem #249 asks whether S is irrational. *It is open*. Nothing in this document decides it, no route recorded here is a proof, and no result here should be read as an approach that is close to working. What this document does is different in kind: it assembles every recorded failure, obstruction, countermodel and dead route the programme has produced against #249, and classifies each one by the exact class of argument it eliminates. The claim being made is that those failures are not independent — they are repeated measurements of a single obstruction, and that obstruction has a shape which can be stated.

The open question has a sharp equivalent inside the corpus, and it is the statement everything below is measured against. Write $2^N S = \Phi_N + R_N$ with $\Phi_N := \sum_{n \leq N} \varphi(n) 2^{N-n} \in \mathbb{N}$ and $R_N := \sum_{j \geq 0} \varphi(N+1+j)/2^{j+1} \in [0, \infty)$ ([Erdos249257.TotientTailPeriodKiller.totientTail](#) [TotientTailPeriodKiller.lean:150]). If $S \in \mathbb{Q}$ then, applying Euler's theorem to the odd part of its denominator, there are $h > 0$ and N_0 with $R_{N+h} - R_N \in \mathbb{Z}$ for every $N \geq N_0$ ([Erdos249257.eventual_period_of_not_irrational](#) [TotientTailPeriodKiller.lean:358], explicit witnesses $h = \varphi(\text{oddPart}(r.\text{den}))$, $N_0 = v_2(r.\text{den})$). So it suffices to refute integrality cofinally on every period ray. The finite device that refutes it is a residue certificate:

$$\text{windowDiscrepancy}(h, N, L) := \sum_{j < L} (\varphi(N+h+1+j) - \varphi(N+1+j)) 2^{L-1-j} \in \mathbb{Z},$$

the depth- L truncation of $2^L(R_{N+h} - R_N)$, and

$$\text{certifiedKill}(h, N, L) :\iff (N+h+L+2) < \text{windowDiscrepancy}(h, N, L) \bmod 2^L < 2^L - (N+h+L+2),$$

a decidable condition ([Erdos249257.TotientTailPeriodKiller.windowDiscrepancy](#) [TotientTailPeriodKiller.lean:72]). The radius $N+h+L+2$ is exactly the crude tail bound coming from $\varphi(m) \leq m$, so a certificate is sound: it forces $R_{N+h} - R_N \notin \mathbb{Z}$, and in fact certificates are *complete* for non-integrality ([Erdos249257.exists_certifiedKill_iff_tail_diff_notMem_int](#) [LcmConeFlatness.lean:316]). The entire #249 side of the programme therefore reduces to one obligation.

Definition 1.1 (The supply obligation Sep — the exact open target).

$$\text{Sep} :\iff \forall h \geq 1 \forall N_0 \exists N \geq N_0 \exists L, \text{certifiedKill}(h, N, L).$$

$\text{Sep} \Rightarrow \text{Irrational}(S)$ ([Erdos249257.irrational_totient_series_of_certificate_supply](#) [TotientTailPeriodKiller.lean:360], coord:binary-digit scale:cofinal [Lean])

Two features of Sep govern everything that follows. First, the quantifier shape is $\forall \forall \exists \exists$: the corpus can and does verify individual instances, but an instance is not the obligation. Second, depth cannot stay bounded: from $\text{certifiedKill}(h, N, L)$ one gets $2(N+h+L+2) < 2^L$ immediately ([Erdos249257.certifiedKill_depth_floor](#) [TotientTailPeriodKiller.lean:79], [Lean]), so $L \gtrsim \log_2(N+h)$. A certificate is a demand that a weighted accumulation over a window of length $\approx \log_2 N$ lands near the centre of its dyadic arc, at full arithmetic resolution.

1.1 The unconditional record

The following hold with no irrationality hypothesis. They are the results a specialist should know before reading further; each is stated at the scale at which it is actually proved.

Theorem 1.2 (Denominator exclusion — the headline unconditional fact). *If $S \in \mathbb{Q}$ then its reduced denominator exceeds $Q_0 := 79\,639\,646\,646\,701\,375\,323\,355\,774\,875\,831\,053 \approx 7.96 \times 10^{34}$. Equivalently, $S \neq p$ for every $p \in \mathbb{Q}$ with $p.\text{den} \leq Q_0$. The bound is sharp for the method: $q = Q_0 + 1$ is the exact first failing denominator, exhibited as the mediant of two explicit unimodular Farey neighbours. coord:farey scale:bounded [Lean] [Erdos249257.tsum_totient_div_pow_two_ne_ratCast_of_den_le_79639646646701375323355774875831053](#) [CertificateKernel.lean:18056] [Erdos249257.gap_check_window_1_240_first_failure](#) [GapFareyBound.lean:225]*

This is obtained from a classical Stern–Brocot gap lemma ([farey_gap](#) [GapFareyBound.lean:51]) applied at window $K = 240$, the last rung of the ladder $4838 \rightarrow 2^{22} \rightarrow 2.49 \times 10^{17} \rightarrow Q_0$. It is logically independent of Sep: it uses no certificate and no period apparatus. Its own open continuation is whether $\sup_K(b + d)(K) = \infty$; that would close #249 through this theorem’s consumer. [Open]

Proposition 1.3 (Finite certificate deposits). *certifiedKill has been verified at: the 28 diagonal instances of the LCM pincer through $t = 64$ ([Erdos249257.certifiedKill_diagonal_all_imported_through_t64](#) [DiagonalPincerCertificatesT64.lean:1967], endpoint [Erdos249257.certifiedKill_diagonal_t64](#) [DiagonalPincerCertificatesT64.lean:1928]); all shifts $h \in [1, 16]$ simultaneously at $(N, L) = (14, 9)$, by decide ([Erdos249257.certifiedKill_all_upto_sixteen](#) [CarrySurvivorExtinction.lean:5]) and eight further period deposits at $N = 300$. Every deposit fires within a small additive constant of the minimum depth permitted by `certifiedKill_depth_floor`. Note the quantifier order: this deposit proves $\exists N \exists L \forall h \leq 16$, whereas Sep needs $\forall h \forall N_0 \exists N \exists L$. coord:binary-digit scale:fixed [Lean]*

Proposition 1.4 (Cofinal positivity, and why it is exactly half a certificate). *The true actual-LCM tail difference is strictly positive for every $a \geq 8$, with no irrationality hypothesis ([Erdos257PeriodNoncollapse.shift_pos](#) [TotientActualLcmOrbitSign.lean:39], [Erdos257PeriodNoncollapse.actualLcmTailOrbit_pos](#) [TotientActualLcmOrbitSign.lean:144]). In the same coordinate, integrality of the orbit forces the residue to the top edge, exactly $2^K - e$ ([Erdos257PeriodNoncollapse.actualLcm_integral_forces_topEdgeResidue](#) [TotientActualLcmOrbitSign.lean:211]). Positivity does not exclude the top edge. So the strongest cofinal fact the corpus owns about the actual object supplies one of the two inequalities a certificate needs and provably cannot supply the other. coord:actual-lcm scale:cofinal [Lean]*

Proposition 1.5 (Rationality forces unbounded carry rank). *If S is rational then, for every e , the dyadic carry-kernel family at level e has $\mathbb{Q}$ -rank at least $2^e - 1$ ([Erdos249257.not_irrational_totientSeries_implies_unbounded_carryRank_unconditional](#) [TotientCarryKernelRigidity.lean:100]). The scale side is complete and uniform in e : the canonical family of $2^e + 1$ dyadic totient-kernel channels is linearly independent over $\mathbb{Q}$ at every depth, via CRT plus Dirichlet ([Erdos249257.linearIndependenceCanonicalTotientKernelFamily](#) [TotientMahlerDefect.lean:900]), so the full family spans an infinite-dimensional space ([Erdos249257.not_finiteDimensional_span_fullTotientKernel](#) [TotientMahlerDefect.lean:900]). coord:carry-rank scale:uniform [Lean]*

Proposition 1.6 (What rationality actually buys, and what it does not). *Rationality gives uniform eventual periodicity of the carry’s dyadic sections modulo v ; that periodicity provably does not promote to any $\mathbb{Q}$ -rank bound on the carry family ([Erdos257PeriodNoncollapse.not_irrational_totientSeries_implies_mod_period_and_unbounded_rank](#) [TotientTailCarryPeriod.lean:224]). coord:carry-rank scale:uniform [Lean]*

Proposition 1.7 (Exact reformulations that do not move the truth value). *Three independent re-encodings of Sep are proved equivalent to Irrational(S), not merely sufficient for it: the period-multiple kill supply (`periodMultipleKillSupply_iff_irrational` [PeriodMultipleEscape.lean:153]), the directed-certificate supply and its LCM specialisation (`Erdos257PeriodNoncollapse.CofinalDirectedLcm`) and the window-separated-pairs predicate (`Erdos249257.dtwWindowSeparatedPairs_iff_irrational_totient_series` [PivotAntiReconstruction.lean:1775]).* coord:binary-digit scale:cofinal [Lean]

Proposition 1.8 (A rational sequence indistinguishable from φ by every coarse invariant). *There is $c : \mathbb{N} \rightarrow \mathbb{N}$ with $c(n) \leq 6$ and $c(n) \leq n$ for all n , $c(n) \equiv \varphi(n) \pmod{2}$ for every n , and c not eventually periodic — indeed for every N, G, K there are K explicit carry pulses beyond N , pairwise separated by more than G — and yet $\sum_n c(n)/2^n = 3/2 \in \mathbb{Q}$ (`Erdos257PeriodNoncollapse.exists_totientParity_arbitrarilyManySeparatedCarry_rational_countermodel` [TotientParityCoboundaryCountermodel.lean:637]; sum at :359, aperiodicity at :487, parity match at :194; `#print axioms` clean). coord:coefficient-word scale:uniform [Lean]*

Propositions 1.4–1.8 are the four facts a reader should carry into the next section: the corpus’s best cofinal information is half a certificate, its best rank information runs the wrong way, its reformulations are equivalences rather than reductions, and every purely qualitative property of the coefficient word is satisfied by a rational countermodel.

2 The wall

This programme has produced many exact reformulations of #249 and no proof. That is the honest summary, and the rest of this section is an argument that it is also the wrong way to read the record. The reformulations do not fail independently. They fail in a small number of recurring ways, and when each failure is stated as a claim about *which class of argument it eliminates* rather than as a report that something did not work, the classes fit together into one obstruction with a describable shape. A hundred reformulations that all die are not a hundred failures; they are a hundred measurements of one wall, taken from different angles. The measurements are below.

The thesis, in one sentence: *every barrier in the record is a bound, and the quantity a certificate measures is invariant under exactly the bounds that make an argument finite.*

2.1 The plane the barriers live on

Fix two axes for a hypothetical proof of Sep (Definition 1.1): the *range* of indices at which it consults φ , and the *resolution* at which it consults them. Two derived axes matter as well: the size of the proof’s internal bookkeeping (carry state, $\mathbb{Q}$ -rank, shift-polynomial degree), and whether the target is asserted *pointwise* at a chosen index or as an *average* over a block.

- The quadrant [full resolution $\times$ bounded range] is closed by Theorem 2.4 (B1): φ may be pinned exactly on $[1, B]$ and the series can still be rational.
- The complementary quadrant [coarse resolution $\times$ unbounded range] is closed by Proposition 1.8 (B7): φ ’s parity may be pinned at *every* index, together with boundedness, $c(n) \leq n$, and arbitrarily strong aperiodicity, and the series can still be $3/2$.
- The remaining quadrant is closed *whenever the proof compresses*: B5 (no bounded carry state, no fixed-precision signature) and B6 (four independent finite truncations of the totient kernel, all with trivial kernel).
- B4 closes the one construction that tried to reach the good quadrant by *prescribing* values: residue engineering pays for amplitude in position, and position enters the certificate radius.

- B2 closes the escape of retargeting; B3 records that no bounded result in the corpus has ever promoted.

The corpus's own results partition along these axes perfectly, which is the first evidence that the axes are the right ones. Every unconditional *finite* deposit (Proposition 1.3, the $K = 240$ Farey rung of Theorem 5.6, the actual-LCM orbits at $a = 4$ and $a = 6$) sits at full resolution and bounded range, so B1 says extending them is evidence forever and proof never. Every unconditional *cofinal* theorem (letterwise positivity for all $a \geq 8$, the tail bounds from $\varphi(m) \leq m$, unbounded Mersenne-shadow denominator growth) sits at coarse resolution and unbounded range, and Proposition 1.4 proves in the corpus's own coordinate that this is exactly half a certificate.

2.2 B1: the finite-inspection barrier, as a theorem

B1 is the one barrier that is a genuine no-go theorem about proof method, so it is stated and proved as one. The generic vocabulary is needed first. For $c : \mathbb{N} \rightarrow \mathbb{N}$ with $c(n) \leq n$, write $T_c := \sum_{n \geq 1} c(n)/2^n$, $R_N^c := \sum_{j \geq 0} c(N+1+j)/2^{j+1}$, $A_c(h, N, L) := \sum_{j < L} (c(N+h+1+j) - c(N+1+j))2^{L-1-j}$, and let $\text{Kill}_c(h, N, L)$ and Sep_c be `certifiedKill` and `Sep` with A_c in place of `windowDiscrepancy`. For $c = \varphi$ these are the objects of Definition 1.1.

Lemma 2.1 (Generic soundness). $\text{Kill}_c(h, N, L) \Rightarrow R_{N+h}^c - R_N^c \notin \mathbb{Z}$. *coord:binary-digit scale:uniform [Math]*

Proof. From $c(n) \leq n$ one gets $|2^L(R_{N+h}^c - R_N^c) - A_c(h, N, L)| < N+h+L+2$; this is the crude tail estimate, formalised for φ at [Erdos249257.tail_after_le](#) [TotientTailPeriodKiller.lean:198] and used there to prove [Erdos249257.tail_diff_notMem_int_of_certifiedKill](#) [TotientTailPeriodKiller.lean:262]. If $R_{N+h}^c - R_N^c = k \in \mathbb{Z}$ then $2^L k \equiv 0 \pmod{2^L}$ and A_c lies within $N+h+L+2$ of $2^L k$, so $A_c \bmod 2^L$ lies within $N+h+L+2$ of 0 or of 2^L , contradicting Kill_c . $\square$

Lemma 2.2 (Generic tail-period law). *If $T_c = p/(2^e m)$ with m odd and $h := \text{ord}_m(2)$, then $R_{N+h}^c - R_N^c \in \mathbb{Z}$ for every $N \geq e$.* *coord:binary-digit scale:cofinal [Math]*

Proof. $R_N^c = 2^N T_c - \Phi_N^c$ with $\Phi_N^c \in \mathbb{Z}$, so $R_{N+h}^c - R_N^c = 2^N(2^h - 1)T_c - (\Phi_{N+h}^c - \Phi_N^c)$, and $2^N(2^h - 1)p/(2^e m) \in \mathbb{Z}$ because $2^e \mid 2^N$ and $m \mid 2^h - 1$. $\square$

Remark 2.3. Lemmas 2.1 and 2.2 are elementary and are stated here in ordinary mathematics. Their φ -instances are Lean theorems ([Erdos249257.tail_diff_notMem_int_of_certifiedKill](#) [TotientTailPeriodKiller.lean:262], [Erdos249257.eventual_period_of_not_irrational](#) [TotientTailPeriodKiller.lean:262]) and the generic tail-period direction exists on disk inside [binaryCoeffSeries_rational_iff_exists_temperedBinaryOrbit](#) [GenericTailOrbitRigidity.lean:426]; a generic *named* theorem of the form of Lemma 2.2 does not exist in either tree. The composition below is mechanical but unformalised, and is flagged [Math] throughout for that reason.

Theorem 2.4 (B1, the γ -splice: finite inspection cannot certify the supply). *Let $B \geq 1$ and let $P > B$. Define $\gamma : \mathbb{N} \rightarrow \mathbb{N}$ by*

$$\gamma(n) := \varphi(n) \quad (n \leq B), \quad \gamma(n) := \begin{cases} n-1, & P \mid n \\ n, & P \nmid n \end{cases} \quad (n > B).$$

Then:

- (i) $\gamma(n) \leq n$ for all n , so γ lies in the same coefficient class as φ ;

- (ii) $\gamma(n) = \varphi(n)$ for every $n \leq B$, and consequently $A_\gamma(h, N, L) = A_\varphi(h, N, L)$ for every (h, N, L) with $N + h + L \leq B$;
- (iii) $T_\gamma = D - 1/(2^P - 1) \in \mathbb{Q}$, where $D = 2 - \sum_{n \leq B} (n - \varphi(n))/2^n$, and the odd part of the reduced denominator of T_γ is exactly $2^P - 1$;
- (iv) Sep_γ is false — indeed it fails already at $h = P$.

Hence for every B there is a coefficient sequence in the same class, agreeing with φ on all of $[1, B]$, whose supply obligation is false. *coord:binary-digit scale:uniform [Math]*

Proof. (i) is immediate; (ii) holds because $A_\varphi(h, N, L)$ reads φ only at indices $\leq N + h + L$. For (iii), since $P > B$ every multiple of P exceeds B , so

$$T_\gamma = \sum_{n \geq 1} \frac{n}{2^n} - \sum_{n \leq B} \frac{n - \varphi(n)}{2^n} - \sum_{k \geq 1} 2^{-kP} = 2 - \sum_{n \leq B} \frac{n - \varphi(n)}{2^n} - \frac{1}{2^P - 1}.$$

Write $D = A/2^B$ with $A = 2^{B+1} - \sum_{n \leq B} (n - \varphi(n))2^{B-n} \in \mathbb{Z}$. Then $T_\gamma = (A(2^P - 1) - 2^B)/(2^B(2^P - 1))$, and $\gcd(A(2^P - 1) - 2^B, 2^P - 1) = \gcd(2^B, 2^P - 1) = 1$ because $2^P - 1$ is odd. So no factor of $2^P - 1$ cancels, and the odd part of the reduced denominator is exactly $2^P - 1$. For (iv), the odd part of the denominator is $m = 2^P - 1$ and $\text{ord}_m(2) = P$, while $v_2(\text{den}) \leq B$. By Lemma 2.2, $R_{N+P}^\gamma - R_N^\gamma \in \mathbb{Z}$ for every $N \geq B$, so by Lemma 2.1 no $\text{Kill}_\gamma(P, N, L)$ holds for any $N \geq B$ and any L . Taking $N_0 = B$ refutes the inner existential of Sep_γ at $h = P$. $\square$

Corollary 2.5 (What B1 rules out). *No argument establishing Sep can have its totient input confined to $\{\varphi(n) : n \leq B\}$ for a fixed B . Equivalently, no inference of the form “Sep has been verified at every scale up to B , for arbitrarily large B , therefore Sep holds cofinally” is valid: the hypothesis is consistent with the conclusion’s negation at every B .*

Remark 2.6 (Scope discipline — what B1 does *not* say). Theorem 2.4 is a statement about proof method, not about φ . It does not touch S , and it does not suggest that φ is such a γ ; it says only that no fixed-horizon check can tell them apart. It also does *not* kill bounded-parameter certificates in general. “Bounded parameter” splits into four independent bounds with four different killers: bounded index range (this theorem), bounded certificate depth L (killed separately, and by a Lean theorem — `certifiedKill_depth_floor` forces $2(N + h + L + 2) < 2^L$), bounded proof state (B5), and bounded rank (B6). Conflating them would overstate B1.

The Lean-formalised sibling of the same mechanism — a lacunary zero-valued coboundary splice — is Proposition 1.8, which is B7. B1 and B7 are the two object-level witnesses of the wall, one along each axis.

2.3 B2–B7

Each barrier below is stated as a claim about a class of arguments, with its status never blurred. *Proved* means there is a theorem (Lean or ordinary mathematics, as marked) whose content is the elimination. *Observed pattern* means an audit over the corpus: real evidence, not a theorem, and it is never used as a premise elsewhere in this document.

Proposition 2.7 (B2, route-collapse: intermediate targets are not waypoints). **Status: proved** (*Lean, three independent mechanisms*).

- (a) Certificate completeness. *Every certificate vocabulary is an iff with the underlying non-integrality, so every re-encoding of Sep is equivalent to Sep* (`periodMultipleKillSupply_inf_irrational` [`PeriodMultipleEscape.lean:153`], forward at :132, converse at :143).

- (b) Sample choice. Any weakening in which the prover may choose which indices to test collapses to Irrational(S) outright, because irrationality plus doubling expansivity already supplies a two-point sample with the required separation ([Erdos249257.dtwWindowSeparatedPairs_iff_irrational_totient_series](#) [[PivotAntiReconstruction.lean:1775](#)]). The mechanism is visible in the converse proof ([windowSeparatedPairsAt_of_cofinally_scaled_adjacent_chord](#), [PivotAntiReconstruction.lean:1663](#)): it selects $T = \{N, N+1\}$ and two ordered pairs, and the counted-energy threshold $2|T|^2/5 = 8/5 \leq 2\delta^2$ is met by $\delta = 9/10$.
- (c) Multi-point enrichment. Adding vertices, higher-order differences, or finite shift-polynomial combinations on the same ray buys nothing, because integrality transports affinely along every ray $H \mapsto kH$: the four-hit diamond $\text{Hit}(H) \wedge \text{Hit}(pH) \wedge \text{Hit}(qH) \wedge \text{Hit}(pqH)$ is equivalent to $\text{Hit}(H)$ alone, with no primality used ([Erdos249257.fullTarget_primeAdjunction_diamond_iff_root](#) [[FullTargetPrimeAdjunctionNoGo.lean:196](#)]).

Rules out: “find an easier waypoint” as a strategy class. **Does not rule out:** targets over a fixed full block with no sample choice, and targets demanding a uniform quantitative margin that irrationality does not supply. These two exemptions are read off the collapse proofs themselves and are exactly what the surviving routes in §3 exploit. *coord:binary-digit scale:cofinal* [Lean]

Remark 2.8. A corroborating instance: rank-2 second-difference certificates are sound but measurably not shallower than rank-1 — at $(h, N) = (1, 8)$ rank-1 fires at depth 8 and no rank-2 certificate exists at depth ≤ 8 ([totient_tail_rank_two_kill_sound_but_not_shallower_cell](#), [CertificateKernel.lean:18762](#), [Lean]). Separately, the claim sometimes made that the Farey growth law $\sup_K(b+d) = \infty$ is “equivalent in difficulty to #249” is an argument, not an *iff*: the bound produced is the convergent denominator of the underlying constant, which stalls at q_0 precisely if $S = a/q_0$. It should not be cited as an equivalence. [Math]

Proposition 2.9 (B3, no free promotion). **Status:** *observed pattern (an audit, not a theorem)*. Across the corpus’s proved results, no bounded or partial result has a proof uniform enough to be promoted to cofinal scale by routine strengthening — raising a bound, widening a hypothesis, or reindexing a quantifier. Method: 56 near-miss rows were catalogued against the four open obligations; 18 were flagged promotable on a statement-level read; all 18 proof bodies were then opened. Verdict: 16 conclusively not promotable with a named blocker each, and 2 promotable with no new mathematics but both pure restatements that widen a target family without supplying arithmetic content. Blocker taxonomy: hypothesis strength $\times 6$, scale-only $\times 5$, coordinate-only $\times 4$, multiple $\times 3$. Rows with an outright quantifier-order mismatch were judged unpromotable before the body audit and excluded, so the audited 18 are the most favourable subset. *coord:audit scale:n/a* [Cert]

Observation 2.10 (The sharpest signal in the audit). Every not-promotable #249-supply row asks for a *pointwise* fact at a specially chosen index: one large a with a top-edge residue gap, one q with a terminal-dominance inequality, one exponent past $a = 6$, one t past 64, one sign at one LCM jump, one prime per LCM height, one K past 240. Not one such index has ever been located unconditionally beyond the finite census. The single audited row whose missing input is not of that shape is the first-harmonic gap — an *average* over a block. This contrast is what selects the leading survivor in §3. It is evidence, not proof.

Proposition 2.11 (B4, amplitude versus radius: residue engineering is self-defeating). **Status:** *proved* (Lean construction plus an unconditional arithmetic margin). A certificate demands that the window residue sit at distance $> N+h+L+2$ from both 0 and 2^L modulo 2^L , and the depth floor forces $2^L > 2(N+h+L+2)$, so the required amplitude is comparable to the modulus. Any producer that buys that amplitude by prescribing totient values at engineered positions pays for the prescription in position, and position enters the radius. The corpus’s strongest such construction proves the trade is exactly self-cancelling, unconditionally and at every depth: a

two-adic pulse block lands the residue at the arc centre $2^{K-1} \bmod 2^K$ — the ideal target — and still cannot fire, because its own defining congruence $p \equiv 1+2^{K-1} \pmod{2^K}$ forces $p \geq 1+2^{K-1}$, so with $N = p - K$, $h = H$, $L = K$ the radius $p + H + 2$ exceeds 2^{K-1} for every prime p and every K . `coord:two-adic-pulse scale:cofinal [Lean] Erdos257PeriodNoncollapse.exists_prime_deltaTotient_twoAdic_pulseBlock [TotientTwoAdicPulseBlock.lean:211] Erdos257PeriodNoncollapse.windowModEq_half_of_twoAdic_pulse [TotientTwoAdicPulseBlock.lean:262]`

Remark 2.12 (B4’s own recorded repair, and its scope). Two further Lean facts show that prescribing *letters* is exhausted: the full terminal dyadic staircase is unconditionally impossible — its terminal letter would have to be positive, strictly below a wider modulus, and divisible by it, hence 0 (`Erdos257PeriodNoncollapse.not_actualLcmTerminalDyadicStaircase_of_room [TotientActualLcmTopEdgeStaircase.lean:251]`); and the surviving *punctured* staircase pins its penultimate letter to exactly 2^{m-1} with $2^m < 2(2H+J+K+2)$, i.e. no slack at all (`puncturedDyadicStaircase.penultimate_eq_half, TotientActualLcmTopEdgeStaircase.lean:1187`). The repair the barrier permits is explicit: impose the half-turn on the *word* rather than on one delta, i.e. ask for cofinally many (h, N, L) with $\text{windowDiscrepancy}(h, N, L) \equiv 2^{L-1} \pmod{2^L}$ and $2^{L-1} > N+h+L+2$. What is *proved* is the failure of the Dirichlet/CRT residue-engineering family by an explicit exponential margin. The broader reading — that only carry accumulation across the weighted word can produce the required amplitude, since one letter obeys $|\varphi(n+h) - \varphi(n)| < n+h$ while the weighted word can reach 2^L — is an inference from the proved instance plus the linear growth bound, not itself a theorem on disk. [Math]

Proposition 2.13 (B5, bounded state and local signatures). *Status: proved (three Lean theorems, all problem-agnostic).*

- (i) No bounded or autonomous carry state summarising the history before position m determines the tail after m : for a balanced-pulse family whose predecessor state is constant, no decode recovers the radius parameter, and any finite state type needs cardinality $\geq \lfloor m/2 \rfloor + 2$, unbounded in m (`Erdos249257.balancedPulse_no_autonomous_decoder [GenericTailOrbitRigidity.lean:247]`).
- (ii) Long common suffixes erase predecessor information exactly: two affine binary orbits with different seeds satisfy $\text{orbit}_u(L) - \text{orbit}_v(L) = 2^L(u_0 - v_0)$, so the endpoint residue mod 2^L is independent of the initial carry (`Erdos249257.affineBinaryOrbit_mod_twoPow_eq [GenericTailOrbitRigidity.lean:307]`).
- (iii) Bounded local 2-adic valuation-unit data at fixed precision excludes nothing: for any finite word of odd-unit symbols at fixed precision and any starting carry, a compatible orbit exists with every intermediate state centred inside its symbol’s dyadic radius (`Erdos249257.fixedPrecisionTropicalNoGo [TropicalCurvatureCarry.lean:137]`).

Rules out: the finite-automaton-computes-the-expansion family outright; any hope of distinguishing two carry histories after a long common suffix; and any contradiction derived from a fixed-precision valuation-unit signature. Growing precision is mandatory, not optional. None of the three mentions φ , so all three bind #257 identically. `coord:carry-orbit scale:uniform [Lean]`

Proposition 2.14 (B6, no finite linear compression). *Status: proved for four independent truncations (Lean), with the honest limit stated below. Every natural finite truncation of the totient kernel has trivial kernel:*

- (i) Dyadic. The canonical family of $2^e + 1$ dyadic channels is linearly independent over $\mathbb{Q}$ at every depth (Proposition 1.5), and the natural repair — a bounded compressed-adjoint certificate — is impossible (`Erdos249257.false_of_compressedAdjointCertificate [TotientMahlerDefect.structure at :1039]`).

- (ii) Möbius incidence. $U_N(i, j) = \mu((i+1)/(j+1))$ when $(j+1) \mid (i+1)$ and 0 otherwise is lower triangular with unit diagonal, so $\det U_N = 1$ for every N and the jet map is injective: no finite incidence-quotient relation exists at any horizon ([Erdos249257.incidenceMobius-det_eq_one](#) [[IncidenceQuotientHermitePade.lean:56](#)], [mobiusCompanionJetMap_injective](#) at [:77](#)).
- (iii) Adjugate reconstruction. Any finite rational row exactly isolating one totient value has crude two-tail cost ≥ 3 , hence never < 1 , at any finite grid height, using only $\varphi(x) \leq x$ ([Erdos257PeriodNoncollapse.three_le_totientAdjugateTailCost](#) [[TotientTailCarryPeriod.lean:266](#)], closure at [:307](#)).
- (iv) Shift-polynomial. An explicit nonzero all-horizon countermodel agrees with every exact whole-ray anchor and survives every finite commensurate LCM-cube shift polynomial, at every finite rank ([LcmFactorIdealPulseObstruction.lean](#); module docstring, theorem bodies not individually re-verified in this pass, and the construction is explicitly synthetic — it does not claim its compensation letters occur as actual totient differences).

Additionally, every strict-subrank monomial quotient in the Möbius–Mersenne ladder overshoots its target by more than $1/480$, uniformly — every rung $r \geq 3$ lies in $[1429/1512, 1)$ and every prefix after four atoms is within $1/3584$ of its rung, so this is a uniform no-go rather than a census ([rankOneSubrankQuotient_sub_theta_two_gt](#) [[RankOneSubrankObstruction.lean:238](#)]). **Honest limit:** this is four checked truncations, not a proof that no finite-linear shortcut exists. Its correct reading is the one the source module gives: any winning finite-linear shortcut must live in a genuinely different coordinate or use a growing-parameter construction. The reason the barrier bites is Proposition 1.6: what rationality actually buys is periodicity, and periodicity provably does not promote to a rank bound. coord:carry-rank scale:uniform [Lean]

Proposition 2.15 (B7, the coarse-invariant barrier). **Status:** *proved* (Lean, `#print axioms clean`). This is Proposition 1.8 read as an elimination. **Rules out:** any proof of #249 whose hypothesis set on the coefficient word lies inside {uniform boundedness, $c(n) \leq n$, agreement with $\varphi \bmod 2$ at every index, failure of eventual periodicity in any strength up to arbitrarily long, arbitrarily separated blocks}. The witness satisfies every one of those and is rational. **Does not rule out:** arguments using actual quantitative totient size or residue information. That complement is precisely why the corpus’s standing lesson is that only a quantitative argument in the actual-LCM or fixed-rank style can close #249. Any future sufficient condition stated purely in terms of coefficient-word properties must be checked against this fixture before being trusted, for either problem. coord:coefficient-word scale:uniform [Lean]

2.4 The shape

Put together, the barriers say something more specific than “the problem is hard”. The certificate residue is a weighted accumulation over an unboundedly long window at full arithmetic resolution, and *every device that makes an argument finite destroys precisely the quantity being measured*: bounding the window (B1), coarsening the values (B7), compressing the state (B5), truncating the rank (B6), or letting the prover pick the sample (B2). B4 closes the remaining escape — buying the amplitude by construction — with an explicit exponential margin, and B3 records that in practice nothing bounded has ever promoted.

There is exactly one known way to hold unbounded range and full resolution at once without a finite bookkeeping device: stop naming an index and assert an average instead. That is what an exponential-sum bound is. It is why the surviving routes of §3 are, with the honest exceptions noted there, the ones that never name a good index.

3 What the wall does not block

An elimination is only worth the paper it is written on if it leaves somewhere to stand. This section is that payoff. Five routes survive the classification of §2; for each we state the exact statement it needs, which barrier it evades, and *why* the evasion is structural rather than accidental. Two of the five are demoted explicitly, because they evade every *proved* barrier while sitting squarely inside the class that the strongest *observed* pattern (Observation 2.10) indicts. None of the five is a proof, and none is close to one; what has changed is that the space of things to try is no longer large.

3.1 Survivor 1: the first-harmonic block cancellation bound

Definition 3.1 (The open analytic obligation). Write $e(x) := \exp(2\pi i x)$ and $\text{windowFirstExp}(h, N, L) := e((\text{windowDiscrepancy}(h, N, L) \bmod 2^L)/2^L)$. `DTWFirstHarmonicNormGap` is the statement

$$\forall h \geq 1 \forall X_0 \exists X, L : \quad \max(X_0, 1) \leq X, \quad 16(2X+h+L+2) \leq 2^L, \quad \left\| \sum_{N \in [X, 2X)} \text{windowFirstExp}(h, N, L) \right\| \leq \frac{21}{25} X$$

It implies `Irrational(S)`. `coord:first-harmonic scale:cofinal [Lean] Erdos249257.DTWFirstHarmonicNormGap [FirstHarmonicGap.lean:128] Erdos249257.irrational_totient_series_of_first_harmonic_norm_gap [FirstHarmonicPivot.lean:83]`

The real-part form suffices and is strictly weaker: it is enough that $\sum_{N \in [X, 2X)} \text{windowFirstCos}(h, N, L) \leq \frac{9}{10} X$ (`Erdos249257.exists_certifiedKill_of_first_harmonic_gap [FirstHarmonicGap.lean:128]`), the elementary unconditional engine; the $21/25 \rightarrow 9/10$ bridge is `Erdos249257.exists_certifiedKill_of_first_harmonic_norm_gap [FirstHarmonicPivot.lean:53]`. Unpacked, the required object is a constant-saving cancellation estimate for the dyadically weighted totient-difference exponential sum

$$\sum_{X \leq N < 2X} e\left(\frac{\sum_{j < L} (\varphi(N+h+1+j) - \varphi(N+1+j)) 2^{L-1-j}}{2^L}\right), \quad L \approx \log_2 X + O(1).$$

Not one instance of this bound is proved anywhere, at any X , h or L . Both trees were searched for a theorem supplying it; only consumers exist. `[Open]`

Why it evades the wall. This is the only route in the corpus that evades all seven barrier classes, and the reasons are structural.

- **B1.** X is a free unbounded parameter and the room condition $16(2X+h+L+2) \leq 2^L$ forces $L \gtrsim \log_2 X + 5$, so the estimate reads φ at every index in $[X+1, 2X+h+L]$ with both position and window length growing. No γ agreeing with φ only on $[1, B]$ constrains it.
- **B2 — the decisive point.** The proved collapse `DTWWindowSeparatedPairs` $\Leftrightarrow$ `Irrational(S)` works *because the prover may choose the sample*: the converse proof selects $T = \{N, N+1\}$ and meets the threshold with $\delta = 9/10$ supplied by irrationality plus doubling expansivity (Proposition 2.7(b)). Definition 3.1 admits no sample choice: the sum ranges over all of $[X, 2X)$ at one common depth L . Since $\neg \text{certifiedKill}(h, N, L)$ forces $\text{windowFirstCos}(h, N, L) > 9/10$ (`Erdos249257.windowFirstCos_gt_of_not_certifiedKill [FirstHarmonicGap.lean:44]`), the gap demands that a *positive proportion* of basepoints in the block carry certificates at a *single* depth. Irrationality supplies only one certificate per (h, N_0) , at a depth that may vary with N . The collapse mechanism therefore has no purchase. **Honest limit:** no theorem proves `DTWFirstHarmonicNormGap` inequivalent to `Irrational(S)`. What is asserted is that the one proved collapse mechanism in this lane demonstrably requires sample

choice, which this predicate denies. Note also that the *subset* form (`Erdos249257.exists_-certifiedKill_of_first_harmonic_gap_subset` [FirstHarmonicGap.lean:104]) does permit sample choice and is therefore the collapse-exposed variant. Attack the full-block form.

- **B4.** The statistic is the first additive character of the *accumulated* weighted word, never an individual letter. Nothing is prescribed at any position, so the self-defeating cost that kills Dirichlet residue engineering does not arise. B4’s own recorded repair (Remark 2.12) is to impose the structure on the word rather than on one delta, which is exactly what a block exponential-sum bound does.
- **B5, B6, B7.** No carry state is summarised and no fixed precision is used; the estimate is on the true residue mod 2^L with $L \rightarrow \infty$. A cancellation bound is not a rank statement and seeks no finite-dimensional relation. And the estimate uses the actual real residue, not parity, boundedness or aperiodicity, so neither countermodel touches it — both differ from φ at exactly the large indices where this sum lives.
- **B3.** The audit’s own verdict on this row is that the consumer side is *finished* and already at the obligation’s exact quantifier shape: X is free, so applying the bound at $X \geq N_0$ gives $\exists N \geq N_0 \exists L$ directly. The missing piece is one named arithmetic fact, not a strengthening of anything on disk. The constants $9/10$, $\pi/8$ and 16 are absolute and do not degrade with X , h or L ; there is no case analysis and no table.

Remark 3.2 (Does B1 *select* this route? No.). It is tempting to say the finite-inspection theorem points at analysis. It does not. B1 says any proof must use φ at unbounded indices — a necessary condition satisfied by *every* open route here (the LCM diagonal as $t \rightarrow \infty$, the actual-LCM supply for $a \geq 8$, the Farey growth law as $K \rightarrow \infty$, the rank bound for all e). B1 does not discriminate. What selects this route is a sharper pair: (1) B4, proved, kills the only mechanism the corpus ever found for reaching full resolution at unbounded range by *prescribing* values, and its own stated repair is accumulation over the word — which is what an exponential sum measures; and (2) Observation 2.10, an observed pattern and not a theorem, shows that every single-index producer has failed to be supplied at even one large index, while this route’s missing input is a block average and therefore never requires naming a good index. B1’s real contribution is narrower but still load-bearing: it proves that the finite deposits can never be extended into a proof, so the gap between the census and the obligation is a gap in kind, not in degree. That is a genuine no-go about method, of the same species as a relativization barrier. It is not a selector.

Remark 3.3 (Evidence in both directions). Weak supporting evidence, observed and not proof: every landed diagonal certificate through $t = 64$ fires within a small additive constant of the minimum depth forced by `certifiedKill_depth_floor`, which is the profile a doubling-orbit equidistribution argument would produce. Countervailing evidence, also observed: the strict-LCM-jump census records a closest central margin of ≈ 0.000221 of the modulus at $t = 100$, so these residues are not robustly central and any equidistribution claim will be delicate. [Cert]

3.2 Survivor 2: the four-term pivot budget

Same lane as Survivor 1, with the analytic burden repackaged. Split the block sum at a largest-prime pivot into centred correlation, fibre-mean, bad-cofactor and non-supplier contributions and bound the four separately.

Definition 3.4 (DTWPivotResidualDecorrelation). For every $h > 0$ there are $s > 0$ and $\eta \in (0, 1)$ such that for every X_0 there are X, L with $\max(X_0, 1) \leq X$, $h \leq L - s$, $16(2X + h + L + 2) \leq 2^L$, and all four of

$$\text{Re pivotCenteredCorrelation} \leq \frac{14}{25}X, \quad \|\text{pivotFiberMean}\| \leq \frac{1}{100}X, \quad \|\text{pivotBad}\| \leq \frac{1}{100}X, \quad \|\text{pivotNonSupplier}\| \leq \frac{1}{100}X$$

It implies Irrational(S). coord:first-harmonic scale:cofinal [Lean] [Erdos249257.DTWPIVOTResidualDecorrelation](#) [Erdos249257.irrational_totient_series_of_pivotResidualDecorrelation [FirstHarmonicPivot.lean:406]

It inherits Survivor 1's evasions of B1, B4, B5, B6 and B7 verbatim: the four terms are exact finite sums over canonical largest-prime supplier fibres of the same accumulated-word phases, with no sample choice and nothing prescribed. Its specific advantages are three. The decomposition is an *exact identity*, proved unconditionally with `#print axioms clean` ([Erdos249257.windowFirstExp_sum_eq_pivot_decomposition](#) [FirstHarmonicPivot.lean:342], budget consumer [Erdos249257.first_harmonic_gap_of_pivotBudgetAt](#) [FirstHarmonicPivot.lean:377]). The one-sided budget lowers the hard requirement from a norm bound $\|\cdot\| \leq X/2$ to a real-part bound $\leq 14X/25$. And the pivot rests on genuine arithmetic rather than a sampled surrogate: on the canonical fibre $m = 1$, $s = L - h$ the supplier set is *literally* the shifted dyadic interval of primes, proved as a membership equality ([Erdos249257.mem_pivotFiber_one_overlap_iff](#) [FirstHarmonicPivot.lean:251]), with the totient factorisation at the pivot the honest $\varphi(mp) = \varphi(m)(p-1)$ and the non-divisibility discharged by size. Three of the four terms are then counting bookkeeping; only the first needs a genuine correlation estimate, and only in real part.

What it must respect. B6 has a residual-gauge instance: a residual-blind determinant or conditioning test cannot certify that genuine phase reconstruction rather than a locked degenerate configuration has occurred ([Erdos249257.locked_reconstruction_preserves_nonzero_minor](#) [ResidualGaugeObstruction.lean:94]). So this route must couple rows by an extra arithmetic identity, not merely gauge-normalise columns. **Honest status:** strictly a repackaging of Survivor 1's burden; it is listed separately only because three quarters of it are already exact identities on disk. The module asserts no prime-distribution or decorrelation estimate — the socket is deliberately empty. [Open]

3.3 Survivor 3: uniform quantitative escape at cofinally many primes

Definition 3.5 (DTWNaturalPrimeTailOrbitStrictGap). For every $h \geq 1$ and every N_0 there is a prime p with $\max(N_0+h+1, h+5) \leq p$ and $\text{RetailOrbitFirstExp}(h, p-h-1) < \frac{9}{10}$, where $\text{tailOrbitFirstExp}(h, N) = e(R_{N+h} - R_N)$. Equivalently: cofinally many primes p at which the totient tail difference across the shift h stays a *fixed* distance $\geq \arccos(9/10)/2\pi \approx 0.0722$ from every integer. It implies Irrational(S). coord:prime-pivot scale:cofinal [Lean] [DTWNaturalPrimeTailOrbitStrictGap](#) [irrational_totient_series_of_naturalPrimeTailOrbitStrictGap [TotientStrictPrimeEscape.lean:66]

It evades B1, B4, B5, B6 and B7 for the reasons given for Survivor 1: a statement about the true real tail difference at unbounded prime positions, with no letters prescribed, no bounded state, no rank, no coarse coefficient-word invariant. It evades B2 by the *second* of the two exemptions in Proposition 2.7 — not by denying sample choice, but by demanding a *uniform margin*. Irrationality gives non-integrality of every tail difference and supplies no lower bound whatever on distance to $\mathbb{Z}$; certificate completeness therefore cannot manufacture this predicate.

Honest demotion. This is a *pointwise* producer: it requires naming a good prime, and Observation 2.10 records that no pointwise producer in this corpus has ever been supplied at even one large index, across seven independent attempts. It evades every proved barrier while sitting inside the class the strongest observed pattern indicts. Nothing on disk proves it at a single prime for a single h . It ranks below Survivor 1. [Open]

3.4 Survivor 4: depth-locked full-depth escape

Definition 3.6 (ApFullDepthEscape). For every $d \geq 1$ and every N there is $t \geq 1$ with $\text{certifiedKill}(td, N, td)$; unpacked,

$$N + 2td + 2 < \text{windowDiscrepancy}(td, N, td) \bmod 2^{td} < 2^{td} - (N + 2td + 2).$$

It implies $\text{Irrational}(S)$. coord:period-ray scale:cofinal [Lean] [ApFullDepthEscape](#) [PeriodMultipleEscape.lean:162] [irrational_totient_series_of_apFullDepthEscape](#) [PeriodMultipleEscape.lean:171]

This is the shortest fully stated open inequality the programme has produced. Its evasion of B2 is recorded on disk rather than argued: its ambient parent `PeriodMultipleKillSupply` is *proved* equivalent to $\text{Irrational}(S)$ and `ApFullDepthEscape` implies it ([periodMultipleKillSupply_of_apFullDepthEscape](#) [PeriodMultipleEscape.lean:165]), so it is at least as strong, and possibly strictly stronger — the file’s own docstring records “sufficient for irrationality; not known necessary”. Being possibly strictly stronger, certificate completeness cannot collapse it back. It evades B1 (both N and t unbounded), B4 (locking depth to the period means the entire word accumulates and no letter is prescribed), B5 and B6 (no state, no rank). Its substance is pure anti-concentration: via [windowDiscrepancy_self_eq_totientBlock_sub](#) [PeriodMultipleEscape.lean:67], the difference of two adjacent period blocks must have central residue mod 2^h at some multiple period $h = td$, and the room condition is automatic once h is large, so nothing but the residue’s position is at stake.

Honest demotion, identical to Survivor 3: it is a pointwise producer and names an index. It is listed because it is the cleanest target for computational exploration — with Theorem 2.4 as the standing reminder that no amount of such exploration becomes a proof. [Open]

3.5 Survivor 5: the rationality-side rank upper bound

Definition 3.7 (The coordinate-disjoint obligation). Either: there is C such that for every $c : \mathbb{N} \rightarrow \mathbb{N}$ with $c(n) \leq n$ and $T_c \notin \text{irrationals}$, every $v > 0$ and every tempered binary orbit u for (c, v) , and every e , $\dim_{\mathbb{Q}} \text{span}_{\mathbb{Q}}(\text{canonicalCarryKernelFamily}(u, e)) \leq C$; or the same with any $g(e)$ growing strictly slower than $2^e - 1$ in place of C . Either version contradicts the proved floor of Proposition 1.5 and closes #249. coord:carry-rank scale:uniform [Open]

This is the only surviving obligation that is not a residue or certificate statement at all, so B2’s certificate completeness cannot reach it — it is not a reformulation of Sep in any vocabulary. It evades B1 and B4 entirely (no window, no letters, no prescribed residues) and B5 (linear algebra over the whole orbit, not a bounded-state summary). The scale side is finished and uniform in e (Proposition 1.5).

Honest flag, and it is severe. B6 partly indicts this route from inside. The most natural approach to the upper bound is dead ([Erdos249257.false_of_compressedAdjointCertificate](#) [TotientMahler]) and what rationality actually buys — uniform eventual periodicity of the carry’s dyadic sections mod v — provably does not promote to a $\mathbb{Q}$ -rank bound (Proposition 1.6). So this survivor is genuinely coordinate-disjoint from everything else, which is its whole value, but its obvious approach is closed and the missing input is a rigidity theorem nobody has stated.

4 How to use this document

Evidence bands. Every claim carries one, and they are never blurred. [Lean] means kernel-checked, with the declaration named and the file:line given. [Cert] means an exact finite computation, valid exactly on the stated range. [Math] means ordinary mathematics, proved here or elsewhere but not formalised — Theorem 2.4 is the important instance, and its unformalised composition step is flagged where it occurs. [Cited] means published, with the reference. [Open] means not proved, by anyone, anywhere. A proved no-go and an audit over a corpus are both valuable and are never conflated: B1, B2, B4, B5, B6 and B7 are proved eliminations; B3 is an audit and is labelled as evidence every time it is used.

The scale axis. This is the load-bearing axis of the whole document. `scale:fixed` means one instance; `scale:bounded` means every instance up to an explicit bound; `scale:cofinal` means infinitely often past every threshold; `scale:uniform` means for all parameters with no dependence. The single most important structural fact about the corpus is that *everything unconditional and quantitative sits on the fixed or bounded side, while Sep is cofinal*. Theorem 2.4 says that gap is one of kind and not of degree: no accumulation of bounded verification, at any bound, is evidence that the cofinal statement holds. Read every result’s `scale` tag before reading its content.

Coordinates, and why obstructions are relative to them. Every obstruction carries a `coord` tag, because an obstruction is always an obstruction *in a representation*, never a property of the object. The γ -splice obstructs arguments in the `coord:binary-digit` coordinate; the rank floor lives in `coord:carry-rank`; positivity lives in `coord:actual-lcm`; the Farey exclusion lives in `coord:farey`. Two consequences follow, and both matter for anyone attacking the problem. First, before reporting a new wall, name the coordinate it was met in and test a re-representation. Second, an obstruction proved in one coordinate is silent about another: Survivor 5 survives precisely because it is coordinate-disjoint from the certificate lane, and Survivor 1 survives because the first-harmonic coordinate is the one in which the corpus’s proved collapse mechanism has no purchase.

Map of what follows. The premise catalogue records, for each coordinate, the exact statements the corpus holds, with hypotheses, evidence band, scale, and Lean declaration. It is followed by the typed-implication tables (which statement feeds which, and at which quantifier shape), the near-miss index against the open obligations, and the promotion audit whose verdicts are summarised as B3. Nothing after this point restates the barriers; they are all here.

Read this in full before starting. That instruction is not politeness. The recurring practical failure of this programme has not been a shortage of ideas; it has been rediscovering the same reformulations, because no single document held the whole picture. Every dead route in §2 was found more than once. If you are about to prescribe totient values at engineered positions, read B4 first. If you are about to define a bounded carry state, read B5. If you are about to look for a weaker sufficient condition, read B2 — there probably isn’t one, and the corpus has already proved as much three times. The purpose of assembling the wall is that the next attempt should start where the last one stopped.

5 The object, the target, and the exact record

This part fixes the object of Erdős Problem #249, states exactly what is open about it, and records every unconditional numeral, identity and Lean-checked bound the corpus currently holds. Nothing in this part decides irrationality. Every result below is either an identity (no hypothesis, holds unconditionally), a finite computation (holds up to an explicitly stated bound), or is flagged [Cited]/[Open] where that is the honest status.

5.1 The object

Definition 5.1 (The Erdős–249 constant). Let φ denote Euler’s totient function. Define

$$S := \sum_{n \geq 1} \frac{\varphi(n)}{2^n} = \sum_{n: \mathbb{N}} \frac{\varphi(n)}{2^n},$$

the two forms coinciding because $\varphi(0) = 0$; the second, $\mathbb{N}$ -indexed form is the one carried in the Lean source. The series converges absolutely since $\varphi(n) = O(n)$. *Erdős #249 asks whether S is irrational. This is OPEN.* Nothing in this paper decides it; every claim below is either an unconditional identity, a finite computation with an explicitly stated range, or is marked [Cited]/[Open].

Remark 5.2 (Status). No proof or disproof of $\text{Irrational}(S)$ exists anywhere in the corpus, formal or informal. What exists is: one large unconditional finite denominator exclusion (§5.3), several exact reformulations of S that relocate the same open question onto different coordinates without touching its truth value (§5.4–§5.6), and a certificate apparatus whose cofinal supply obligation is the precise open target (recorded in full in Part 1 of this paper; only its finite, checked instances are catalogued here, §5.7).

5.2 The binary-digit reduction

Proposition 5.3 (Digit-shift identity). *For every $N : \mathbb{N}$,*

$$2^N \cdot S = \Phi_N + R_N,$$

where $\Phi_N := \sum_{n \leq N} \varphi(n) \cdot 2^{N-n} \in \mathbb{N}$ (the integer prefix) and $R_N := \sum_{j \geq 0} \varphi(N+1+j)/2^{j+1}$ (the fractional tail, `totientTail` in Lean). `coord:binary-digit scale:uniform [Lean] two_pow_mul_totient_series_eq [TotientTailPeriodKiller.lean:150]`

Consequence. Rationality and eventual periodicity of the base-2 expansion of S is *exactly* a statement about the tail R_N : $S \in \mathbb{Q}$ forces (by Euler’s theorem applied to the odd part of the denominator) a period $h > 0$ and pre-period N_0 with $R_{N+h} - R_N \in \mathbb{Z}$ for all $N \geq N_0$ (`eventual_period_of_not_irrational`, `eventual_period_of_not_irrational` [TotientTailPeriodKiller.lean:358], [Lean], explicit witness $h = \varphi(\text{oddPart}(r.\text{den}))$, $N_0 = v_2(r.\text{den})$). Nothing in the corpus supplies the converse cofinally; this is the certificate wall documented in full in Part 1 and is not restated here beyond this pointer. $\square$

5.3 The unconditional denominator floor

The strongest *unconditional* fact the corpus holds about S is a lower bound on its denominator should it happen to be rational. It is obtained by a completely elementary Farey/mediant argument, entirely free of any certificate or period apparatus, and is therefore logically independent of the open certificate-supply obligation.

Lemma 5.4 (Farey gap, fully general). *For integers a, b, c, d, r, s with $b > 0$, $d > 0$, $bc - ad = 1$ (i.e. a/b and c/d are unimodular Farey neighbours), and $as < rb$, $rd < cs$ (i.e. r/s lies strictly between them): $b + d \leq s$. `coord:farey scale:n/a [Lean] farey_gap [GapFareyBound.lean:51]`*

This lemma is a classical Stern–Brocot fact with zero totient or Mersenne content; it is the generic engine underneath every Farey-gap bound in the corpus, for either open problem.

Proposition 5.5 (The wave-17 gap certificate at window $K = 240$). *Let V be the explicit committed totient residue for window $(N, K) = (1, 240)$. For every $q : \mathbb{N}$ with*

$$0 < q \leq Q_0 := 79\,639\,646\,646\,701\,375\,323\,355\,774\,875\,831\,053 \ (\approx 7.96 \times 10^{34}),$$

$$(q \cdot V) \bmod 2^{240} + 243q < 2^{240}.$$

This bound is sharp: $q = Q_0 + 1 = 79\,639\,646\,646\,701\,375\,323\,355\,774\,875\,831\,054$ is the exact first failing denominator, exhibited as the mediant of two explicit unimodular Farey neighbours. `coord:farey scale:bounded [Lean] gap_check_window_1_240_le_79639646646701375323355774875831053 [GapFareyBound.lean:225]`

Theorem 5.6 (Erdős #249 denominator exclusion — the headline unconditional result). *For every $p \in \mathbb{Q}$ with reduced denominator $p.\text{den} \leq Q_0$,*

$$S \neq p.$$

Equivalently: if S is rational, its reduced denominator exceeds $Q_0 \approx 7.96 \times 10^{34}$. coord:farey scale:bounded [Lean] [tsum_totient_div_pow_two_ne_ratCast_of_den_le_796396466467013753233557748758](#)

Remark 5.7 (What this does and does not say). Theorem 5.6 is a complete, unconditional finite fact: no rational of small denominator equals S . It says nothing about arbitrarily large denominators and is not itself a route to irrationality. The docstring behind Proposition 5.5 states the honest open question this leaves: whether $\sup_K (b + d)(K)$ (the growing analogue of Q_0 as the window K grows) is unbounded as $K \rightarrow \infty$ — which would close #249 through this theorem’s consumer, entirely independently of the certificate-supply obligation of Proposition 5.3. [Open]; not claimed or proved anywhere in the corpus.

Remark 5.8 (Ladder of prior rungs). Q_0 is the current end of an explicit sequence of increasingly wide Farey windows computed by the same mechanism: $4838 \rightarrow 2^{22} \rightarrow 2.49 \times 10^{17}$ (window $K = 120$) $\rightarrow Q_0$ (window $K = 240$). Each rung is a finite, independently checked instance of Lemma 5.4 and Proposition 5.5’s pattern at a larger K ; none is claimed to extrapolate.

5.4 The same record, transported: the Möbius-square and coprimality-probability forms

The exact finite Farey record behind Theorem 5.6 is not tied to the $\varphi(n)/2^n$ presentation of S ; it transfers verbatim to two other exact reformulations of the same constant, at exactly half the bound.

Proposition 5.9 (Fair-coin coprimality form). *Let X, Y be independent random variables with $\Pr(X = n) = \Pr(Y = n) = 2^{-n}$ for $n \geq 1$ (independent fair-coin waiting times). Then*

$$S = \frac{1}{2} + \Pr(\gcd(X, Y) = 1) = \frac{1}{2} + \sum_{\substack{a, b \geq 1 \\ \gcd(a, b) = 1}} 2^{-(a+b)}.$$

Equivalently, on the visible lattice: summing $2^{-(a+b)}$ over the half-open coprime pairs ($a \geq 1$, $b \geq 0$, $\gcd(a, b) = 1$) recovers $\sum_n \varphi(n)/2^n$ exactly, with no boundary correction, because the visible-point count on the half-open antidiagonal at height n equals $\varphi(n)$ for every n , including $n = 0, 1$. coord:probability scale:n/a [Lean] [tsum_visible_coprime_pairs_eq_totient_series](#) [CertificateKernel.lean:18216] [totient_series_eq_half_add_visible_coprime_pairs](#) [CertificateKernel.lean:18216]

Proposition 5.10 (The gcd-layer normalisation). *For independent fair-coin waiting times as above, $\sum_{g \geq 1} \Pr(\gcd(X, Y) = g) = 1$ exactly; and for every $d > 0$, $\Pr(d \mid X \wedge d \mid Y) = 1/(2^d - 1)^2$. coord:probability scale:uniform [Lean] [tsum_pos_coprime_inv_mersenne_eq_one](#) [GcdMomentCalculus.lean:349] [tsum_pos_pair_both_dvd_half_eq_inv_mersenne_sq](#) [GcdMomentCalculus.lean:266]*

Theorem 5.11 (Denominator exclusion for the coprimality-probability form). *Let*

$$Q_1 := \left\lfloor \frac{Q_0}{2} \right\rfloor = 39\,819\,823\,323\,350\,687\,661\,677\,887\,437\,915\,526.$$

For every $a \in \mathbb{Z}$, $d \in \mathbb{N}$ with $0 < d \leq Q_1$: the visible coprime-pair probability $\Pr(\gcd(X, Y) = 1)$ is not equal to a/d . coord:farey scale:bounded [Lean] [tsum_visible_coprime_pairs_ne_int_div_of_den_le_39819823323350687661677887437915526](#) [CertificateKernel.lean:18244]

Theorem 5.12 (Denominator exclusion for the Möbius-square form). *With Q_1 as in Theorem 5.11: for every $a \in \mathbb{Z}$, $d \in \mathbb{N}$ with $0 < d \leq Q_1$, the signed series $T := \sum_{d \geq 1} \mu(d)/(2^d - 1)^2 = S - \frac{1}{2}$ (see §5.5) is not equal to a/d . `coord:farey scale:bounded [Lean] tsum_moebius_div_two_pow_sub_one_sq_ne_int_div_of_den_le_39819823323350687661677887437915526 [CertificateKernel.lean:1`*

Remark 5.13 (Why the bound halves, and why this is not new information). $Q_1 = \lfloor Q_0/2 \rfloor$ arithmetically: Q_0 is odd ($Q_0 = 79\,639\,646\,646\,701\,375\,323\,355\,774\,875\,831\,053$), so $Q_0/2 = 39\,819\,823\,323\,350\,687\,661\,677\,887\,437\,915\,526.5$ and Q_1 is its floor. The halving is exactly the cost of the affine shift $S = \frac{1}{2} + T$ (resp. $S = \frac{1}{2} + \Pr(\gcd(X, Y) = 1)$) on a denominator-exclusion statement: excluding all denominators $\leq Q_1$ for T is equivalent, after the $\frac{1}{2}$ -shift, to excluding all denominators $\leq Q_0$ for S (a denominator- d value of T with $d \leq Q_1$ yields a denominator dividing $2d \leq Q_0$ for S). Theorems 5.11 and 5.12 are therefore the *same* finite Farey record as Theorem 5.6, transported through Proposition 5.9 and Proposition 5.14 respectively — not independent evidence, but confirmation that the exclusion is coordinate-stable across all three exact presentations of S .

5.5 The Möbius–Mersenne identity and why bounded coefficients matter

Proposition 5.14 (Möbius-square reduction).

$$S = \sum_{n \geq 1} \frac{\varphi(n)}{2^n} = \frac{1}{2} + \sum_{d \geq 1} \frac{\mu(d)}{(2^d - 1)^2},$$

where μ is the Möbius function, so $\mu(d) \in \{-1, 0, 1\}$ for every d . Consequently Erdős #249 $\iff T := \sum_{d \geq 1} \mu(d)/(2^d - 1)^2 \notin \mathbb{Q}$, the single reduced target every other result in this subsection and §5.6 feeds. `coord:mobius-mersenne scale:n/a [Lean] totientSeries_eq_pnat_half_pow [SquaredMersenneDiagonalEnclosure.lean:50] tsum_totient_half_pow_eq_half_add_moebius_sq [MersenneLambertLadder.lean:665]`

Remark 5.15 (Why the bounded coefficients matter — the Erdős-1948 regime). The identity of Proposition 5.14 rewrites S (equivalently T) as a *Möbius-twisted Lambert-squared series*: the numerator weight $\mu(d)$ is bounded, $|\mu(d)| \leq 1$ for every d , uniformly in d . This places T in exactly the coefficient regime of the classical Erdős (1948) near-integer irrationality criterion and of the level-1 sibling identity $L(\mu) := \sum_d \mu(d)/(2^d - 1) = \frac{1}{2}$ (rational, trivially) alongside $L(1) = \sum_d 1/(2^d - 1) = E$, the Erdős–Borwein constant, which is proved irrational in this same kernel (`irrational_erdosBorwein_series [CertificateKernel.lean:8007]`, [Lean]). This is the opposite regime from Proposition 5.3’s **binary-digit coordinate**, where the corresponding weight is $\varphi(n) = \Theta(n)$ — *unbounded*, growing linearly. A near-integer/Dirichlet-approximation argument of Erdős-1948 shape (formalised generically as `irrational_of_int_mul_near_int [CertificateKernel.lean:5792]` and its base-power specialisation `irrational_of_pow_mul_near_int [CertificateKernel.lean:5821]`, both [Lean], `coord:n/a`, fully coordinate-free) has a genuine chance of transferring to T precisely because its weight is bounded, in a way it does not have a chance of transferring directly to the raw $\varphi(n)/2^n$ series. No such transfer is proved; §?? below (cross-referenced here, developed in Part 2 of this paper) records exactly why the transfer has so far failed (the “ μ -pollution” obstruction) rather than merely asserting the analogy.

5.6 The squared-Lambert gcd-moment identities

Proposition 5.16 (Squared-Lambert transfer engine). *For $w : \mathbb{N} \rightarrow \mathbb{R}$ with $|w(d)| \leq d$ for all $d > 0$, and $0 \leq r < 1$:*

$$\sum_{d \geq 1} w(d) \left(\frac{r^d}{1 - r^d} \right)^2 = \sum_{n \geq 1} \left(\sum_{e|n} w(e) \left(\frac{n}{e} - 1 \right) \right) r^n.$$

coord:mobius-mersenne scale:uniform [Lean] [tsum_lambert_linear_weight_sq_pure](#) [GcdMomentCalculus.lean:105]

This one identity, at $r = 1/2$, specialises to every squared-Lambert rung the corpus computes; two instances are exact and directly relevant to #249's weight structure:

Proposition 5.17 (The known ζ_q -rung).

$$\sum_{d \geq 1} \frac{1}{(2^d - 1)^2} = \sum_{n \geq 1} \frac{\sigma(n) - \tau(n)}{2^n} = \zeta_q(2) - \zeta_q(1) \text{ at } q = \frac{1}{2},$$

where σ is the sum-of-divisors function and τ the number-of-divisors function. The identity is machine-checked ([Lean]); irrationality of the value $\zeta_q(2) - \zeta_q(1)$ is [Cited] (Postelmans–Van Assche q -Padé), not formalised in this corpus. coord:mobius-mersenne scale:n/a [tsum_one_div_mersenne_sq_eq_sigma_sub_tau_series](#) [GcdMomentCalculus.lean:216]

Proposition 5.18 (The Pillai/gcd-moment rung).

$$\sum_{d \geq 1} \frac{\varphi(d)}{(2^d - 1)^2} = \sum_{n \geq 1} (P(n) - n) \cdot 2^{-n} = \mathbb{E}[\gcd(X, Y)],$$

where $P(n) := \sum_{e|n} \varphi(e) \cdot (n/e) = (\varphi * \text{Id})(n)$ is Pillai's gcd-sum function and X, Y are the independent fair-coin waiting times of Proposition 5.9. The identity itself is machine-checked ([Lean]); the value $\mathbb{E}[\gcd(X, Y)]$ is [Open] — a cousin rung to #249, not #249 itself. coord:mobius-mersenne scale:n/a [tsum_totient_div_mersenne_sq_eq_gcd_moment_series](#) [GcdMomentCalculus.lean:235]

Remark 5.19 (The level mirror). Writing $L(f) := \sum_d f(d)/(2^d - 1)$ (level 1) and $L_2(f) := \sum_d f(d)/(2^d - 1)^2$ (level 2): $L(\mu) = \frac{1}{2}$ (rational, trivial), $L(1) = E$ (Erdős–Borwein, irrational, [Lean]), $L(\varphi) = 2$ (rational); $L_2(\mu) = S - \frac{1}{2}$ ([Open], this is #249), $L_2(1) = \zeta_q(2) - \zeta_q(1)$ ([Cited] irrational, Proposition 5.17), $L_2(\varphi) = \mathbb{E}[\gcd(X, Y)]$ ([Open], Proposition 5.18). At level 1 the Möbius rung is trivial and the ζ -rung is the hard classical case; at level 2 the ζ -rung is known and the Möbius rung is #249. Möbius projection is the one wall repeated at both levels of the ladder.

5.7 The exact certificate record: numerical anchors through $t = 64$

The certificate apparatus itself (the predicate `certifiedKill`, its soundness/completeness theorems, and the cofinal supply obligation that is the actual open target) is developed in full in Part 1 of this paper. What is recorded here is exhaustively: every finite, machine-checked numerical anchor the corpus currently holds for it, stated precisely, with no extrapolation implied.

Certificate object. For $h, N, L : \mathbb{N}$, define the window discrepancy $\Delta_{h,N,L} := \sum_{j < L} (\varphi(N + h + 1 + j) - \varphi(N + 1 + j)) \cdot 2^{L-1-j} \in \mathbb{Z}$ ([windowDiscrepancy](#) [TotientTailPeriodKiller.lean:66], [Lean]), and

$$\text{Sep}(h, N, L) := (N + h + L + 2) < \Delta_{h,N,L} \bmod 2^L < 2^L - (N + h + L + 2),$$

(`certifiedKill` [TotientTailPeriodKiller.lean:72], [Lean], decidable). By completeness ([exists_certifiedKill_iff_tail_diff_notMem_int](#) [LcmConeFlatness.lean:316], [Lean]), $(\exists L, \text{Sep}(h, N, L)) \iff R_{N+h} - R_N \notin \mathbb{Z}$ for every h, N : an unbounded (cofinal, over h and N) supply of Sep is exactly equivalent to Irrational(S). The table below records every (h, N, L) -shaped or t -shaped instance that has actually been checked; none of it is cofinal, and none of it is presented as deciding #249.

Anchor	Exact statement of what was checked	Scale	Site
Fixed-window deposit, depth 16	$\text{Sep}(h, 12, 16)$ holds for every $h \in \{1, \dots, 8\}$ (by <code>decide</code> , 256 totient values below 37); consequently $S \neq r$ for every $r \in \mathbb{Q}$ with $1 \leq h \leq 8$ and $r.\text{den} \mid 2^{12}(2^h - 1)$.	scale:fixed	certifiedKill_all_small [TotientTailPeriodKiller.lean] totient_series_ne_rat_of_den_dvd [TotientTailPeriodKiller.lean]
Fixed-window deposit, depth 14, wider net	$\text{Sep}(h, 14, \text{depth } 14)$ -shaped exclusion extends to every $h \in \{1, \dots, 16\}$: $S \neq r$ for every $r \in \mathbb{Q}$ with $1 \leq h \leq 16$ and $r.\text{den} \mid 2^{14}(2^h - 1)$.	scale:fixed	totient_series_ne_rat_of_den_dvd_pow_two_mul_mersenne_upto_sixteen [CertificateKernel.lean] certifiedKill_diagonal_all_imported [DiagonalPincerCertificateSc] diagonalPincerKillDepth [Diag
Diagonal-pincer certificates, base 12 scales	$\text{Sep}(H_t, H_t, D(t))$, $H_t := \text{lcm}(1, \dots, t)$, holds (by explicit <code>decide/norm_num</code> on checked <code>Nat.totient</code> values, via factored prime-power blocks with Lucas-primality certificates) for $t \in \{1, 2, 3, 4, 5, 7, 8, 9, 11, 13, 16, 17\}$ with certificate depths $D(t) \in \{6, 5, 7, 7, 9, 14, 15, 14, 21, 22, 23, 26\}$ respectively (in the same order).	scale:fixed	certifiedKill_diagonal_all_imported [DiagonalPincerCertificateSc] diagonalPincerKillDepth [Diag
Diagonal-pincer certificates, extended scales through $t = 64$	The same predicate $\text{Sep}(H_t, H_t, D(t))$ is additionally checked, one sibling module per scale, at $t \in \{19, 23, 25, 27, 29, 31, 32, 37, 41, 43, 47, 49, 53, 59, 61, 64\}$ (16 further explicit values). Together with the base 12 scales above this is 28 explicit values of t in total , the complete list being $\{1, 2, 3, 4, 5, 7, 8, 9, 11, 13, 16, 17, 19, 23, 25, 27, 29, 31, 32, 37, 41, 43, 47, 49, 53, 59, 61, 64\}$. This does <i>not</i> establish $\text{Sep}(H_t, H_t, \cdot)$ for infinitely many t ; it is the concrete finite floor of the diagonal sequence, exactly what an unbounded-supply argument would need to extend.	scale:fixed	DiagonalPincerCertificatesT...T61.lean (one file per scale, each named DiagonalPincerCertificatesT endpoint certifiedKill_diagonal_ 3741 [DiagonalPincerCertificateSc, ffl, cde] T64
Farey denominator floor	Every $q \in \mathbb{N}$ with $0 < q \leq Q_0 = 79\,639\,646\,646\,701\,375\,323\,355\,774\,875\,831\,053$ satisfies the window- $(N, K) = (1, 240)$ gap certificate; $q = Q_0 + 1$ is the exact first failure.	scale:bounded	gap_check_window_1_240_le_796396466467013753233557748

Anchor	Exact statement of what was checked	Scale	Site
Coprimalty-probability / Möbius-square Farey floor	Every $d \in \mathbb{N}$ with $0 < d \leq Q_1 = 39\,819\,823\,323\,350\,687\,661\,677\,887\,437\,915\,526$ (exactly $\lfloor Q_0/2 \rfloor$) is excluded as a denominator of $\Pr(\gcd(X, Y) = 1)$ and, separately, of $T = \sum_d \mu(d)/(2^d - 1)^2$.	scale:bounded	tsum_visible_coprime_pairs_ne_int_div_of_den_le_39819823323350687661677887437915526_tsum_moebius_div_two_pow_sub_one_sq_ne_int_div_of_den_le_39819823323350687661677887437915526

Remark 5.20 (What this table is not). No row above is cofinal in its indexing parameter (h , t , or the denominator bound), and no row is claimed to extrapolate. The certificate depths in the diagonal-pincer row are strictly increasing and irregular (6, 5, 7, 7, 9, 14, 15, 14, 21, 22, 23, 26, ...); no closed-form growth rate for $D(t)$ is proved or conjectured in the corpus. This table is the complete inventory of finite Erdős-#249 certificate evidence as of this writing; the cofinal supply obligation it falls short of is the open target developed in Part 1.

6 Catalogue of usable premises

This catalogue renders every producer and converter/identity result found for Erdős #249 ($S = \sum_{n \geq 1} \varphi(n)/2^n$) in the certificate-kernel lane (`TotientTailPeriodKiller.lean`, `CertificateKernel.lean`, and the lcm/diagonal/cone reduction family) and the Möbius–Mersenne / squared-Lambert lane (`GcdMomentCalculus.lean`, `RepunitMobiusNumerator.lean` through `PrimePowerJumpDynamics.lean`, `MersenneLambertLadder.lean`). Each entry is a self-contained premise: exact hypotheses, exact conclusion, the coordinate it was proved in, and the Lean site. **#249 is OPEN**; nothing below decides it. Entries labelled **consumer**-shaped (an implication whose hypothesis is an unsupplied cofinal predicate) are marked explicitly as conditional the hypothesis itself carries `[Open]` and is never claimed as proved. Site paths are relative to `Erdos249257/` in `/Users/willcook/src/plectis-lean-work/Erdos249257/` (mirrored from `/Users/willcook/src/plectis-`

6.1 Producers

Producers conclude an existence or a supply: a witnessed object, a witnessed finite family, or an unconditional dimension/growth lower bound obtained by exhibiting witnesses (CRT, Dirichlet primes in AP, Bertrand’s postulate, explicit `decide` computation). Sorted by scale: uniform, then bounded, then fixed; no producer in this lane is cofinal (the cofinal-scale reduction theorems are consumer-shaped and are catalogued as converters below, since each moves a cofinal hypothesis across a coordinate boundary to the target conclusion).

Theorem 6.1 (`cert:a9 — eventual_period_of_not_irrational`, THE TAIL-PERIOD LAW). *If S is rational then a period exists: $\neg \text{Irrational}(S) \rightarrow \exists h : \mathbb{N}, 0 < h \wedge \exists N_0 : \mathbb{N}, \forall N \geq N_0, \text{totientTail}(N + h) - \text{totientTail}(N) \in \text{range}((\uparrow) : \mathbb{Z} \rightarrow \mathbb{R})$. The witness is explicit: $h = \varphi(\text{oddPart}(r.\text{den}))$, $N_0 = v_2(r.\text{den})$, from Euler’s theorem applied to the odd part of the hypothetical denominator.*

[Lean] `scale:uniform coord:binary-digit` [eventual_period_of_not_irrational](#) [`TotientTailPeriodKiller.lean:358`]

Theorem 6.2 (`cert:d5 — not_irrational_totientSeries_implies_unbounded_carryRank_unconditional`). $\neg \text{Irrational}(S) \rightarrow \exists v > 0, \exists u : \mathbb{N} \rightarrow \mathbb{Z}, \text{IsTemperedBinaryOrbit } \varphi v u \wedge$

$\forall e, 2^e - 1 \leq \text{finrank}_{\mathbb{Q}}(\text{span}(\text{range}(\text{canonicalCarryKernelFamily } u e)))$. *Rationality of S forces its associated tempered integral binary-carry orbit to have unboundedly rich dyadic-section rank; a second, coordinate-independent necessary condition on rationality, parallel to `cert:a9` but in the carry-kernel-rank coordinate rather than the binary-digit-periodicity coordinate. Explicitly flagged in its own docstring as not by itself an irrationality proof: a separate rationality-side rank-compression theorem (contradicting `cert:d4` below) would still be needed.*

[Lean] scale:uniform coord:other:carry-kernel-rank [not_irrational_totientSeries_implies_unbounded_carryRank_unconditional](#) [TotientCarryKernelRigidity.lean:284]

Theorem 6.3 (`cert:d4` — `linearIndependent_canonicalTotientKernelFamily`). *For every $e : \mathbb{N}$, the canonical dyadic totient-kernel family $\text{canonicalTotientKernelFamily}(e) : \text{TotientCanonicalIndex}(e) \rightarrow \mathbb{N} \rightarrow \mathbb{Q}$, which has exactly $2^e + 1$ channels, is linearly independent over $\mathbb{Q}$. Proved unconditionally by constructing, via CRT and Dirichlet's theorem on primes in arithmetic progression (`PrimesCongruentOne/PrimesInAP` from `Mathlib`), an explicit evaluation point at which one channel becomes prime and every other channel picks up a fresh prime $\equiv 1$ modulo a large power of 2 — a genuine witnessed producer, not merely a dimension count. Consequently $\neg \text{FiniteDimensional } \mathbb{Q}(\text{span } \mathbb{Q}(\text{range fullTotientKernelFamily}))$. Self-flagged: this shows the dyadic-kernel side is infinite-rank; it is not itself an irrationality proof.*

[Lean] scale:uniform coord:other:dyadic-kernel-rank [card_totientCanonicalIndex](#) [TotientMahlerDefect.lean:51] [linearIndependent_canonicalTotientKernelFamily](#) [TotientMahlerDefect.lean:900]

Proposition 6.4 (`cert:a8` — `tail_diff_int_of_den_dvd`). *If $S = (r : \mathbb{R})$ for $r : \mathbb{Q}$ and $r.\text{den} \mid 2^N \cdot (2^h - 1)$, then $\text{totientTail}(N + h) - \text{totientTail}(N) \in \text{range}((\uparrow) : \mathbb{Z} \rightarrow \mathbb{R})$: a rational value of S with a denominator of this shape forces the shifted tail difference to be an integer. The sharp contrapositive engine underlying `cert:a9` and `cert:a6`.*

[Lean] scale:uniform coord:binary-digit [tail_diff_int_of_den_dvd](#) [TotientTailPeriodKiller.lean:327]

Proposition 6.5 (`mob:b5` — `top fibre survives, T3, mobiusNumerator_gcd_cyclotomicValue`). *For r squarefree: $\text{gcd}(|\text{mobiusNumerator}(r)|, \text{cyclotomicValue}(r)) = 1$, hence $\text{cyclotomicValue}(r) \mid \text{baseMobiusShadow}(r).\text{den}$. The top cyclotomic channel $|\Phi_r(2)|$ is produced as a survivor: it can never cancel from the numerator and is exhibited as an unconditional divisor of the reduced denominator at every squarefree r .*

[Lean] scale:uniform coord:cyclotomic [mobiusNumerator_gcd_cyclotomicValue](#) [CyclotomicProjectionOfShadow.lean:389] [cyclotomicValue_dvd_baseMobiusShadow_den](#) [CyclotomicProjectionOfShadow.lean:389]

Proposition 6.6 (`mob:b6` — `upper-half prime channel survival, T4`). *For a scale coprime to $C = \prod_{p \in \text{upperHalfPrimes}(t)} \text{mersenne}(p)$ (or with prime support $\leq t$), the whole finite product C over the explicit upper-half prime set $\text{upperHalfPrimes}(t) = \{p \text{ prime} : t/2 < p \leq t\}$ divides the reduced denominator of the scaled Möbius shadow at LCM height t , unscaled. A genuine finite family of surviving Mersenne-prime channels is exhibited explicitly at every t , not merely bounded below.*

[Lean] scale:uniform coord:cyclotomic [upperHalfPrimes](#) [MersenneShadowCyclotomicNoncollapse.lean:914] [upperHalfChannel_product_dvd_den_of_scale_primeFactors_le](#) [MersenneShadowCyclotomicNoncollapse.lean:914] [lcmHeight_upperHalf_product_dvd_den](#) [MersenneShadowCyclotomicNoncollapse.lean:983]

Proposition 6.7 (`mob:b7a` — `denominator growth lower bound`). *For $t \geq 5$ (Bertrand's postulate supplies nonemptiness of $\text{upperHalfPrimes}(t)$): $2^{t/2} \leq \prod_{p \in \text{upperHalfPrimes}(t)} \text{mersenne}(p) \leq (\text{lcmHeight}(t) \cdot \text{numericMobiusShadow}(\text{lcmHeight}(t))).\text{den}$. An exponential-in- $t/2$ growth lower bound for the reduced denominator at every LCM height, produced from `mob:b6`'s explicit surviving channel product. Denominator-only: does not by itself rule out cancellation by a foreign-defect term, hence does not by itself prove #249.*

[Lean] scale:uniform coord:mobius-mersenne [upperHalfMersenneProduct_lower_bound](#) [MersenneShadowDenominatorGrowth.lean:1000]

Proposition 6.8 (mob:b7b — exact denominator value). *For every t (no lower bound on t needed for this direction): $(\text{lcmHeight}(t) \cdot \text{numericMobiusShadow}(\text{lcmHeight}(t))).\text{den} = \text{mersenne}(\text{lcmRadical}(t) | \text{oddJordanScalar}(\text{lcmRadical}(t)))$. A fully closed form for the reduced denominator at every scale, produced (not merely bounded) as an explicit rational function of t .*

[Lean] scale:uniform coord:mobius-mersenne [lcmHeight_scaledMobiusShadow_den_exact](#) [MersenneShadowDenominator]

Proposition 6.9 (mob:d3 — signed dyadic sum nonvanishing, scaled_dyadic_sum_ne_zero). *If a finite signed sum $\sum_{i \in s} u(i) \cdot 2^{e(m)-e(i)}$ (clearing dyadic denominators to a common exponent $e(m)$) has one index $m \in s$ with strictly maximal exponent $e(m)$ and odd coefficient $u(m)$, while every other index has strictly smaller exponent, then the cleared sum is $\equiv 1 \pmod{2}$, hence nonzero. A positive finite-nonvanishing engine, produced via a rectangular Cauchy–Binet determinant expansion (`det_mul_rectangular`) applied to the signed-Hankel/ q -moment construction (`hankelDet`, `truncatedMoment`); the underlying arithmetic is a $q = 1/2$ Hankel-determinant framework for a possible Padé route to #249. No supply theorem is proved that such a unique-terminal configuration exists cofinally this is a per-instance producer, not a cofinal producer.*

[Lean] scale:bounded coord:p-adic [scaled_dyadic_sum_ne_zero](#) [SignedQMomentObstruction.lean:95] [det_mul_rectangular](#) [SignedQMomentObstruction.lean:28]

Proposition 6.10 (cert:a11 — certifiedKill_all_small). $\forall h \in [1, 8]$, certifiedKill h 12 16 a finite, fully unconditional **decide**-checked family of 8 certificate witnesses at the fixed point $N = 12$, $L = 16$ (256 explicit totient values below 37 evaluated). Consequence, via cert:a6: $\forall r : \mathbb{Q}, 1 \leq h \leq 8, r.\text{den} \mid 2^{12} \cdot (2^h - 1) \rightarrow S \neq r$.

[Cert] scale:fixed coord:other:binary-window [certifiedKill_all_small](#) [TotientTailPeriodKiller.lean:404] [totient_series_ne_rat_of_den_dvd](#) [TotientTailPeriodKiller.lean:416]

Proposition 6.11 (cert:a12 — upto-sixteen deposit). $\forall h \in [1, 16]$ at exponent 14: a wider, fully unconditional **decide**-checked family of certificate witnesses, subsuming cert:a11 on both axes (larger h -range, larger 2-adic exponent). Consequence: $\forall r : \mathbb{Q}, 1 \leq h \leq 16, r.\text{den} \mid 2^{14} \cdot (2^h - 1) \rightarrow S \neq r$.

[Cert] scale:fixed coord:other:binary-window [totient_series_ne_rat_of_den_dvd_pow_two_mul_mersenne_upto_sixteen](#) [CertificateKernel.lean:18562]

Proposition 6.12 (cert:b11 — diagonal pincer finite deposits, THE DIAGONAL-PINCER IDENTITY). `certifiedKill_diagonal_all_imported` : $\forall t \in \{1, 2, 3, 4, 5, 7, 8, 9, 11, 13, 16, 17, \dots\}$, `certifiedKill`(periodLcm(t)) with explicit certificate depths $[6, 5, 7, 7, 9, 14, 15, 14, 21, 22, 23, 26, \dots]$, extended by sibling modules through $t = 64$ (28 explicit deposits total, endpoint `certifiedKill_diagonal_t64`). Each instance is a finite **decide**/`norm_num` computation on explicit `Nat.totient` values, factored via checked prime-power blocks (`FactorBlock`, `totient_factorBlocks`, `prime_dvd_factorBlocks`) with Lucas-primality certificates for the primes involved. This is the concrete finite floor of the single-parameter diagonal sequence $P(t) := \exists L, \text{certifiedKill}(\text{periodLcm}(t))(\text{periodLcm}(t)) L$ (cert:b3 below): it produces $P(t)$ for these 28 explicit t , and does not establish $P(t)$ for infinitely many t quantifier order is exact: a finite witnessed list, not a cofinal supply.

[Cert] scale:fixed coord:other:lcm-diagonal [certifiedKill_diagonal_all_imported](#) [DiagonalPincerCertificates.lean:2870] [diagonalPincerKillDepth](#) [DiagonalPincerCertificates.lean:2870] [certifiedKill_diagonal_t64](#) [DiagonalPincerCertificateT64Endpoint.lean:1928]

6.2 Converters and exact identities

Converters and identities move a statement between coordinates without changing its truth value: definitions, exact algebraic identities, iff-characterisations, and the implication theorems that convert a supply hypothesis in one coordinate into the target conclusion `Irrational(S)`. This subsection carries the certificate-kernel core (the `Sep(h,N,L)` predicate, its completeness

iff, and every known reformulation of the open supply obligation) and the Möbius–Mersenne / squared-Lambert identity family, including the Mersenne–Lambert five-row status ladder. Sorted by scale: foundational (n/a) definitions and identities first, then uniform, then cofinal, then bounded, then fixed.

6.2.1 Foundational definitions and identities (scale n/a)

Definition 6.13 (cert:a1 — `totientTail`). $\text{totientTail}(N) := \sum'_{j \geq 0} \varphi(N + 1 + j) / 2^{j+1}$, the fractional layer of $2^N \cdot S$; well-defined for every N .

[Lean] scale:n/a coord:binary-digit `totientTail` [TotientTailPeriodKiller.lean:57]

Definition 6.14 (cert:a3 — `windowDiscrepancy`). $\text{windowDiscrepancy}(h, N, L) := \sum_{j < L} (\varphi(N + h + 1 + j) - \varphi(N + 1 + j)) \cdot 2^{L-1-j} \in \mathbb{Z}$, the depth- L truncation of $2^L \cdot (\text{totientTail}(N + h) - \text{totientTail}(N))$; computable and decidable given h, N, L .

[Lean] scale:n/a coord:other:binary-window `windowDiscrepancy` [TotientTailPeriodKiller.lean:66]

Definition 6.15 (cert:a4 — `certifiedKill`, THE Sep(h,N,L) PREDICATE). $\text{certifiedKill}(h, N, L) := (N + h + L + 2 : \mathbb{Z}) < \text{windowDiscrepancy}(h, N, L) \bmod 2^L < 2^L - (N + h + L + 2)$ the residue of the window discrepancy modulo 2^L avoids the shrinking radius- $(N + h + L + 2)$ neighbourhood of 0. This is exactly the object named $\text{Sep}(h, N, L)$: purely finite arithmetic, decidable, with no analytic hypothesis. This is the certificate kernel's core object; every theorem below is either a hypothesis-shape wrapping it or an unconditional finite instance of it.

[Lean] scale:n/a coord:other:binary-window `certifiedKill` [TotientTailPeriodKiller.lean:72]

Proposition 6.16 (cert:a2 — shift identity, `two_pow_mul_totient_series_eq`). $2^N \cdot \left(\sum'_{n \geq 0} \varphi(n) / 2^n \right) = \text{totientPrefix}(N) + \text{totientTail}(N)$, where $\text{totientPrefix}(N) = \sum_{n \leq N} \varphi(n) \cdot 2^{N-n} \in \mathbb{N}$ is an exact integer. Rationality/periodicity of S 's binary expansion reduces exactly to a statement about totientTail . Template shape (constant · prefix + tail split) reusable for any $\sum f(n) / 2^n$.

[Lean] scale:uniform coord:binary-digit `two_pow_mul_totient_series_eq` [TotientTailPeriodKiller.lean:150]

Lemma 6.17 (cert:a5 — certificate depth floor, `certifiedKill_depth_floor`). $\text{certifiedKill}(h, N, L) \rightarrow 2 \cdot (N + h + L + 2) < 2^L$. A structural corollary of the kernel definition: the certificate depth L cannot stay bounded while $N + h \rightarrow \infty$; L must grow at least logarithmically with $N + h$. Necessary context for reading every fixed/bounded-scale kernel instance below.

[Lean] scale:uniform coord:other:binary-window `certifiedKill_depth_floor` [TotientTailPeriodKiller.lean:79]

Proposition 6.18 (cert:a6 — certificate soundness, the kernel's converter direction, `tail_diff_notMem_int_of_certifiedKill`). $\text{certifiedKill}(h, N, L) \rightarrow \text{totientTail}(N + h) - \text{totientTail}(N) \notin \text{range}((\uparrow) : \mathbb{Z} \rightarrow \mathbb{R})$. This is the kernel's soundness converter: it moves a finite, decidable, binary-window fact into a real-analytic non-integrality fact. Composes with cert:a9 (tail-period law) by contradiction to kill a hypothetical rational's period.

[Lean] scale:uniform coord:other:binary-window `tail_diff_notMem_int_of_certifiedKill` [TotientTailPeriodKiller

Theorem 6.19 (cert:a7 — certificate completeness, `exists_certifiedKill_iff_tail_diff_notMem_int`). $(\exists L, \text{certifiedKill}(h, N, L)) \iff \text{totientTail}(N + h) - \text{totientTail}(N) \notin \text{range}((\uparrow) : \mathbb{Z} \rightarrow \mathbb{R})$, for all h, N . Certificates are complete receipts of non-integrality, not merely sufficient: the certificate vocabulary is dispensable, and the real target of the whole kernel is exactly the right-hand real-analytic statement. Converts the open #249 obligation freely between "supply of certificates" language and "supply of non-integral tail differences at arbitrarily large scale" language (cert:b8 below uses the latter form). Any independent non-integrality proof from any coordinate automatically yields a certificate by this iff, and vice versa.

[Lean] scale:uniform coord:other:binary-window `exists_certifiedKill_iff_tail_diff_notMem_int` [LcmConeFlatness.lean:316]

Definition 6.20 (cert:b1 — `periodLcm`, the universal period ray). $\text{periodLcm}(0) = 1$, $\text{periodLcm}(t+1) = \text{lcm}(\text{periodLcm}(t), t+1)$, i.e. $\text{periodLcm}(t) = \text{lcm}(1, \dots, t)$; $t \leq \text{periodLcm}(t)$; every primitive period $h_0 \leq t$ divides $\text{periodLcm}(t)$. Pure number theory about $\text{lcm}(1..t)$, zero totient content; directly reusable for #257 or any period-search problem "stand on the universal-period ray to remove one free parameter" is a general reduction technique.

[Lean] scale:n/a coord:other:lcm-period-ray `periodLcm` [CarrySurvivorExtinction.lean:515] `le_periodLcm` [CarrySurvivorExtinction.lean:515]

Proposition 6.21 (cert:c1 — `farey_gap`, the mediant lemma). For $a, b, c, d, r, s : \mathbb{Z}$ with $b > 0$, $d > 0$, unimodular neighbours $bc - ad = 1$, and r strictly between a/b and c/d ($a \cdot s < r \cdot b$, $r \cdot d < c \cdot s$): $b + d \leq s$. Any rational strictly between two unimodular Farey neighbours has denominator at least the sum of the neighbours' denominators. Fully problem-agnostic pure Stern–Brocot/Farey fact, zero totient content, directly reusable for #257.

[Lean] scale:n/a coord:farey `farey_gap` [GapFareyBound.lean:51]

Proposition 6.22 (cert:d1 — `irrational_of_den_mul_abs_sub_tendsto_zero`, generic Dirichlet-gap criterion). If $u : \mathbb{N} \rightarrow \mathbb{Q}$ is eventually never equal to $x : \mathbb{R}$, and $\text{den}(u(k)) \cdot |x - u(k)| \rightarrow 0$, then $\text{Irrational}(x)$. Classical Dirichlet-approximation irrationality criterion, zero totient/Mersenne content, already reused in this corpus to prove full-support Erdős–Borwein irrationality (#257-shaped). Apply to any explicit sequence of convergents to S with a provable denominator-gap $\rightarrow 0$ bound.

[Lean] scale:n/a coord:n/a `irrational_of_den_mul_abs_sub_tendsto_zero` [CertificateKernel.lean:5371]

Proposition 6.23 (cert:d2 — `irrational_of_int_mul_near_int`, classical near-integer criterion). If $\forall q > 0, \exists m, z : \mathbb{Z}, 0 < |m \cdot \xi - z| < 1/q$, then $\text{Irrational}(\xi)$. The classical Erdős-1948-shape criterion behind digit/carry irrationality proofs. Base-power specialisation `irrational_of_pow_mul_near_int` (witnesses of form $b^n \cdot \xi$) directly matches a digit/carry construction in any base b , hence directly usable for #257's $b^n - 1$ denominators. Together with cert:d1 these are the only two general-purpose irrationality criteria in the whole kernel; everything else exists to supply or substitute for their hypotheses.

[Lean] scale:n/a coord:n/a `irrational_of_int_mul_near_int` [CertificateKernel.lean:5792] `irrational_of_pow_mul_near_int` [CertificateKernel.lean:5821]

Proposition 6.24 (cert:d3 — `SeparatedMinorCertificate`, linear independence from a nonzero minor). For an indexed family $\text{family} : \iota \rightarrow \mathbb{N} \rightarrow \mathbb{Q}$: a `SeparatedMinorCertificate` (an explicit finite evaluation-point assignment $\text{rowIndex} : \iota \rightarrow \mathbb{N}$ with $\det(\text{family}(j)(\text{rowIndex}(i)))_{i,j} \neq 0$) implies `LinearIndependent` $\mathbb{Q}$ family. Fully generic finite-dimensional linear algebra; family is a free variable, nothing here mentions totient or Mersenne structure. Reusable for any finite-rank/linear-independence obstruction in either open problem.

[Lean] scale:n/a coord:n/a `SeparatedMinorCertificate` [TotientMahlerDefect.lean:73] `linearIndependent_of_separatedMinorCertificate` [TotientMahlerDefect.lean:81]

Proposition 6.25 (cert:d9 — `positive_rational_difference_lower_bound`). For $\text{pfx} < \text{whole} : \mathbb{Q}$ (strict): $1/(\text{whole.den} \cdot \text{pfx.den}) \leq (\text{whole} : \mathbb{R}) - (\text{pfx} : \mathbb{R})$. A positive rational difference is bounded below by the reciprocal of the product of the two actual reduced denominators, not a displayed/guessed one. Fully general; feeds `prefixDenominator_shell_power_bound_of_rational_difference` and `nextSupport_power_bound_of_rational_difference`, converting any analytic upper bound on a rational gap into a denominator-growth lower bound plug in any tail estimate from either open problem's coordinate.

[Lean] scale:n/a coord:n/a `positive_rational_difference_lower_bound` [PrimitiveRationalGapSupply.lean:31]

Proposition 6.26 (mob:a1a — the Möbius-squared-Mersenne identity). $S = \sum_{n \geq 1} \varphi(n)/2^n = \sum'_{d \geq 1} \mu(d) \cdot 2^d / (2^d - 1)^2 = \frac{1}{2} + \sum'_{d \geq 1} \mu(d) / (2^d - 1)^2$. Unconditional identity of convergent real series; this is the reduced target every Möbius–Mersenne-lane result in this catalogue feeds.

[Lean] scale:n/a coord:mobius-mersenne `tsum_totient_half_pow_eq_half_add_moebius_sq` [SquaredMersenneDiagonalEnclosure.lean:50] `mobius_square_series_eq_partial_add_tail` [SquaredMersenneDiagonalEnclosure.lean:72]

Corollary 6.27 (mob:a1b — #249 restated in the squared-Lambert coordinate). *Immediate from mob:a1a: $\text{Irrational}(S) \iff \sum'_{d \geq 1} \mu(d)/(2^d - 1)^2 \notin \mathbb{Q}$. Denote the right-hand series $L_2(\mu) := S - 1/2$ (mob:a3 below). Every subsequent Möbius-coordinate obstruction or identity in this catalogue is measured against this exact restatement, not the original totient series.*

[Lean] scale:n/a coord:mobius-mersenne `tsum_totient_half_pow_eq_half_add_moebius_sq` [SquaredMersenneDiagonalEnclosure.lean:72]

Proposition 6.28 (mob:a2 — squared-Lambert transfer engine, `tsum_lambert_linear_weight_sq_pure`). *For $w : \mathbb{N} \rightarrow \mathbb{R}$ with $|w(d)| \leq d$ for $d > 0$, and $0 \leq r < 1$: $\sum'_{d:\mathbb{N}^+} w(d) \cdot (r^d / (1 - r^d))^2 = \sum'_{n:\mathbb{N}^+} \left(\sum_{e|n} w(e) \cdot (n/e - 1) \right) \cdot r^n$. Converts any linear-growth-bounded squared-Lambert series into a divisor-convolution power series the single reusable brick for the whole level-2 Möbius-Lambert ladder below. Pure Dirichlet-convolution/Lambert-series algebra, no Mersenne-specific structure; instantiate at $w = \mu, 1, \varphi$ to obtain mob:a3–a5, and directly reusable for a weighted or squared variant of #257's series.*

[Lean] scale:uniform coord:mobius-mersenne `tsum_lambert_linear_weight_sq_pure` [GcdMomentCalculus.lean:105]

Definition 6.29 (mob:a3 — $L_2(\mu)$ is exactly #249). $L_2(\mu) := \sum'_{d:\mathbb{N}^+} \mu(d)/(2^d - 1)^2 = S - 1/2$. The open #249 atom, restated as the Möbius rung of the level-2 (squared) Lambert ladder; consequence of mob:a1a and mob:a2 at $w = \mu$.

[Cited] scale:n/a coord:mobius-mersenne `tsum_totient_half_pow_eq_half_add_moebius_sq` [SquaredMersenneDiagonalEnclosure.lean:72]

Proposition 6.30 (mob:a4 — $L_2(1)$, the known q -zeta anchor rung, `tsum_one_div_mersenne_sq_eq_sigma_sub_tau_series`). $\sum'_{d:\mathbb{N}^+} 1/(2^d - 1)^2 = \sum'_{n:\mathbb{N}^+} (\sigma(n) - \tau(n)) \cdot (1/2)^n = \zeta_q(2) - \zeta_q(1)$ at $q = 1/2$. The level-2 ζ -rung, exactly evaluated in Lean. Irrationality of this value is cited (Postelmans–Van Assche q -Padé), not formalised only the identity itself is machine-checked. This is the "known" sibling rung mirroring mob:a3 in the level-mirror table mob:a6.

[Lean] scale:n/a coord:mobius-mersenne `tsum_one_div_mersenne_sq_eq_sigma_sub_tau_series` [GcdMomentCalculus.lean:105]

Proposition 6.31 (mob:a5 — $L_2(\varphi)$, the Pillai gcd-moment rung, `tsum_totient_div_mersenne_sq_eq_gcd_moment_series`). $\sum'_{d:\mathbb{N}^+} \varphi(d)/(2^d - 1)^2 = \sum'_{n:\mathbb{N}^+} (P(n) - n) \cdot (1/2)^n$, where $P = \varphi * \text{Id}$ (Pillai's gcd-sum function). Equals $\mathbb{E}[\text{gcd}(X, Y)]$ for independent fair-coin waiting times X, Y (probabilistic coordinate, mob:a7–a9). A cousin rung, not #249 itself; status open.

[Lean] scale:n/a coord:mobius-mersenne `tsum_totient_div_mersenne_sq_eq_gcd_moment_series` [GcdMomentCalculus.lean:105]

Observation 6.32 (mob:a6 — the level-mirror table). Level 1 ($L(f) := \sum f(d)/(2^d - 1)$): $L(\mu) = 1/2$ (rational, trivial); $L(1) = \mathcal{E}$, the Erdős–Borwein constant (irrational, Erdős 1948, machine-checked in this kernel, cert:d7c below); $L(\varphi) = 2$ (rational). Level 2 ($L_2(f) := \sum f(d)/(2^d - 1)^2$): $L_2(\mu) = S - 1/2$ (mob:a3, OPEN, = #249); $L_2(1) = \zeta_q(2) - \zeta_q(1)$ (mob:a4, irrational, cited); $L_2(\varphi) = \mathbb{E}[\text{gcd}]$ (mob:a5, open, Pillai). At level 1 the Möbius rung is trivial and the ζ -rung is hard; at level 2 the ζ -rung is known and the Möbius rung is #249 Möbius projection is the single wall at both levels. This mirror-structure phenomenon is a general observation about Dirichlet-convolution ladders, potentially informative for #257's Mersenne-shifted sums too.

[Lean] scale:n/a coord:mobius-mersenne

Proposition 6.33 (mob:a7 — gcd-divisibility factorises, `tsum_pos_pair_both_dvd_half_eq_inv_mersenne_sq`). *For independent fair-coin waiting times X, Y ($P(X = n) = 2^{-n}$) and $d > 0$: $P(d \mid X \wedge d \mid Y) = 1/(2^d - 1)^2$. The foundation stone for reading $L_2(f)$ as $\mathbb{E}[(f * \zeta)(\text{gcd}(X, Y))]$. Pure probability/geometric-series fact about independent geometric random variables, zero #249-specific content; directly reusable for a two-coordinate gcd structure in #257.*

[Lean] scale:uniform coord:probability `tsum_pos_pair_both_dvd_half_eq_inv_mersenne_sq` [GcdMomentCalculus.lean:105]

Proposition 6.34 (mob:a8 — reduced-direction law, `tsum_pos_coprime_inv_mersenne_eq_one`). $\sum'_{(a,b): a,b \geq 1, \gcd(a,b)=1} 1/(2^{a+b} - 1) = 1$: the sum over exact bipartite coprime pairs. Every positive coprime pair carries a slope mass $1/(2^{a+b} - 1)$, and these mass exactly one the root cylinder $M(1,1) = 1$ of the Stern–Brocot tree, mob:a9. Pure coprimality/geometric-series law, applies verbatim to any base- b analogue.

[Lean] scale:n/a coord:probability [tsum_pos_coprime_inv_mersenne_eq_one](#) [GcdMomentCalculus.lean:349]

Proposition 6.35 (mob:a9a — Stern–Brocot cylinder recursion, `cylinderMass_split`). $M(a,b) := 1/((2^a - 1)(2^b - 1))$ satisfies the exact telescoping identity $M(a,b) = 1/(2^{a+b} - 1) + M(a+b,b) + M(a,a+b)$ for $a,b : \mathbb{N}^+$: an exact Markov measure on the Stern–Brocot median tree, whose closed form is a product of two independent-coordinate divisor probabilities $P(a | X) \cdot P(b | Y)$ (mob:a7).

[Lean] scale:uniform coord:other:stern-brocot [cylinderMass_split](#) [GcdMomentCalculus.lean:474]

Proposition 6.36 (mob:a9b — depth- d convergence rate, `sternBrocotDepthMass_error / tendsto_sternBrocotDepthMass`). Children of a Stern–Brocot cylinder carry at most $2/3$ of the parent’s mass; the depth- d finite unfolding of mob:a9a converges to $M(a,b)$ at explicit rate $|M(a,b) - M_d(a,b)| \leq (2/3)^d \cdot M(a,b)$. The “recursion limit = subtree mass” identification is explicitly not claimed beyond this rate bound (flagged as a wave-21 gap in the source). All-left cusp cylinders $M(N,1) = 1/(2^N - 1)$ are near-Mersenne-reciprocal, one exponential order closer to the Erdős–Borwein engine than raw $\varphi(n)$ coefficients an advisory bridge candidate to Erdős-1948-style arguments, not a proved route.

[Lean] scale:uniform coord:other:stern-brocot [sternBrocotDepthMass_error](#) [GcdMomentCalculus.lean:525]
[tendsto_sternBrocotDepthMass](#) [GcdMomentCalculus.lean:560]

Proposition 6.37 (mob:b1 — repunit gcd word, T1, `mobiusNumeratorPolynomial_eq_gcdWord`). For r squarefree, the divisor-signed polynomial `mobiusNumeratorPolynomial(r) :=  $\sum_{d|r} \mu(d) \cdot (r/d) \cdot \text{spacedRepunit}(d, r/d)$` equals `gcdWord(r)`, whose X^k coefficient ($k < r$) is `gcdWordCoeff(r, k) =  $(r/\gcd(r, k)) \cdot \varphi(\gcd(r, k))$` , strictly positive for $k < r$ and zero for $k \geq r$. The signed repunit numerator is exactly a positive gcd word; the classical polynomial identity underlying it (signed spaced repunit = gcd word) is base-independent, though the application here is Mersenne-specific.

[Lean] scale:n/a coord:cyclotomic [mobiusNumeratorPolynomial_eq_gcdWord](#) [RepunitMobiusNumerator.lean:205]
[mobiusNumeratorPolynomial_coeff](#) [RepunitMobiusNumerator.lean:217] [mobiusNumeratorPolynomial_coeff_pos](#) [RepunitMobiusNumerator.lean:234]

Proposition 6.38 (mob:b2 — evaluation at 2 recovers the integer numerator, `mobiusNumeratorPolynomial_eval_two`). `mobiusNumeratorPolynomial(r).eval 2 = mobiusNumerator(r)` for r squarefree, where `mobiusNumerator(r) :=  $\sum_{s \subseteq \text{primeFactors}(r)} (-1)^{|s|} \cdot (r/d) \cdot (\text{mersenne}(r)/\text{mersenne}(d))$` , $d = \prod s$. Bridges the polynomial (mob:b1) world to the integer arithmetic used by every denominator-survival theorem below.

[Lean] scale:uniform coord:cyclotomic [mobiusNumeratorPolynomial_eval_two](#) [RepunitMobiusNumerator.lean:320]
[mobiusNumerator](#) [RadicalMobiusShadow.lean:101]

Proposition 6.39 (mob:b3 — radical shadow scale decomposition). `baseMobiusShadow(r) := mobiusNumerator(r)/(2^r - 1)` (unscaled); `numericMobiusShadow(H) := baseMobiusShadow(rad(H))/rad(H)`; exactly $H \cdot \text{numericMobiusShadow}(H) = (H/\text{rad}(H)) \cdot \text{baseMobiusShadow}(\text{rad}(H))$ for $H > 0$. Reduced-denominator identity for the unscaled shadow: `baseMobiusShadow(r).den = mersenne(r)/gcd(|mobius` for $r > 0$ exact and generic, no coprimality assumed, no channel-survival hidden.

[Lean] scale:uniform coord:mobius-mersenne [scaledMobiusShadow_eq_radicalBase](#) [RadicalMobiusShadow.lean:125]
[baseMobiusShadow_den](#) [RadicalMobiusShadow.lean:150]

Proposition 6.40 (mob:b4 — cyclotomic congruence, T2, `cyclotomic_dvd_mobiusNumeratorPolynomial_sub`). For r squarefree, $m | r$: $\Phi_m | (\text{mobiusNumeratorPolynomial}(r) - C(\mu(m) \cdot J_2(r/m)))$ in

$\mathbb{Z}[X]$, where $J_2 = \mu * \text{id}^2$ is the Jordan totient. Evaluated: $\text{cyclotomicEval}(m) \mid \text{mobiusNumerator}(r) - \mu(m) \cdot J_2(r/m)$. Every cyclotomic fibre $\Phi_m(2)$ of the numerator is congruent to an explicit constant depending only on $\mu(m)$ and the Jordan totient of the cofactor. Setting $m = r$ gives mob:b5 (top fibre survives).

[Lean] scale:uniform coord:cyclotomic [cyclotomic_dvd_mobiusNumeratorPolynomial_sub](#) [CyclotomicProjectionOfShadow.lean:310]
[mobiusNumerator_mod_cyclotomicEval](#) [CyclotomicProjectionOfShadow.lean:310]

Proposition 6.41 (mob:b8a — prime-power jump recurrence, T5, new fibre). *Adjoining a new prime $p \nmid r$: on a genuinely new fibre $m \cdot p$ (with $m \mid r$), $\Phi_{mp} \mid \text{mobiusNumeratorPolynomial}(rp) + \text{expand}_p(\text{mobiusNumeratorPolynomial}(r))$ the new fibre picks up a sign-flipped p -expansion of the old numerator. Holds for p prime, $p \nmid r$, $m \mid r$; the underlying recurrence needs no squarefreeness of r .*

[Lean] scale:uniform coord:cyclotomic [cyclotomic_dvd_primeJump_new_fibre](#) [PrimePowerJumpDynamics.lean:281]
[cyclotomic_dvd_primeJump_new_fibre_constant](#) [PrimePowerJumpDynamics.lean:347]

Proposition 6.42 (mob:b8b — prime-power jump recurrence, T5, old fibre). *Adjoining a new prime $p \nmid r$: on an old fibre m (with $m \mid r$), $\Phi_m \mid \text{mobiusNumeratorPolynomial}(rp) - C(p^2 - 1) \cdot \text{mobiusNumeratorPolynomial}(r)$ old fibres scale by exactly $p^2 - 1$. Requires r squarefree (needed for the T2 constant-fibre step mob:b4 , not for the underlying recurrence itself). An inductive tool for building explicit denominator/numerator values one prime at a time; a candidate base case for an induction proving mob:b7a/b7b ’s growth bound tight.*

[Lean] scale:uniform coord:cyclotomic [cyclotomic_dvd_primeJump_old_fibre](#) [PrimePowerJumpDynamics.lean:310]

Proposition 6.43 (cert:d7 — the Mersenne–Lambert ladder identities). *Define $L(f) := \sum_{n \geq 1} f(n)/(2^n - 1)$ (level 1, no square). Writing $A := \varphi * \mu$ (the primitive Euler weight, α in prose): the Dirichlet-convolution identities $A * \zeta = \varphi$ and $\varphi * \zeta = \text{Id}$ separate five exact values of L along one ladder, each obtained by factoring $L(f)$ as a divisor-transform $(f * 1)$ followed by binary evaluation, $L(f) = \sum_n (f * 1)(m)/2^m$. S sits inside this ladder as $L(A)$: this is #249 restated in positive Erdős–Borwein form.*

[Lean] scale:n/a coord:mobius-mersenne [tsum_totient_div_pow_two_eq_pnat_half_pow](#) [CertificateKernel.lean:18063]

Table 2: The Mersenne–Lambert ladder five-row status table (docstring @ CertificateKernel.lean:18063–18083, body MersenneLambertLadder.lean).

Weight f	Value $L(f)$	Status
μ	$L(\mu) = 1/2$	rational, trivial
φ	$L(\varphi) = 2$	rational
Id	$L(\text{Id}) = \sum_m \sigma(m)/2^m$	transcendental (Nesterenko 1996, [Cited], not formalised)
1	$L(1) = \mathcal{E}$, Erdős–Borwein constant	irrational, [Lean] (irrational_erdosBorwein_series , CertificateKernel.lean:8007) matches #257’s full-supply
$A = \varphi * \mu$	$L(A) = S$	OPEN this is #249

Remark 6.44 (cert:d7 — reading the ladder). This is the single clearest bridge placing #249 inside #257’s native Mersenne–Lambert coordinate ($\sum f(n)/(b^n - 1)$, here $b = 2$): if the #257 lane’s Mersenne-coordinate machinery (achievement sets, greedy orbits) can say anything about the sign or density structure of $A = \varphi * \mu$ specifically, it transfers here for free via this ladder identity. Conversely, any #249 result about $L(A)$ phrased purely in Lambert-series terms (not totient terms) is directly #257-lane-portable. The ‘evaluated’ identity confirming S ’s $\mathbb{N}$ -indexed form equals its $\mathbb{N}^+$ /Lambert-indexed form is the one row here checked directly against source

([tsum_totient_div_pow_two_eq_pnat_half_pow](#) [CertificateKernel.lean:18087]); the body machinery in `MersenneLambertLadder.lean` establishing the other four rows was read only via this docstring and is flagged [Cited]/[Math] accordingly except where a specific declaration is named above.

Proposition 6.45 (mob:e3 — joint-35 cone annihilator, `oldChannel_affine_moment_annihilation` / `joint35_oldChannel_zero`). *The four-vertex affine annihilator $q(X, Y) = XY - 3X - 2Y + 4$ ($q(1, 1) = q(3, 5) = 0$, $q(9, 25) = 152$) kills every “old” divisor channel exactly at any LCM height: for $d \mid H$, $d > 0$, $\text{transportResidueKernel}(d, 15H) - 3 \cdot \text{transportResidueKernel}(d, 3H) - 2 \cdot \text{transportResidueKernel}(d, 5H) + 4 \cdot \text{transportResidueKernel}(d, H) = 0$. General form (`oldChannel_affine_moment_annihilation`): any finite affine annihilator with $\sum c_i = 0$, $\sum c_i \cdot m_i = 0$ kills every old residue channel, independent of the particular $(3, 5)$ choice fully coordinate-free finite linear algebra. Sharp cone radius $19H + 5L + 5$ (`sharpJoint35ConeRadius`). An unbounded certificate supply built from this (not proved) would close #249 via the same cone-flatness route as mob:e2/cert:b6.*

[Lean] scale:uniform coord:other:lcm-diagonal [oldChannel_affine_moment_annihilation](#) [JointExponentTransport.lean:71] [joint35_oldChannel_zero](#) [JointExponentTransport.lean:71] [sharpJoint35ConeRadius](#) [JointExponentTransport.lean:71]

Proposition 6.46 (mob:f1 — composite dilation defect identity, `supportCoeff_mul_eq_add_defect`). *For an abstract support set $A \subseteq \mathbb{N}$, $\text{supportCoeff}_A(n) := \#\{a \in A : a \mid n\}$: for $a \in A$, $a > 0$, $x > 0$, $\text{supportCoeff}_A(ax) = \text{supportCoeff}_A(x) + [a \nmid x] + \text{compositeDilationDefect}_A(a, x)$. On prime-only support (every element of A prime) the defect is identically zero, recovering the classical $\text{supportCoeff}_A(px) = \text{supportCoeff}_A(x) + [p \nmid x]$. Pure divisor combinatorics over an abstract support set, no 2^n -specific structure a shared #249/#257 substrate identity: instantiate $A = \mathbb{N}$ weighted by φ for #249, or A the chosen infinite subset for #257's $\sum_{n \in A} 1/(2^n - 1)$.*

[Lean] scale:uniform coord:other:support-divisor-counting [supportCoeff_mul_eq_add_defect](#) [CompositeDilationDefect.lean:103] [compositeDilationDefect_eq_zero_of_prime_support](#) [CompositeDilationDefect.lean:103] [supportCoeff_mul_prime_support](#) [CompositeDilationDefect.lean:119]

6.2.2 Cofinal-scale converters the certificate-kernel reduction ladder

Every entry in this block is an implication whose *conclusion* is `Irrational(S)` (#249 solved) and whose *hypothesis* is a cofinal supply predicate. **None of these hypotheses is supplied anywhere in the corpus.** Each theorem itself is [Lean]-checked; the hypothesis it consumes carries [Open]. These are listed here, not as producers, because each one is exactly a converter: it moves an unsupplied cofinal statement in the binary-window/lcm-cone coordinate across to the target statement about S , and (via cert:a7's iff) several of them are genuine logical equivalences, not mere one-directional implications.

Theorem 6.47 (cert:a10 — `irrational_totient_series_of_certificate_supply`, WAVE-21 REDUCTION, THE WALL). $(\forall h : \mathbb{N}, 0 < h \rightarrow \forall N_0 : \mathbb{N}, \exists N \geq N_0, \exists L, \text{certifiedKill}(h, N, L)) \rightarrow \text{Irrational}(S)$. *The hypothesis is exactly $\text{Sep}(h, N, L)$ quantified as $\forall h \geq 1 \forall N_0 \geq 0 \exists N \geq N_0 \exists L$. **This hypothesis is nowhere proved in the corpus; this is the central open supply obligation.** Every other reduction theorem in this block is a re-derivation of an equivalent-or-weaker sufficient condition for the same missing supply, never a proof of it.*

[Lean] (implication proved; hypothesis [Open]) scale:cofinal coord:other:binary-window [irrational_totient_series_of_certificate_supply](#) [TotientTailPeriodKiller.lean:394]

Theorem 6.48 (cert:b2 — multiple-period collapse, wave-22, `irrational_totient_series_of_multiple_certificate_supply`). $(\forall h_0 > 0, \forall N_0, \exists m > 0, \exists N \geq N_0, \exists L, \text{certifiedKill}(m \cdot h_0, N, L)) \rightarrow \text{Irrational}(S)$. *Weaker hypothesis than cert:a10: for every primitive period it suffices to certify some multiple. Proof idea (`tail_diff_mul`, telescoping an $m \cdot h$ -difference into*

a sum of m shifted h -differences) is problem-agnostic; logically equivalent-strength weakening of `cert:a10`, still open.

[Lean] (implication proved; hypothesis [Open]) `scale:cofinal coord:other:lcm-period-multiple irrational_totient_series_of_multiple_certificate_supply` [CarrySurvivorExtinction.lean:502]

Theorem 6.49 (`cert:b3` — lcm-diagonal collapse, wave-23, the canonical single-parameter form, `irrational_totient_series_of_lcm_diagonal_certificate_supply`). $(\forall t_0 : \mathbb{N}, \exists t \geq t_0, \exists L, \text{certifiedKill}(\text{periodLcm}(t), \text{periodLcm}(t), L)) \rightarrow \text{Irrational}(S)$. Standing at $N = h = \text{periodLcm}(t)$ beats every hypothetical rational simultaneously (any $t \geq \max(h_0, N_0)$ works for both parameters at once). **This is the canonical restatement with only one quantified unknown t** , and the best target for search/heuristics; the finite floor is `cert:b11` above (28 explicit values, not a cofinal supply).

[Lean] (implication proved; hypothesis [Open]) `scale:cofinal coord:other:lcm-diagonal irrational_totient_series_of_lcm_diagonal_certificate_supply` [LcmDiagonalReduction.lean:112]

Lemma 6.50 (`cert:b4` — lcm-window structure, `eq_prime_pow_of_not_dvd_periodLcm`). For $0 < j < 2t$: if $j \nmid \text{periodLcm}(t)$ then $\exists$ prime p, k with $j = p^k \wedge t < j$ below $2t$, every non-divisor of $\text{lcm}(1..t)$ is a bare prime power exceeding t . Pure elementary number theory about $\text{lcm}(1..t)$; structural input for anyone trying to search for a diagonal certificate narrows where the "noise" in φ on the window comes from.

[Lean] `scale:uniform coord:other:lcm-window eq_prime_pow_of_not_dvd_periodLcm` [CertificateKernel.lean:18613]

Proposition 6.51 (`cert:b5` — lcm-ray window totient factorisation, `totient_periodLcm_ray_split`). On a clean divisor $j \mid \text{periodLcm}(t)$ (every prime factor of j still divides $\text{periodLcm}(t)/j$): $\varphi(q \cdot \text{periodLcm}(t) + j) = \varphi(j) \cdot \varphi(q \cdot (\text{periodLcm}(t)/j) + 1)$. An exact multiplicative split of window totient values into a known local part and a cofactor forced $\equiv 1$ modulo every exhausted prime the exact algebraic handle needed to attempt an unconditional supply proof at `cert:b3`.

[Lean] `scale:uniform coord:other:lcm-window-multiplicative totient_periodLcm_ray_split` [LcmDiagonalReduction.lean:18613]

Theorem 6.52 (`cert:b6` — lcm-cone flatness law, wave-24, `rational_totient_series_forces_lcm_cone_flatness`). $\neg \text{Irrational}(S) \rightarrow \exists t_1, \forall t \geq t_1, \forall q, m : \mathbb{N}, 0 < q \rightarrow \text{totientTail}(q \cdot \text{periodLcm}(t) + m \cdot \text{periodLcm}(t)) - \text{totientTail}(q \cdot \text{periodLcm}(t)) \in \text{range}((\uparrow) : \mathbb{Z} \rightarrow \mathbb{R})$. Rationality forces one fractional constant on the entire lcm cone $\{k \cdot \text{periodLcm}(t) : k \geq 1\}$ at every scale $t \geq t_1$, not just the diagonal pair a strict generalisation of `cert:a9`'s hypothesis-generating side. Any certificate anywhere on the cone kills #249 (`cert:b7`).

[Lean] `scale:uniform coord:other:lcm-cone rational_totient_series_forces_lcm_cone_flatness` [CertificateKernel.lean:18613]

Theorem 6.53 (`cert:b7` — cone collapse, wave-24, annihilator umbrella, `irrational_totient_series_of_lcm_cone_window_kill_supply`). $(\forall t_0 : \mathbb{N}, \exists t \geq t_0, \exists q, m, L : \mathbb{N}, 0 < q \wedge \text{certifiedKill}(m \cdot \text{periodLcm}(t), q \cdot \text{periodLcm}(t), L)) \rightarrow \text{Irrational}(S)$. One certified kill anywhere on the two-multiplier lcm cone, at arbitrarily large t , suffices. Diagonal (`cert:b3`) is the cell $q = m = 1$; q -ray steps are $m = 1$; prime-jump pairs are $(q, m) = (1, p - 1)$. The widest known target still logically equivalent-in-strength to `cert:a10`.

[Lean] (implication proved; hypothesis [Open]) `scale:cofinal coord:other:lcm-cone irrational_totient_series_of_lcm_cone_window_kill_supply` [CertificateKernel.lean:18686]

Corollary 6.54 (`cert:b8` — pure non-integrality frontier forms, no certificate vocabulary). Diagonal: $(\forall t_0, \exists t \geq t_0, \text{totientTail}(2 \cdot \text{periodLcm}(t)) - \text{totientTail}(\text{periodLcm}(t)) \notin \text{range}((\uparrow) : \mathbb{Z} \rightarrow \mathbb{R})) \rightarrow \text{Irrational}(S)$; cone form is the (q, m) generalisation identical in shape. Via `cert:a7`'s iff these hypotheses are exactly equivalent to `cert:b3`/`cert:b7` respectively. **The frontier of #249 stripped of certificate vocabulary:** does $\text{totientTail}(2H_t) - \text{totientTail}(H_t) \notin \mathbb{Z}$ for infinitely many t , $H_t = \text{lcm}(1..t)$?

[Lean] (implication proved; hypothesis [Open]) scale:cofinal coord:other:real-analytic-nonintegrality [irrational_totient_series_of_lcm_diagonal_nonintegrality_supply](#) [CertificateKernel.lean:18706] [irrational_totient_series_of_lcm_cone_nonintegrality_supply](#) [CertificateKernel.lean:18718]

Proposition 6.55 (cert:b9a — second-difference certificates, sound but measured not shallower, `second_diff_notMem_int_of_certifiedRank2Kill`). $\text{certifiedRank2Kill}(h, N, L) \rightarrow (\text{totientTail}(N + 2h) - \text{totientTail}(N + h)) - (\text{totientTail}(N + h) - \text{totientTail}(N)) \notin \text{range}((\uparrow) : \mathbb{Z} \rightarrow \mathbb{R})$. Sound second-difference non-integrality via a doubled band radius. Measured cell $(h, N) = (1, 8)$: rank-1 fires at depth 8, no rank-2 certificate exists at depth ≤ 8 , rank-2 first fires at depth 9 a probe over $t \leq 20$ finds rank-1 at least as shallow in 30/40 cells. This route is empirically not a shortcut over rank-1; flagged do-not-re-attempt without new information.

[Lean] scale:uniform coord:other:binary-window [second_diff_notMem_int_of_certifiedRank2Kill](#) [CertificateKernel.lean:18718] (measured verdict: [totient_tail_rank_two_kill_sound_but_not_shallower_cell](#) [CertificateKernel.lean:18718])

Theorem 6.56 (cert:b10a — cone non-flatness menu refuter, wave-25, sharper than pairwise, `exists_nonintegral_pair_of_coneNonflatCert`). For a nonempty menu Q of positive vertex multipliers with the one-sided floor $\forall q \in Q, q \cdot H + L + 2 < 2^L$ (half the pairwise floor of `certifiedKill`): $\text{coneNonflatCert}(H, L, Q) \rightarrow \exists q_i, q_j \in Q, \text{totientTail}(q_j \cdot H) - \text{totientTail}(q_i \cdot H) \notin \text{range}((\uparrow) : \mathbb{Z} \rightarrow \mathbb{R})$. Proved by an argmin/Helly-avoidance argument over one-sided arcs: if all vertices shared one fractional part, the minimal-deep-tail vertex would be a common left endpoint of every arc, which `coneNonflatCert` denies. Information-theoretically half the depth floor of pairwise `certifiedKill`; the reusable combinatorial-geometry technique transplants to any modular-residue pincer with more than two points.

[Lean] scale:uniform coord:other:lcm-cone-menu [exists_nonintegral_pair_of_coneNonflatCert](#) [CertificateKernel.lean:18718]

Theorem 6.57 (cert:b10b — cone non-flat supply, wave-25, `irrational_totient_series_of_lcm_cone_nonflat_supply`). If `coneNonflatCert` fires (via `cert:b10a`) on a menu whose scale is unbounded, then $\text{Irrational}(S)$. The sharpest known certificate-depth-reduced restatement of the wall; for $|Q| \geq 3$ genuinely joint (menu inconsistent while every pair consistent) this is the best-known target for a search-based attempt at supplying `cert:a10/cert:b3/cert:b7`.

[Lean] (implication proved; supply [Open]) scale:cofinal coord:other:lcm-cone-menu [irrational_totient_series_of_lcm_cone_nonflat_supply](#) [CertificateKernel.lean:18812]

Proposition 6.58 (cert:b12 — survivorKill, alternative certificate via bounded carry orbit). $\text{survivorKill}(h, N, K) := \forall j < 2(N + h + 1) + 1, \exists i \leq K, \text{carryOrbit}(h, N, j - (N + h + 1), i)$ escapes the strip $|\cdot| \leq N + i + h + 2$. Soundness: $\text{survivorKill}(h, N, K) \rightarrow \text{totientTail}(N + h) - \text{totientTail}(N) \notin \text{range}((\uparrow) : \mathbb{Z} \rightarrow \mathbb{R})$ a bounded-orbit argument, alternative to `cert:a4/cert:a6`'s residue-window argument, exhausting all $2(N + h + 1) + 1$ integer box candidates and checking each provably escapes within K steps. `cert:a4` (window) and this (orbit) are two independently complete coordinates for the same non-integrality target (via `cert:a7`'s iff): the "enumerate all integer candidates in a shrinking box, verify each provably diverges" pattern is a general technique for irrationality-by-integer-orbit arguments.

[Lean] scale:uniform coord:other:carry-orbit [survivorKill](#) [CarrySurvivorExtinction.lean:412] [tail_diff_notMem_int_of_survivorKill](#) [CarrySurvivorExtinction.lean:428]

Theorem 6.59 (mob:e1 — first-harmonic norm-gap supply, `irrational_totient_series_of_first_harmonic_norm_gap`). Hypothesis $\text{DTWFirstHarmonicNormGap} := \forall h > 0, \forall X_0, \exists X \geq \max(X_0, 1), \exists L, 16(2X + h + L + 2) \leq 2^L \wedge \|\sum_{N \in [X, 2X]} \text{windowFirstExp}(h, N, L)\| \leq (21/25) \cdot X$. If this holds then $\text{Irrational}(S)$. A genuinely cofinal first-harmonic exponential-sum (Weyl-sum) cancellation statement is sufficient for #249; unconditional companion `exists_certifiedKill_of_first_harmonic_gap` shows any constant-saving first-harmonic gap on one dyadic block forces a finite kill certificate (via $\cos(\pi/8) > 9/10$ and a pigeonhole/averaging argument).

[Lean] (conditional theorem proved; hypothesis [Open]) scale:cofinal coord:binary-digit [irrational_totient_series_of_first_harmonic_norm_gap](#) [FirstHarmonicPivot.lean:83] [exists_certifiedKill_of_first_harmonic_gap](#) [FirstHarmonicGap.lean:128]

Theorem 6.60 (mob:e2 — prime-jump sharp-kill supply, irrational_totient_series_of_primeJumpSharpKill_supply). $\text{primeJumpTailCommutator}(H, p) := D(pH) - p \cdot D(H)$, $D(H) = R_{2H} - R_H$, has an exact partial/tail split with sharp radius $3pH + (p+1)(L+2)$ (tighter than the earlier $4pH$ two-cell-disjunction radius). If $\forall t_0 \exists t \geq t_0 \exists p, L > 0$, $\text{primeJumpSharpKill}(\text{periodLcm}(t), p)$ then $\text{Irrational}(S)$. One concrete deposit is proved unconditionally: $\text{primeJumpSharpKill}(12, 5, 15)$, kernel-decided. Proof strategy: assume S rational $\Rightarrow$ cone flatness (cert:b6) $\Rightarrow$ contradiction via the sharp prime-jump kill.

[Lean] (conditional theorem + [Cert] witness; supply [Open]) scale:cofinal coord:other:lcm-diagonal [irrational_totient_series_of_primeJumpSharpKill_supply](#) [PrimeJumpWindow.lean:193] [primeJumpSharpKill_twelve_five](#) [PrimeJumpWindow.lean:186]

6.3 Consumers

This subsection lists every premise in the corpus whose conclusion is *conditional* on an unsupplied hypothesis, with that hypothesis rendered in display maths so an agent can recognise it as a search target. All targets below are logically downstream of the single exact equivalence that defines the open problem:

Proposition 6.61 (The exact target: #249 = cofinal actual-orbit nonintegrality). *Unconditionally,*

$$\text{Irrational}\left(\sum'_n \varphi(n)/2^n\right) \iff \forall a_0, \exists a \geq a_0, \text{actualLcmTailOrbit } a \notin \text{range}(\mathbb{Z} \rightarrow \mathbb{R}).$$

Nothing beyond a supply of the right-hand side is needed in principle; every other proposition in this subsection is a route to it. scale:n/a [Lean] coord:mobius-mersenne

[irrational_totientSeries_iff_actualLcmOrbitNonintegralitySupply](#) [TotientActualLcmOrbitNonintegralitySupply.lean:2107]

Proposition 6.62 (Short-window arithmetic-kill supply). *If*

$$\forall a_0, \exists a \geq a_0, \exists L < 2 \cdot 2^a, \text{LcmDiagonalArithmeticKill}(2^a, L)$$

holds, then #249 follows (via [actualLcmOrbitNonintegralitySupply_of_shortArithmeticKill](#) then Prop. 6.61). The predicate is exactly the residue-band exclusion of Appendix A4/A6: [lcmDiagonalArithmeticWord](#) at scale 2^a escapes a shrinking central arc mod 2^L . Only two instances are proved (Prop. 6.67's base cases); the cofinal supply is the open trigger of the entire private batch. scale:cofinal [Open] coord:mobius-mersenne

[PowerTwoActualLcmShortArithmeticKillSupply](#) [TotientActualLcmOrbitArithmetic.lean:2107]

Proposition 6.63 (Diophantine separation supply). *If, at canonically-guarded odd ranks q , cofinally many a satisfy*

$$|\text{actualLcmTailOrbit } a - z| > \frac{1}{32} + (\text{explicit error radius}) \quad \forall z \in \mathbb{Z},$$

then #249 follows via the landed signed-margin producer. This restates the target as effective irrationality-measure / anti-concentration for the actual orbit rather than exact residue exclusion, using the explicit approximant of Prop. ???. scale:cofinal [Open] coord:mobius-mersenne

[irrational_totientSeries_of_actualLcmOrbitSeparationSupply](#) [TotientActualLcmOrbitSeparation.lean:308]

Proposition 6.64 (One-sided top-edge residue-gap supply). *If, for the room bound of Prop. 6.84 ($a \geq 8, J + K + (a + 6) < 2 \cdot 2^a$),*

$$\text{ActualLcmTopEdgeResidueGap } a \ J \ K \ m \quad : \iff \quad m \leq K \wedge (\text{residue of windowDiscrepancy at scale } m) \leq$$

holds cofinally, then #249 follows (`actualLcmTailDiff_notMem_int_of_topEdgeResidueGap`, sufficiency at line 1260), because Prop. 6.84/6.85 already exclude the negative-side residue independently only the positive arc needs excluding. This is a strictly weaker target than the old symmetric `certifiedKill` band and is the genuinely easier open target of the whole batch.

`scale:cofinal [Open] coord:mobius-mersenne`
[PowerTwoActualLcmTopEdgeResidueGapSupply](#) [`TotientActualLcmTopEdgeStaircase.lean:900,1260`]

Proposition 6.65 (The five-link equivalence chain below the residue-gap target). *Each of the following cofinal supplies is proved sufficient for Prop. 6.64’s target, chained in decreasing strength:*

`PowerTwoActualLcmTopEdgeResidueGapSupply` $\Leftarrow$ `PowerTwoAdjacentSuffixMidbandSupply`
 $\Leftarrow$ `PowerTwoOddGuardTopEdgeHalfWordBandSupply` $\iff$ `PowerTwoActualFinalTopEdgeMagnitudeSupply`
 $\Leftarrow$ `PowerTwoFlexibleActualTopEdgeMagnitudeSupply` $\Leftarrow$ `PowerTwoFlexibleActualTerminalDominanceSupply`
 $\Leftarrow$ `PowerTwoFlexibleActualTerminalCarryCorridorEscapeSupply`

All six named predicates are open; none is proved. Because the chain is implication-only-downward, proving the single weakest link (`TerminalCarryCorridorEscapeSupply`) closes the entire cluster and hence #249. This is the true minimal remaining target of `TotientActualLcmTopEdgeStaircase`.

`scale:cofinal [Open] coord:mobius-mersenne`
[powerTwoActualLcmTopEdgeResidueGapSupply_of_adjacentSuffixMidband](#) [`TotientActualLcmTopEdgeStaircase.lean:900,1260`]

Proposition 6.66 (The exact closed-form escape identity). *Under integrality of the actual orbit (representative z) and the half-cell fit condition $2(H+q+2) \leq 4^a$ ($a \geq 8$, room bound as above),*

$$2 \cdot \text{actualOddHalfCenteredLift } a \ q = \text{diagonalWindowIncrement}(2^a)(2q+2) - \text{carryOrbit } H \ H \ z \ (2q+1)$$

holds exactly. Escaping either side of this named open interval (the “terminal/carry corridor”) is both necessary and sufficient for non-integrality at rank q (`actualLcmTailOrbit_notMem_int_of_actualTerminalCarryCorridorEscape`). This is the sharpest fully explicit statement of “what remains to prove” anywhere in the batch.

`scale:bounded [Lean] coord:mobius-mersenne`
[two_mul_actualOddHalfCenteredLift_eq_terminal_sub_trueCarry](#) [`TotientActualLcmTopEdgeStaircase.lean:900,1260`]

Proposition 6.67 (Extend the short-kill stub past $a_0 = 6$). *Currently proved only for $a_0 \leq 6$:*

$$\forall a_0 \leq 6, \exists a \ L, a_0 \leq a \wedge L < 2 \cdot 2^a \wedge \text{LcmDiagonalArithmeticKill}(2^a) \ L.$$

The module’s own comment states plainly: “the remaining endpoint gap is now precisely the unbounded continuation beyond this finite prefix.” Supplying a single further instance at $a_0 = 7$ or $a_0 = 8$ already extends the stub and is independently interesting evidence toward Prop. 6.62’s full cofinal supply; the two base cases $a = 4, 6$ trace to `certifiedKill_diagonal_t16/t64`.

`scale:bounded [Lean (finite prefix only)] coord:mobius-mersenne`
[powerTwoActualLcmShortArithmeticKillSupply_through_six](#) [`TotientActualLcmShortKill.lean:54`]

Proposition 6.68 (Fixed-rank extremal-ordering supply). *Define `fixedRankSecondDifference` $H \ j := \varphi(3H+j) - 2\varphi(2H+j) + \varphi(H+j)$. If, at $H = \text{periodLcm}(2^a)$ and some fixed small j ,*

$$\text{MiddleRankTotientExtremal } H \ j \quad : \iff \quad \varphi(2H+j) \text{ is a strict min or max among } \{\varphi(H+j), \varphi(2H+j), \varphi(3H+j)\}$$

then `fixedRankSecondDifference` $H \ j \neq 0$, with sign matching the extremum. Notably an ordering suffices no quantitative gap is required. This is a genuinely different coordinate from the sliding LCM window: j fixed and small, only three fixed ranks examined. The remaining open step is `PowerTwoLcmMiddleRankExtremalSupply`: does this ordering hold cofinally in a .

`scale:uniform [Lean] coord:other:fixed-rank-curvature`
[MiddleRankTotientExtremal](#) [`TotientFixedRankLcmAsymptotic.lean:256,267,275,291,307`]

Proposition 6.69 (The directed/LCM-specialised certificate supply). `directedCertifiedKill` h N L is sound and complete:

$$(\exists L, \text{directedCertifiedKill } h \ N \ L) \iff \text{totientTail}(N+h) - \text{totientTail}(N) \notin \text{range}(\mathbb{Z} \rightarrow \mathbb{R}),$$

exactly, no gap; and $\text{Irrational}(S) \iff \text{CofinalDirectedLcmCertificateSupply}$ (the LCM-diagonal specialisation). The asymmetric strip is a genuine finite-depth improvement over the symmetric certificate (kills $t = 3$ at depth 6, one level earlier than the symmetric one), but the file itself notes the improvement does not turn into an independent sieve theorem: supplying the cofinal predicate is exactly as hard as #249 itself. scale:uniform [Lean] coord:seam-integer
[CofinalDirectedLcmCertificateSupply](#) [TotientTailCarryPeriod.lean:766,829,875,845]

Proposition 6.70 (Pulse-restricted survivor-search supply). Given the cofinal mod-4 pulse supply of Prop. 6.91, it suffices to kill only the 2 mod 4-class candidate states (a fourfold reduction of the initial search strip) at one cofinal arithmetic-pulse prime per putative period, to conclude #249:

$$\text{modFourPulseSurvivorKill} \implies \text{Irrational}(S).$$

The reduction technique (use a cofinal totient-specific pulse to legally restrict the survivor search to one residue class mod a small modulus) transfers wherever an analogous cofinal pulse can be built on the #257 side. scale:cofinal [Lean] coord:mobius-mersenne

[irrational_totientSeries_of_cofinal_modFourPulseSurvivorKill](#) [TotientTailCarryPeriod.lean:906,973,1

Proposition 6.71 (The wave-21 wall: certificate supply over all periods).

$$(\forall h \geq 1, \forall N_0, \exists N \geq N_0, \exists L, \text{certifiedKill } h \ N \ L) \implies \text{Irrational}(S).$$

This is exactly the quantifier structure $\forall h \geq 1 \ \forall N_0 \ \exists N \geq N_0 \ \exists L \ \text{Sep}(h, N, L)$. Nothing in the public or private corpus supplies this hypothesis; every other reduction in Part B of the certificate bank is a logically equivalent-or-weaker reformulation of this same missing supply, never independent progress on it. scale:cofinal [Open] coord:mobius-mersenne

[irrational_totient_series_of_certificate_supply](#) [TotientTailPeriodKiller.lean:394]

Proposition 6.72 (Diagonal collapse one free parameter).

$$(\forall t_0, \exists t \geq t_0, \exists L, \text{certifiedKill}(\text{periodLcmt})(\text{periodLcmt}) \ L) \implies \text{Irrational}(S).$$

Standing at $N = h = \text{periodLcmt}$ beats every hypothetical rational simultaneously (any $t \geq \max(h_0, N_0)$ serves both parameters at once), collapsing Prop. 6.71's two unbounded parameters to one. This is the canonical single-unknown restatement, the best target for search or heuristics. scale:cofinal [Open] coord:mobius-mersenne

[irrational_totient_series_of_lcm_diagonal_certificate_supply](#) [LcmDiagonalReduction.lean:112]

Proposition 6.73 (Cone collapse annihilator umbrella).

$$(\forall t_0, \exists t \geq t_0, \exists q \ m \ L, 0 < q \wedge \text{certifiedKill}(m \cdot \text{periodLcmt})(q \cdot \text{periodLcmt}) \ L) \implies \text{Irrational}(S).$$

One certified kill anywhere on the two-multiplier LCM cone, at arbitrarily large t , suffices; the diagonal (Prop. 6.72) is the cell $q = m = 1$. This is the widest known target still logically equivalent-in-strength to Prop. 6.71. scale:cofinal [Open] coord:mobius-mersenne

[irrational_totient_series_of_lcm_cone_window_kill_supply](#) [CertificateKernel.lean:18686]

Proposition 6.74 (Menu non-flatness supply sharper than pairwise). For a nonempty menu Q of positive vertex multipliers with one-sided floor $\forall q \in Q, qH + L + 2 < 2^L$ (half the pairwise floor of `certifiedKill`),

$$\text{coneNonflatCert } H \ L \ Q \implies \exists q_i, q_j \in Q, \text{totientTail}(q_j H) - \text{totientTail}(q_i H) \notin \text{range}(\mathbb{Z} \rightarrow \mathbb{R}).$$

If this fires at unbounded scale over a menu with $|Q| \geq 3$ genuinely joint (menu inconsistent while every pair is separately consistent), #249 follows via Prop. 6.73. The argmin/Helly-avoidance proof technique is a reusable combinatorial-geometry pattern for any modular-residue pincer with more than two points. scale:cofinal [Open] coord:mobius-mersenne

[irrational_totient_series_of_lcm_cone_nonflat_supply](#) [CertificateKernel.lean:18801,18812]

Proposition 6.75 (Unbounded Farey growth a second independent wall). *The Farey-gap denominator bound at window K is currently $\sim 7.96 \times 10^{34}$ at $K = 240$ (Prop. 6.99). If*

$$\sup_K (b + d)(K) = \infty$$

(the bound growing without limit as $K \rightarrow \infty$), then #249 follows via the C3-style denominator-exclusion consumer, logically independently of Prop. 6.71's certificate-supply wall. No unbound-ness claim is proved or attempted in the corpus. scale:cofinal [Open] coord:other:farey-gap

[gap_check_window_1_240_le_79639646646701375323355774875831053](#) [GapFareyBound.lean:176,225]

Proposition 6.76 (Finite-rank compression a third independent wall). *By Prop. 6.98, the canonical dyadic totient-kernel family is unconditionally $(2^e + 1)$ -dimensional at every level e . Rationality of S forces an associated tempered carry orbit with $\mathbb{Q}$ -rank $\geq 2^e - 1$ at every level (Prop. 6.90). A theorem*

bounding the dyadic-section rank of every rationality-supplied tempered carry

would contradict the unconditional lower bound and close #249 by a linear-algebraic coordinate wholly distinct from the binary-window certificate coordinate of Props. 6.71-6.74. scale:cofinal [Open] coord:other:carry-kernel-rank

[not_irrational_totientSeries_implies_unbounded_carryRank_unconditional](#) [TotientCarryKernelRigidity.lean:100,101]

Proposition 6.77 (Alternative coordinate: bounded carry-orbit survivor supply). *survivorKill h N K enumerates all $2(N + h + 1) + 1$ integer box candidates and checks each provably diverges from the strip $|\cdot| \leq N + i + h + 2$ within K steps; soundness gives $\text{survivorKill } h \ N \ K \implies \text{totientTail}(N+h) - \text{totientTail}(N) \notin \text{range}(\mathbb{Z} \rightarrow \mathbb{R})$, proving the same target as Prop. 6.71/6.61 from a different coordinate (bounded dynamics, not residue windows). Per the doctrine “obstructions are coordinate-relative,” a case hard in the window coordinate may be easy here. scale:n/a [Lean] coord:other:carry-orbit-dynamics*

[tail_diff_notMem_int_of_survivorKill](#) [CarrySurvivorExtinction.lean:412,428]

6.4 Invariants

Necessary conditions that any successful (or unsuccessful) approach must be consistent with facts a solution cannot contradict, whichever way #249 eventually resolves.

Proposition 6.78 (Exact quotient-scale digit closed form, no cleanliness hypothesis).

$\text{lcmRayArithmeticLetter } t \ j = \text{deltaTotient}(\text{periodLcmt})(\text{periodLcmt}+j) = \text{diagonalWindowIncrement}$

valid for both divisor and non-divisor offsets j , with no side condition on which primes of j survive in H/j . Every downstream sign, positivity, or residue statement about the diagonal word must factor through this identity; it generalises the older `deltaTotient_periodLcm_ray_split`, which needed j 's primes to survive. scale:uniform [Lean] coord:mobius-mersenne

[lcmRayArithmeticLetter_eq_deltaTotient](#) [TotientActualLcmOrbitArithmetic.lean:291,298,310]

Proposition 6.79 (Uniform 3/4-density retained on rough integers). *For $a \geq 8$, $t = 2^a$, every prime factor of $n > 0$ exceeding t , and $n < 2^{2 \cdot 2^a}$: $n.\text{primeFactors}.card < 2^a/4$, hence $(3/4) \cdot n <$*

$\varphi(n)$ over $\mathbb{Q}$. Any argument bounding φ on the short window below height $\text{periodLcm}(2^a)^2$ must respect this floor; the counting mechanism (union-bound-in-product form, then convert via `Nat.totient_eq_mul_prod_factors`) is reusable for #257's Mersenne-rough objects at the analogous height. *scale:bounded [Lean] coord:mobius-mersenne*
[three_quarters_mul_lt_totient_of_rough_lt_two_pow](#) [TotientActualLcmOrbitArithmetic.lean:722,768]

Proposition 6.80 (Every short-window digit sign is fixed to +). For $a \geq 8$, $0 < j < 2 \cdot 2^a$: $0 < \text{lcmRayArithmeticLetter}(2^a) j$. Every coefficient of the actual diagonal word in the entire short window is strictly positive divisor offsets via a quantitative $H < 4jc$ bound, foreign prime powers via predecessor descent plus Prop. 6.79's rough-density bracket ($H/4 < c < 5H/2$), exponent-one new primes via $H/4 < c$. This unconditional fact (no irrationality hypothesis) is what makes every sign theorem in the batch Prop. 6.84 and the staircase-impossibility of Prop. 6.107 provable at all. *scale:bounded [Lean] coord:mobius-mersenne*

[lcmRayArithmeticLetter_pos_of_lt_two_mul](#) [TotientActualLcmOrbitArithmetic.lean:1703]

Proposition 6.81 (Window word = truncated discrepancy; kill $\iff$ certificate). $\text{lcmDiagonalArithmeticWord}(\text{windowDiscrepancy}(\text{periodLcm} t)(\text{periodLcm} t) L, \text{and } \text{lcmDiagonalArithmeticKill } t L \iff \text{certifiedKill}(\text{periodLcm} t)(\text{periodLcm} t) L$. Any residue-band existence claim about the actual diagonal word converts directly to a certifiedKill/non-integrality fact through this iff it is the fixed bridge every consumer in this subsection implicitly uses. *scale:n/a [Lean] coord:mobius-mersenne*

[lcmDiagonalArithmeticKill_iff_certifiedKill](#) [TotientActualLcmOrbitArithmetic.lean:1998,2045]

Proposition 6.82 (Exact global-to-local bridge).

$$\text{actualLcmTailOrbit } a = 2^H(2^H - 1) \left(\sum'_n \varphi(n)/2^n \right) - (\text{totientPrefix}(2H) - \text{totientPrefix}(H)), \quad H = \text{periodLcm}(2^a)$$

Every local orbit statement is secretly a statement about the global series value minus a computable finite prefix; any rational-approximation or continued-fraction style separation argument for #249 must express its target through this exact affine image. *scale:n/a [Lean] coord:mobius-mersenne*

[actualLcmTailOrbit_eq_scaled_totientSeries_sub_prefix](#) [TotientActualLcmOrbitSeparation.lean:68]

Proposition 6.83 (Explicit computable approximant with explicit error radius). For all a, q : $|\text{actualLcmTailOrbit } a - \text{actualLcmRawApprox } a q| < (2H + 2q + 3)/2^{2q+1}$, where $\text{actualLcmRawApprox } a q := \text{diagonalAdjacentSuffixRawBlock}(2^a, 0, 2q + 1)/2^{2q+1}$. Any separation-from-integers argument (Prop. 6.63) is forced to work with this finite, computable block rather than the infinite tail, at this exact error rate. *scale:uniform [Lean] coord:mobius-mersenne*

[abs_actualLcmTailOrbit_sub_rawApprox_lt](#) [TotientActualLcmOrbitSeparation.lean:141]

Proposition 6.84 (Unconditional positive-sign corridor a real theorem, not a supply). For $a \geq 8$, $J + (a + 6) < 2 \cdot 2^a$:

$$0 < \text{totientTail}(2H + J) - \text{totientTail}(H + J), \quad H = \text{periodLcm}(2^a),$$

with no irrationality hypothesis the true, infinite, real translated tail difference is strictly positive throughout almost the entire short window, proved unconditionally from Prop. 6.80 plus a directed one-sided tail bound. Any argument about the sign of the actual orbit (not merely its residue mod 2^L) must agree with this; specialised at $J = 0$ this gives $0 < \text{actualLcmTailOrbit } a$. *scale:bounded [Lean] coord:mobius-mersenne*

[actualLcmTailDiff_shift_pos](#) [TotientActualLcmOrbitSign.lean:39,144]

Proposition 6.85 (Integrality forces the exact top-edge residue names the obstruction). Under the room bound of Prop. 6.84 plus $2H + J + K + 2 < 2^K$: integrality of the actual orbit forces

$$\text{windowDiscrepancy } H (H+J) K \bmod 2^K = 2^K - e$$

(the top-edge representative), provably outside the central arc required by `directedCertifiedKill`. This documents precisely why the sign theorem alone cannot close #249: any contradiction needs an independent exclusion of this specific top-edge boundary band, not a re-derivation of the carry reset. The named residue $2^K - e$ ($e > 0$ small) is exactly what Prop. 6.64 sets out to exclude. *scale:bounded [Lean] coord:mobius-mersenne*

[actualLcm_integral_forces_topEdgeResidue](#) [TotientActualLcmOrbitSign.lean:211]

Proposition 6.86 (Punctured staircase is pinned to the half-turn). *Under the room bound and the punctured-staircase hypothesis (all but the last letter vanish at growing dyadic weight; the last letter retained below the top-edge carry band),*

$$\text{lcmRayArithmeticLetter}(2^a)(J+K-1) = 2^{m-1} \quad \text{exactly, and} \quad 2^m < 2(2H + J + K + 2).$$

The modulus must be the first dyadic scale above the room bound; any extra bit of modulus makes even the punctured route empty. Any attempt to revive a partial-staircase strategy must land exactly here. *scale:bounded [Lean] coord:mobius-mersenne*

[puncturedDyadicStaircase_penultimate_eq_half](#) [TotientActualLcmTopEdgeStaircase.lean:1187]

Proposition 6.87 (Any certificate reduces to a two-bit test universal normal form). *Unconditionally, for arbitrary h, N (no totient content in the proof):*

$$(\exists L, \text{certifiedKill } h \ N \ L) \iff \text{GuardCylinderWitness } h \ N,$$

where the witness is a logarithmic-depth socket $(b+1)$ or a two-bit mixed-guard cylinder at scale $b = \log_2(N+h+L+2)+1$. Any certificate-search algorithm or complexity bound for either problem's tail differences must respect this compression an unbounded-depth search is never truly necessary once b is fixed. *scale:uniform [Lean] coord:seam-integer*

[exists_certifiedKill_iff_guardCylinderWitness](#) [TotientActualLcmTopEdgeStaircase.lean:844]

Proposition 6.88 (Primitive kernel factor of the fixed-rank curvature). *On the square-root clean window $j^2 \leq 2^a$, $a \geq 4$: $2\varphi(j) \mid \text{fixedRankSecondDifference}(\text{periodLcm}(2^a)) \ j$. The exact local factor $2\varphi(j)$ comes from the affine-rank-3 kernel $(1, -2, 1)$ combined with the parity of three odd rough cofactors, and is provably tight Prop. 6.108 shows no bare application of the kernel can force one more factor of 2. Any curvature-based attack at fixed rank must land inside this exact divisibility ceiling. *scale:bounded [Lean] coord:other:fixed-rank-curvature**

[two_mul_totient_dvd_fixedRankSecondDifference](#) [TotientFixedRankLcmAsymptotic.lean:519]

Proposition 6.89 (Carry displacement $\iff$ integral tail difference central plumbing). *For a positive-multiplier tempered totient carry u (`IsTemperedBinaryOrbit` $\varphi \ v \ u$, $v > 0$):*

$$(v : \mathbb{Z}) \mid u(N+k) - u(N) \iff \text{totientTail}(N+k) - \text{totientTail}(N) \in \text{range}(\mathbb{Z} \rightarrow \mathbb{R}).$$

Every statement in this batch that converts between “carry orbit divisibility” and “real tail difference is an integer” is an instance of this identity the underlying machinery (`IsTemperedBinaryOrbit`, `binaryCoeffTail`) is generic over any f with $f(n) \leq n$, so it transfers to #257's own coefficient function verbatim. *scale:uniform [Lean] coord:seam-integer*

[carryShift_dvd_iff_tailDiff_mem_int](#) [TotientTailCarryPeriod.lean:77]

Proposition 6.90 (What rationality does and does not buy the exact frontier). *If $\neg \text{Irrational}(S)$, there exist $v > 0$ and a tempered orbit u such that for every level e , the $\mathbb{Q}$ -rank of the canonical carry-kernel-family span is $\geq 2^e - 1$ (torsion-free rank grows exponentially in level), yet u 's dyadic sections are uniformly eventually periodic modulo v (`CarrySectionsEventuallyPeriodicMod`). Rationality buys quotient-mod- v periodicity but does not buy any bound on $\mathbb{Q}$ -rank this is precisely the boundary any Mersenne-specific residue argument (mod-4 pulse, two-adic pulse) exists to cross, and it is the exact hypothesis feeding Prop. 6.76's third independent wall. *scale:uniform [Lean] coord:other:carry-kernel-rank**

[not_irrational_totientSeries_implies_mod_period_and_unbounded_rank](#) [TotientTailCarryPeriod.lean:22]

Proposition 6.91 (Cofinal mod-4 residue pulse genuinely totient-specific). *For any $h > 0$ and bound B , there is a prime $p > B$ with $\text{deltaTotient}(4h) \ p \equiv 2 \pmod{4}$, constructed via Dirichlet on $p \equiv (3r - H) \pmod{4r}$ for a fresh prime $r \equiv 1 \pmod{4}$. Unlike the Mersenne-primitivity killer of Prop. 6.111, this is a genuinely totient-specific residue supplied cofinally beyond every threshold any argument invoking Prop. 6.70 inherits this as its forcing mechanism. The fresh-prime CRT template is scaled to arbitrary depth in Prop. 6.92 below. scale:cofinal [Lean] coord:other:dirichlet-crt*
[exists_prime_deltaTotient_four_mul_mod_four_two](#) [TotientTailCarryPeriod.lean:384,553]

Proposition 6.92 (Arbitrary-depth zero-prefix-then-pulse construction, and its transfer). *For $K \geq 2$, $H > K$, B : cofinally many primes $p > B$ satisfy $p \equiv 1 + 2^{K-1} \pmod{2^K}$, $2^K \mid \varphi(p+H)$, and $2^K \mid \varphi(p-j) \wedge 2^K \mid \varphi(p-j+H)$ for every $1 \leq j < K$ an entire length- $(K-1)$ zero prefix followed by a half-turn terminal pulse, at arbitrary two-adic depth (Prop. 6.91's $K = 2$ case generalised to every K , via $K-1$ fresh Dirichlet primes glued by CRT). This forces*

$$\text{windowDiscrepancy } H \ (p-K) \ K \equiv 2^{K-1} \pmod{2^K},$$

and under eventual integrality transfers to $\exists z, (z : \mathbb{R}) = \text{totientTail}(p+H) - \text{totientTail}(p) \wedge z \equiv 2^{K-1} \pmod{2^K}$. Any solution attempt must be consistent with the existence of this pulse family at every depth K . scale:cofinal [Lean] coord:p-adic
[exists_prime_totient_twoAdic_pulse_divisors](#) [TotientTwoAdicPulseBlock.lean:54,211,262,301,327]

Proposition 6.93 (Exact Möbius-inversion tail identity). $\text{totientTail } N = \sum'_d \mu(d) \cdot 2^{d-r_d(N)} (q_d(N)/(2^d - 1) + 1/(2^d - 1)^2)$, with $r_d(N) = d - N \bmod d$ (forward shift to the next multiple of d), $q_d(N) = \lfloor N/d \rfloor + 1$ exact, not a definition-by-subtraction from a target. Any Möbius/Lambert-series manipulation of totientTail in this batch or in #257 factors through this identity and its fully generic regrouping primitive $\text{tsum_lambert_pair_regroup_if}$ (proved for arbitrary weight functions $|w(d)| \leq d, |v(m)| \leq m$). scale:uniform [Lean] coord:cyclotomic
[totientTail_eq_tsum_shiftedMobiusPulse](#) [TotientShiftedMobiusPulse.lean:535,112]

Proposition 6.94 (The rare unconditional deposits necessary base cases). $\text{LcmDiagonalArithmeticKill}(2^4) \ 2$ and $\text{LcmDiagonalArithmeticKill}(2^6) \ 93$ are proved, routed through pre-existing compressed certificates $\text{certifiedKill_diagonal_t16/t64}$, giving $\text{actualLcmTailOrbit } 4, 6 \notin \text{range}(\mathbb{Z} \rightarrow \mathbb{R})$ the only two unconditional actual-orbit nonintegrality facts currently proved. Any induction or bootstrapping attempt at Prop. 6.62 must reproduce these as base cases, and they double as a computational sanity check that the $\text{lcmRayArithmeticLetter}$ machinery agrees with the older compressed-certificate route. scale:fixed [Cert] coord:mobius-mersenne
[actualLcmTailOrbit_four_and_six_notMem_int](#) [TotientActualLcmShortKill.lean:18,27,35]

Proposition 6.95 (Certificate depth floor). $\text{certifiedKill } h \ N \ L \implies 2(N+h+L+2) < 2^L$. Certificate depth L cannot stay bounded while $N+h \rightarrow \infty$: L must grow at least logarithmically. Any complexity argument about certificate search inherits this floor, and it is exactly what Prop. 6.87 compresses to a two-bit test at the corresponding logarithmic scale. scale:n/a [Lean] coord:seam-integer
[certifiedKill_depth_floor](#) [TotientTailPeriodKiller.lean:79]

Proposition 6.96 (The tail-period law necessary consequence of rationality). $\neg \text{Irrational}(S) \implies \exists h : \mathbb{N}, 0 < h \wedge \exists N_0, \forall N \geq N_0, \text{totientTail}(N+h) - \text{totientTail}(N) \in \text{range}(\mathbb{Z} \rightarrow \mathbb{R})$, with explicit witnesses $h = \varphi(\text{oddPart}(r.\text{den}))$, $N_0 = v_2(r.\text{den})$ for $S = r$. This is the “only if” half of the master equivalence: any rationality-refutation strategy must eventually contradict some period h and preperiod N_0 produced this way. Composes with Prop. 6.81/soundness to give Prop. 6.71. scale:uniform [Lean] coord:other:euler-theorem
[eventual_period_of_not_irrational](#) [TotientTailPeriodKiller.lean:358]

Proposition 6.97 (Rationality flattens the whole LCM cone). $\neg \text{Irrational}(S) \implies \exists t_1, \forall t \geq t_1, \forall q m : \mathbb{N}, 0 < q, \text{totientTail}(q \cdot \text{periodLcm } t + m \cdot \text{periodLcm } t) - \text{totientTail}(q \cdot \text{periodLcm } t) \in \text{range}(\mathbb{Z} \rightarrow \mathbb{R})$. Rationality forces ONE fractional constant on the entire cone $\{k \cdot \text{periodLcm } t : k \geq 1\}$ at every scale $t \geq t_1$, not just the diagonal pair a strict generalisation of Prop. 6.96. Any certificate anywhere on the cone therefore kills #249 (Prop. 6.73). scale:n/a [Lean] coord:mobius-mersenne

[rational_totient_series_forces_lcm_cone_flatness](#) [CertificateKernel.lean:18674]

Proposition 6.98 (Unconditional (2^e+1) -dimensional dyadic kernel the rank floor). The canonical dyadic-kernel family `canonicalTotientKernelFamily e` has exactly $2^e + 1$ channels, unconditionally linearly independent for every e (CRT + Dirichlet: one channel made prime, every other channel gets a fresh prime $\equiv 1 \pmod a$ large power of 2). This is a proved fact, needing no rationality hypothesis at all, and it is the fixed lower bound that Prop. 6.76 would have to contradict. scale:n/a [Lean] coord:other:carry-kernel-rank

[card_totientCanonicalIndex](#) [TotientMahlerDefect.lean:51]

Proposition 6.99 (Sharp Farey rung concrete necessary floor at $K = 240$). For all $q : \mathbb{N}$, $0 < q \leq 79639646646701375323355774875831053$ ($\sim 7.96 \times 10^{34}$): $(qV) \bmod 2^{240} + 243q < 2^{240}$, where V is the explicit committed totient residue for window $(N, K) = (1, 240)$; this is SHARP $q = 79639646646701375323355774875831054$ is the exact first failing denominator (an explicit Farey mediant). Any candidate rational value for S must have reduced denominator strictly larger than this bound (Prop. 6.100 below). scale:fixed [Cert] coord:other:farey-gap

[gap_check_window_1_240_first_failure](#) [GapFareyBound.lean:176,225]

Proposition 6.100 (Denominator exclusion headline unconditional consequence). $\forall p : \mathbb{Q}, p.\text{den} \leq 79639646646701375323355774875831053 \implies S \neq (p : \mathbb{R})$. If S is rational, its reduced denominator exceeds $\sim 7.96 \times 10^{34}$ an unconditional necessary condition on any hypothetical rational value, logically independent of the certificate-supply wall (Prop. 6.71); it excludes small denominators outright, without needing a period at all. scale:fixed [Lean] coord:other:farey-gap

[tsum_totient_div_pow_two_ne_ratCast_of_den_le_79639646646701375323355774875831053](#) [CertificateKernel.lean:18674]

Proposition 6.101 (The two foundational irrationality engines). Two, and only two, general-purpose irrationality criteria exist in the kernel, and any successful proof of #249 must ultimately instantiate one of them: (i) if $u : \mathbb{N} \rightarrow \mathbb{Q}$ is eventually never x and $\text{den}(u_k) \cdot |x - u_k| \rightarrow 0$, then $\text{Irrational}(x)$; (ii) if $\forall q > 0, \exists m, z : \mathbb{Z}, 0 < |m\xi - z| < 1/q$, then $\text{Irrational}(\xi)$ (with a base-power specialisation $b^n\xi$ matching any digit/carry construction in base b). Every certificate, cone, or Farey argument in this catalogue exists either to supply one of these two hypotheses or to substitute for them via the `certifiedKill` route. scale:n/a [Lean] coord:other:dirichlet-approximation

[irrational_of_den_mul_abs_sub_tendsto_zero](#) [CertificateKernel.lean:5371,5792,5821]

Proposition 6.102 (S sits on the Mersenne-Lambert ladder). Writing $L(f) := \sum_{n \geq 1} f(n)/(2^n - 1)$: $L(\mu) = 1/2$, $L(\varphi) = 2$ (exactly rational, machine-checked), $L(1) = E$ = the Erdős-Borwein constant, proved **irrational** in this same kernel; $L(A) = S$ for $A = \varphi * \mu$ is exactly #249 restated in “positive Erdős-Borwein form,” still open; $L(\text{Id}) = \sum \sigma(m)/2^m$ is transcendental by Nesterenko 1996 (cited, not formalised). Any proof of #249 sits on this Dirichlet-convolution ladder next to a proved irrational neighbour ($L(1)$) and a proved rational neighbour ($L(\varphi)$); this is the clearest bridge into #257’s native Lambert-series coordinate. scale:n/a [Lean, Cited] coord:mobius-mersenne

[tsum_totient_div_pow_two_eq_pnat_half_pow](#) [CertificateKernel.lean:18063-18087]

Proposition 6.103 (Generic rational-gap adapter). For $\text{pfx} < \text{whole} : \mathbb{Q} : 1/(\text{whole.den} \cdot \text{pfx.den}) \leq \text{whole} - \text{pfx}$ (as reals). Any analytic upper bound on a totient-tail gap converts, via this lemma, into a denominator-growth lower bound extending Prop. 6.99’s Farey rungs without redoing the Farey-neighbour search from scratch a clean, fully general adapter usable on either

problem's tail estimate. scale:n/a [Lean] coord:other:farey-gap

[positive_rational_difference_lower_bound](#) [PrimitiveRationalGapSupply.lean:31]

Proposition 6.104 (Finite pincer floor concrete necessary evidence through $t = 64$). `certifiedKill_diagonal`

Prop. 6.72's predicate Pt holds for $t \in \{1, 2, 3, 4, 5, 7, 8, 9, 11, 13, 16, 17, \dots\}$ (28 explicit cases through $t = 64$), each a finite `decide/norm_num` computation on explicit φ values via checked prime-power blocks with Lucas-primality certificates. This does not establish Pt for infinitely many t and hence does not close Props. 6.71/6.72/6.73 it is exactly the finite floor any unbounded-supply argument must extend. scale:fixed [Cert] coord:mobius-mersenne

[certifiedKill_diagonal_all_imported](#) [DiagonalPincerCertificates.lean:2842-2870]

Proposition 6.105 (The fair-coin coprimality reformulation of S). $\#\{(a, b) : a+b = n, a > 0, \gcd(a, b) = 1\} = \varphi(n)$ for every n (uniform, $\varphi(0) = 0$, $\varphi(1) = 1$ included automatically). At $r = 1/2$, for independent fair-coin waiting times X, Y ($P(X=n) = 2^{-n}$, $n \geq 1$): $P(\gcd(X, Y) = 1) = S - 1/2$. #249's constant is, up to the additive constant $1/2$, a literal lattice-point probability. Any digit argument for #249 necessarily also carries a positional meaning about visible lattice points on the addition antidiagonal a reformulation any proof attempt should keep in view even though it is not itself a route to a proof. scale:n/a [Lean] coord:other:visible-lattice-points

[tsum_pos_coprime_pair_pow](#) [GeometricCoprimality.lean:120,182,56]

6.5 Killers and countermodels

Adversarial constructions and no-go results, kept with the mechanism that closes each route so the machinery outlives the framing. Per the corpus's own doctrine (project memory `feedback_invent_dont_audit`): obstructions below are coordinate-relative, and each entry names exactly which coordinate or proof-template it excludes, not the underlying object.

Observation 6.106 (Nonnegative-branch elimination for the true survivor). If the translated actual orbit is integral at the start of Prop. 6.84's positive corridor, then for every later depth K with room $J+K+(a+6) < 2 \cdot 2^a$, the negated carry-orbit trajectory `-carryOrbit H (H+J) d K` is negative and is an `endpointSurvivor` at every depth in the corridor. This eliminates only the nonnegative branch of the true survivor's sign it is emphatically *not* a nonintegrality claim; spurious survivors of either sign may still exist. Any strategy hoping to conclude non-integrality purely from this sign fact fails: Prop. 6.85 names exactly what remains. scale:bounded [Lean] coord:mobius-mersenne

[actualLcm_trueEndpointSurvivor_neg](#) [TotientActualLcmOrbitSign.lean:172]

Observation 6.107 (Total dyadic staircase is impossible route-pruning). For $a \geq 8$, room bound, and modulus wide enough ($2H+J+K+2 < 2^m$): `ActualLcmTerminalDyadicStaircase a J K m` is **false** a terminal window where every one of the last m letters is divisible by its own growing power of two ($2^{r+1} \mid \text{letter}_r$) cannot occur. The last letter would have to be positive (Prop. 6.80), strictly below the wide modulus, and divisible by it, forcing it to be exactly 0 contradicting positivity. **Kills outright:** any strategy aiming for total dyadic annihilation of the terminal suffix. The generic mechanism ($0 < e < 2^m \wedge 2^m \mid e \implies e = 0$) is reusable wherever a positive quantity is asked to vanish mod a wider-than-itself modulus. The surviving route is the *punctured* staircase (Prop. 6.86) or the residue-gap producers (Prop. 6.64). scale:bounded [Lean] coord:mobius-mersenne

[not_actualLcmTerminalDyadicStaircase_of_room](#) [TotientActualLcmTopEdgeStaircase.lean:251]

Observation 6.108 (The bare 3-rank kernel is exactly tight a sharpness no-go). `fixedRankSecondDifference(2^n)` $2(2^{n+1} \cdot 3) = -2^{n+1}$ exactly, for all n (a closed-form computed fixture). The two-adic valuation gained by Prop. 6.88 is exactly tight the normalised 1×1 minor is odd at every depth, so the bounded-height primitive kernel alone can **never** force one additional factor

of 2. **Kills:** any attempt to squeeze more 2-adic information out of the bare 3-rank curvature kernel; further progress needs new arithmetic input, matching the file’s own open target `PowerTwoLcmMiddleRankExtremalSupply`. `scale:uniform [Cert] coord:other:fixed-rank-curvature`
[fixedRankSecondDifference_dyadic_fixture_exact](#) [TotientFixedRankLcmAsymptotic.lean:462]

Observation 6.109 (The parity coboundary countermodel object). `parityCoboundaryWeight n := parityBaseWeight n + 2 · largePowerTwoBit n - 4 · largePowerTwoBit (n - 1)`, where `parityBaseWeight` is the eventually-constant word $0, 1, 1, 2, 4, 4, 4, \dots$ and `largePowerTwoBit n = 1` iff $n = 2^{k+3}$ (lacunary spikes starting at 8). This is a hand-built adversary sequence containing *no* φ in its definition at all except via its parity. The construction technique `add a zero-valued sparse binary coboundary  $2 \cdot 2^{-m} - 4 \cdot 2^{-(m+1)} = 0$  at lacunary ranks to an eventually-periodic base word, destroying periodicity while preserving the rational sum` is a completely general recipe, directly portable to an analogous #257 adversary matched to $1/(2^n - 1)$ -parity patterns. `scale:n/a [Lean] coord:other:binary-digit`

[parityCoboundaryWeight](#) [TotientParityCoboundaryCountermodel.lean:37,41,46,59]

Observation 6.110 (What the countermodel refutes the flagship kill of the batch). There is $c: \mathbb{N} \rightarrow \mathbb{N}$ with: (a) arbitrarily many, arbitrarily-separated, cofinal explicit “6,0” carry pairs (for every N, G, K a block of K such pairs beyond N , pairwise separated by $> G$); (b) $\forall n, c(n) \leq 6$; (c) $\forall n, c(n) \leq n$; (d) $\forall n, c(n) \equiv \varphi(n) \pmod{2}$ (exact parity agreement with Euler’s totient); (e) c is **not** eventually periodic; yet (f) $\sum'_n c(n)/2^n = 3/2$, a rational number. **Kills:** any #249 proof strategy whose only inputs are uniform boundedness, the trivial growth bound $c(n) \leq n$, exact φ -parity agreement, and failure of eventual periodicity *even strengthened* to cofinal, arbitrarily-separated, arbitrarily-long-block non-periodicity. The obstruction is structural, not a shortage of non-periodicity witnesses: a rational coboundary can carry unlimited *visible* aperiodic structure while summing to a fixed rational. **Doctrine consequence** (matches project memory `feedback_erdos_reductions_rejected_bank_real_results`): only arguments using actual quantitative totient/Mersenne size or residue information (as in the `TotientActualLcm*/TotientFixedRank*` families) can possibly close #249; pure symbolic-word arguments cannot. This countermodel is also the ready-made stress-test fixture for any future sufficient-condition candidate stated purely in coefficient-word terms (boundedness/growth/parity/periodicity), on either #249 or #257. `scale:cofinal [Cert] coord:other:binary-digit`

[exists_totientParity_arbitrarilyManySeparatedCarry_rational_countermodel](#) [TotientParityCoboundaryCountermodel.lean:342,320]

Observation 6.111 (Primitive Mersenne-prime factors alone force nothing). For the completely-multiplicative control $c(n) = n$ (zero totient content: `binaryCoeffTail id N = N + 2`): if $q > K > 0$ and $q \mid 2^K - 1$ (e.g. q a primitive prime factor of the homogeneous Mersenne multiplier), the shift is integral at every N (value exactly K) but $q \nmid K$. **Kills:** “a large primitive Mersenne prime factor alone forces a contradiction” as a proof strategy this is a route-pruning warning applicable *verbatim* to #257, whose denominators $2^n - 1$ are literally the object $q \mid 2^K - 1$ tested here. Any successful use of such a factor needs genuinely totient-specific residue/size input beyond bare Mersenne primitivity contrast with the genuinely totient-specific pulse of Prop. 6.91. `scale:bounded [Lean] coord:mobius-mersenne`

[idCoeff_mersenneModulus_does_not_annihilate_tailShift](#) [TotientTailCarryPeriod.lean:342,320]

Observation 6.112 (Universal no-go for absolute-adjugate coefficient reconstruction). For any finite rational row $w: \iota \rightarrow \mathbb{Q}$, integer targets $x: \iota \rightarrow \mathbb{N}$ that *exactly* isolate one totient channel ($\sum_i w_i \varphi(x_i) = 1$): the crude two-tail cost $\sum_i |w_i| \cdot (2(x_i+1) + (x_i+2))$ is ≥ 3 , hence never < 1 . **Kills:** the strategy of recovering $\varphi(x)$ from $2R_{x-1} - R_x$ and bounding each tail termwise by $R_M \leq M+2$ it can *never* reach the strict < 1 tail-error threshold, at any finite grid height, independent of matrix height, row translation, determinant, or target channel. The proof only uses $\varphi(x) \leq x$, so the same triangle-inequality floor transfers *verbatim* to any #257 reformulation with $c(n) \leq n$. `scale:uniform [Lean] coord:other:adjugate-linear-algebra`

[three_le_totientAdjugateTailCost](#) [TotientTailCarryPeriod.lean:266,307]

Observation 6.113 (Rank-2 second-difference certificates: sound but measured not shallower). $\text{certifiedRank2Kill } h \ N \ L \implies$ the second difference $(\text{totientTail}(N+2h) - \text{totientTail}(N+h)) - (\text{totientTail}(N+h) - \text{totientTail } N) \notin \text{range}(\mathbb{Z} \rightarrow \mathbb{R})$, sound at doubled band radius versus rank 1. Measured at $(h, N) = (1, 8)$: rank-1 fires at depth 8, no rank-2 certificate exists at depth ≤ 8 , rank-2 first fires at depth 9 empirically **not** a shortcut over rank-1 (rank-1 at least as shallow in 30/40 probed cells for $t \leq 20$). **Kills**: expecting a depth improvement from moving to second differences. Explicitly flagged as a dead end do not re-attempt this exact refinement without new information. scale:n/a [Cert] coord:mobius-mersenne
[totient_tail_rank_two_kill_sound_but_not_shallower_cell](#) [CertificateKernel.lean:18729,18762]

Observation 6.114 (Unit-gap strengthening rescues at most one lattice point). $\text{ReducedDenominatorUnitGapCert} \forall t \in [1, L], \neg \text{Coprime}(J+t, u)$ (nonunits-only refinement of the Farey-gap consumer). At a prime-power reduced denominator p^e : $\text{ReducedDenominatorUnitGapCert}(p^e) \ N \ K \iff L=0 \vee (L=1 \wedge p \mid (J+1))$ the unit-gap strengthening can rescue **at most one** additional candidate lattice point beyond the ordinary Farey-gap certificate. **Kills**: expecting this refinement to unboundedly strengthen Prop. 6.99/6.100's Farey rungs it is self-flagged in its own docstring as “a strict certificate-level strengthening, not a supply theorem.” A known dead end; do not re-attempt expecting more than +1 lattice point per prime power. scale:bounded [Lean] coord:other:farey-gap
[reducedDenominatorCandidateCount_prime_pow_le_one](#) [PrimitiveWeightCertificate.lean:30,54,111]

Observation 6.115 (Synthetic all-horizon countermodel to homogeneous-factor-only proof strategies). There is an explicit nonzero “all-horizon countermodel” sequence built from multiples of $\varphi(\text{periodLcm } t)$ that (i) agrees with every exact whole-ray anchor $\text{deltaTotient } H \ (qH) = \varphi(H)$ for $2 \leq q < t$; (ii) stays inside the natural diagonal bounds; and (iii) **survives every finite integer shift polynomial** every commensurate finite-rank LCM cube, via the normal form $P_m(E_H) \cdot \prod (E_H^n - 1)$. Explicitly flagged synthetic: it does not claim the compensation letters occur as actual totient differences. **Kills**: the strategy of “retain only homogeneous LCM-ray factors” at *every* finite rank, not just rank 2 or 3. This is a representation-level exclusion (a particular factor-ideal projection throws away information that can be adversarially reconstructed), not an exclusion of φ itself an actual proof must control the fresh Möbius channel, per [totient_eq_sum_mobiusTotientChannel](#) in the same file. scale:n/a [Lean (docstring-sourced)] coord:mobius-mersenne
[LcmFactorIdealPulseObstruction](#) [LcmFactorIdealPulseObstruction.lean:1-24,328]

Observation 6.116 (Fixed-precision local valuation-unit signatures never obstruct). For any finite word of odd-valuation-unit symbols at fixed local 2-adic precision $u > 0$, and any starting carry state e , there **exists** a compatible carry orbit realising that exact word with every intermediate state centred in its dyadic interval ($|e'| \leq \text{vuRadius } u \ \sigma$). **Kills**: any proof strategy trying to derive a contradiction purely from “the local valuation-unit signature at fixed precision u is incompatible with X ” such signatures are *always* realisable by some carry orbit. Framed against #249 in the docstring, but the theorem itself carries no totient- or Mersenne-specific content: a proof needs growing precision or extra arithmetic coupling, not a fixed-window local signature. scale:n/a [Lean] coord:p-adic
[fixedPrecisionTropicalNoGo](#) [TropicalCurvatureCarry.lean:137,69,115]

Observation 6.117 (Square-CRT correction-suppression is independent of nonvanishing). Square-CRT correction suppression (fixing a cofactor mod a prime square to remove a weighted correction term on a finite horizon) is achievable, **but** suppression alone does not force a nonzero coefficient: the smallest returned countermodel has the whole two-step finite block vanish identically, while a separate clean witness shows it can also be nonzero “clean” (correction-suppressed) is consistent with *both* vanishing and nonvanishing. **Kills**: “achieve a clean square-CRT horizon” as a sufficient condition by itself for a nonvanishing correction term; an additional anti-concentration or residue producer is still needed, exactly as the module's own docstring states.

scale:n/a [Lean (docstring-sourced)] coord:other:square-crt-cocycle

[squareCRT_clean_block_can_vanish](#) [SquareCRTCube.lean:285,294,303,311]

Observation 6.118 (No fixed integer clears every normalised primitive-Euler coordinate). $\forall D > 0, \neg \forall n > 0, \exists z \in \mathbb{Z}, D \cdot (A(n)/n) = z$, where $A = \varphi * \mu$ is the Mersenne-Lambert primitive weight (so $S = L(A)$, Prop. 6.102). Any D clearing normalised coordinates through horizon N must be divisible by every odd prime $p \leq N$, every $p^2 \leq N$, and by 4 once $N \geq 4$ so no fixed D works for all n . **Kills:** any strategy seeking a single fixed integer denominator that simultaneously integralises every primitive-Euler coordinate $A(n)/n$ a finite no-lift theorem in the integral Euler/Witt-coordinate category (weaker than the general Dieudonné-Dwork theorem, and explicitly *not* itself an irrationality proof). An independent-coordinate no-go, orthogonal to the binary-window (Prop. 6.71) and carry-rank (Prop. 6.76) obstructions a third distinct coordinate attacking the same open problem. scale:n/a [Lean] coord:mobius-mersenne

[no_fixed_integral_primitive_euler_index](#) [PrimitiveDeterminantLift.lean:169,148]

Observation 6.119 (Finite certificates only ever prove exclusion, never membership). Certified-death/kill families (finite decidable checks at a given depth) are explicitly **one-sided**: a found certificate proves exclusion, but failure to find one, or survival through any finite probed depth, proves *nothing* about membership or rationality. This is a structural caveat recurring across every certificate family in this catalogue (`certifiedKill`, `directedCertifiedKill`, `survivorKill`, `LcmDiagonalArithmeticKill`): none of them can ever certify membership/rationality by finite search, only non-membership/non-integrality. **Kills:** treating an unsuccessful finite search over any of the certificate families above as evidence toward rationality, or treating a finite prefix of confirmed kills (Prop. 6.94, Prop. 6.104) as anything more than a floor to extend. scale:n/a [Lean (structural caveat, cross-problem)] coord:n/a

[CertifiedGreedyMersenneDeath](#) [GreedyAchievementSet.lean:1720,1742-1744]

6.6 Recently published declarations

This subsection catalogues material that entered the public record after the rest of this paper was drafted: declarations union-merged into modules that were already public (`SignedQMomentObstruction`, `GenericTailOrbitRigidity`, `DiagonalFreshLossBridge`, `SquaredMersenneDiagonalEnclosure`, `ActualForeignResidueProjection`, `DyadicPrefixCompression`, `RepunitMobiusNumerator`, `GeometricCoprimality`, `LcmDiagonalReduction`, `PivotAntiReconstruction`, all in `Erdos249257/`), and the newly published per-problem tree `ErdosProblems/Erdos249/`. Every declaration below was read directly from source; none is inferred from a name. Several of these files contain long chains of “`XSply` $\rightarrow$ irrational” implications whose hypothesis is an unproved cofinal predicate. Per the standing rule of this paper, those chains are named as exactly what they are — reductions, not results — and are not restated as if they narrowed the open problem. The module `SquareCRTCube` is deliberately excluded: its private additions did not compile even in the private tree and were reverted, so nothing from it beyond the existing public file is cited here.

6.6.1 The Möbius–Mersenne ladder: unconditional log-concavity

`SignedQMomentObstruction` adds an infinite ladder $\Theta_r := \sum_{n \geq 0} \mu(n+1)/(2^{n+1}-1)^r$ built from the same Möbius–Mersenne atoms as the rest of the corpus, together with a full unconditional order-two Hankel (log-concavity) theorem for every rung.

Definition 6.120. $\Theta_r := \sum_{n \geq 0} \mu(n+1)/(2^{n+1}-1)^r$, split exactly into the first two atoms $\text{TwoAtom}(r) := 1-3^{-r}$ and the tail $\text{TailAfterTwo}(r) := \sum_{n \geq 0} \mu(n+3)/(2^{n+3}-1)^r$. [Erdos249257.SignedQMomentObstruction.mobiusMersenneTwoAtom](#) [SignedQMomentObstruction.lean:140] [Erdos249257.SignedQMomentObstruction.mobiusMersenneTailAfterTwo](#) [SignedQMomentObstruction.lean:144] [Lean] scale:uniform coord:mobius-mersenne.

Theorem 6.121 (Two low rungs are exact rationals tied to #249). $\Theta_1 = 1/2$, and $\Theta_2 = (\sum_{n \geq 1} \varphi(n)2^{-n}) - 1/2$ — the second rung is literally $S - 1/2$. [Erdos249257.SignedQMomentObstruction.mobiusMersenneTheta_two_eq_totient_offset](#) [SignedQMomentObstruction.lean:197] [Lean] scale:fixed coord:mobius-mersenne.

Theorem 6.122 (Two-atom exact Hankel gap). $\text{TwoAtom}(r+1)^2 - \text{TwoAtom}(r)\text{TwoAtom}(r+2) = 4/3^{r+2}$, hence the two-atom truncation alone is strictly log-concave at every r . [Erdos249257.SignedQMomentObstruction.hankelGap](#) [SignedQMomentObstruction.lean:311] [Lean] scale:uniform coord:mobius-mersenne.

Theorem 6.123 (Full ladder: unconditional strict log-concavity, every rung). For every $r \geq 1$, $\Theta_r \Theta_{r+2} < \Theta_{r+1}^2$: the shifted 2×2 Hankel determinant of the whole ladder is negative at every rung, not just asymptotically. The proof propagates the exact two-atom gap above against a geometric tail-error budget that contracts by a factor $1/4$ per rung against a gap that only contracts by $1/3$, giving an inductive floor from rung 5 on (kernel-checked base cases below). [Erdos249257.SignedQMomentObstruction.mobiusMersenneTheta_strict_logConcave](#) [SignedQMomentObstruction.lean:534] [Erdos249257.SignedQMomentObstruction.mobiusMersenneTheta_strict_logConcave_of_five_le](#) [SignedQMomentObstruction.lean:534] [Erdos249257.SignedQMomentObstruction.mobiusMersenneTheta_hankel_two_neg](#) [SignedQMomentObstruction.lean:740] [Lean] scale:uniform coord:mobius-mersenne.

Observation 6.124. This is a genuine unconditional analytic fact about the ladder, but it is not by itself an irrationality obstruction or a step toward one: negative Hankel determinants say the sequence (Θ_r) is not the moment sequence of a positive measure, which is unrelated to whether any single Θ_r (in particular $\Theta_2 = S - 1/2$) is rational. It is recorded here as exactly what it is — a structural fact about the ladder — and no stronger claim is made.

The file also supplies the underlying algebraic infrastructure: a rectangular Cauchy–Binet expansion derived from the Leibniz formula ([Erdos249257.SignedQMomentObstruction.det_mul_rectangular](#) [SignedQMomentObstruction.lean:29], [Lean], scale:uniform, coord:other:hankel-determinant, needed because Mathlib’s pinned determinant API does not expose rectangular Cauchy–Binet directly), and a finite unique-terminal-dyadic-exponent parity lemma showing that clearing a common denominator by its uniquely largest power of two preserves oddness of the numerator ([Erdos249257.SignedQMomentObstruction.scaled_dyadic_sum_odd](#) [SignedQMomentObstruction.lean:78], [Lean], scale:uniform, coord:other:dyadic-parity).

6.6.2 Generic tail-orbit rigidity: a self-labelled non-claim

`GenericTailOrbitRigidity` is explicit in its own header that it asserts no novelty and no priority, and contains an explicit `NON_CLAIM` guard against a superseded “positive orbit” route; it is a formal algebra/analysis interface, reproduced here only because it entered the public tree after the earlier parts of this paper were drafted.

Theorem 6.125 (T7: tempered-orbit equivalence, abstract binary series). For any $c : \mathbb{N} \rightarrow \mathbb{N}$ with $c(n) \leq n$, writing $X_c := \sum_{n \geq 1} c(n)/2^n$, X_c is rational iff there exist $v > 0$ and an integer sequence u with $u(N+1) = 2u(N) - v c(N+1)$ and $u(N)/2^N \rightarrow 0$; when it exists such an orbit is rigid, $u(N) = v \cdot T_c(N)$ for every N , where T_c is the scaled tail. [Erdos249257.binaryCoeffSeries_rational_iff_exists_temperedBinaryOrbit](#) [GenericTailOrbitRigidity.lean:426] [Erdos249257.temperedBinaryOrbit_eq_scaledTail](#) [GenericTailOrbitRigidity.lean:339] [Lean] scale:uniform coord:other:tail-orbit-rigidity.

Proposition 6.126 (Rigidity engine). A real sequence d with $d(N+1) = 2d(N)$ and $d(N) = o(2^N)$ is identically zero. [Erdos249257.doublingOrbit_eq_zero_of_tempered](#) [GenericTailOrbitRigidity.lean:339] [Lean] scale:uniform coord:other:tail-orbit-rigidity.

Observation 6.127 (Finite-state no-go for successor decoders). For fixed $m \geq 2$, the “balanced-pulse” family $c_{m,r}$ (mass r moved from digit m to digit $m+1$, $0 \leq r \leq \lfloor (m+1)/2 \rfloor$) has the same value and the same complete pre- m history for every r , yet the parameter r is recovered exactly from the first post-pulse digit. Consequently no state that identifies all members of one such family can be decoded by any autonomous map, and the fan-out of any correct decoder is unbounded in m . [Erdos249257.balancedPulse_no_autonomous_decoder](#) [GenericTailOrbitRigidity.lean:247] [Erdos249257.balancedPulse_label_card_lower_bound](#) [GenericTailOrbitRigidity.lean:228] [Lean] scale:uniform coord:other:tail-orbit-rigidity. This is a genuine barrier of the class the brief asks for: it rules out exactly the class of arguments that try to predict/decode the exact tail orbit from a state depending only on the pre-pulse history, because the family exhibited is a literal counterexample generator for any such decoder. It says nothing about the totient-specific series itself.

Two “rational control models” are included as honesty checks, not irrationality statements: the pair-balanced family $c(2k)=k-a(k)$, $c(2k+1)=2a(k)$ has tempered orbit for every bounded payload a and a fixed marked value $4/9$ regardless of a ([Erdos249257.globalBalancedCoeff_value](#) [GenericTailOrbitRigidity.lean:545], [Lean], scale:uniform, coord:other:tail-orbit-rigidity); and the identity coefficients $c(n) = n$ have the explicit multiplier-one tempered orbit $u(N) = N + 2$ ([Erdos249257.idCoeff_temperedOrbit](#) [GenericTailOrbitRigidity.lean:564], [Lean], scale:fixed, coord:other:tail-orbit-rigidity).

6.6.3 Squared-Mersenne diagonal enclosure

`SquaredMersenneDiagonalEnclosure` spends the exactly-summable first-order Lambert term of the actual #249 diagonal exactly, leaving only a squared-Mersenne tail to bound.

Proposition 6.128 (Exact rational centre, direct). `scaleDiagonalTailDifference(H)–lambertProjectedDiagonalCH·mobiusSquareTail(D)` for every H, D — no residue split and no side condition on D versus H . [Erdos249257.SquaredMersenneDiagonalEnclosure.scaleDiagonalTailDifference_sub_lambertProjectedDiagonal](#) [SquaredMersenneDiagonalEnclosure.lean:138] [Lean] scale:uniform coord:other:squared-mersenne-tail.

Proposition 6.129 (Sharp geometric tail bound). $|\text{mobiusSquareTail}(D)| \leq 4/(3(2^{D+1} - 1)^2)$. [Erdos249257.SquaredMersenneDiagonalEnclosure.abs_mobiusSquareTail_le](#) [SquaredMersenneDiagonalEnclosure.lean:145] [Lean] scale:uniform coord:other:squared-mersenne-tail.

Observation 6.130 (Directed vs. symmetric enclosure). When the first nonzero Möbius channel past D is known, its sign directs a one-sided interval half the width of the symmetric bound ([Erdos249257.SquaredMersenneDiagonalEnclosure.mobiusSquareTail_directed_of_first_support](#) [SquaredMersenneDiagonalEnclosure.lean:313], [Lean], scale:uniform, coord:other:squared-mersenne-tail), and either enclosure composes with an integer-lattice separation hypothesis into a consumer theorem ([Erdos249257.SquaredMersenneDiagonalEnclosure.scaleFullTarget_miss_of_lambert_projected_separation](#) [SquaredMersenneDiagonalEnclosure.lean:426]). The separation hypothesis itself is exactly the remaining open content; the enclosure is real but does not supply it.

6.6.4 Old/foreign Möbius channel split (DiagonalFreshLossBridge)

The 4243-line addition to `DiagonalFreshLossBridge` isolates, at the level of individual Möbius-totient channels, which channels are “old” (index divides the LCM height H) and which are “foreign” (index does not), and proves exact identities for both. Most of the file’s bulk beyond what is recorded here is a long ladder of `...Supply-conditional “ $\Rightarrow$  irrational”` implications at successively more specialised hypotheses; those are reductions and are not restated as results.

Proposition 6.131 (Exact old-channel value: a positive gcd word, scaled). *For $H > 0$, the sum of totient forward-differences over exactly the divisor channels of H equals $(H/\text{rad}(H)) \cdot \text{gcdWordCoeff}(\text{rad}(H), s)$, where gcdWordCoeff is the positive gcd-word coefficient of §6.6.5 below.* [Erdos249257.DiagonalFreshLossBridge.oldMobiusIncrement_eq_scale_gcdWordCoeff](#) [DiagonalFreshLossBridge.lean:828] [Lean] scale:uniform coord:other:gcd-word.

Proposition 6.132 (Exact old/foreign split of the diagonal height increment). $\text{diagonalHeightIncrement}(H, s) = \text{oldMobiusIncrement}(H, s) + \text{finiteForeignChannelIncrement}(H, s)$, a literal finite-sum identity, not an estimate. [Erdos249257.DiagonalFreshLossBridge.diagonalHeightIncrement_eq_oldMobius_add_finiteForeign](#) [DiagonalFreshLossBridge.lean:829] [Lean] scale:uniform coord:other:gcd-word.

Theorem 6.133 (Exact doubling law for the old-channel increment). *If H and r are both even, $\text{diagonalHeightIncrement}(2H, 2r) = 2 \cdot \text{diagonalHeightIncrement}(H, r)$; if H is even and r odd, the same doubled height increment equals $\text{diagonalHeightIncrement}(H, r)$ unchanged.* [Erdos249257.DiagonalFreshLossBridge.diagonalHeightIncrement_two_mul_even](#) [DiagonalFreshLossBridge.lean:830] [Erdos249257.DiagonalFreshLossBridge.diagonalHeightIncrement_two_mul_odd](#) [DiagonalFreshLossBridge.lean:831] [Lean] scale:uniform coord:other:gcd-word.

Observation 6.134 (Every nonzero foreign phase term is squarefree-supported). A “foreign” channel $d \nmid H$ contributes a nonzero phase term to the literal increment only if d is squarefree and d divides exactly one of the two window endpoints $2H+s$, $H+s$ (never both, since $d \nmid H$ forces the two endpoints incongruent mod d). [Erdos249257.DiagonalFreshLossBridge.squarefree_of_foreignChannelPhaseTerm_ne_zero](#) [DiagonalFreshLossBridge.lean:909] [Erdos249257.DiagonalFreshLossBridge.endpoint_support_disjoint](#) [DiagonalFreshLossBridge.lean:852] [Lean] scale:uniform coord:other:gcd-word. This is a real structural narrowing — it rules out non-squarefree indices and simultaneous double support as sources of foreign contribution — but it narrows a finite bookkeeping decomposition, not the analytic separation obligation itself.

Proposition 6.135 (Low/top echo doubling). *If a foreign channel d has a lower-endpoint hit at offset s , its value there is $-\mu(d) \cdot \lfloor (H+s)/d \rfloor$, and its top-endpoint value at the doubled offset $2s$ is exactly twice that quantity in absolute value (with sign flipped): $\text{foreignChannelPhaseTerm}(d, H, 2s) = 2(\mu(d) \lfloor (H+s)/d \rfloor)$.* [Erdos249257.DiagonalFreshLossBridge.foreignChannelPhaseTerm_low_double_echo](#) [DiagonalFreshLossBridge.lean:941] [Lean] scale:uniform coord:other:gcd-word.

6.6.5 Repunit Möbius numerator is a positive gcd word (T1)

Theorem 6.136 (T1). *For squarefree r , the signed repunit numerator polynomial $\sum_{d|r} \mu(d)(r/d)(1 + X^d + \dots + X^{r-d})$ equals the “gcd word” whose coefficient at X^k ($k < r$) is $(r/\text{gcd}(r, k)) \cdot \varphi(\text{gcd}(r, k))$ — strictly positive for $k < r$, zero for $k \geq r$. The squarefree hypothesis is kept explicit and is not promoted to arbitrary r .* [Erdos249257.RepunitMobiusNumerator.mobiusNumeratorPolynomial_eq_gcdWord](#) [RepunitMobiusNumerator.lean:205] [Erdos249257.RepunitMobiusNumerator.gcdWordCoeff](#) [RepunitMobiusNumerator.lean:206] [Lean] scale:uniform coord:other:gcd-word.

Corollary 6.137. *Evaluation at $X = 2$ recovers the integral Möbius–Mersenne numerator already owned by `RadicalMobiusShadow`, on the same squarefree boundary.* [Erdos249257.RepunitMobiusNumerator.eval_two](#) [RepunitMobiusNumerator.lean:445] [Lean] scale:uniform coord:other:gcd-word.

6.6.6 Dyadic prefix compression: exact greedy-carry arithmetic

`DyadicPrefixCompression` adds roughly 3200 lines of exact-arithmetic infrastructure for a greedy half-orbit carry construction (Mersenne-weight achievement sets); the bulk is machinery for a specific producer hypothesis. Two pieces of genuinely reusable exact arithmetic:

Definition 6.138. For a displayed residual $p/(2L)$, the integer excess numerator above the next dyadic point $2^{-(n+1)}$ is $\text{nextDyadicExcessIntNumerator}(p, n, L) := 2^n p - L$, chosen so the skipped-branch comparison is an exact Diophantine inequality rather than a real-valued phase estimate; it obeys the exact doubling recurrence $E(p, n+1, L) = 2E(p, n, L) + L$. [Erdos249257.nextDyadicExcessIntNumerator_succ](#) [DyadicPrefixCompression.lean:161] [Lean]
scale:uniform coord:other:dyadic-carry.

Proposition 6.139 (Exact geometric tail bound for the Mersenne-weight remainder). *For $n \geq 2$, $\text{mersenneWeightRemainder}(n) \leq (4/3)(1/8)^n$, and the corresponding infinite tail from depth $m \geq 1$ is $\leq (4/21)(1/8)^m$.* [Erdos249257.mersenneWeightRemainder_le_four_thirds](#) [DyadicPrefixCompression.lean:1024]
[Erdos249257.mersenneWeightRemainderTail_le_four_twentyone](#) [DyadicPrefixCompression.lean:1024] [Lean]
scale:uniform coord:other:dyadic-carry.

6.6.7 Pivot-fibre anti-reconstruction: an exact energy identity

PivotAntiReconstruction adds roughly 1800 lines building a finite-fibre variance/energy machinery around the “first-harmonic” producer hypothesis, ending in another chain of DTW...Supply $\Leftrightarrow$ irrationality equivalences (again reductions, flagged as such). The exact algebraic core underneath is reusable:

Proposition 6.140 (Anchor defect is a squared complex distance). $\text{firstHarmonicAnchorDefect}(h, L, T) = \sum_{N \in T} \|\text{windowFirstExp}(h, N, L) - 1\|^2$, and it equals $2|T| - 2 \sum_{N \in T} \text{windowFirstCos}(h, N, L)$. [Erdos249257.TotientTailPeriodKiller.firstHarmonicAnchorDefect_eq_sum_norm_sq](#) [PivotAntiReconstruction.lean:551]
[Erdos249257.TotientTailPeriodKiller.firstHarmonicAnchorDefect_eq](#) [PivotAntiReconstruction.lean:551] [Lean]
scale:uniform coord:other:pivot-fiber.

Lemma 6.141 (Separated-pairs energy floor). *For a finite family $z : T \rightarrow \mathbb{C}$ and any set of pairs $P \subseteq T \times T$ each separated by $\geq \delta$, $|P| \cdot \delta^2 \leq \sum_{i,j \in T} \|z_i - z_j\|^2$.* [Erdos249257.TotientTailPeriodKiller.card_mul_sq_le_sum_pairwise_norm_sq_of_separatedPairs](#) [PivotAntiReconstruction.lean:724] [Lean]
scale:uniform coord:other:pivot-fiber.

6.6.8 Actual foreign-residue projection

ActualForeignResidueProjection is explicit that it is “the proof consumer” for a receipt whose analytic kernel identity remains a separate Lean validation lane; it supplies the finite bridge, not the estimate.

Proposition 6.142 (Explicit shadow is exactly the divisor-channel sum). *For $H > 0$, $\text{scaleExplicitShadow}(H) = \sum_{d|H} \text{residueIncrement}(d, H)$.* [Erdos249257.ActualForeignResidueProjection.scaleExplicitShadow_eq_sum_divisors_residueIncrement](#) [ActualForeignResidueProjection.lean:338] [Lean]
scale:uniform coord:other:foreign-residue.

Proposition 6.143 (Complement-noncancellation consumer). *If a projection’s error against the true foreign defect is controlled by the closed geometric budget $\text{foreignComplementBound}(H, D)$, and the finite rational state $\text{scaleExplicitShadow}(H) + \text{projectedForeignDefect}(H, D)$ is farther from every integer than that budget, then $S \neq \text{fullTargetHit}$ at scale H .* [Erdos249257.ActualForeignResidueProjection.miss_of_projected_separation](#) [ActualForeignResidueProjection.lean:414] [Lean]
scale:uniform coord:other:foreign-residue. *The control and separation hypotheses are exactly the two remaining open obligations; the consumer is a real theorem, not a claim they hold.*

6.6.9 Geometric coprimality: a coordinate that provably cannot decide #249 alone

GeometricCoprimalty relocates #249 onto the visible-lattice-point mass of coprime pairs, and then proves that this specific coordinate change, by itself, cannot decide irrationality at

any rational parameter — a genuine proved barrier against an entire family of approaches.

Theorem 6.144 (Visible-point count is the totient, uniformly). *For every $n \in \mathbb{N}$, $\#\{(a, b) : a + b = n, 0 < a, \gcd(a, b) = 1\} = \varphi(n)$, with no case split at $n = 0, 1$.* [GeometricCoprimalty.card_antidiagonal_filter_pos_coprime](#)[\[GeometricCoprimalty.lean:56\]](#) [\[Lean\]](#) [scale:uniform](#) [coord:other:coprime-lattice](#).

Proposition 6.145 (Coprime-pair lattice mass bridges). *For $0 \leq r < 1$: $\sum_{(a,b) \text{ coprime}, a \geq 1} r^{a+b} = \sum_n \varphi(n) r^n$; over strictly positive coprime pairs the same sum is $\sum_n \varphi(n) r^n - r$; and layering by $\gcd = g$ recovers the full positive-quadrant mass $\sum_g [\text{layer } g] = (r/(1-r))^2$, which equals 1 exactly at $r = 1/2$ (two independent fair-coin waiting times have a finite gcd almost surely).* [GeometricCoprimalty.tsum_coprime_pair_pow_eq_tsum_totient_mul_pow](#)[\[GeometricCoprimalty.lean:120\]](#) [GeometricCoprimalty.tsum_pos_coprime_pair_pow](#)[\[GeometricCoprimalty.lean:182\]](#) [GeometricCoprimalty.gcd_layer_pos_coprime_pow](#)[\[GeometricCoprimalty.lean:328\]](#) [\[Lean\]](#) [scale:uniform](#) [coord:other:coprime-lattice](#).

Theorem 6.146 (Rational geometric-model falsifier — a proved barrier). *For every $0 \leq r < 1$, $\sum_{(a,b) \text{ coprime}, a, b \geq 1} \frac{r^{a+b}}{1 - r^{a+b}} = \left(\frac{r}{1-r}\right)^2$, an elementary rational function of r , hence rational at every rational r including $r = 1/2$.* [GeometricCoprimalty.tsum_pos_coprime_lambert_eq_sq](#)[\[GeometricCoprimalty.lean:407\]](#) [\[Lean\]](#) [scale:uniform](#) [coord:other:coprime-lattice](#).

Observation 6.147. This is a proved barrier, not an observed pattern: it names exactly the class of argument it rules out. Coprime-pair restriction, exact Stern–Brocot-type splitting by gcd, and geometric cylinder decay of the summand under the map $n \mapsto r^n/(1-r^n)$, used together and evaluated at a rational r , cannot imply irrationality of the resulting sum, because the identical construction is provably rational at *every* rational r in $[0, 1)$ — the mechanism has zero sensitivity to whether r itself is rational. Any future argument built on this specific coordinate must use some further feature of $r = 1/2$ beyond coprimality, gcd-layering, and geometric decay; the coordinate change is real and interesting, but this particular route through it is closed.

6.6.10 The lcm-diagonal collapse: reduced to one $\mathbb{N}$ -indexed predicate

`LcmDiagonalReduction` (module `TotientTailPeriodKiller`, “wave 23”) removes the second free parameter from the wave-22 one-parameter reduction, producing a single decidable $\mathbb{N}$ -indexed statement equivalent to #249. The module’s own header states plainly that the resulting supply hypothesis “is the open content of #249 and is NOT claimed”; the equivalence itself is a reduction, reported as such.

Observation 6.148 (Diagonal collapse — reduction, not a result). *irrational(S) follows from: for every t_0 , some $t \geq t_0$ admits a certified kill at the diagonal point (H_t, H_t) , $H_t := \text{lcm}(1, \dots, t)$. This is proved as an implication only; the antecedent is the unproved open content.* [Erdos249257.TotientTailPeriodKiller.totient_series_of_lcm_diagonal_certificate_supply](#)[\[LcmDiagonalReduction.lean:112\]](#) [\[Lean\]](#) (implication proved; antecedent [\[Open\]](#)) [scale:cofinal](#) [coord:other:lcm-diagonal](#).

Theorem 6.149 (Window structure of the lcm ray — unconditional). *Below $2t$, the only indices j that fail to divide H_t are bare prime powers exceeding t ; every $j \leq t$ divides H_t outright. On a “clean” divisor $j \mid H_t$ (every prime of j still divides H_t/j) the window totient factors exactly: $\varphi(qH_t + j) = \varphi(j) \cdot \varphi(q(H_t/j) + 1)$.* [Erdos249257.TotientTailPeriodKiller.eq_prime_pow_of_not_dvd_periodLcm](#)[\[LcmDiagonalReduction.lean:137\]](#) [Erdos249257.TotientTailPeriodKiller.totient_periodLcm_ray_split](#)[\[LcmDiagonalReduction.lean:208\]](#) [\[Lean\]](#) [scale:uniform](#) [coord:other:lcm-diagonal](#).

Observation 6.150 (Diagonal deposits). The kernel decides $P(t)$ (a certified diagonal kill) unconditionally for every $1 \leq t \leq 8$ (totient arguments stay ≤ 130), at tabulated depths.

[Erdos249257.TotientTailPeriodKiller.certifiedKill_periodLcm_diagonal_upto_six](#) [LcmDiagonalReduct
[Erdos249257.TotientTailPeriodKiller.certifiedKill_periodLcm_diagonal_at_eight](#) [LcmDiagonalReduct
[Cert] scale:fixed coord:other:lcm-diagonal — a finite floor, not evidence toward the cofinal supply.

6.6.11 Cyclotomic anchored kills: unconditional prime support, plus new exclusion certificates

The newly published `ErdosProblems/Erdos249/CyclotomicAnchoredKill.lean` (3222 lines, namespace `ErdosProblems.Erdos249.CyclotomicAnchoredKill`) discharges the abstract order-consumer producer of §6.6.12 completely for the concrete polynomial $X - 2$, and separately deposits new kernel-checked denominator exclusions. Its middle ≈ 1600 lines are a further ladder of `CyclotomicPrime...CarryKillSupply` $\Leftrightarrow$ irrationality equivalences; those are reductions and are reported only as architecture, not as narrowing progress.

Theorem 6.151 (Unconditional unbounded prime support for the Mersenne layer). *Every prime divisor p of $2^q - 1$, for prime q , satisfies $q \mid p - 1$ (the order of 2 mod p is exactly q , by Fermat/Lagrange in $(\mathbb{Z}/p)^\times$); consequently the prime divisors appearing in the layers $\{2^n - 1\}$ are unbounded, unconditionally, with no cyclotomic resultant hypothesis left open.*

[ErdosProblems.Erdos249.CyclotomicAnchoredKill.prime_index_dvd_pred](#) [CyclotomicAnchoredKill.lean:333]
[ErdosProblems.Erdos249.CyclotomicAnchoredKill.mersenneLayer_unboundedPrimeDivisorSupply](#) [CyclotomicAnchoredKill.lean:334]
[Lean] scale:uniform coord:other:cyclotomic-anchor.

Theorem 6.152 (Unconditional clean cyclotomic anchor existence). *For every period $h > 0$ and threshold N_0 , there exist a prime q and a prime factor p of $|\Phi_{hq}(2)|$ (the binary cyclotomic layer) with p coprime to hq , $hq \mid p - 1$, and $p - 1 \geq N_0$. The characteristic-prime exceptional case in the cyclotomic order decomposition is eliminated directly, by choosing $q > 2^h$ (rules out $p = q$) and $q > h$ (rules out $p \mid h$). [ErdosProblems.Erdos249.CyclotomicAnchoredKill.exists_clean_binaryCyclotomicAnchor](#) [CyclotomicAnchoredKill.lean:1464] [ErdosProblems.Erdos249.CyclotomicAnchor.prime_order_decomposition](#) [CyclotomicAnchoredKill.lean:1390] [Lean] scale:uniform coord:other:cyclotomic-anchor.*

Observation 6.153. These two theorems are genuine unconditional number theory — an exact-order argument and a Dirichlet-plus-cyclotomic-root existence construction — and they close the abstract producer hypotheses of `PrimeRayCyclotomicCurvature` (§6.6.12) for the $X - 2$ layer with no residual conditional. Neither result touches the irrationality question itself; they supply arithmetic support for a *different*, still-open cofinal predicate about certified kills at cyclotomic-anchored periods.

Observation 6.154 (New certified exclusions at period 30). The kernel decides two independent certificates at the composite cyclotomic anchor period $h = 30$ (natural basepoint $N = 300$, and the prime-anchored basepoint $N = 330$), yielding a genuine new denominator exclusion: $S \neq r$ for every $r \in \mathbb{Q}$ with $r.\text{den} \mid 2^{300}(2^{30} - 1)$. [ErdosProblems.Erdos249.CyclotomicAnchoredKill.certifiedKill_cyclotomic_30_331_natural](#) [CyclotomicAnchoredKill.lean:3167] [ErdosProblems.Erdos249.CyclotomicAnchor.series_ne_rat_of_den_dvd_30_300](#) [CyclotomicAnchoredKill.lean:3188] [Cert] scale:fixed coord:other:cyclotomic-anchor — a finite exclusion, not evidence toward the cofinal supply.

6.6.12 Prime-ray cyclotomic curvature: the abstract order-consumer producer

Theorem 6.155 (Bounded-degree order realisability forces unbounded prime support). *If every prime divisor of a layer $C(mq)$ has order at most d in the relevant residue group (`BoundedDegreeOrderConsumer`), then for every fixed finite set of primes S , all sufficiently large prime indices q avoid S entirely on layer $C(mq)$; combined with a nontrivial-layer hypothesis this forces arbitrarily large new prime*

divisors on cofinally many prime indices. [ErdosProblems.Erdos249.PrimeRayCyclotomicCurvature.finite_of_orderConsumer](#) [[PrimeRayCyclotomicCurvature.lean:116](#)] [ErdosProblems.Erdos249.PrimeRayCyclotomicCurvature.finite_of_orderConsumer](#) [[PrimeRayCyclotomicCurvature.lean:172](#)] [Lean] scale:uniform coord:other:cyclotomic-anchor. This is the abstract producer instantiated unconditionally for $X - 2$ in §6.6.12’s sibling result above; polynomial resultant realisability and Archimedean growth for other layers remain explicit upstream obligations, not proved here.

6.6.13 Rank-one subrank obstruction: a uniform proved barrier

`RankOneSubrankObstruction` is, together with the coprimality falsifier above, the sharpest new barrier in this batch: it names an entire family of candidate linear-form constructions and proves, uniformly, that none of them can work.

Theorem 6.156 (Uniform rank-one no-go). *Let Θ_r be the Möbius–Mersenne ladder rung and let $t(Y, r)$ be its first Y atoms. For every $e \geq 1$ and $Y \geq 4$,*

$$t(Y, e+2)^2 / t(Y, 2e+2) - \Theta_2 > 1/480.$$

Every rank-one monomial Schur quotient built this way overshoots $\Theta_2 = S - 1/2$ by a fixed positive margin, uniformly in e and Y ; the bound uses only two interval facts (every rung $r \geq 3$ lies in $[1429/1512, 1)$, and every length- ≥ 4 prefix is within $1/3584$ of its rung). [ErdosProblems.Erdos249.RankOneSubrankObstruction.sub_theta_two_gt](#) [[RankOneSubrankObstruction.lean:238](#)] [ErdosProblems.Erdos249.RankOneSubrankObstruction.ge_alpha](#) [[RankOneSubrankObstruction.lean:163](#)] [Lean] scale:uniform coord:other:rank-one-subrank.

Proposition 6.157 (The gap survives positive averaging). *For any nonempty finite positively-weighted family of admissible quotients, the weighted average still overshoots Θ_2 by more than $1/480$; and every primitive integer linear form $q\Theta_2 - p$ realising such a quotient satisfies $|q\Theta_2 - p| > q/480$. [ErdosProblems.Erdos249.RankOneSubrankObstruction.positive_direct_sum_sub_theta_two_gt](#) [[RankOneSubrankObstruction.lean:300](#)] [ErdosProblems.Erdos249.RankOneSubrankObstruction.form_abs_gt](#) [[RankOneSubrankObstruction.lean:341](#)] [Lean] scale:uniform coord:other:rank-one-subrank.*

Observation 6.158. This is a proved barrier stated at the required precision: the class of argument it rules out is exactly “a primitive rational linear form for Θ_2 (equivalently for S) obtained as a rank-one strict-subrank monomial quotient of finite Möbius–Mersenne prefixes,” and the reason is the explicit uniform lower bound above, not an empirical failure report. It does not touch, and is not claimed to touch, linear forms built by any other mechanism.

6.6.14 Period-multiple escape: the nesting identity and eight new exclusions

`PeriodMultipleEscape` proves the period-multiple kill supply is *exactly equivalent* to irrationality of S (both directions, sufficiency by telescoping the tail-period law, necessity by certificate completeness) — reported here as an equivalence-class reduction, not progress — and separately deposits genuinely new denominator exclusions.

Theorem 6.159 (Nesting: the cyclotomic fan collapses onto the dyadic tower). $Q_{a+b, N} = 2^b Q_{a, N} + Q_{b, N+a}$ (block concatenation), so the order-4 cyclotomic channel at height h is literally the order-2 channel at height $2h$: the 2–3–4 fan of channels is a nested family along the tower $h, 2h, 4h, \dots$, not three unrelated moduli. [ErdosProblems.Erdos249.PeriodMultipleEscape.totientBlock_add](#) [[PeriodMultipleEscape.lean:77](#)] [ErdosProblems.Erdos249.PeriodMultipleEscape.totientBlock_two_mul](#) [[PeriodMultipleEscape.lean:94](#)] [Lean] scale:uniform coord:other:lcm-period-multiple.

Observation 6.160 (Supply $\Leftrightarrow$ irrational — reduction, not a result). [ErdosProblems.Erdos249.PeriodMultipleEscape.iff_irrational](#) [[PeriodMultipleEscape.lean:153](#)] [Lean] (equivalence proved; both directions of

the underlying predicate are exactly as open as #249 itself) `scale:cofinal coord:other:lcm-period-multiple`. The paper’s existing diagonal bank already certifies kills at H_t through $t = 64$; this module’s equivalence contributes no new information about which t beyond 64 will certify.

Observation 6.161 (Eight new certified denominator exclusions past the 64-smooth diagonal bank). The kernel certifies kills, unconditionally, at the eight prime-power periods the diagonal bank could not reach (67, 81, 97, 101, 121, 125, 127, 128, all at basepoint $N = 300$), each yielding $S \neq r$ for every r with $r.\text{den} \mid 2^{300}(2^h - 1)$ — new odd denominator classes, including the Cole factors of $2^{67} - 1$ and the Mersenne prime $2^{127} - 1$. One of the eight is certified at its own locked depth $L = h = 67$, the concrete instance of the sufficient (not known necessary) depth-equals-period form `ApFullDepthEscape`. [ErdosProblems.Erdos249.PeriodMultipleEscape.certifiedKill_67_300](#) [`PeriodMultipleEscape.lean:192`] [ErdosProblems.Erdos249.PeriodMultipleEscape.certifiedKill_127_300](#) [`PeriodMultipleEscape.lean:211`] [ErdosProblems.Erdos249.PeriodMultipleEscape.certifiedKill_67_300_fullDepth](#) [`PeriodMultipleEscape.lean:219`] [ErdosProblems.Erdos249.PeriodMultipleEscape.totient_series_ne_rat_of_den_dvd_127_300](#) [`PeriodMultipleEscape.lean:255`] [`Cert`] `scale:fixed coord:other:lcm-period-multiple` — eight finite exclusions, not evidence toward the cofinal supply.

6.6.15 Strict prime-orbit escape: a sharper reduction

`TotientStrictPrimeEscape` sharpens the legacy first-harmonic producer’s threshold from a 4/5 gap to a strict 9/10 gap with an adaptive truncation budget, and proves the sharper predicate still closes #249 through the existing singleton-certificate endpoint. This is, exactly as its own comment states, a reduction: “the producer itself remains unproved.” [ErdosProblems.Erdos249.irrational_totient_series_of_naturalPrimeTailOrbitStrictGap](#) [`TotientStrictPrimeEscape.lean:66`] [`Lean`] (implication proved; antecedent [`Open`]) `scale:cofinal coord:other:first-harmonic`.

6.6.16 Finite Euler-sieve algebra

`FiniteEulerSieve` records the elementary finite-stage identities behind the squared-Möbius Euler factor, with no transcendence claim asserted: $(1 - 2/p + 1/p^2) = (1 - 1/p)^2$ and its degree-two analogue, and the second finite difference of $1 + p + \dots + p^e$ recovering $p^{e+1}(p - 1)$, the prime-power totient row. [ErdosProblems.Erdos249.FiniteEulerSieve.muSqEulerFactor_one](#) [`FiniteEulerSieve.lean:28`] [ErdosProblems.Erdos249.FiniteEulerSieve.sigmaPrimePow_secondDiff](#) [`FiniteEulerSieve.lean:51`] [`Lean`] `scale:uniform coord:other:euler-sieve`.

7 The scale ladder

Every statement in the Erdős #249 corpus that this paper draws on is tagged with a *scale*: `scale:fixed` (a finite, explicitly enumerated set of parameter values, typically closed by `decide/interval_cases`), `scale:bounded` (holds for all parameter values on one side of a threshold, e.g. $\forall a \geq 8$, but the proof or the constants inside it do not survive removing the threshold), `scale:uniform` (holds unconditionally for every value of every free parameter, with no scale restriction at all), and `scale:cofinal` (an existential claim of the shape $\forall N_0 \exists N \geq N_0, P(N)$ infinitely often, arbitrarily far out). The source tables underlying this ladder also mark a handful of pure identities and converters `n/a` when they are not indexed by any problem-scale parameter at all (they are definitions or unconditional equivalences); for ladder purposes these are folded into `scale:uniform`, since an unconditional statement is, if anything, stronger than a uniform one. This convention is applied uniformly below and is stated here once rather than re-flagged on every row.

#249’s own supply obligation is *exactly* cofinal. The reduction chain (§??, and see [irrational_totientSeries_iff_actualLcmOrbitNonintegralitySupply](#) [`TotientActualLcmOrbitNonintegrality.lean:37`])

shows

$$\text{Irrational}\left(\sum_{n \geq 1} \varphi(n)/2^n\right) \iff \forall a_0 \exists a \geq a_0, \text{actualLcmTailOrbit } a \notin \text{range}(\mathbb{Z} \rightarrow \mathbb{R}),$$

an **cofinal** statement in the exponent a that indexes $H = \text{periodLcm}(2^a)$. Nothing weaker in scale can close #249; the entire question is whether the corpus’s uniform, bounded, and fixed machinery can be pushed to cofinal. The table below lists every catalogued #249 result, grouped by scale, with a horizontal rule separating everything that is *proved* (uniform, bounded, fixed) from the handful of statements that are themselves the *cofinal target* or consumers of it (below the rule). Quantifier prefixes are written out explicitly and exactly as recorded against the Lean source; nothing is compressed to “ $\forall \dots$ ” where the source specifies bounds.

Table 3: The #249 scale ladder. Every row above the rule is proved; every row below it is the cofinal target or a direct consumer of it.

Result	Lean site	Scale	Quantifier prefix	Coordinate
– <i>scale:uniform</i> (includes source-tagged <i>n/a</i> unconditional identities/converters) –				
totientTail defined	well- totientTail [TotientTailPeriodKiller.lean:57]	uniform	$\forall N$	binary-digit
$2^N \cdot S$ split	two_pow_mul_-totient_series_-eq [TotientTailPeriodKiller.lean:150]	uniform	$\forall N$	binary-digit
windowDiscrepancy (def)	windowDiscrepancy [TotientTailPeriodKiller.lean:66]	uniform	$\forall h \forall N \forall L$	other-binary-window
certifiedKill Sep(h, N, L) (def)	certifiedKill [TotientTailPeriodKiller.lean:72]	uniform	$\forall h \forall N \forall L$	other-binary-window
certificate floor	certifiedKill_-depth_-floor [TotientTailPeriodKiller.lean:79]	uniform	$\forall h \forall N \forall L$, given $\text{certifiedKill } h \ N \ L$	other-binary-window
kill engine soundness	tail_diff_-notMem_int_of_-certifiedKill [TotientTailPeriodKiller.lean:262]	uniform	$\forall h \forall N \forall L$, given certifiedKill	other-binary-window
certificates are complete receipts	exists_-certifiedKill_-iff_tail_-diff_notMem_-int [LcmConeFlatness.lean:316]	uniform	$\forall h \forall N$	other-binary-window
forced integrality under den. divisibility	tail_diff_-int_of_den_-dvd [TotientTailPeriodKiller.lean:327]	uniform	$\forall N \forall h \forall r:\mathbb{Q}$, given $S = r, r.\text{den} \mid 2^N(2^h - 1)$	binary-digit
tail-period law	eventual_-period_of_not_-irrational [TotientTailPeriodKiller.lean:358]	uniform	$\neg \text{Irrational } S \rightarrow \exists h > 0 \exists N_0 \forall N \geq N_0$	binary-digit
periodLcm facts	periodLcm [CarrySurvivorDef.lean:515]	uniform	$\forall n$	other-lcm-period-ray

continued on next page

Result	Lean site	Scale	Quantifier prefix	Coordinate
non-divisor < 2t is a bare prime power	eq_prime_pow_of_not_dvd_periodLcm [CertificateKernel.lean:18613]	uniform	$\forall t \forall j (0 < j < 2t), \text{ given } j \nmid \text{periodLcm } t$	other-lcm-window
LCM-ray multiplicative split	totient_periodLcm_ray_split [LcmDiagonalReduction.lean:208]	uniform	$\forall t \forall j \forall q, \text{ given } j \mid \text{periodLcm } t, \text{ clean-divisor}$	other-lcm-window-multiplicative
rationality flattens the whole LCM cone	rational_totient_series_forces_lcm_cone_flatness [CertificateKernel.lean:18674]	uniform	$\neg \text{Irrational } S \rightarrow \exists t_1 \forall t \geq t_1 \forall q > 0 \forall m$	other-lcm-cone
second-difference kill engine	second_diff_notMem_int_of_certifiedRank2Kill [CertificateKernel.lean:18729]	uniform	$\forall h \forall N \forall L, \text{ given certifiedRank2Kill}$	other-binary-window
cone-menu nonintegral pair	exists_nonintegral_pair_of_coneNonflatCert [CertificateKernel.lean:18801]	uniform	$\forall H \forall L \forall Q \neq \emptyset, \text{ given coneNonflatCert}$	other-lcm-cone-menu
survivorKill engine (carry-orbit route)	tail_diff_notMem_int_of_survivorKill [CarrySurvivorExtinction.lean:428]	uniform	$\forall h \forall N \forall K, \text{ given survivorKill}$	other-carry-orbit
Farey neighbour denominator law	farey_gap [GapFareyBound.lean:51]	uniform	$\forall a b c d r s : \mathbb{Z}, b > 0, d > 0, bc - ad = 1, \text{ unimodular-between}$	farey
Dirichlet near-integer criterion	irrational_of_den_mul_abs_sub_tendsto_zero [CertificateKernel.lean:5371]	uniform	$\forall u : \mathbb{N} \rightarrow \mathbb{Q} \forall x : \mathbb{R}$	n/a (generic)
Erdős 1948 near-integer criterion	irrational_of_int_mul_near_int [CertificateKernel.lean:5792]	uniform	$\forall \xi : \mathbb{R}, \forall q > 0 \exists m z 0 < m\xi - z < 1/q$ (generic)	n/a (generic)
finite-minor $\Rightarrow$ linear independence	linearIndependent_of_separatedMinorCertificate [TotientCarryDefect.lean:81]	uniform	$\forall \text{family} : \iota \rightarrow \mathbb{N} \rightarrow \mathbb{Q}, \text{ given SeparatedMinorCertificate}$	n/a (generic)
dyadic totient-kernel rank = $2^e + 1$	linearIndependent_canonicalTotientKernelFamily [TotientMahlerDefect.lean:90]	uniform	$\forall e$	other-dyadic-kernel-rank
rationality forces unbounded carry-kernel rank	not_irrational_totientSeries_implies_unbounded_carryRank_unconditional [TotientCarryKernelRigidity.lean:284]	uniform	$\neg \text{Irrational } S \rightarrow \exists v > 0 \exists u \forall e$	other-carry-kernel-rank
no fixed D clears all primitive Euler jets	no_fixed_integral_primitive_euler_index [PrimitiveDeterminantLift.lean:169]	uniform	$\forall D > 0$	mobius-mersenne

continued on next page

Result	Lean site	Scale	Quantifier prefix	Coordinate
MersenneLambert ladder value identities	tsum_totient_- div_pow_two_- eq_pnat_half_- pow [CertificateKernel.lean:18063]	uniform	value identities, $L(\mu)=\frac{1}{2}$, $L(1)=\text{Erdős-Borwein}$	mobius- mersenne
prime-power reduced-denom. unit-gap ceiling	reducedDenominatorUnitGapCeiling prime_pow_- iff [PrimitiveWeightCertificate.lean:54]	uniform	$\forall p \text{ prime } \forall e > 0$	farey
positive rational- difference lower bound	positive_- rational_- difference_- lower_- bound [PrimitiveRationalGapSupply.lean:31]	uniform	$\forall \text{ whole pfx: } \mathbb{Q}, \text{ pfx} < \text{whole}$	n/a (generic)
Möbius-square identity ($S = \frac{1}{2} + \Sigma$)	tsum_totient_- half_pow_- eq_half_- add_moebius_- sq [SquaredMersenneDiagonalEnclosure.lean:50]	uniform	unconditional	mobius- mersenne
squared-Lambert transfer engine	tsum_lambert_- linear_- weight_sq_- pure [GcdMomentCalculus.lean:105]	uniform	$\forall w: \mathbb{N} \rightarrow \mathbb{R} (w(d) \leq d) \forall r \in [0, 1)$	mobius- mersenne
$L_2(\mu) = S - \frac{1}{2}$	prose:L2_of_mu_ is_249	uniform	unconditional	mobius- mersenne
$L_2(1) = \zeta_q(2) - \zeta_q(1)$ identity	tsum_one_div_- mersenne_sq_eq_- sigma_sub_tau_- series [GcdMomentCalculus.lean:216]	uniform	unconditional	mobius- mersenne
$L_2(\varphi) = \text{gcd moment}$	tsum_totient_- div_mersenne_sq_- eq_gcd_moment_- series [GcdMomentCalculus.lean:235]	uniform	unconditional	mobius- mersenne/probability
level-mirror table	prose:level_mirror_ table	uniform	unconditional	mobius- mersenne
gcd-divisibility fac- torizes	tsum_pos_pair_- both_dvd_half_- eq_inv_mersenne_- sq [GcdMomentCalculus.lean:266]	uniform	$\forall d: \mathbb{N}, d > 0$	probability
reduced-direction law $\Sigma = 1$	tsum_pos_- coprime_inv_- mersenne_eq_- one [GcdMomentCalculus.lean:349]	uniform	unconditional (sum over all coprime pairs)	probability
Stern-Brocot cylinder recursion	cylinderMass_- split [GcdMomentCalculus.lean:474]	uniform	$\forall a, b: \mathbb{N}^+ \forall \text{depth } d$	other-stern- brocot
repunit gcd-word T1	mobiusNumeratorPolynomial eq_- gcdWord [RepunitMobiusNumerator.lean:205]	uniform	$\forall r \text{ (Squarefree } r) \forall k < r$	cyclotomic
eval-at-2 recovers numerator	mobiusNumeratorPolynomial eval_- two [RepunitMobiusNumerator.lean:320]	uniform	$\forall r \text{ (Squarefree } r)$	cyclotomic

continued on next page

Result	Lean site	Scale	Quantifier prefix	Coordinate
radical-shadow scale decomposition	scaledMobiusShadow_eq_- radicalBase [RadicalMobiusShadow.lean:125]	uniform	$\forall H > 0 \forall r > 0$	mobius-mersenne
cyclotomic congruence T2	cyclotomic_dvd_- mobiusNumeratorPolynomial_- sub [CyclotomicProjectionOfShadow.lean:299]	uniform	$\forall r$ (Squarefree r) $\forall m (m r)$	cyclotomic
top-fibre survives T3	mobiusNumerator_- gcd_- cyclotomicValue [CyclotomicProjectionOfShadow.lean:350]	uniform	$\forall r$ (Squarefree r)	cyclotomic
upper-half prime channel survival T4	upperHalfChannel_- product_dvd_- den_of_scale_- primeFactors_- le [MersenneShadowCyclotomicNoncollapse.lean:809]	uniform	$\forall t \forall p$ prime $\in (t/2, t]$, scale-coprime-to- C	cyclotomic
denominator lower bound / exact value	lcmHeight_- scaledMobiusShadow_- den_- exact [MersenneShadowDenominatorGrowth.lean:147]	uniform	$\forall t$ (exact-value); $t \geq 5$ (lower bound)	mobius-mersenne
prime-power jump recurrence T5	cyclotomic_dvd_- primeJump_new_- fibre [PrimePowerJumpDynamics.lean:281]	uniform	$\forall p$ prime ($p \nmid r$) $\forall m (m r)$	cyclotomic
squared-Mersenne diagonal tail enclosure	mobius_square_- series_eq_- partial_add_- tail [SquaredMersenneDiagonalEnclosure.lean:72]	uniform	$\forall D: \mathbb{N}$	mobius-mersenne
(3, 5) joint annihilator	joint35_- oldChannel_- zero [JointExponentTransport.lean:71]	uniform	$\forall d > 0 (d H)$; general: finite affine annihilator	other-lcm-diagonal
composite-dilation defect identity	supportCoeff_- mul_eq_add_- defect [CompositeDilationDefect.lean:30]	uniform	$\forall a \in A (a > 0) \forall x > 0$	other-divisor-support-coeff
sublogarithmic zero-window T11	supportCoeffZeroWindow_- length_le_- eps_logb_- add [SublogDivisorCoverage.lean:392]	uniform	$\forall \varepsilon > 0 \exists B$, given rationality of the base-2 support series	other-support-divisor-counting
campbell shift $\leftrightarrow$ Mersenne endpoint (shared namespace, #257-flavored)	shiftWindowZero_- iff_half_mem_- mersenneAchievementSet [CampbellShiftSynchronization.lean:486]	uniform	iff, no extra parameter	other-greedy-mersenne-achievement
totient overlap-factor multiplicativity	totient_mul_eq_- overlapFactor_- mul [TotientActualLcmOrbitArithmetic.lean:43]	uniform	$\forall x > 0 (j \text{ fixed})$	other-totient-arithmetic
totient relative-Euler-product split	totient_mul_- eq_totient_mul_- relativeEulerProduct [TotientActualLcmOrbitArithmetic.lean:410]	uniform	$\forall j x > 0$	other-totient-arithmetic

continued on next page

Result	Lean site	Scale	Quantifier prefix	Coordinate
lcmRayArithmeticLetter = deltaTotient = diagonalWindowIn- crement	lcmRayArithmeticLetter eq_- deltaTotient [TotientActualLcmOrbitArithmetic.lean:298]	uniform	$\forall t \forall j$	mobius- mersenne
lcmDiagonalArithmeticWindow = windowDiscrep- ancy	lcmDiagonalArithmeticWindow eq_- windowDiscrepancy [TotientActualLcmOrbitArithmetic.lean:1998]	uniform	$\forall t \forall L$	mobius- mersenne
#249 $\iff$ cofinal actual-orbit nonin- tegrality	irrational_- totientSeries_- iff_- actualLcmOrbitNonintegralitySupply [TotientActualLcmOrbitNonintegrality.lean:37]	uniform	iff	mobius- mersenne
actualLcmTailOrbit = scaled series - prefix	actualLcmTailOrbit_- eq_scaled_- totientSeries_- sub_- prefix [TotientActualLcmOrbitSeparation.lean:68]	uniform	$\forall a$	mobius- mersenne
explicit finite-block approximation	abs_- actualLcmTailOrbit_- sub_rawApprox_- lt [TotientActualLcmOrbitSeparation.lean:141]	uniform	$\forall a \forall q$	mobius- mersenne
guard-cylinder nor- mal form	exists_- certifiedKill_- iff_- guardCylinderWitness [TotientActualLcmTopEdgeStaircase.lean:844]	uniform	$\forall h \ N$, iff	seam- integer
fixedRankSecondDiffer- ence sign under extremal- ity	fixedRankSecondDifference [TotientFixedRankLcmAsymptotic.lean:256]	uniform	$\forall h \ j$, given $\text{rankTotientExtremal}$	fixed-rank- curvature
dyadic fixture exactness (kernel- valuation sharpness)	fixedRankSecondDifference dyadic_fixture_- exact [TotientFixedRankLcmAsymptotic.lean:462]	uniform	$\forall n$ (closed-form)	other- fixed-rank- curvature
parityCoboundaryWeight (def)	parityCoboundaryWeight [TotientParityCoboundaryCountermodel.lean:37]	uniform	$\forall n$	binary-digit
totientTail = pos- itive foreign-residue kernel sum	totientTail_- eq_tsum_- positiveForeignResidueKernel [TotientShiftedMobiusForeignBridge.lean:52]	uniform	$\forall N$	cyclotomic
totientTail = shifted Möbius-pulse sum	totientTail_- eq_tsum_- shiftedMobiusPulse [TotientShiftedMobiusPulse.lean:535]	uniform	$\forall N$	cyclotomic
Lambert double- sum regroup engine	tsum_lambert_- pair_regroup_- if [TotientShiftedMobiusPulse.lean:112]	uniform	$\forall w \ v:\mathbb{N} \rightarrow \mathbb{R} (w(d) \leq d, v(m) \leq m)$	cyclotomic
carryShift- integrality iff	carryShift_- dvd_iff_- tailDiff_mem_- int [TotientTailCarryPeriod.lean:77]	uniform	$\forall N \ k$, given tempered orbit	seam- integer

continued on next page

Result		Lean site	Scale	Quantifier prefix	Coordinate
rationality	$\Rightarrow$	not_irrational_-	uniform	$\neg \text{Irrational } S \rightarrow \exists v u \forall e$	seam-integer
periodicity & unbounded rank		totientSeries_- implies_mod_- period_and_- unbounded_- rank [TotientTailCarryPeriod.lean:224]			
adjugate tail-cost floor ≥ 3		three_le_- totientAdjugateTailCost [TotientTailCarryPeriod.lean:1266]	uniform	$\forall \text{ finite } w:\iota \rightarrow \mathbb{Q} x:\iota \rightarrow \mathbb{N} \sum_{i \in w} \text{period}(x_i) \geq 3$	other-linear-algebra
directedCertifiedKill exact iff		directedCertifiedKill [TotientTailCarryPeriod.lean:766]	uniform	$\forall n \in \mathbb{N} \exists L \forall c \in \mathbb{N} \rightarrow \mathbb{N} (c(n) \leq n), \text{ iff}$	seam-integer
tempered orbit rigidity iff		binaryCoeffSeries_- rational_- iff_exists_- temperedBinaryOrbit [GenericTailOrbitRigidity.lean:425]	uniform	$\forall c:\mathbb{N} \rightarrow \mathbb{N} (c(n) \leq n), \text{ iff}$	binary-digit
doubling-orbit rigidity		doublingOrbit_- eq_zero_of_- tempered [GenericTailOrbitRigidity.lean:314]	uniform	$\forall d:\mathbb{N} \rightarrow \mathbb{R}, \text{ doubling} + \text{tempered} \rightarrow d \equiv 0$	other-abstract-recursion
finite-state (unbounded state)	no-go	balancedPulse_- no_autonomous_- decoder [GenericTailOrbitRigidity.lean:246]	uniform	$\forall m \forall \text{State}, \text{ collapsing-state} \rightarrow \text{no decoder}$	binary-digit
fixed-depth reset	affine	affineBinaryOrbit_- mod_twoPow_- eq [GenericTailOrbitRigidity.lean:306]	uniform	$\forall a:\mathbb{N} \rightarrow \mathbb{Z} \forall u_0 v_0 \forall L$	binary-digit
signed engine (full-block certs)	dichotomy (full-block)	int_coeff_- series_- irrational_- or_bpow_mul_- eq_intCast_- of_full_block_- certificates [CertificateKernel.lean:13606]	uniform	$\forall b \geq 2 \forall c \forall q, \text{ given full-block certificate at } q$	binary-digit
Lambert sum regroup (dyadic forward-difference calc.)	double-	dvd_- iteratedForwardDifference [LcmFactorIdealPulseObstruction.lean:211]	uniform	$\forall f, c:\mathbb{N} \rightarrow \mathbb{Z} \forall \text{ shift-lists}$	binary-digit
fair-coin coprimality bridge		tsum_coprime_- pair_pow_- eq_tsum_- totient_mul_- pow [GeometricCoprimalty.lean:120]	uniform	$\forall n; \forall r \in [0, 1)$	probability
linear-descender rigidity		linearDescender_- eq_smul_- eval [AdelicHeightObstruction.lean:120]	uniform	$\forall V, W \forall \text{ev} \forall L, \text{ker}(\text{ev}) \leq \text{ker}(L)$	p-adic
rational denominator survival law		divisor_- dvd_divInt_- den [RationalDenominatorSurvival.lean:17]	uniform	$\forall D > 0 \forall m D \forall a$	p-adic

continued on next page

Result	Lean site	Scale	Quantifier prefix	Coordinate
scalar-localization	scalarLocalization_-	uniform	$\forall x:\mathbb{Q} \forall H (H x.\text{den}) \forall c ((cx) \text{pad} H)$	padic
complement-dvd	complement_-			
(adelic height tax)	dvd [AdelicHeightObstruction.lean:23]			
signed Hankel	scaled_-	bounded	$\exists m \in s$ (strictly max-	p-adic
terminal-parity	dyadic_sum_ne_-		imal, odd coeff),	
engine	zero [SignedQMomentObstruction.lean:95]		$\forall m e(i) < e(m)$	
residual gauge ob-	det_-	uniform	$\forall d \forall e: \text{Fin } d \rightarrow \mathbb{N} \forall z: \text{Fin } d \rightarrow \mathbb{C}$	other-first-
struction (locked	residualMonomialMatrix [ResidualGaugeObstruction.lean:47]			harmonic-
minors)				phase
incidence-quotient	incidenceMobius_-	uniform	$\forall N:\mathbb{N}$	mobius-
no compression	det_eq_-			mersenne
	one [IncidenceQuotientHermitePade.lean:56]			
– scale:bounded –				
Farey window	gap_check_-	bounded	$\forall q:\mathbb{N}, 0 < q \leq 79,639,646,646,701,375,323,355,774,875,$	
$K=240$ denomina-	window_1_240_le_-			
tor bound	79639646646701375323355774875831053 [GapFareyBound.lean:176]			
denominator lower	tsum_totient_-	bounded	$\forall p:\mathbb{Q}, p.\text{den} \leq Q_0$	farey
bound $S \neq p/q,$	div_pow_two_-			
$q \leq Q_0$	ne_ratCast_-			
	of_den_le_-			
	79639646646701375323355774875831053 [CertificateKernel.lean:18056]			
foreign-residue tail-	controlledForeignProb_	bounded	$\forall H D$ with $2H \leq D$,	mobius-
limit converter	of_tail_-		given tail-limit conver-	mersenne
	limit [LambertDiagonalEnclosure.lean:60]			
fixed-precision trop-	fixedPrecisionTropical	bounded	$\forall n \geq 0$ (fixed precision) $\forall n$	p-adic
ical no-go			finite carry word	
μ -clean- Padé feasi-	prose:mu_pollu-	bounded	claimed $\forall d$; machine-	mobius-
bility wall	tion_qpade_wall		audited only $d \leq 12$	mersenne
				(q- Padé)
iterated pullback /	iteratedDilationDiff_	bounded	given $\Omega(a) \leq K \forall a \in A$	other-
bounded- Ω vanish-	eq_zero_of_-			support-
ing	cardFactors_-			divisor-
	le [SupportDilationDifferences.lean:250]			counting
rough-density $\Rightarrow$ to-	three_quarters_-	bounded	$\forall a \geq 8 \forall n > 0$ (n is t -	mobius-
tient $\geq \frac{3}{4}n$	mul_lt_totient_-		rough, $t=2^a, n < 2^{2 \cdot 2^a}$)	mersenne
	of_rough_lt_two_-			
	pow [TotientActualLcmOrbitArithmetic.lean:768]			
short-window sign	lcmRayArithmeticLetto_	bounded	$\forall a \geq 8 \forall j (0 < j < 2 \cdot 2^a)$	mobius-
fixed positive	pos_of_lt_two_-			mersenne
	mul [TotientActualLcmOrbitArithmetic.lean:1703]			
unconditional lower-	actualLcmTailDiff_-	bounded [†]	$\forall a \geq 8 \forall J (J + (a+6) < 2 \cdot$	mobius-
half positivity	shift_-		$2^a)$, no rationality	mersenne
	pos [TotientActualLcmOrbitSign.lean:139]		hypothesis	
top-edge residue	actualLcm_-	bounded	$\forall a \geq 8 \forall J K$ (room	mobius-
survivor negative	trueEndpointSurvivor_-	bound)		mersenne
	neg [TotientActualLcmOrbitSign.lean:172]			
integrality forces	actualLcm_-	bounded	$\forall a \geq 8 \forall J K$ (room,	mobius-
top-edge residue	integral_forces_-		$2H + J + K + 2 < 2^K$)	mersenne
	topEdgeResidue [TotientActualLcmOrbitSign.lean:211]			

continued on next page

Result	Lean site	Scale	Quantifier prefix	Coordinate
short-kill supply through $a \leq 6$	powerTwoActualLcmShortKill boundedMeta supply_through_ six [TotientActualLcmShortKill.lean:54]		$\forall a \leq 6$	mobius-mersenne
terminal staircase impossible with room	not_ actualLcmTerminalDyadicStaircase modulus_of_ room [TotientActualLcmTopEdgeStaircase.lean:251]	bounded	$\forall a \geq 8 \forall J K m$ (room, modulus-wide)	mobius-mersenne
punctured staircase penultimate pin	puncturedDyadicStaircase penultimate_eq_ half [TotientActualLcmTopEdgeStaircase.lean:1187]	bounded	$\forall a \geq 8 \forall J K m$ (room + punctured hyp.)	mobius-mersenne
top-edge residue-gap $\Rightarrow$ nonintegral	actualLcmTailDiff_ notMem_int_of_ topEdgeResidueGap [TotientActualLcmTopEdgeStaircase.lean:1260]	bounded	$\forall a \geq 8 \forall J K m$ (room, one-sided)	mobius-mersenne
odd-rank centred-lift = terminal – true carry	two_mul_ actualOddHalfCenteredLift_ eq_terminal_sub_ trueCarry [TotientActualLcmTopEdgeStaircase.lean:1678]	bounded	$\forall a \geq 8 \forall q$ (room, fit bound $2(H+q+2) \leq 4^q$)	mobius-mersenne
prime-power kernel valuation floor	two_mul_ totient_dvd_ fixedRankSecondDifference [TotientFixedRankLcmAsymptotic.lean:1019]	bounded	$\forall a \geq 4 \forall j > 0 (j^2 \leq 2^a)$	other-fixed-rank- kernel
finite rational-separation $\Rightarrow \neg$ hit (#257-flavored)	renormalizedResidueApprox of_twice_ le [TotientShiftedMobiusForeignBridge.lean:329]	bounded	$\forall H > 0 \forall D \geq 2H$, given finite separation	mobius-mersenne
homogeneous Mersenne multiplier does not annihilate	idCoeff_ mersenneModulus_ does_not_ annihilate_ tailShift [TotientTailCarryPeriod.lean:342]	bounded	$\forall K, q (K > 0, K < q, q 2^K - 1)$	mobius-mersenne
numerator-gap $\Rightarrow \neg$ hit (#257-flavored)	scaleFullTarget_ miss_of_lambert_ projected_num_ gap [LambertProjectedNumeratorGap.lean:23]	bounded	$\forall H, D$ fixed, given uniform numerator-gap bound	mobius-mersenne
periodic-weight Lambert dichotomy	irrational_ or_bpow_mul_ eq_intCast_ intWeightedErdosSeries_ periodic [CertificateKernel.lean:13847]	bounded	$\forall b \geq 2 \forall m > 0 \forall w$ ($w(n+m) = w(n)$)	mobius-mersenne
non-negative periodic $\Rightarrow$ irrational (unconditional)	irrational_ intWeightedErdosSeries_ periodic_of_ coeff_nonneg_of_ frequently_ne_ zero [CertificateKernel.lean:14255]	bounded	$\forall b \geq 2 \forall m > 0, w$ periodic, ≥ 0 , frequently $\neq 0$	mobius-mersenne
– <i>scale:fixed</i> – all $h \in [1, 8]$ kill certificate at depth 16, $N=12$	certifiedKill_ all_ small [TotientTailPeriodKiller.lean:1404]	fixed	$\forall h \in [1, 8]$, certifiedKill h 12 16 (104 decide)	other-binary-window

continued on next page

Result	Lean site	Scale	Quantifier prefix	Coordinate
$h \in [1, 16]$, exponent 14 kill table	totient_series_-ne_rat_of_den_-dvd_pow_two_mul_-mersenne_upto_-sixteen [CertificateKernel.lean:18562]	fixed	$\forall h \in [1, 16]$, denominator exponent 14 (by decide)	other-binary-window
rank-2 kill not shallower at $(h, N)=(1, 8)$	totient_tail_-rank_two_kill_-sound_but_-not_shallower_-cell [CertificateKernel.lean:18762]	fixed	measured over $t \leq 20$ (30/40 cells)	other-binary-window
28-point diagonal pincer table through $t=64$	certifiedKill_-diagonal_all_-imported [DiagonalPincerCertificateKernel.lean:231]	fixed	$t \in \{1, 2, 3, 4, 5, 7, 8, 9, 11, 13, 16, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63\}$ (28 explicit values)	other-lcm-diagonal
short kill at $a=4$ and $a=6$	lcmDiagonalArithmeticFixed1_-two_pow_-four [TotientActualLcmShortKill.lean:18]	fixed	$a=4, a=6$ (concrete, kernel-checked)	mobius-mersenne
square-CRT clean block: vanishing / nonvanishing witnesses	squareCRT_-clean_block_can_-vanish [SquareCRTCube.lean:285]	fixed	witness $n=52$ (vanishes); witness $n=27$ (nonzero)	other-crt-dyadic-residue
echo-vs-height two-front wall	prose:echo_versus_-height_two_front_-wall	fixed	empirical over tested families only (not universal)	mobius-mersenne (q-Padé, quantitative)
period-4 zeroset escape witness	intWeightedCoeff_-periodFourSignWeight_-eq_zero_of_-mod_four_eq_-three [CertificateKernel.lean:13898]	fixed	$\forall n, n \equiv 3 \pmod{4}$ (fixed weight w)	mobius-mersenne
— frontier: everything below this line is the cofinal target or a direct consumer of it —				
– scale:cofinal (OPEN) –				
THE WALL certificate supply $\Rightarrow$ Irrational S	irrational_-totient_series_-of_certificate_-supply [TotientTailPeriodKiller.lean:394]	cofinal	$\forall h \geq 1 \forall N_0 \geq 0 \exists N \geq N_0 \exists L$, certifiedKill $h N L$	other-binary-window
multiple-certificate supply (weakened)	irrational_-totient_series_-of_multiple_-certificate_-supply [CarrySurvivorExtinction.lean:502]	cofinal	$\forall h_0 > 0 \forall N_0 \exists m > 0 \exists N \geq N_0 \exists L$	other-lcm-period-multiple
LCM-diagonal one-parameter restatement	irrational_-totient_series_-of_lcm_diagonal_-certificate_-supply [LcmDiagonalReduction.lean:112]	cofinal	$\forall t_0 \exists t \geq t_0 \exists L$	other-lcm-diagonal

continued on next page

Result	Lean site	Scale	Quantifier prefix	Coordinate
LCM-cone window-kill supply (widest target)	irrational_-totient_series_of_lcm_cone_window_kill_supply [CertificateKernel.lean:18686]	cofinal	$\forall t_0 \exists t \geq t_0 \exists q > 0 \exists m \exists L$	other-lcm-cone
LCM diagonal/cone nonintegrality restatement	irrational_-totient_series_of_lcm_diagonal_nonintegrality_supply [CertificateKernel.lean:18706]	cofinal	$\forall t_0 \exists t \geq t_0, \text{totientTail}(2H_t) - \text{totientTail}(H_t) \notin \mathbb{Z}$	other-real-nonintegrality
LCM cone-nonflat supply (sharpest depth-reduced form)	irrational_-totient_series_of_lcm_cone_nonflat_supply [CertificateKernel.lean:18812]	cofinal	$\exists$ unbounded-scale menu Q with coneNonflatCert firing	other-lcm-cone-menu
first-harmonic norm-gap supply	irrational_-totient_series_of_first_harmonic_norm_gap [FirstHarmonicPivot.lean:83]	cofinal	$\forall h > 0 \forall X_0 \exists X \geq \max(X_0, 1) \exists L$ room $16(2X + h + L + 2) \leq 2^L$	binary-digit
prime-jump sharp-kill supply	irrational_-totient_series_of_prime_jump_sharp_kill_supply [PrimeJumpWindow.lean:193]	cofinal	$\forall t_0 \exists t \geq t_0 \exists p, L > 0$	binary-digit
a -exponent short-arithmetic kill supply (the open trigger)	PowerTwoActualLcmShortArithmeticKillSupply [PowerTwoActualLcmShortArithmeticKillSupply.lean:229]	cofinal	$\forall a_0 \exists a \geq a_0 \exists L$	mobius-mersenne
actual-orbit separation supply	irrational_-totientSeries_of_actualLcmOrbitSeparationSupply [TotientActualLcmOrbitSeparation.lean:305]	cofinal	cofinal 1/32+error separation at guarded odd ranks	mobius-mersenne
top-edge-residue-gap sufficiency chain (5 links, weakest first)	irrational_-totientSeries_of_flexibleActualTerminalDominanceSupply [TotientActualLcmTopEdgeStaircase.lean:222]	cofinal	$\forall a_0 \exists a \geq a_0 \exists$ (supply predicate)	mobius-mersenne
rational countermodel refutes {bound, parity, aperiodicity}	exists_-totientParity_of_arbitrarilyManySeparatedCarryCountermodel [TotientParityCoboundaryCountermodel.lean:637]	cofinal	$\exists c: \mathbb{N} \rightarrow \mathbb{N}, c \leq 6, c \leq n, c \equiv \varphi \pmod{2},$ arbitrarily-separated blocks	binary-digit
mod-4 pulse co-final supply (via CRT+Dirichlet)	exists_prime_deltaTotient_four_mul_mod_four_two [TotientTailCarryPeriod.lean:384]	cofinal	$\forall h > 0 \forall B \exists \text{prime } p > B$	mobius-mersenne

continued on next page

Result	Lean site	Scale	Quantifier prefix	Coordinate
mod-4 pulse survivor kill (fourfold reduction)	irrational_- totientSeries_- of_cofinal_- modFourPulseSurvivorKill [TotientTailCarryPeriod.lean:1066]	cofinal	given CP-05's cofinal supply	mobius-mersenne
two-adic pulse at arbitrary depth K	exists_prime_- totient_- twoAdic_pulse_- divisors [TotientTwoAdicPulseBlock.lean:54]	cofinal	$\forall K \geq 2 \forall H > K \forall B \exists$ cofinally many primes $p > B$	p-adic
two-adic pulse transfer (needs eventual integrality)	eventual_- integral_- tailDiff_- has_cofinal_- twoAdic_half_- pulse [TotientTwoAdicPulseBlock.lean:327]	cofinal	$\forall K \geq 2$, given pulse + eventual integrality	TA-01 p-adic

[†] The source normal-form table tags `actualLcmTailDiff_shift_pos` `scale:bounded` because the window parameter J is finite-range relative to $2 \cdot 2^a$; the exponent a itself is completely unbounded ($\forall a \geq 8$), and this row's constants (4, 8, 32) do not degrade with a . It is flagged here, not silently re-tagged, because it is exactly the kind of row an automated promotion sweep must not misclassify see the audit below.

7.1 Checked negative result: no free promotion from bounded to cofinal

A ladder like the one above invites an obvious question: is any `scale:bounded` result actually uniform in disguise, so that removing its threshold is a routine generalisation rather than new mathematics? This was checked directly rather than assumed. An audit read the Lean proof body behind every bounded-scale result in the #249 corpus eighteen candidates in total, spanning the near-miss list against the #249-supply obligation catalogued in the interface index (fourteen rows, each independently checked against the exact cofinal target it is closest to) together with the remaining bounded/fixed rows in the certificate, Möbius–Lambert, and private-totient banks above and found **no** bounded result whose proof is uniform enough in its threshold parameter to promote to cofinal or uniform for free. Every candidate failed for one of three reasons, and the failure mode is worth recording because it is a checked negative result, not an absence of search:

1. **Finite certificate tables.** Results such as [certifiedKill_diagonal_all_imported](#) [DiagonalPincerCertificate.lean:404] (28 explicit values of t through $t=64$), [certifiedKill_all_small](#) [TotientTailPeriodKiller.lean:404] ($h \in [1, 8]$ at fixed depth 16), and [totient_series_ne_rat_of_den_dvd_pow_two_mul_mersenne_upto_sixteen](#) [CertificateKernel.lean:18562] ($h \in [1, 16]$ at exponent 14) are each a finite *list* of independently verified rows, not a single argument instantiated at a free parameter. There is no uniform proof underneath to strip the bound from; each new row is a fresh finite computation.
2. **interval_cases/decide over a bounded range.** Results such as [three_quarters_mul_lt_totient_of_rough_lt_two_pow](#) [TotientActualLcmOrbitArithmetic.lean:768] ($a \geq 8$) and the entire [actualLcmTailDiff_shift_pos](#) [TotientActualLcmOrbitSign.lean:39]/top-edge-staircase family (room conditions of the shape $J + (a+6) < 2 \cdot 2^a$) are proved by case-splitting a residue or a divisor structure that is only exhaustively enumerable inside the stated range; the mathematical content genuinely narrows as the range widens, so `decide` cannot simply be re-run at a larger bound without an exponential blow-up in the search space it certifies.

3. **Constants that degrade with the parameter.** The clearest instance is the echo-versus-height wall (prose:echo_versus_height_two_front_wall, §5e–5f of the ambitious-modular-route note): height-optimal ladder families pay quadratic height $P(1) \sim 2^{K^2/2}$ matching `lcmHeight_scaledMobiusShadow_den_exact` [MersenneShadowDenominatorGrowth.lean:147]’s own growth rate, while height-minimal integer-relation minimizers regrow the echo term E_R to match for every tested family, the two costs cannot both be driven to zero as the scale parameter grows. The interface index records the same phenomenon for `HalfRung(J)`-shaped truncation rungs: the certificate window $B(J)$ is driven by $L_J = \text{lcm}(2, \dots, J)$, which grows super-exponentially in J , so the per- J constant degrades with the very parameter the cofinal claim needs to range over “the hallmark of a proof that does not survive bound removal” (interface index, row d-3a/d-3b).

None of the fourteen near-miss rows against the #249-supply obligation is closed by a scale argument alone either: reading each one’s own recorded `exact_mismatch` shows the residual gap is a missing *arithmetic input* (a constant-saving Weyl-sum bound for `e1-companion`; a one-sided top-edge residue gap for `TE-04/TE-05-weakest`; an unbounded finite-rank compression on the rationality side for `d4/d5`), never a generalisable proof technique sitting one `omega` call away from cofinal. The one row where the source tag itself is imprecise `SGN-01`, marked `scale:bounded` even though its underlying exponent a is unbounded, see [†] above is precisely the kind of false positive the audit was built to catch, and even there what remains open is the *other* half of the certificate band (the top-edge residue), not a scale defect in `SGN-01` itself.

8 Coordinate atlas and transport maps

#249 is not proved or disproved in a single representation. The corpus expresses $S = \sum_{n \geq 1} \varphi(n)/2^n$ in at least eight distinct coordinates, and the central methodological lesson of the whole programme flagged repeatedly in the source banks as AP17/AP18, “obstructions are coordinate-relative” is that a mechanism which kills a proof strategy in one coordinate can be silent, or even actively helpful, in another. This section lists every coordinate used, gives at least one exact transport map out of it into another coordinate with its Lean site, and then makes the coordinate-relativity claim precise for the one pair where it matters most: binary-digit versus Möbius–Mersenne.

binary-digit

S as the literal base-2 expansion object: `totientTail`, `windowDiscrepancy`, `certifiedKill`.

Transport to mobius-mersenne: the core identity

$$S = \sum_{n \geq 1} \frac{\varphi(n)}{2^n} = \frac{1}{2} + \sum_{d \geq 1} \frac{\mu(d)}{(2^d - 1)^2}$$

proved by `tsum_totient_half_pow_eq_half_add_moebius_sq` [SquaredMersenneDiagonalEnclosure.lean:50] (consumed at `mobius_square_series_eq_partial_add_tail` [SquaredMersenneDiagonalEnclosure.lean:72]) turns every binary-digit statement about S into a statement about a squared Möbius–Mersenne Lambert series, and back.

mobius-mersenne

S as $\frac{1}{2} + \sum_d \mu(d)/(2^d - 1)^2$, the Lambert ladder $L(w) = \sum_n w(n)/(2^n - 1)$ at $w = \mu, 1, \varphi$, and the whole cyclotomic denominator-survival chain built on it. *Transport to cyclotomic:* evaluating the Möbius-numerator polynomial at $x = 2$, `mobiusNumeratorPolynomial_eval_two` [RepunitMobiusNumerator.lean:320] together with the definition `mobiusNumerator` [RadicalMobiusSha] moves the squared-denominator question into finite cyclotomic-polynomial arithmetic over $\mathbb{Z}$.

probability/coprime-pair

$S - \frac{1}{2} = \Pr(\gcd(X, Y) = 1)$ for independent fair-coin waiting times X, Y . *Transport from mobius-mersenne*: the gcd-divisibility factorisation `tsum_pos_pair_both_dvd_half_eq_inv_mersenne_sq` [GcdMomentCalculus.lean:266] ($\Pr(d \mid X \wedge d \mid Y) = 1/(2^d - 1)^2$) together with the reduced-direction law `tsum_pos_coprime_inv_mersenne_eq_one` [GcdMomentCalculus.lean:349] converts the squared-Lambert sum into a probability mass, and the independently-built `tsum_coprime_pair_pow_eq_tsum_totient_mul_pow` [GeometricCoprimality.lean:120] gives the same bridge from the totient generating function directly.

cyclotomic

The T1–T6 chain: `mobiusNumeratorPolynomial`, its coefficients (an exact divisor-counting formula, `mobiusNumeratorPolynomial_coeff` [RepunitMobiusNumerator.lean:217]), cyclotomic-value survival (`cyclotomicValue_dvd_baseMobiusShadow_den` [CyclotomicProjectionOfShadow.lean:389]), and prime-power jump dynamics. *Transport to mobius-mersenne (denominator growth)*: the upper-half prime-channel survival theorem `upperHalfChannel_product_dvd_den_of_scale_primeFactors_le` [MersenneShadowCyclotomicNoncollapse.lean:809] feeds directly into the exact denominator-growth formula `lcmHeight_scaledMobiusShadow_den_exact` [MersenneShadowDenom.lean:103] in the Möbius–Mersenne coordinate.

p-adic

Fixed and growing 2-adic valuation-unit dynamics: the adelic height/denominator-tax law, the tropical fixed-precision no-go, the signed-Hankel terminal-parity engine, the two-adic pulse-block construction. *Transport to farey*: the scalar-localisation complement-dvd law `scalarLocalization_complement_dvd` [AdelicHeightObstruction.lean:23] together with the Mersenne-height corollary `positiveRat_mersenne_height` [AdelicHeightObstruction.lean:103] ($(2^r \mid |x.\text{num}| \wedge x < 2/(2^n - 1) \Rightarrow 2^r(2^n - 1) < 2 \cdot x.\text{den})$) is exactly a denominator lower bound of the same shape the Farey coordinate proves directly.

farey

Unimodular denominator exclusion: `farey_gap` [GapFareyBound.lean:51] and the committed $K=240$ window certificate `gap_check_window_1_240_le_79639646646701375323355774875831053` [GapFareyBound.lean:225] giving $S \neq p/q$ for every $q \leq Q_0 \approx 7.96 \times 10^{34}$. *Transport to binary-digit*: the window bound is built directly from a committed 2^{240} -scale totient residue V for the window $(N, K) = (1, 240)$ a binary-digit object via `gap_check_window_1_240_first_failure` [GapFareyBound.lean:225], which shows the bound is sharp at exactly the mediant $b_K + d_K$.

lcm-orbit

The periodLcm/LCM-diagonal/LCM-cone family: `periodLcm` [CarrySurvivorExtinction.lean:515], `eq_prime_pow_of_not_dvd_periodLcm` [CertificateKernel.lean:18613], the multiplicative ray split, and the cone-flatness/cone-nonflat producers. *Transport to binary-digit*: the identity `lcmRayArithmeticLetter_eq_deltaTotient` [TotientActualLcmOrbitArithmetic.lean:298] and `diagonalWindowIncrement_eq_lcmRayArithmeticLetter` [TotientActualLcmOrbitArithmetic.lean:310] identifies the LCM-ray letter literally with a `deltaTotient/windowDiscrepancy` value, so every LCM-orbit statement is, term for term, a binary-digit statement in disguise.

first-harmonic/exponential-sum

Weyl-sum block cancellation over the totient window discrepancy: `irrational_totient_series_of_first_harmonic_norm_gap` [FirstHarmonicPivot.lean:83] and the unconditional companion at `exists_certifiedKill_of_first_harmonic_gap` [FirstHarmonicGap.lean:128] (any constant-saving cancellation, via $\cos(\pi/8) > 9/10$ plus pigeonhole, forces a finite kill certificate). *Transport to binary-digit*: by construction the exponential sum is literally $\sum_N \cos(2\pi \cdot (\text{windowDiscrepancy } h \ N \ L \bmod 2^L)/2^L)$ a first-harmonic statistic of the

binary-digit residue, so a norm-gap in this coordinate is definitionally a statement about `windowDiscrepancy`.

8.1 Obstructions are coordinate-relative: the binary-digit vs. Möbius–Mersenne case

The clearest instance of coordinate-relative obstruction in the whole #249 corpus is the fate of the classical Erdős (1948) near-integer digit method. That method needs, in essence, a finite automaton: a bounded amount of state carried forward from one digit block to the next, so that the value of a far-away digit block can be recovered from a bounded summary of everything before it. In the binary-digit coordinate this is exactly what fails, and it fails as a proved Lean theorem, not a heuristic remark: `balancedPulse_no_autonomous_decoder` [`GenericTailOrbitRigidity.lean:246`] shows that for a balanced-pulse family of radius m , any finite `State` type collapsing the family to a single autonomous summary must satisfy $|\text{State}| \geq \lfloor m/2 \rfloor + 2$ unbounded in m so no finite-state decoder can recover the shift r from its collapsed state ([`Lean`], scale:`uniform`, coordinate `binary-digit`). This is the formal shadow of the classical fact that Euler’s totient function has $\varphi(n) = \Theta(n)$ (average order $\frac{6}{\pi^2}n$, elementary; [`Math`]) rather than being bounded or sparse: the digit-window discrepancy `windowDiscrepancy` $h N L$ genuinely needs $\Omega(m)$ bits of state to track as N grows, which is exactly why #249’s own binary-digit engine is forced into a *cofinal certificate supply* (`irrational_totient_series_of_certificate_supply` [`TotientTailPeriodKiller.lean:394`]) rather than a single finite-state argument: the same shape of no-go recurs at `det_residualMonomialMatrix` [`ResidualG`] (locked gauge defeats residual-blind rank certificates for the first-harmonic pivot) and at `linearIndependent_canonicalTotientKernelFamily` [`TotientMahlerDefect.lean:900`] (the dyadic totient-kernel span is genuinely infinite-rank, not compressible to any fixed dimension).

The Möbius–Mersenne coordinate does not carry this obstruction, because it is built on a coefficient sequence, $\mu(d) \in \{-1, 0, 1\}$, that is bounded rather than growing with d . The squared-Lambert transfer engine `tsum_lambert_linear_weight_sq_pure` [`GcdMomentCalculus.lean:105`] takes any weight with $|w(d)| \leq d$ μ trivially qualifies with room to spare and converts it into a divisor-convolution power series for free; no finite-state decoder is ever needed because the relevant recursion (the cyclotomic T1–T6 chain: `mobiusNumeratorPolynomial_eq_gcdWord` [`RepunitMobiusNumerator.lean:2`] through `cyclotomic_dvd_primeJump_new_fibre` [`PrimePowerJumpDynamics.lean:281`]) is a finite, exact polynomial identity at every squarefree radical r , uniform in r , with no growth-driven state explosion. This is precisely why the corpus’s strongest *unconditional* denominator result—the exact-value and lower-bound formulas of `lcmHeight_scaledMobiusShadow_den_exact` [`MersenneShadowDenominatorGrowth.lean:147`] and `upperHalfMersenneProduct_lower_bound` [`MersenneShadow`]—lives entirely in the Möbius–Mersenne coordinate and is scale-`uniform`, while the sharpest *conditional* route in the binary-digit coordinate is scale-`cofinal` and unclosed. The two coordinates are not equivalent representations of the same difficulty: the wall each one hits is a fact about that coordinate’s own state-growth, not about #249 itself, which is exactly the AP17/AP18 discipline this corpus was built to enforce before reporting an obstruction, name the coordinate it was measured in, and test whether a re-representation removes it.

9 Interface index: what stands between the corpus and a proof

This section is the join layer of the paper. Parts I–II inventory what the corpus proves; this part inventories, obligation by obligation, exactly what is *missing* to turn each proved reduction into a proof of Erdős #249, in a form usable as a premise without opening the Lean tree. Erdős #249 asks whether $S = \sum_{n \geq 0} \varphi(n)/2^n$ is irrational. It is OPEN. Nothing in this section decides it. Every row below either (a) states a proved theorem exactly, or (b) states an unproved but well-posed proposition (an *open producer* or *open supply*) together with the exact proved consumer

that would turn it into irrationality of S .

9.1 Reading this index

Every entry follows the same discipline. **Yields**: the exact statement proved, with quantifiers in their proved order. **Exact mismatch**: precisely where the proved statement falls short of the open obligation it is compared against — never “it doesn’t work,” always the named axis of shortfall. **What would close it**: the exact remaining proposition, stated in full, whose proof (via the cited Lean consumer, already on disk) proves $\text{Irrational}(S)$.

Each numbered claim carries three tags. `scale:fixed` means proved at finitely many explicit numerals with no argument in the growing parameter; `scale:bounded` means proved on a bounded range of a parameter that is itself unbounded elsewhere in the statement; `scale:uniform` means proved for *every* value of the relevant parameter (the strongest positive tag short of matching the open target); `scale:cofinal` means the statement itself has the shape $\forall a_0 \exists a \geq a_0$, i.e. it is exactly the quantifier shape the open obligation needs; `scale:n/a` applies to structural non-existence results with no growing parameter. The evidence tags are `[Lean]`, `[Cert]`, `[Math]`, `[Cited]`, `[Open]`, used exactly as defined in Part I.

The canonical open obligation for #249, proved in Part II to be sufficient (`irrational_totient_series_of_certificate_supply` `[TotientTailPeriodKiller.lean]`), is the **249-supply** obligation:

$$\forall h \geq 1, \forall N_0, \exists N \geq N_0, \exists L, \text{certifiedKill } h \ N \ L.$$

Fourteen near-miss rows are catalogued against it below: one headline row (the sharpest reduction in the corpus, given its own subsection) and thirteen further rows, each attacking the same obligation from a different coordinate.

9.2 The headline: #249 reduces to a single unsupplied exponential-sum bound

The following two definitions, verified against the live Lean tree in this session, are the sharpest statement of the #249 frontier that the corpus contains.

Definition 9.1 (Window discrepancy). `windowDiscrepancy` `[TotientTailPeriodKiller.lean:66]`. For $h, N, L \in \mathbb{N}$,

$$A_{h,N,L} = \sum_{j=0}^{L-1} (\varphi(N+h+1+j) - \varphi(N+1+j)) \cdot 2^{L-1-j} \in \mathbb{Z}.$$

This is the depth- L truncation of $2^L \cdot (R_{N+h} - R_N)$, where $R_N = \sum_{j \geq 1} \varphi(N+j)/2^j$ is the local totient tail (`totientTail` `[TotientTailPeriodKiller.lean:57]`).

Definition 9.2 (Certified kill). `certifiedKill` `[TotientTailPeriodKiller.lean:72]`.

$$\text{certifiedKill } h \ N \ L :\Leftrightarrow (N+h+L+2) < A_{h,N,L} \bmod 2^L < 2^L - (N+h+L+2).$$

That is: the residue of $A_{h,N,L}$ modulo 2^L avoids the radius- $(N+h+L+2)$ neighbourhood of 0. `certifiedKill_depth_floor` `[TotientTailPeriodKiller.lean:79]` records the necessary room condition this forces: $2(N+h+L+2) < 2^L$.

Definition 9.3 (First-harmonic real and complex characters). `windowFirstCos` `[FirstHarmonicGap.lean:27]` and `windowFirstExp` `[FirstHarmonicPivot.lean:38]`:

$$\text{windowFirstCos } h \ N \ L = \cos\left(2\pi \frac{A_{h,N,L} \bmod 2^L}{2^L}\right), \quad \text{windowFirstExp } h \ N \ L = \exp\left(i 2\pi \frac{A_{h,N,L} \bmod 2^L}{2^L}\right).$$

`windowFirstExp_re` [FirstHarmonicPivot.lean:42] records $\text{Re}(\text{windowFirstExp}) = \text{windowFirstCos}$ and `norm_windowFirstExp` [FirstHarmonicPivot.lean:48] that $\|\text{windowFirstExp } h \ N \ L\| = 1$: this is literally the first additive character of the endpoint discrepancy modulo 2^L , a standard exponential-sum object.

Theorem 9.4 (Real-part certificate consumer). `exists_certifiedKill_of_first_harmonic_gap` [FirstHarmonicGap.lean:128]. [Lean]. *scale:uniform. coord:first-harmonic. For all h, X, L with $0 < X$ and the room condition $16(2X+h+L+2) \leq 2^L$, if*

$$\sum_{N=X}^{2X-1} \text{windowFirstCos } h \ N \ L \leq \frac{9}{10} X,$$

then $\exists N \in [X, 2X)$ with `certifiedKill` $h \ N \ L$.

Theorem 9.5 (Subset consumer — no density or partition hypothesis). `exists_certifiedKill_of_first_harmonic_gap_subset` [FirstHarmonicGap.lean:104]. [Lean]. *scale:uniform. coord:first-harmonic. For any nonempty finite $T \subseteq \mathbb{N}$ with $T \subset [0, 2X)$ and the same room condition, if*

$$\sum_{N \in T} \text{windowFirstCos } h \ N \ L \leq \frac{9}{10} |T|,$$

then $\exists N \in T$ with `certifiedKill` $h \ N \ L$. This is strictly stronger than Theorem 9.4: T can be any explicitly chosen nonempty finite subset of the dyadic block, not the whole block, and the proof (an averaging pigeonhole, §below) never uses that T has positive density or comes from a partition.

Theorem 9.6 (Complex norm consumer and the open producer). `exists_certifiedKill_of_first_harmonic_norm_gap` [FirstHarmonicPivot.lean:53] and `irrational_totient_series_of_first_harmonic_norm_gap` [FirstHarmonicPivot.lean:83]. [Lean] for both implications; the hypothesis `DTWFirstHarmonicNormGap` is [Open]. *The complex norm bound is strictly stronger than the real-part bound ($|z| \geq \text{Re}(z)$), and $21/25 < 9/10$ absorbs the slack), so it composes through Theorem 9.4 to the same certificate. Define*

`DTWFirstHarmonicNormGap` $:\Leftrightarrow \forall h > 0 \ \forall X_0 \ \exists X, L, \max(X_0, 1) \leq X \wedge 16(2X+h+L+2) \leq 2^L \wedge$

$$\left\| \sum_{N=X}^{2X-1} \text{windowFirstExp } h \ N \ L \right\| \leq \frac{21}{25} X.$$

coord:first-harmonic. scale:cofinal (this is the open target itself). Then

$$\text{DTWFirstHarmonicNormGap} \implies \text{Irrational} \left(\sum_{n \geq 0} \frac{\varphi(n)}{2^n} \right),$$

proved in full, with no further gap, by chaining Theorem 9.6 $\rightarrow$ Theorem 9.4 $\rightarrow$ `irrational_totient_series_of_certificate_supply` [TotientTailPeriodKiller.lean].

Observation 9.7 (The exact negative). No instance of `DTWFirstHarmonicNormGap`'s hypothesis, nor of `hgap` in Theorems 9.4–9.5, is proved anywhere in the corpus, at any h, X, L . [Open] is exact, not conservative rounding. The two consumer theorems that *use* the subset form (Theorem 9.5) are conditional consumers, not a supply: `AdjacentPhaseSeparation.lean` [AdjacentPhaseSeparation.lean] instantiates T as the explicit two-element set $\{N, M\}$ (apply `exists_certifiedKill_of_first_harmonic_gap_subset` ($\{N, M\} : \text{Finset } \mathbb{N}$)), and both call sites in `PivotAntiReconstruction.lean` and `PivotAntiReconstruction.lean` [PivotAntiReconstruction.lean:1541] likewise apply it to a finite explicit pair, never to an infinite or growing family. There is no theorem anywhere in the corpus that discharges `hgap` at a single instance, let alone cofinally.

Remark 9.8 (Why this is the sharpest reduction, and why it is tractable). This reduction converts the open producer for #249 from certificate-supply language — “exhibit a certified kill (N, L) ” — into a *constant-saving cancellation estimate for the first additive character of the totient window discrepancy over a dyadic block*: exactly the shape of a classical exponential-sum bound in analytic number theory (a Weyl-type estimate for $\sum_N e(\theta_N)$). Three features make this genuinely the narrowest gap in the corpus.

First, the saving required is a *constant*, not $o(X)$: $21/25$ (complex norm) or $9/10$ (real part), fixed independent of h, X, L . Any nontrivial cancellation estimate — even one far weaker than square-root cancellation — would suffice; this is not asking for GRH-strength input.

Second, the room condition $16(2X+h+L+2) \leq 2^L$ forces only $L \gtrsim \log_2 X + 5$, which is satisfiable for every X by simply taking L large enough (the depth floor grows logarithmically in X , not polynomially); no scale obstruction of the kind documented elsewhere in this paper (e.g. the $\sqrt{2\log_2 N}$ silent-position defect for #257, Part I) stands in the way of the producer’s own room requirement.

Third, the subset form (Theorem 9.5) removes even the requirement that the cancellation happen on the whole dyadic block or a positive-density subset of it: any single explicitly named finite T with the averaged bound would do. In particular a supplier-fibre or sparse-subsequence argument — e.g. picking N to be one less than a prime, as the exact pivot algebra in [FirstHarmonicPivot.lean](#) [FirstHarmonicPivot.lean] already sets up via `pivotArgument`, `pivotPrime`, `pivotSupplierBases` — is a legitimate route to this obligation and does not need to control the discrepancy on a full interval.

9.3 Row-by-row: the remaining near misses against 249-supply

The headline (§9.2) is row **e1-companion** of the source interface catalogue in full. The remaining thirteen rows below attack the same 249-supply obligation from independent coordinates in the Lean tree: the actual-LCM sign corridor, the top-edge staircase, the raw rational approximant, the short-window arithmetic word, the diagonal pincer certificate bank, the h -uniform single-window certificate, the Farey/continued-fraction denominator growth law, the 2-adic pulse block, the LCM-jump slack scalar, the prime-jump commutator, the Möbius–Mersenne denominator channel, and the carry-kernel rank. None of them is closed; each is recorded with its exact remaining content.

9.3.1 SGN-01 — the positive-sign half of the certified-kill band (coord:actual-lcm-sign)

Definition 9.9. The *actual LCM tail orbit* at exponent a is `actualLcmTailOrbit` $a = R_{2H+2H} - R_{H+\dots}$, precisely `totientTail(2H) - totientTail(H)` where $H = \text{periodLcm}(2^a)$ ([actualLcmTailOrbit_pos](#) [TotientActualLcmOrbitSign.lean:144]).

Theorem 9.10. [actualLcmTailDiff_shift_pos](#) [TotientActualLcmOrbitSign.lean:39] and [actualLcmTailOrbit_pos](#) [TotientActualLcmOrbitSign.lean:144]. [Lean]. *scale:uniform. For every $a \geq 8$ and every J with $J + (a+6) < 2 \cdot 2^a$,*

$$0 < \text{totientTail}(2H+J) - \text{totientTail}(H+J), \quad H = \text{periodLcm}(2^a).$$

Hence $0 < \text{actualLcmTailOrbit } a$ for every $a \geq 8$. The proof is genuinely uniform in a : it rests on two facts proved for all $a \geq 8$ by a two-case structural split (divisor letter vs. foreign prime power), with absolute constants 4, 8, 32 and no lookup table — [lcmRayArithmeticLetter_pos_of_lt_two_mul](#) [TotientActualLcmOrbitArithmetic.lean:1703] and [periodLcm_lt_eight_mul_t_mul_lcmRayArithmeticLetter](#) [TotientActualLcmOrbitArithmetic.lean:1906].

Remark 9.11 (Exactly half the certificate band, and where the other half goes). `certifiedKill` needs the residue to avoid a *symmetric* radius- $(N+h+L+2)$ neighbourhood of 0 on both sides. SGN-01 gives positivity only. Its companion `actualLcm_integral_forces_topEdgeResidue` [TotientActualLcm0] proves that, under integrality, the consequence is not a kill but the opposite: the residue is forced to the *top edge*, exactly $2^K - e$ where e is the true carry orbit (`carryOrbit` []). So the lower half of the certified-kill band is discharged unconditionally and cofinally by SGN-01; the upper half is untouched, and what remains open is a genuinely different, one-sided statement about the top edge — row TE-04 below. The corpus’s own normal-form tables tag this row `scale:bounded`; that tag is misleading, since only the window offset J is bounded relative to the height, and the height $2 \cdot 2^a$ itself is unbounded: the correct tag is `scale:uniform`.

9.3.2 TE-04 — the one-sided top-edge staircase (coord:actual-lcm-top-edge)

This is the sharpest *landed weakening* of the 249-supply obligation in the corpus: the symmetric certified-kill band is replaced by a single upper inequality, halving the information the open producer must supply.

Theorem 9.12 (Consumers, fully proved). `actualLcmTailDiff_notMem_int_of_topEdgeResidueGap` [TotientActualLcmTailDiff_notMem_int_of_topEdgeResidueGap] [TotientActualLcmTopEdgeStaircase.lean:1314], `irrational_totientSeries_of_topEdgeResidueGapSupply` [TotientActualLcmTopEdgeStaircase.lean:2184]. [Lean]. *scale:uniform*. For every $a \geq 8$ and every J, K, m inside the sign corridor $(J+K+(a+6) < 2 \cdot 2^a)$, a one-sided residue gap at precision $m \leq K$ — room $2H+J+K+2 < 2^m$ and $A_{H,H+J,K} \bmod 2^m \leq 2^m - (2H+J+K+2)$ — already forces `totientTail(2H+J) - totientTail(H+J)  $\notin \mathbb{Z}$` . No lower margin at all is demanded. The proof chain is complete: the theorem holds for every $a \geq 8$, and `PowerTwoActualLcmTopEdgeResidueGapSupply` $\Rightarrow$ `Irrational S` is proved.

Definition 9.13 (The open producer). `PowerTwoActualLcmTopEdgeResidueGapSupply` [TotientActualLcmTopEdgeResidueGapSupply] [Open]. *scale:cofinal*.

$\forall a_0 \exists a, K, m, a_0 \leq a \wedge 8 \leq a \wedge K+(a+6) < 2 \cdot 2^a \wedge \text{ActualLcmTopEdgeResidueGap } a \ 0 \ K \ m.$

No unconditional statement anywhere in the corpus locates the residue of the diagonal word $A_{H,H,K}$ (where $H = \text{periodLcm}(2^a)$) below the top strip at even one large a . `actualLcmTopEdgeResidueGap_iff_terminal` [TotientActualLcmTopEdgeStaircase.lean:909] further reduces the condition to the last m arithmetic letters of the word alone.

Proposition 9.14 (The five strictly weaker sufficient links — proving any one closes #249). All five are proved sufficient for `PowerTwoActualLcmTopEdgeResidueGapSupply` [TotientActualLcmTopEdgeStaircase.lean:2184] in the same module, and each is strictly weaker in the sense that it drops a hypothesis, widens a band, or asks only for magnitude rather than a signed inequality.

1. `PowerTwoAdjacentSuffixMidbandSupply` [TotientActualLcmTopEdgeStaircase.lean:1334] ([Open], *scale:cofinal*):

$$\forall a_0 \exists a, m, a_0 \leq a \wedge 8 \leq a \wedge m+1+(a+6) < 2 \cdot 2^a \wedge 2H+m+3 < 2^m \wedge$$

$$2H+m+2 \leq \text{diagonalAdjacentSuffixResidue}(2^a) \ 0 \ m \leq 2^m - (2H+m+2)$$

(a two-sided band on the adjacent-suffix residue directly, one candidate depth m).

2. `PowerTwoOddGuardTopEdgeHalfWordBandSupply` [TotientActualLcmTopEdgeStaircase.lean:1349] ([Open], *scale:cofinal*): at the odd guarded depth $2q+1$, a two-sided band of half-width $H+q+2$ on the half-word residue $\bmod 4^q$ — “substantially weaker than the older fixed $1/32$ central band.”

3. [PowerTwoActualFinalTopEdgeMagnitudeSupply](#) [TotientActualLcmTopEdgeStaircase.lean:1471] ([Open], scale:cofinal), proved equivalent to the previous one via [topEdgeHalfWordBandSupply_iff_actualFinalCenteredMagnitudeSupply](#) [TotientActualLcmTopEdgeStaircase.lean:1479]:

$$\forall a_0 \exists a, q, \max(14, a_0) \leq a \wedge \text{oddGuardedCanonicalAdjacentSuffixDepth}(2^a) = 2q+1 \wedge H+q+2 \leq |\text{actual}0$$

4. [PowerTwoFlexibleActualTopEdgeMagnitudeSupply](#) [TotientActualLcmTopEdgeStaircase.lean:1927] ([Open], scale:cofinal): the same magnitude bound with the depth restriction relaxed from “canonical guarded” to any odd $2q+1$ satisfying the half-cell fit $2(H+q+2) \leq 4^q$ and the sign-corridor room $2q+2+(a+6) < 2 \cdot 2^a$.
5. [PowerTwoFlexibleActualTerminalDominanceSupply](#) [TotientActualLcmTopEdgeStaircase.lean:1908] — the weakest of all five; treated separately as row TE-05-weakest below.

Proving any one of these five closes Erdős #249; the corpus has already proved all the implications from each to [PowerTwoActualLcmTopEdgeResidueGapSupply](#) [TotientActualLcmTopEdgeStaircase.lean:1325] (chain: midband $\rightarrow$ residue-gap at [TotientActualLcmTopEdgeStaircase.lean:2109]; half-word-band $\rightarrow$ midband at [TotientActualLcmTopEdgeStaircase.lean:2058]; final-magnitude $\Leftrightarrow$ half-word-band at [TotientActualLcmTopEdgeStaircase.lean:1479]; flexible-magnitude $\rightarrow$ midband at [TotientActualLcmTopEdgeStaircase.lean:1993]).

9.3.3 TE-05-weakest — one scalar inequality at cofinally many odd ranks (coord:actual-lcm-top-edge)

This is the weakest landed link in the whole #249 sufficiency lattice: a single comparison between one terminal arithmetic letter and twice a centred lift.

Definition 9.15. [PowerTwoFlexibleActualTerminalDominanceSupply](#) [TotientActualLcmTopEdgeStaircase.lean:1908] ([Open]. scale:cofinal. Writing $H = \text{periodLcm}(2^a)$,

$$\forall a_0 \exists a, q, a_0 \leq a \wedge 8 \leq a \wedge 2q+2+(a+6) < 2 \cdot 2^a \wedge 2(H+q+2) \leq 4^q \wedge$$

$$\text{diagonalWindowIncrement}(2^a)(2q+2) \leq 2 \cdot \text{actualOddHalfCenteredLift } a \ q.$$

Theorem 9.16 (Consumer). [actualLcmTailOrbit_notMem_int_of_actualTerminalDominance](#) [TotientActualLcmTopEdgeStaircase.lean:1908] and [irrational_totientSeries_of_flexibleActualTerminalDominanceSupply](#) [TotientActualLcmTopEdgeStaircase.lean:1908] [Lean]. scale:uniform. The dominance hypothesis at a single odd rank already excludes integrality of $\text{actualLcmTailOrbit } a$, and the supply predicate composes to $\text{Irrational}(S)$.

Observation 9.17 (The exact identity, and the internal tension it exposes). [two_mul_actualOddHalfCenteredLift_eq_terminal_sub_trueCarry](#) [TotientActualLcmTopEdgeStaircase.lean:1678]. [Lean]. scale:uniform. For $a \geq 8$, room $2q+1+1+(a+6) < 2 \cdot 2^a$, half-cell fit $2(H+q+2) \leq 4^q$, and any integral representative z with $z = \text{actualLcmTailOrbit } a$:

$$2 \cdot \text{actualOddHalfCenteredLift } a \ q = \text{diagonalWindowIncrement}(2^a)(2q+2) - \text{carryOrbit } H \ H \ z \ (2q+1).$$

This is an *exact equality*, not a bound. Comparing it against the dominance inequality above, the dominance inequality is literally equivalent to $\text{carryOrbit } H \ H \ z \ (2q+1) \leq 0$. But SGN-02 ([actualLcm_trueEndpointSurvivor_neg](#) [TotientActualLcmOrbitSign.lean:172], [Lean], scale:uniform) proves that under integrality the true carry orbit is *strictly positive* throughout this exact corridor: for $a \geq 8$ and $J+K+(a+6) < 2 \cdot 2^a$,

$$0 < \text{carryOrbit } H \ (H+J) \ d \ K$$

whenever d is the real-valued representative of the translated tail difference. So the two branches of the corridor-escape disjunction that [actualLcmTailOrbit_notMem_int_of_actualTerminalCarryCorridor](#)

offers are *not* symmetric: SGN-02 has already eliminated the branch that the identity most naturally supplies (dominance, $\text{carryOrbit} \leq 0$), and the surviving branch is the *lower* escape

$$2 \cdot \text{actualOddHalfCenteredLift } a \ q \leq \text{diagonalWindowIncrement}(2^a)(2q+2) - (2H+2q+3).$$

This is stated precisely so the reader can check it: it is neither a refutation of the dominance route (SGN-02 constrains the sign of carryOrbit , it does not itself bound $\text{diagonalWindowIncrement}$ or $\text{actualOddHalfCenteredLift}$ against 0, so dominance is not shown *false*, only shown to entail a carry-orbit sign the census never violates) nor a proof (no theorem excludes dominance outright). It is the precise reason the finite census keeps landing just inside the corridor rather than outside it, and it identifies the lower-escape branch as the coordinate where a producer is actually needed.

Proposition 9.18 (What would close it). *Either (i) the lower-escape branch cofinally — as displayed just above — or (ii) the two-sided magnitude form [PowerTwoFlexibleActualTopEdgeMagnitudeSupply](#) [item 4 of Proposition 9.14], which asks only $H+q+2 \leq |\text{actualOddHalfCenteredLift } a \ q|$ and which the file proves ([flexibleActualTerminalCarryCorridorEscapeSupply_of_magnitude](#) [TotientActualLcmOrbitSeparation.lean:141] implies corridor escape via a clean sign split (positive branch escapes above the terminal letter, negative branch escapes below the directed bound), sidestepping the SGN-02 tension entirely because it does not commit to a sign for the centred lift.*

9.3.4 SEP-02 — separation from an explicit rational approximant (coord:raw-approximant)

Theorem 9.19. [abs_actualLcmTailOrbit_sub_rawApprox_lt](#) [TotientActualLcmOrbitSeparation.lean:141] and [irrational_totientSeries_of_actualLcmOrbitSeparationSupply](#) [TotientActualLcmOrbitSeparation.lean:141] [Lean]. scale:uniform. Unconditionally, for every a and q ,

$$|\text{actualLcmTailOrbit } a - \text{actualLcmRawApprox } a \ q| < \frac{4H + 2(2q+1) + 4}{2^{2q+2}},$$

where $\text{actualLcmRawApprox } a \ q$ is an explicit finite computable rational block. The error radius is uniform in a and q and decays like $H/4^q$: the analytic tail is fully discharged.

Remark 9.20. Consequently a cofinal $1/32$ -separation of the raw approximant from every integer suffices for #249. The corpus verifies the analogous kill *unconditionally* at exactly two exponents, $a = 4$ and $a = 6$ ([actualLcmTailOrbit_four_and_six_notMem_int](#) [TotientActualLcmShortKill.lean:35], [Cert], scale:fixed, via [certifiedKill_diagonal_t16](#) [] and [certifiedKill_diagonal_t64](#) [DiagonalPincerCertifiedKill.lean:35]). Nothing beyond $a = 6$ is proved. The depth q is not free: the supply predicate pins $2q+1 = \text{oddGuardedCanonicalAdjacentSuffixDepth}(2^a)$, so per scale a there is exactly one admissible depth, not a search over depths.

Definition 9.21 (What would close it). $\text{PowerTwoActualLcmOrbitSeparationSupply}$: [Open], scale:cofinal.

$$\forall a_0 \ \exists a \geq \max(2, a_0) \ \exists q, \ \text{oddGuardedCanonicalAdjacentSuffixDepth}(2^a) = 2q+1 \wedge$$

$$\forall z \in \mathbb{Z}, \ \frac{1}{32} + \text{actualLcmRawErrorRadius } a \ q \leq |\text{actualLcmTailOrbit } a - z|.$$

Since the error radius is explicit, this reduces to a distance-to-nearest-integer lower bound for one explicitly computable rational per scale — the cleanest reduction of the #249 obligation to a purely finite, computable question in the whole corpus.

9.3.5 SK-02 — the short-window arithmetic kill, verbatim, truncated at $a_0 \leq 6$ (coord:short-window-arithmetic)

Theorem 9.22. [powerTwoActualLcmShortArithmeticKillSupply_through_six](#) [TotientActualLcmShortKill.lean:63] and [irrational_totientSeries_of_shortArithmeticKillSupply](#) [TotientActualLcmShortKill.lean:63]. [Cert] for the witness, [Lean] for the consumer. scale:bounded.

$$\forall a_0 \leq 6, \exists a, L, a_0 \leq a \wedge L < 2 \cdot 2^a \wedge \text{LcmDiagonalArithmeticKill}(2^a) L$$

— literally the open cofinal supply predicate below with its universal quantifier truncated at $a_0 \leq 6$. This is the purest scale:fixed-vs-scale:cofinal row in the corpus: the proof term is 6, 93, ha, by norm_num, lcmDiagonalArithmeticKill_two_pow_six — a single hard-coded witness $(a, L) = (6, 93)$, itself discharged from [certifiedKill_diagonal_t64](#) [DiagonalPincerCertificateT64Endpoint.lean:1928] (a norm_num evaluation over an explicit table of φ values). There is no argument in a whatsoever; removing the bound $a_0 \leq 6$ requires an entirely new proof, not a re-run.

Definition 9.23 (What would close it). PowerTwoActualLcmShortArithmeticKillSupply: [Open], scale:cofinal.

$$\forall a_0 \exists a, L, a_0 \leq a \wedge L < 2 \cdot 2^a \wedge \text{LcmDiagonalArithmeticKill}(2^a) L.$$

The short-window restriction $L < 2 \cdot 2^a$ buys extra structure — every non-divisor offset in that window is a bare prime power, by eq_prime_pow_of_not_dvd_periodLcm — so this is a better-equipped target than the raw SEP-02 supply even though it is formally a stronger statement (it implies certified-kill directly, without the separation-and-round step).

9.3.6 b11 — the diagonal pincer certificate bank (coord:diagonal-pincer)

Theorem 9.24. [certifiedKill_diagonal_all_imported](#) [DiagonalPincerCertificates.lean:2860], with [diagonalPincerCertificateScales](#) [DiagonalPincerCertificates.lean:2842] and [diagonalPincerKillDepth](#) [Cert]. scale:fixed.

$$\forall t \in \{1, 2, 3, 4, 5, 7, 8, 9, 11, 13, 16, 17\}, \exists L, \text{certifiedKill}(\text{periodLcm } t)(\text{periodLcm } t) L$$

at depths $\{6, 5, 7, 7, 9, 14, 15, 14, 21, 22, 23, 26\}$ respectively, extended by the separate T19...T64 modules to 28 values through $t = 64$ (the $t = 64$ endpoint is [certifiedKill_diagonal_t64](#) [DiagonalPincerCertificateT64Endpoint.lean:1928]). Every witness is a norm_num evaluation over a hard-coded block of φ values — e.g. the $t = 17$ certificate lists $\varphi(12252241), \dots, \varphi(24504506)$ explicitly — so nothing in the proof is a function of t .

Remark 9.25 (A stronger signal in the depth table than the theorem states). [certifiedKill_depth_floor](#) [TotientTailPeriodKiller.lean:79] forces $2(2H_t + L + 2) < 2^L$, i.e. $L \gtrsim \log_2(4 \cdot \text{periodLcm } t)$ at any certified depth. Comparing that floor against diagonalPincerKillDepth shows every landed certificate fires within a small additive constant of the theoretical minimum depth, at every tested scale — a numerically supported ([Cert], not [Math]) anti-concentration observation, not a theorem.

Proposition 9.26 (What would close it).

$$\exists C \forall t_0 \exists t \geq t_0 \exists L \leq \log_2(4 \cdot \text{periodLcm } t) + C, \text{certifiedKill}(\text{periodLcm } t)(\text{periodLcm } t) L.$$

This is an anti-concentration statement about the diagonal word's residue at the first admissible depth — exactly the shape a doubling-orbit equidistribution argument would produce.

9.3.7 a12 — one window, sixteen periods, wrong-way quantifiers (coord:h-uniform-certificate)

Theorem 9.27. [certifiedKill_all_upto_sixteen](#) [CertificateKernel.lean:18556] and [totient_series_ne_rat_of_den_dvd_pow_two_mul_mersenne_upto_sixteen](#) [CertificateKernel.lean:18562] (shallower sibling [certifiedKill_all_small](#) [TotientTailPeriodKiller.lean:404]). [Cert]. scale:fixed.

$$\forall h \in [1, 16], \quad \text{certifiedKill } h \ 14 \ 9$$

— a single position $N = 14$ and a single depth $L = 9$ that simultaneously certify every period h up to 16. Consequence: $S \neq r$ for every rational r with $r.\text{den} \mid 2^{14}(2^h - 1)$, $1 \leq h \leq 16$. Proof is *decide* on a 9-bit window; no part of it is a function of N or of the h -range.

Remark 9.28 (Quantifier inversion — the only h -uniform row in the corpus). The 249-supply obligation is $\forall h \forall N_0 \exists N \geq N_0 \exists L$. This row proves $\exists N \exists L \forall h \in [1, 16]$: the quantifiers are inverted, and the h -range is finite. The inversion helps in one direction (one (N, L) covers a whole block of periods, more than the obligation asks) and hurts in the other (N is pinned at 14, not cofinal). This is the only row in the corpus with h -uniformity, and it is the reason a scale-uniform version would be unusually strong.

Proposition 9.29 (What would close it). A scale-uniform version of the same shape: $\exists f : \mathbb{N} \rightarrow \mathbb{N}$ with $f(N) \rightarrow \infty$ such that $\forall N_0 \exists N \geq N_0 \exists L$ with [certifiedKill](#) $h \ N \ L$ for all $h \leq f(N)$. This implies the obligation immediately (fix h , take N_0 large enough that $f(N) \geq h$) and, unlike the obligation, is a statement about one window at a time rather than a per-period search.

9.3.8 c3 — the Farey denominator floor and its stalled growth law (coord:farey-window)

Theorem 9.30. [tsum_totient_div_pow_two_ne_ratCast_of_den_le_79639646646701375323355774875831053](#) built from the classical mediant lemma [farey_gap](#) [GapFareyBound.lean:51] and the window sharpness certificate [gap_check_window_1_240_le_79639646646701375323355774875831053](#) [GapFareyBound.lean:17] / [gap_check_window_1_240_first_failure](#) [GapFareyBound.lean:225]. [Lean] for the general mediant lemma; [Cert] for the $K = 240$ window instantiation. scale:fixed. If S is rational, its reduced denominator exceeds $7.9639646646701375323355774875831053 \times 10^{34}$ (the exact numeral in the declaration name). This is the strongest unconditional statement about S in the corpus, logically independent of the certificate-supply reduction.

Remark 9.31 (Why re-running the same window buys nothing). The cofinal upgrade of this statement — $S \neq p/q$ for every bound q , not just $q = 7.96 \times 10^{34}$ — is literally Irrational(S), so the target coordinate is right. What blocks promotion is proved on disk: [gap_check_window_1_240_first_failure](#) [GapFareyBound.lean:225] establishes that the $K = 240$ window bound is *sharp*, at exactly the mediant $b+d$, so re-running the same argument at the same window buys nothing further. Each new K needs (i) a freshly committed 2^K -scale totient residue V_K and (ii) fresh continued-fraction convergents of $V'_K/2^K$, both hard-coded numerals in the current proofs. The growth of the bound b_K+d_K is governed by the convergent denominators of the underlying constant, which stall at q_0 precisely if $S = a/q_0$: the growth law is equivalent in difficulty to #249 itself.

Observation 9.32 (A closed sub-route). [ReducedDenominatorUnitGapCert](#) [PrimitiveWeightCertificate.lean:30] and [PrimitiveReducedDenominatorUnitGapSupply](#) [PrimitiveWeightCertificate.lean:47], the corpus's own attempt at strengthening this window refinement via a unit-modulo-odd-part criterion, prove a hard ceiling in [reducedDenominatorUnitGapCert_prime_pow_iff](#) [PrimitiveWeightCertificate.lean:110] and [reducedDenominatorCandidateCount_prime_pow_le_one](#) [PrimitiveWeightCertificate.lean:111] ([Lean], scale:n/a): at a prime-power denominator, the strengthening rescues at most *one* extra lattice point. That route is closed — it cannot be the source of a growth law.

Proposition 9.33 (What would close it). *A proved growth law for the window family: a function g with $g(K) \rightarrow \infty$ and a proof that for every K the $(N=1, K)$ gap check passes for all $q \leq g(K)$. Equivalently, a lower bound on the convergent denominators of the totient-window constant, uniform in K .*

9.3.9 TA-01/TA-02 — the two-adic pulse block reaches the arc centre and fails by an exponential (coord:two-adic-pulse)

Theorem 9.34. *`exists_prime_totient_twoAdic_pulse_divisors` [TotientTwoAdicPulseBlock.lean:54], `exists_prime_deltaTotient_twoAdic_pulseBlock` [TotientTwoAdicPulseBlock.lean:211], `windowDiscrepancy_modEq_half_of_twoAdic_pulse` [TotientTwoAdicPulseBlock.lean:262], `eventual_integral_tailDiff_has_cofinal_twoAdic_half_pulse` [TotientTwoAdicPulseBlock.lean:327]. [Lean]. scale:uniform (the theorem is unconditional and holds for every K, H, B , not merely cofinally many). For every $K \geq 2$, every $H > K$, and every bound B , there are primes $p > B$ with a length- $(K-1)$ zero prefix and a terminal half-turn, giving*

$$A_{H,p-K,K} \equiv 2^{K-1} \pmod{2^K}.$$

Under eventual integrality this transfers to an integer z with $(z : \mathbb{R}) = \text{totientTail}(p+H) - \text{totientTail}(p)$ and $z \equiv 2^{K-1} \pmod{2^K}$.

Remark 9.35 (Lands at the exact centre of the arc, fails by an exponential — and why). This lands the residue at the exact centre of the certified-kill arc: 2^{K-1} is maximally far from 0 modulo 2^K , precisely the coordinate the obligation wants. Taking $N=p-K$, $h=H$, $L=K$ gives radius $N+h+L+2 = p+H+2$, so `certifiedKill` requires $2^{K-1} > p+H+2$. But $z \equiv 2^{K-1} \pmod{2^K}$ forces $|z| \geq 2^{K-1}$, while the directed tail bound forces $|z| < p+H+2$; and the construction's own congruence $p \equiv 1+2^{K-1} \pmod{2^K}$ (equivalently $v_2(p-1) = K-1$) forces $p \geq 1+2^{K-1}$. So $p+H+2 > 2^{K-1}$ always, and no choice of p rescues it. The quantifier order is inverted: the theorem gives, for fixed K , cofinally many *large* p ; the certificate needs a p *small* relative to 2^{K-1} , impossible for a single letter since $|\varphi(N+H) - \varphi(N)| < N+H$. Depth-promotion is not the issue: it has already been performed, generalizing an exponent-2 construction to every K uniformly by CRT-gluing over a $\text{Unit} \oplus \text{Fin}(K-1) \oplus \text{Fin}(K-1)$ family.

Proposition 9.36 (What would close it). *The same half-turn residue realised by an accumulated window rather than a single terminal letter: cofinally many (h, N, L) with $A_{h,N,L} \equiv 2^{L-1} \pmod{2^L}$ and $2^{L-1} > N+h+L+2$. Only the weighted sum $\sum \Delta\varphi \cdot 2^{L-1-j}$ can carry magnitude 2^L ; the per-letter pulse provably cannot. Concretely: a pulse-block construction whose zero prefix and half-turn are imposed on the word, not on one delta.*

9.3.10 d-3a/d-3b — cofinal jump positions, one unproved slack sign (coord:lcm-jump-slack)

Theorem 9.37 (Cofinal producer, fully proved). *`exists_periodLcm_strict_jump_ge` [DiagonalFreshLossBridge.lean:248]. scale:cofinal (proved, not open): $\forall t_0 \exists t \geq t_0$ with `periodLcm` $t < \text{periodLcm}(t+1)$, witnessed by $p-1$ for any prime $p > t_0$. The power-of-two specialization `CanonicalAdjacentSuffixPowerTwoPost` shows the positions $t = 2^a - 1$ already suffice, so no position search is needed at all.*

Definition 9.38 (The slack scalar and the open supply). *`canonicalAdjacentSuffixCentralSlack` [DiagonalFreshLossBridge.lean:249]*

$$\text{canonicalAdjacentSuffixCentralSlack } t = \min(d - 2^{m-5}, (2^m - 2^{m-5}) - d),$$

with m the canonical adjacent-suffix depth and d the residue at that depth. The open producers, both [Open] scale:cofinal, are `CanonicalAdjacentSuffixPostJumpSlackSupply` [DiagonalFreshLossBridge.lean:249] (any strict jump) and `CanonicalAdjacentSuffixPowerTwoPostJumpSlackSupply` [DiagonalFreshLossBridge.lean:250]

(power-of-two positions), each asking $0 \leq \text{canonicalAdjacentSuffixCentralSlack}(t+1)$ cofinally, and both compose to $\text{Irrational}(S)$ at `irrational_totientSeries_of_canonicalAdjacentSuffixPostJ` and `irrational_totientSeries_of_canonicalAdjacentSuffixPowerTwoPostJumpSlackSupply` [DiagonalFre

Remark 9.39 (Census evidence only). [Cert], scale:fixed. All 40 strict jumps up to endpoint 113 pass, closest margin ≈ 0.000221 of the modulus at $t = 100$; the power-of-two endpoints 4, 8, 16, 32 pass with slack fractions 0.41, 0.036, 0.40, 0.19. Census, not theorem. The structural handle nobody has used: `periodLcm_succ_eq_prime_mul_of_strict_jump` (`periodLcm_succ_eq_prime_mul_of_strict_jump` [DiagonalFreshLossBridge.lean:2134]) says a strict jump at t means $t+1 = p^k$ and $\text{periodLcm}(t+1) = p \cdot \text{periodLcm}(t)$ — so the required statement is a transfer lemma for how the adjacent-suffix residue at height H moves under $H \mapsto pH$. `canonicalAdjacentSuffixCentralSlack_eq_of_periodLcm_eq` [DiagonalFreshLossBridge.lean:2460] already proves the slack is constant across each LCM plateau, so only jump indices matter.

Proposition 9.40 (What would close it). $0 \leq \text{canonicalAdjacentSuffixCentralSlack}(2^a)$ for cofinally many a .

9.3.11 e2 — the prime-jump commutator, one fixed witness (coord:prime-jump-commutator)

Theorem 9.41. `primeJumpSharpRadius` [PrimeJumpWindow.lean:37] and `primeJumpTailCommutator_notMem_int_of_sharpKill` [PrimeJumpWindow.lean:178]. [Lean]. scale:uniform. Uniform consumer for all H, p, L : a residue of the four-vertex commutator $J(H, p)$ outside the sharp radius $3pH + (p+1)(L+2)$ forces $J(H, p) \notin \mathbb{Z}$. This is tighter than the earlier $4pH$ two-cell disjunction.

Theorem 9.42 (The one witness). `primeJumpSharpKill_twelve_five` [PrimeJumpWindow.lean:186]. [Cert]. scale:fixed. `primeJumpSharpKill 12 5 15` (i.e. $H = \text{periodLcm } 4 = 12$, $p = 5$, $L = 15$), proved by *decide* on an explicit 15-bit window with no dependence on t or p ; only a single instance exists, at $t = 4$, the smallest nontrivial height. The full endpoint composition is `irrational_totient_series_of_primeJumpSharpKill_supply` [PrimeJumpWindow.lean:193].

Remark 9.43. The supply hypothesis $\forall t_0 \exists t \geq t_0 \exists p, L > 0$ with `primeJumpSharpKill`(`periodLcm` t) p L is a genuinely cheaper target than the raw SEP-02 supply: it asks for one fresh prime p per LCM height rather than a full central-arc certificate. Because the consumer route is `rational_totient_series_forces_lcm_cone_flatness` → contradiction, the natural attack is to pick p as the fresh prime introduced at the next strict LCM jump (linking this row to d-3a), where the commutator’s old-channel contributions are annihilated by `oldChannel_affine_moment_annihilation` [JointExponentTransport.lean:36].

9.3.12 b7 — the Möbius–Mersenne denominator channel: unbounded, but off-coordinate (coord:mobius-mersenne)

Theorem 9.44. `upperHalfMersenneProduct_lower_bound` [MersenneShadowDenominatorGrowth.lean:60], `lcmHeight_scaledMobiusShadow_den_lower_bound` [MersenneShadowDenominatorGrowth.lean:85], `exists_upperHalf_mersenne_channel_dvd_den_with_bounds` [MersenneShadowDenominatorGrowth.lean:99]. [Lean]. scale:uniform. For every $t \geq 5$, uniformly in t :

$$2^{t/2} \leq \prod_{p \in \text{upperHalfPrimes}(t)} \text{mersenne}(p) \leq \text{den}(\text{lcmHeight}(t) \cdot \text{numericMobiusShadow}(\text{lcmHeight}(t))).$$

An unbounded denominator lower bound at every LCM height, with an individual surviving channel isolated ($2^{t/2} \leq \text{mersenne}(p) < 2^t$). The proof is genuinely uniform in t (Bertrand’s postulate via `upperHalfPrimes_nonempty`, plus $2^{p-1} \leq 2^p - 1$ factorwise — no table, no constant degrading with t).

Remark 9.45 (The corpus’s only proved unbounded-growth quantity in this coordinate, but the wrong coordinate for #249). This lives in a different coordinate from the 249-supply obligation: it bounds the reduced denominator of the *scaled* shadow at LCM height t , and there is no landed transport from a denominator lower bound to `certifiedKill` or to a totient-tail non-integrality. Two further honest caveats are recorded in the module’s own docstring: it does not rule out cancellation by the foreign-defect term, and the growth rate $2^{t/2}$ is far below the scale $\text{lcmHeight}(t) \approx 2^{1.44t}$ against which the enclosure consumers measure error. The existing consumers built on it (`scaleFullTarget_miss_of_lambert_projected_separation`, `scaleFullTarget_miss_of_lambert_projected_num_gap`) are #257-facing (`ScaleFullTargetHit`), not #249-facing.

Proposition 9.46 (What would close it). *Either (i) a Dirichlet-criterion bridge in the shape of `irrational_of_den_mul_abs_sub_tendsto_zero` [`CertificateKernel.lean:5371`]: a sequence of rationals u_t with $u_t \neq S$ and $\text{den}(u_t) \cdot |S - u_t| \rightarrow 0$, which requires the approximation error to beat the denominator, not merely the denominator to grow; or (ii) a transport map from “channel of size $\geq 2^{t/2}$ survives in the denominator” to “`windowDiscrepancy` residue avoids the arc,” which does not exist in the corpus. Route (i) needs the growth rate raised from $2^{t/2}$ to beat the shadow’s own truncation error.*

9.3.13 d4/d5 — the parallel rank obligation, and a corrected pair of citations (`coord:carry-rank`)

This row runs a second, independent open obligation in parallel to 249-supply, in the carry-kernel *rank* coordinate rather than the residue coordinate. **Correction:** the source index attributed the two theorems below to swapped files; direct reads this session confirm the correct attribution is as given.

Theorem 9.47. `linearIndependent_canonicalTotientKernelFamily` [`TotientMahlerDefect.lean:900`]. [`Lean`]. *scale:uniform. Unconditional and uniform in e : the dyadic totient-kernel family is linearly independent at every depth e (dimension exactly 2^e+1 , via CRT and Dirichlet on primes in arithmetic progression).*

Theorem 9.48. `not_irrational_totientSeries_implies_unbounded_carryRank_unconditional` [`TotientMahlerDefect.lean:900`]. [`Lean`]. *scale:uniform. If S is rational there is a tempered integral carry orbit u with*

$$2^e - 1 \leq \text{finrank}_{\mathbb{Q}} \text{span}(\text{canonicalCarryKernelFamily } u \ e) \quad \text{for every } e.$$

Remark 9.49 (The scale side is finished; the missing direction is proved dead on one route). So: rationality forces *unbounded* carry-kernel rank, and the 2^e-1 floor holds for all e with a uniform proof. What is missing is the opposite inequality — a rationality-side rank upper bound, which would contradict the floor and close #249. The corpus proves one natural route to it is dead: `false_of_compressedAdjointCertificate` [`TotientMahlerDefect.lean:1051`] ([`Lean`], *scale:n/a*) shows no `CompressedAdjointCertificate` [`TotientMahlerDefect.lean:1039`] ($Q \cdot v \cdot A = \text{boundary}$ with $|\text{boundary}| < Q \cdot v$ and $A \neq 0$) can exist, and `not_finiteDimensional_span_fullTotientKernel` (immediately above it in the same file) shows the full family spans an infinite-dimensional space. Cross-checked against a separate row, `lcm_factorIdeal_finiteRank_shiftAlgebra` a single explicit countermodel defeats every finite-rank shift-polynomial observation simultaneously, so finite-rank amplification per se is not the missing rigidity.

Proposition 9.50 (What would close it). *A rationality-side rank upper bound: $\exists C$ such that any tempered integral binary carry orbit for a rational `binaryCoeffSeries` has $\text{finrank}_{\mathbb{Q}} \text{span}(\text{canonicalCarryKernelFamily } C \ e) \leq C$ for all e , or any bound growing slower than 2^e-1 . `not_irrational_totientSeries_implies_mod_period_and_unbounded_rank` (`TotientTailCarryPeriod.lean:224`) pins the obstacle precisely: rationality buys uniform eventual periodicity of u ’s dyadic sections mod v , and that periodicity provably does not promote to a $\mathbb{Q}$ -rank bound without extra arithmetic input.*

9.4 Synthesis: where the frontier actually is

Collecting the fourteen rows, three facts stand out as loci for future effort, stated with the same precision as the rows themselves rather than as summary rhetoric.

First, the single closest approach to 249-supply is the first-harmonic reduction of §9.2: a constant-saving ($21/25$ or $9/10$) exponential-sum cancellation estimate, with a satisfiable room condition and, via the subset consumer (Theorem 9.5), no requirement that the saving hold on a full interval or a positive-density subset. This is the only row in the index whose remaining content is recognisably a single classical estimate rather than a compound arithmetic-geometric statement.

Second, the top-edge staircase (TE-04 through TE-05-weakest, coord:actual-lcm-top-edge) is the only row where the corpus has both halved the obligation (one-sided instead of symmetric) and exposed, via the exact identity of Observation 12.9, precisely which of two logically symmetric escape branches survives the sign machinery already proved elsewhere in the same file family. That five strictly weaker equivalent or sufficient forms are already proved inter-derivable (Proposition 9.14) means effort here is not fragmented across restatements: proving any one closes the others automatically.

Third, the d4/d5 rank obligation is structurally independent of the residue-coordinate rows above: it is a second, self-contained sufficient condition for #249 (an upper rank bound contradicting the proved $2^e - 1$ lower bound), in a coordinate (carry-kernel rank) where a natural finite-rank strengthening is already proved impossible. A rank upper bound, if found, would not need to route through `certifiedKill` at all.

No row in this index is a solution, a partial solution in the sense of resolved cases, or evidence that #249 is likely true or false. Every open producer listed is exactly as open as the original Erdős–Borwein question it reduces to; what the index adds is the exact remaining content, verified against the live Lean tree, of each reduction.

10 Closed routes, with the mechanism that closed them

This section catalogues every no-go, countermodel, obstruction, and refuted proof strategy found in the #249/#257 corpus that bears on Erdős #249 ($S := \sum_{n \geq 1} \varphi(n)/2^n$ irrational). Erdős #249 is [Open]; nothing in this section decides it. What follows is a machinery inventory: for each dead route, the exact statement that closed it, the exact scope of the closure, and the exact machinery salvaged from the wreckage. The organising warning, stated once here and applicable to every item below: several of these “obstructions” are statements about a *representation* of the problem (a proof strategy, a certificate family, a coordinate system) and not about the object S itself. Confusing the two is the single most common error this catalogue exists to prevent.

10.1 Meta-level: finite inspection cannot certify the supply (the γ -splice remark)

(a) **Route as conceived.** Every unconditional route to #249 in this corpus bottoms out in a *certificate-supply* obligation of the shape $\forall h \geq 1 \forall N_0 \exists N \geq N_0 \exists L, \text{Sep}(h, N, L)$ ([Erdos249257.LcmConeFlatnes](#) the exact open-obligation form quoted throughout the existing manuscript). A natural hope is that verifying `Sep` or its diagonal specialisation at every scale up to some large, explicit bound B is evidence that the supply holds, and that pushing B far enough would eventually amount to a proof.

(b) **Exact mechanism that closed it.** The manuscript’s own remark ([Math], prose-only, not formalised in Lean: §5.4, “*finite inspection cannot establish the supply*”) gives an explicit

splice construction. Take any coefficient stream $\gamma : \mathbb{N} \rightarrow \mathbb{N}$ that agrees with φ on every index $\leq B$, and alter γ at one residue class beyond B so that the resulting binary series $\sum \gamma(n)/2^n$ is forced *rational*. Such a γ exists for every B (this is the same coboundary-splice technique that produces §10.2’s `parityCoboundaryWeight` witness, generalised: insert a lacunary zero-valued coboundary edit $2/2^m - 4/2^{m+1} = 0$ at some $m > B$). Because γ agrees with φ on every index $\leq B$, γ passes every finite certificate that only inspects indices $\leq B$ in particular every instance of $\text{Sep}(h, N, L)$ with $N + L \leq B$ that φ itself passes or fails. Yet $\sum \gamma(n)/2^n \in \mathbb{Q}$ by construction.

(c) Precise scope. This is a statement about *proof method*, not about φ : it does not touch S at all. What it excludes is the inference “Sep verified up to bound B , for arbitrarily large B , therefore Sep holds cofinally.” The scope is exactly `scale:bounded` versus `scale:cofinal`: any finite- B verification, however large, is compatible with both outcomes, because a rational stream can be built to survive any *fixed* inspection horizon. It says nothing about whether φ itself is such a γ only that no finite-horizon check can distinguish φ from a γ that is.

(d) Salvage. The splice construction is exactly the mechanism underlying §10.2’s Lean-formalised countermodel below, so its *content* is not lost even though this specific remark is unformalised: everywhere a proof strategy in this corpus proposes to certify a supply by exhaustive finite search (the diagonal deposits at $t \in \{1, \dots, 64\}$, [Erdos249257.DiagonalPincerCertificates](#) [DiagonalPincer], the row-200,000 sqrt-escape census on the #257 side), this remark is the standing reason such a search, however large, is evidence and not proof. [Math] `scale:n/a` `coord:binary-digit`.

10.2 The parity-aperiodicity countermodel (`TotientParityCoboundaryCountermodel`)

(a) Route as conceived. A natural strengthening of #249 attempts: show that any coefficient word $c : \mathbb{N} \rightarrow \mathbb{N}$ that (i) is uniformly bounded, (ii) satisfies the trivial linear-growth bound $c(n) \leq n$, (iii) matches $\varphi(n) \bmod 2$ exactly, and (iv) is not eventually periodic even strengthened to *cofinally, arbitrarily separated, arbitrarily long blocks* of non-periodicity must have irrational binary series $\sum c(n)/2^n$. This is the natural target for anyone trying to use only φ ’s coarse parity/aperiodicity profile.

(b) Exact mechanism. The construction `parityCoboundaryWeight(n) := parityBaseWeight(n) + 2 · largePowerTwoBit(n) - 4 · largePowerTwoBit(n - 1)`, where `parityBaseWeight` is the eventually-constant word $0, 1, 1, 2, 4, 4, 4, \dots$ and `largePowerTwoBit(n) = 1` iff $n = 2^{k+3}$, is a lacunary zero-valued coboundary splice on top of an eventually-constant base. The strongest of five landed theorems,

Proposition 10.1 (`Totient-parity arbitrarily-many-separated-carry rational countermodel`). *There exists $c : \mathbb{N} \rightarrow \mathbb{N}$ such that: $c(n) \leq 6$ for all n ; $c(n) \leq n$; $c(n) \equiv \varphi(n) \pmod{2}$ for every n ; for every N, G, K there is a block of K explicit $(6, 0)$ carry-pulse pairs beyond N , each pair separated by more than G ; and $\sum_n c(n)/2^n = 3/2$.*

[Erdos257PeriodNoncollapse.exists_totientParity_arbitrarilyManySeparatedCarry_rational_countermodel](#) [TotientParityCoboundaryCountermodel.lean:637] (sum computation `tsum_parityCoboundaryWeight_eq_three_halves` [TotientParityCoboundaryCountermodel.lean:359]; non-periodicity `parityCoboundaryWeight_not_eventually_periodic` [TotientParityCoboundaryCountermodel.lean:48]; exact parity match `parityCoboundaryWeight_mod_two_eq_totient` [TotientParityCoboundaryCountermodel.lean:48]) [Lean] `scale:cofinal` `coord:binary-digit`.

(c) Precise scope. This is an *object-level* existence witness ($c \neq \varphi$, only $c \equiv \varphi \pmod{2}$) that closes a *proof-route*: it proves that the hypothesis set {bounded, linear growth, φ -parity match, non-eventual-periodicity}, no matter how strongly the non-periodicity clause is strengthened (up to the arbitrarily-separated arbitrarily-numerous form above), can never entail irrationality of the associated binary series, because c satisfies every hypothesis and is rational. It says nothing about φ itself the witness sequence c is a different, hand-built sequence. Any

future sufficient-condition candidate stated purely in terms of {coefficient boundedness, growth, parity, periodicity} must be checked against this countermodel before being trusted, for *either* #249 or #257.

(d) **Salvage.** The lacunary coboundary-splice technique itself inserting zero-valued edits $2 \cdot 2^{-m} - 4 \cdot 2^{-(m+1)} = 0$ at sparse ranks to destroy periodicity while preserving the rational sum is a fully general recipe for building rational-valued, non-eventually-periodic, bounded coefficient sequences matching *any* parity template, and is literally the mechanism instantiated informally in §??’s γ -splice remark above. It is also a ready-made stress-test fixture for any future parity-based sufficient condition proposed anywhere in the corpus. Confirms the corpus-wide lesson (project memory `feedback_erdos_reductions_rejected_bank_real_results`): only arguments using *actual* quantitative totient/Mersenne size or residue information as in the `TotientActualLcm*`, `TotientFixedRank*` families below can possibly close #249; pure symbolic-word arguments cannot.

10.3 Two scoped #249 no-go countermodels: square-CRT correction suppression is ambiguous

(a) **Route as conceived.** The totient prime-dilation cocycle $\varphi(pm) = (p-1)\varphi(m) + [p \mid m]\varphi(m)$ carries a correction term whenever $p \mid m$. A natural strategy imposes a square congruence $n \equiv A + pa \pmod{p^2}$ with a “clean” shift h ($p \nmid a + h$) to suppress that correction on a finite horizon, then hopes that correction-suppression (“cleanliness”) is itself enough to guarantee that the resulting finite-difference cube is *nonzero* i.e. that cleaning away the correction term automatically leaves behind a genuine, nonvanishing arithmetic signal usable in a curvature argument.

(b) **Exact mechanism that closed it.** Two explicit, opposite finite computations refute the hoped-for entailment simultaneously:

- *Vanishing witness.* The smallest returned countermodel has the *entire* two-step finite block `actualOneCubeCoeff` equal to zero at $n = 52$, $r_0 = 13$, $r_1 = 13/18$ (`squareCRT_clean_block_can_vanish` [SquareCRTCube.lean:285], kernel-decided; cleanliness witnessed by `squareCRT_vanishing_countermodel_is_clean` [SquareCRTCube.lean:294]).
- *Nonvanishing witness.* A separate, nearby clean configuration at $n = 27$ has the same block *nonzero* (`squareCRT_clean_block_can_be_nonzero` [SquareCRTCube.lean:303]; cleanliness witnessed by `squareCRT_nonzero_countermodel_is_clean` [SquareCRTCube.lean:311]).

Both witnesses are exhibited alongside the positive achievability result `exists_squareCRT_clean_horizon` [SquareCRTCube.lean:198] (a bounded common base satisfying prescribed CRT anchors/residues does exist). [Cert] (theorem bodies not independently re-verified beyond signatures and docstring flagged `DOCSTRING-SOURCED` in the source bank) scale:fixed coord:other:square-crt-cocycle.

(c) **Precise scope.** These are two #249-scoped countermodels the coordinate is the totient prime-dilation cocycle specifically, not a problem-agnostic phenomenon. Together they show that *correction-suppression is logically independent of the sign/nonvanishing question it was hoped to settle*: “clean” (i.e. correction-free) is consistent with *both* outcomes. This is a statement about a specific finite-difference construction (the representation), not about φ : it does not exclude a nonvanishing finite-difference cube in general, only the inference from cleanliness *alone* to nonvanishing.

(d) **Salvage.** The general cocycle identity $\varphi(pm) = (p-1)\varphi(m) + [p \mid m]\varphi(m)$ (`totient_mul_prime_cocycle` [SquareCRTCube.lean]) and the CRT-witness apparatus (`exists_squareCRT_base` [SquareCRTCube.lean], `exists_large_squareCRT_base` [SquareCRTCube.lean]) remain fully reusable for *any* residue-forcing construction, in any coordinate. More valuably, the abstract cube-cancellation lemma extracted from the same file, `alternating_powerset_sum_eq_zero_of_`

[insert_invariant](#) [SquareCRTCube.lean:264] “if inserting one coordinate never changes the value of f , the alternating powerset sum of f vanishes” is completely divorced from totient/CRT content and is directly reusable for any Hilbert-cube or higher-difference argument on either problem. What remains as the honest open gap: an independent anti-concentration or residue-producer argument for why the clean residual does *not* vanish, which this module explicitly does not supply.

10.4 The adelic height obstruction

(a) **Route as conceived.** Several proof strategies attempt to *clear* an unwanted denominator factor from a rational value x by multiplying by a small integer coefficient c , hoping to discard the complementary denominator information entirely (e.g. localising a diagonal tail difference to one LCM ray channel while dropping the rest).

(b) **Exact mechanism.**

Lemma 10.2 (Scalar-localisation complement divisibility). *For $x : \mathbb{Q}$, $c : \mathbb{Z}$, $H : \mathbb{N}$: if $H \mid x.\text{den}$ and $(c \cdot x).\text{den} \mid H$, then $x.\text{den}/H \mid c.\text{natAbs}$.*

[Erdos249257.AdelicHeightObstruction.scalarLocalization_complement_dvd](#) [AdelicHeightObstruction.lean:56]. The equality strengthening, $\exists t : \mathbb{Z}, H \cdot c \cdot x = t \cdot x.\text{num}$, is [Erdos249257.AdelicHeightObstruction.scalarLocalization_eq_mul_num](#) [AdelicHeightObstruction.lean:56]. The Mersenne-scale corollary for positive x , if $2^r \mid x.\text{num.natAbs}$ and $x < 2/(2^n - 1)$, then $2^r(2^n - 1) < 2 \cdot x.\text{den}$ is [Erdos249257.AdelicHeightObstruction.mersenne_height](#) [AdelicHeightObstruction.lean:103] (generic form [positiveRat_numDivisor_mul_lt_two_mul_den](#) [AdelicHeightObstruction.lean:77]). A companion rigidity fact rules out smuggling the discarded information back in through a linear channel: if a linear map $\Lambda : V \rightarrow W$ factors through a surjective evaluation ev ($\ker \text{ev} \leq \ker \Lambda$), then Λ is *exactly* $\text{ev}(\cdot) \bullet w_0$ for a single fixed w_0 [Erdos249257.AdelicHeightObstruction.linearDescender_eq_smul_eval](#) [AdelicHeightObstruction.lean:120]. The upstream primitive these import, [Erdos249257.RationalDenominatorSurvival.dvd_divInt_den](#) [RationalDenominatorSurvival.lean:17] and [survivingDivisor_dvd_scaled_divInt_den](#) [RationalDenominatorSurvival.lean:38], gives the exact survival/cancellation law: a divisor $m \mid D$ of a displayed denominator survives reduction of a/D iff m is coprime to a ; after scaling the numerator by h , a surviving coprime divisor $C \mid D$ shrinks to exactly $C/\gcd(C, h)$, never further. [Lean] scale:uniform coord:other:rational-height.

(c) **Precise scope.** This is pure $\mathbb{Q}$ -arithmetic with zero Mersenne or totient content it is not a statement about S at all, but about *any* rational-height bookkeeping argument. Its exact content: scalar denominator-clearing never *erases* the complementary denominator; it moves that complement into the coefficient’s (Archimedean) size. Any construction that tries to shrink a displayed denominator down to one surviving channel pays for it in coefficient growth. This directly explains *why* denominator-compression strategies are structurally hard: the Mersenne-shadow denominator lower bound $2^{t/2} \leq \prod_{p \in \text{upperHalfPrimes } t} \text{mersenne}(p)$ ([upperHalfMersenneProduct_lower_bound](#) [MersenneShadowDenominatorGrowth.lean:60]) forces a quadratic-in-scale coefficient cost via this lemma if one tries to localise to a single channel.

(d) **Salvage.** Fully problem-agnostic and directly reusable for #257 or any other Lambert-type denominator-survival argument as-is. It is the general mechanism underlying the corpus’s Farey-gap denominator-exclusion family (Part C of the certificate bank, [tsum_totient_div_pow_two_ne_ratCast_of_den_le_79639646646701375323355774875831053](#) [CertificateKernel.lean:18056]), the strongest current unconditional statement about S : if S is rational its reduced denominator exceeds $79,639,646,646,701,375,323,355,774,875,831,053 \approx 7.96 \times 10^{34}$).

10.5 The signed q -moment machinery a positive tool, not an obstruction

(a) **Route as conceived.** A “signed Hankel determinant” / Hermite–Padé style route to #249 would exhibit a finite signed linear combination of dyadically-cleared totient terms and show it is provably nonzero, giving a nonvanishing certificate for a first-harmonic or Möbius-companion construction.

(b) **Exact mechanism (a nonvanishing *engine*, not a no-go).**

Lemma 10.3 (Unique-terminal parity nonvanishing). *If a finite signed sum $\sum_{i \in s} u(i) \cdot 2^{e(m)-e(i)}$, clearing dyadic denominators to a common exponent $e(m)$, has one index $m \in s$ with strictly maximal exponent $e(m)$ and odd coefficient $u(m)$, while every other index has strictly smaller exponent, then the whole cleared sum is $\equiv 1 \pmod{2}$, hence nonzero.*

`scaled_dyadic_sum_ne_zero` [SignedQMomentObstruction.lean:95], built from a fully generic rectangular Cauchy–Binet-style determinant-of-product expansion `det_mul_rectangular` [SignedQMomentObstruction.lean:107], `hankelDet/truncatedMoment` defs at `SignedQMomentObstruction.lean` [SignedQMomentObstruction.lean:107, 111] [Lean] scale:bounded coord:p-adic.

(c) **Precise scope the point this section exists to make.** Despite living in a file named alongside the corpus’s “obstruction” modules, this is explicitly *not* phrased as a no-go in its own docstring: it is the finite nonvanishing engine that a Hankel–Padé construction for #249 would *consume*. The parity lemma and the Cauchy–Binet expansion use no totient or Mersenne structure at all pure linear algebra, fully problem-agnostic, directly transplantable to any base- q dyadic-clearing argument including #257’s series family. What is missing, and what prevents this from closing anything, is a *supply* theorem: no result in the corpus proves that a unique-terminal configuration of this shape exists cofinally for the totient-derived family. The obstruction, such as it is, is not in this lemma but in the absence of its hypothesis’s cofinal supply exactly the same shape of gap as §10.8’s Mahler defect below.

(d) **Salvage.** The entire lemma is reusable machinery, not wreckage nothing here needs salvaging because nothing here failed. It stands ready for whichever future construction can supply the missing cofinal unique-terminal-configuration existence.

10.6 The residual-gauge obstruction

(a) **Route as conceived.** A “first-harmonic pivot” strategy for #249 (via exponential/character-sum cancellation, `FirstHarmonicPivot.lean` [FirstHarmonicPivot.lean]) might try to certify a nonzero determinant or full-rank minor of a residual-weighted monomial matrix as sufficient evidence that a genuine phase reconstruction (as opposed to a degenerate locked configuration) has occurred.

(b) **Exact mechanism.**

Proposition 10.4 (Locked reconstruction preserves a nonzero minor). *For a monomial matrix $\text{residualMonomialMatrix}(e, r, z)(i, j) = r(i, j) \cdot z(j)^{e(i)}$ with all residual entries $z(j) \neq 0$ and $e(i) = 1$ for a distinguished row i : $\det(\text{phasePowerMatrix}) \neq 0 \implies \det(\text{residualMonomialMatrix}) \neq 0$, and the distinguished row is identically 1 regardless.*

`Erdos249257.locked_reconstruction_preserves_nonzero_minor` [ResidualGaugeObstruction.lean:94] (row-dependent relaxation `rowResidualMonomialMatrix_reconstructs_phasePowerMatrix` [ResidualGaugeObstruction.lean:107]) the column-gauge identity underlying it, $\det(\text{residualMonomialMatrix}) = \det(\text{phasePowerMatrix}) \cdot \prod_j W(j)$, is `det_residualMonomialMatrix` [ResidualGaugeObstruction.lean:47]; a row-dependent relaxation still fails at `rowReconstruction_zero_row_one` [ResidualGaugeObstruction.lean:145]). [Lean] scale:uniform coord:other:first-harmonic-phase.

(c) **Precise scope.** Pure linear algebra over $\mathbb{C}$, fully generic in dimension d and exponent function e , zero #249-specific content this is a statement about *residual-weighted monomial*

matrices as a representation, not about the totient object. It rules out exactly one class of certificate: a residual-blind rank/determinant/conditioning test, used alone, cannot distinguish genuine phase reconstruction from an explicit “locked” degenerate configuration in which a residual weighting scales every column by $W(j) = z(j)^{-1}$, collapsing the exponent-one row to the constant row $(1, \dots, 1)$ while the determinant stays nonzero whenever the base (Vandermonde-shaped) phasePowerMatrix determinant is nonzero. It does *not* rule out determinants carrying an additional arithmetic coupling identity between rows.

(d) **Salvage.** A hard requirement for any future first-harmonic determinant construction (Part E of the certificate bank, [irrational_totient_series_of_first_harmonic_norm_gap](#) [FirstHarmonicPivot.lean:83]): the construction *must* couple rows by an extra arithmetic identity, not merely gauge-normalise columns. The diagonal-gauge-action technique itself is reusable for any determinant-based nonvanishing strategy in any coordinate.

10.7 Cone flatness and cone nonflatness

(a) **Route as conceived.** The diagonal certificate-supply obligation ([Erdos249257.LcmConeFlatness](#) [LcmConeFlatness.lean]) concerns only the pair $(H, 2H)$ at LCM height $H = \text{periodLcm}(t)$. Two natural strengthenings: (i) show rationality of S is even more constrained than a single diagonal pair it flattens an *entire cone* of ratios; (ii) conversely, exhibit *nonflatness* on a wider “menu” of cone vertices as a cheaper sufficient certificate than a single pairwise certifiedKill.

(b) **Exact mechanism flatness (necessary consequence of rationality).**

Proposition 10.5 (Wave-24 LCM-cone flatness). $\neg \text{Irrational}(S) \implies \exists t_1, \forall t \geq t_1, \forall q, m : \mathbb{N}, 0 < q \implies \text{totientTail}(q \cdot \text{periodLcm}(t) + m \cdot \text{periodLcm}(t)) - \text{totientTail}(q \cdot \text{periodLcm}(t)) \in \text{range}((\uparrow) : \mathbb{Z} \rightarrow \mathbb{R})$.

[rational_totient_series_forces_lcm_cone_flatness](#) [CertificateKernel.lean:18674] (body [LcmConeFlatness.lean](#)). [Lean] scale:uniform (for all sufficiently large t , *given* rationality) coord:other:lcm-ray-totient. Rationality forces one fractional constant on the entire LCM cone $\{k \cdot \text{periodLcm}(t) : k \geq 1\}$ at every scale $t \geq t_1$, not merely the single diagonal pair the umbrella collapse theorem [irrational_totient_series_of_lcm_cone_window_kill_supply](#) [CertificateKernel.lean:18686] shows any certifiedKill *anywhere* on the two-multiplier cone, at arbitrarily large t , forces Irrational(S).

(b') **Exact mechanism nonflatness (a sharper sufficient certificate).**

Proposition 10.6 (Wave-25 cone-nonflat menu refuter). *For a nonempty menu Q of positive vertex multipliers with a one-sided floor $\forall q \in Q, q \cdot H + L + 2 < 2^L$ (half the pairwise floor of certifiedKill): if coneNonflatCert $H L Q$ fires (an argmin/Helly-avoidance argument: the vertices cannot all share one fractional part, else the minimal-deep-tail vertex would be a common left endpoint of every arc), then $\exists q_i, q_j \in Q, \text{totientTail}(q_j H) - \text{totientTail}(q_i H) \notin \text{range}((\uparrow) : \mathbb{Z} \rightarrow \mathbb{R})$.*

[exists_nonintegral_pair_of_coneNonflatCert](#) [CertificateKernel.lean:18801] (supply theorem [irrational_totient_series_of_lcm_cone_nonflat_supply](#) [CertificateKernel.lean:18812]). [Lean] scale:bounded (proved for a given finite menu Q ; an *unbounded*-scale menu is the open sufficient target) coord:other:lcm-ray-totient.

(c) **Precise scope.** Flatness is a genuine *necessary consequence* of rationality it is a statement about what rationality of S would force, conditional on rationality, not an unconditional fact about S . Nonflatness is an unconditional *sufficient* certificate mechanism, information-theoretically half the depth floor of the pairwise certifiedKill (one-sided versus two-sided radius charging), but it has been proved only for individual finite menus Q ; no menu of unbounded scale has been supplied. Neither direction decides #249: flatness gives a contrapositive route (exhibit *any* cone nonintegrality at arbitrarily large t and rationality is refuted) and nonflatness sharpens the certificate needed, but the underlying producer obligation a nonintegral tail

difference at cofinally many scales is unchanged. Note also the coordinate warning from the manuscript’s own Appendix C ([res:flat](#) [LcmConeFlatness.lean:90]): cone flatness is a *necessary* consequence, and only the sharper diagonal-pincer / full-target-avoidance normal forms ([DiagonalPincerDecomposition.lean](#) [DiagonalPincerDecomposition.lean:215,290]) are an exact iff with irrationality.

(d) Salvage. The argmin/Helly-avoidance combinatorial-geometry technique behind cone-nonflatness is a reusable pattern for any modular-residue pincer argument with more than two points, in any coordinate. The rank-2 second-difference certificate family built on the same window apparatus was independently tested and found *not* to be a shortcut over rank 1 (measured, not merely conjectured: at $(h, N) = (1, 8)$, rank-1 fires at depth 8 while no rank-2 certificate exists at depth ≤ 8 ; [totient_tail_rank_two_kill_sound_but_not_shallower_cell](#) [CertificateKernel.lean:18762]) an explicitly flagged dead end not to re-attempt without new information.

10.8 The Mahler defect: dyadic totient-kernel finite rank per level, infinite rank overall

(a) Route as conceived. A finite-linear-algebra shortcut to #249: find a bounded-depth linear relation among the dyadic totient-kernel channels $n \mapsto \varphi(2^j n + r)$ that would compress the infinite family to a finite-dimensional space, from which a rationality-forcing contradiction (or a genuine rank obstruction) might be extracted.

(b) Exact mechanism.

Theorem 10.7 (Infinite dyadic totient-kernel rank). *For every depth $e \geq 0$, the canonical family of $2^e + 1$ dyadic totient-kernel channels (`card_totientCanonicalIndex`, [TotientMahlerDefect.lean](#) [TotientMahlerDefect.lean:900]) is linearly independent over $\mathbb{Q}$, proved via a *SeparatedMinorCertificate* (an explicit finite evaluation-point assignment with nonzero determinant, forced by CRT + Dirichlet’s theorem on primes in arithmetic progressions one channel made prime, every other channel forced through a fresh prime $\equiv 1 \pmod{a}$ large power of 2).*

[linearIndependent_canonicalTotientKernelFamily](#) [TotientMahlerDefect.lean:900]; consequently the full infinite family spans an infinite-dimensional $\mathbb{Q}$ -space, [not_finiteDimensional_span_fullTotientKernel](#) [TotientMahlerDefect.lean:1014]. A companion impossibility result closes the natural repair attempt directly: a bounded *compressed-adjoint certificate* a triple $(Q, A, \text{boundary})$ with $Q \cdot v \cdot A = \text{boundary}$, $|\text{boundary}| < Q \cdot v$, $A \neq 0$ is provably impossible, [false_of_compressedAdjointCertificate](#) [TotientMahlerDefect.lean:1051]. [Lean] `scale:uniform` (holds for every e ; existence side is unconditional, via Mathlib’s CRT + primes-in-AP machinery) `coord:binary-digit` (dyadic totient-kernel *not* the Möbius coordinate).

(c) Precise scope coordinate-relative, stated explicitly by the source module. This does *not* show irrationality of S . It proves the dyadic-kernel *side* is infinite-rank; the module’s own docstring names the missing input as “a rationality-side finite-rank compression, or equivalent contradiction.” The companion necessary-consequence-of-rationality theorem, $\neg \text{Irrational}(S) \implies \exists v > 0, \exists U : \mathbb{N} \rightarrow \mathbb{Z}, \text{IsTemperedBinaryOrbit}(\varphi, v, U) \wedge \forall e, 2^e - 1 \leq \text{finrank}_{\mathbb{Q}} \text{span}(\text{range}(\text{canonicalCarryKernelFamily}(U, e)))$ ([not_irrational_totientSeries_implies_unbounded_carryRank_unconditional](#) [TotientCarryKernelRigidity.lean:284]), shows the SCALE side is finished the $2^e - 1$ floor holds for all e with a uniform proof but the opposite inequality (a rationality-side rank *upper* bound) is exactly what is missing, and its absence is the entire open content here. A *third*, independent coordinate exhibits the same coordinate-relative phenomenon: the Möbius-incidence companion matrix $U_N(i, j) = \mu((i + 1)/(j + 1))$ if $(j + 1) \mid (i + 1)$ else 0 is lower triangular with diagonal 1, hence $\det U_N = 1$ for every N , hence its jet-evaluation map is injective a finite “incidence quotient” compression is

impossible at any finite horizon in the Möbius coordinate too ([incidenceMobius_det_eq_one](#) [IncidenceQuotientHermitePade.lean:56]). Both natural finite truncations dyadic-residue and Möbius-incidence turn out to have no nontrivial kernel, in two completely different coordinates. That is strong evidence any winning finite-linear-algebra shortcut, if one exists, must live in a genuinely third coordinate or use an infinite/growing-parameter construction, not evidence that no shortcut exists at all.

(d) **Salvage.** [SeparatedMinorCertificate](#) [TotientMahlerDefect.lean:73] and [linearIndependent_of_separatedMinorCertificate](#) [TotientMahlerDefect.lean:81] are fully generic finite-rank linear-independence infrastructure the indexed family is a free variable, reusable for any finite-rank obstruction argument on either problem (e.g. testing independence of Mersenne-indexed sequences for #257 via an explicit nonzero minor). The CRT+Dirichlet evaluation-forcing technique (make one channel prime, force every other channel through a fresh arithmetic-progression prime) is a reusable construction pattern independent of the totient specifics.

10.9 Further closed routes, catalogued for completeness

The following additional no-gos and route-pruning results were read in full and are recorded here in compressed form; each follows the same (a)/(b)/(c)/(d) discipline as above but is presented as a table row for space.

Route	Mechanism & site	Scope / salvage
Full terminal dyadic staircase (annihilate every suffix letter mod growing powers of 2)	$a \geq 8$, room bound $\Rightarrow$ ActualLcmTerminalDyadicStaircase is false: the terminal letter would have to be positive (by short-window positivity) and strictly below a wider-than-itself modulus while divisible by it, forcing it to 0, contradicting positivity. not_of_room [TotientActualLcmTopEdgeStaircase.lean:1251] [Lean] scale:uniform coord:binary-digit	Kills the FULL terminal-staircase proof strategy outright and unconditionally; the <i>punctured</i> staircase (all but the last letter vanish) survives and is pinned exactly to the half-turn value 2^{m-1} at the penultimate letter (puncturedDyadicStaircase_actualLcmTerminalDyadicStaircase_penultimate_eq_of_room [TotientActualLcmTopEdgeStaircase.lean:1251]). The generic mechanism $(0 < e < 2^m \wedge 2^m \mid e \Rightarrow e = 0)$ is reusable anywhere a positive quantity is asked to vanish mod a wider modulus.
LCM factor-ideal retention (keep only the homogeneous “factor-only” part of an LCM-ray decomposition)	An explicit nonzero all-horizon countermodel, built from multiples of $\varphi(\text{periodLcm}(t))$, agrees with every exact whole-ray anchor $\Delta\varphi(H, qH) = \varphi(H)$ for $2 \leq q < t$ and survives every finite commensurate LCM-cube shift-polynomial, at every finite rank. LcmFactorIdealPulseObstruction.lean:1-24 (module docstring; theorem bodies [Cert], DOCSTRING-SOURCED). scale:uniform coord:other:lcm-ray-totient	This is a <i>representation</i> -level exclusion, not an object-level one: it shows the homogeneous factor-ideal projection discards fresh Möbius-channel information that can be adversarially reconstructed. Explicitly flagged SYNTHETIC no claim that the construction is as actual totient differences. Salvage: a future factor-ideal argument must control the <i>fresh</i> channel via totient_eq_sum_mobiusTotientChannel [LcmFactorIdealPulseObstruction.lean:1-24] not just the old one.

Route	Mechanism & site	Scope / salvage
Two-prime “full-target diamond” (four simultaneous ray hits certify more than one)	The four-hit diamond $\text{Hit}(H) \wedge \text{Hit}(pH) \wedge \text{Hit}(qH) \wedge \text{Hit}(pqH)$ is logically equivalent to $\text{Hit}(H)$ alone, no primality of p, q needed hits transport multiplicatively for free because $\text{Hit}(H) \iff \text{IsIntegralValue}(\text{scaleDiagonalTailDifference}(\text{Hit}(H)))$ and integrality transports affinely along every ray $H \mapsto kH$. Erdos249257.fullTarget_primeAdjunction_diamond_iff_root [FullTargetPrimeAdjunctionNoGo.lean:196]. [Lean] scale:uniform coord:other:mobius-mersenne	The 4-condition diamond carries zero information beyond its base point a “curvature-zero collapse” of a multi-point certificate. Any future positive-holonomy argument needs an independently- $\text{Hit}(H)$ projection of the foreign state <i>plus</i> control of the discarded complement; a bare 2/4-point diamond cannot supply new information beyond its own scale.
Fixed-precision local valuation-unit signature (2-adic “tropical curvature carry” attack)	For any finite word of odd-unit-part 2-adic symbols at fixed precision $u > 0$ and any starting carry state e , there exists a compatible carry orbit realising the word with every intermediate state centred inside its symbol’s dyadic radius. Erdos249257.TotientTailPeriodKiller.isRequiredPrecision2-adicDynamics [TropicalCurvatureCarry.isRequiredPrecision2-adicDynamics] [Lean] scale:bounded coord:p-adic	Bounded LOCAL valuation-unit data at FIXED precision can never exclude all finite centred carry completions no obstruction is derivable from a fixed-window local signature alone; growing precision, or extra arithmetic coupling, is required. Preson 2-adic dynamics, zero totient content, reusable against any similarly-shaped fixed-window carry-certificate attack.
Fixed-depth affine carry reset (distinguishing two carry histories after a long common tail)	For the affine binary orbit $\text{orbit}(0) = u_0$, $\text{orbit}(n+1) = 2\text{orbit}(n) - a(n+1)$ with common forcing word a : $\text{orbit}_u(L) - \text{orbit}_v(L) = 2^L(u_0 - v_0)$ exactly. Erdos249257.affineBinaryOrbit_mod_twoPow_eq [GenericTailOrbitRigidity.lean:306]. [Lean] scale:uniform coord:binary-digit	After L common steps the end-point residue mod 2^L is <i>independent</i> of the initial carry long common tails erase predecessor information exactly, not approximately. Directly relevant to any argument hoping to distinguish two carry histories from a shared long suffix; the terminal forcing word alone determines the residue.
Bounded finite-state / autonomous successor decoder for the binary carry	For a balanced-pulse family at location m whose predecessor State is constant across the whole family, no decode : $\text{State} \rightarrow \mathbb{N}$ recovers the radius parameter; any finite Fintype State needs $\text{card} \geq \lfloor m/2 \rfloor + 2$, unbounded in m . Erdos249257.balancedPulse_no_autonomous_decoder [GenericTailOrbitRigidity.lean:246]. [Lean] scale:uniform coord:binary-digit	Rules out, unconditionally, ANY proof strategy that defines a bounded or autonomous “carry state” summarising pre- m history and claims it exactly determines the post- m tail independent of whether the coefficients are φ or a Möbius-support indicator. A hard stop against finite-automaton- $\text{computes-the-expansion}$ style attacks on either problem.

Route	Mechanism & site	Scope / salvage
Period-4 sign weight as a route to A7's "frequently nonzero" hypothesis for free	The divisor coefficient of the period-4 sign weight ($w(a)=1$ if $a \equiv 1, -1$ if $a \equiv 3$, else 0, mod 4) vanishes identically at every $n \equiv 3 \pmod{4}$, via the divisor-pairing involution $d \mapsto n/d$. Erdos249257.intWeightedCoeff_-periodFourSignWeight_-eq_zero_of_mod_four_eq_-three [CertificateKernel.lean:13898]. [Lean] scale:fixed coord:mobius-mersenne	Witnesses that Dirichlet-convolution coefficients CAN vanish identically on an arithmetic progression, so the "frequently nonzero" hypothesis in the nonnegative-periodic irrationality closure (irrational_-intWeightedErdosSeries_-periodic_of_coeff_-nonneg_of_frequently_ne_-zero [CertificateKernel.lean:14255]) is not free it is why the corpus lands a dichotomy for general periodic signed weights rather than an unconditional supply theorem.
Certified finite-depth "death" certificates as a route to membership proofs (both problems)	CertifiedGreedyMersenneDeath (x , level, StochasticRead) is a decidable, finite-depth certificate proving non-membership only (greedyMersenneDeath_not_-mem [GreedyAchievementSet.lean:1720]; witness three_fourths_-certifiedGreedyMersenneDeath [GreedyNonmembershipSet.lean:1742] 3/4 certified dead at level 1). [Cert] scale:bounded coord:greedy-orbit	StochasticRead one-sidedness, recurring across the WHOLE certified-kill family (TotientTailPeriodKiller [TotientTailPeriodKill], LcmConeFlatness [LcmConeFlatness.lean], this one): certified-death $\Rightarrow$ non-membership. Hence of a found certificate, or survival through any finite depth, proves <i>nothing</i> about membership or rationality.
Reconstructing one totient value from a two-tail absolute-value linear combination	For any finite $w : \iota \rightarrow \mathbb{Q}$, $x : \iota \rightarrow \mathbb{N}$ with $\sum w_i \varphi(x_i) = 1$ (exact isolation), the crude two-tail cost $\sum w_i (2(x_i + 1) + (x_i + 2)) \geq 3$, using only $\varphi(x) \leq x$. Erdos257PeriodNoncollapse.three_-le_totientAdjugateTailCost [TotientTailCarryPeriod.lean:307]. (closure not_-totientAdjugateTailCost_lt_-one [TotientTailCarryPeriod.lean:307]). [Lean] scale:uniform coord:other:totient-adjugate-linear-algebra	The strategy needs cost < 1 and gets ≥ 3 at ANY finite grid height kills the entire "isolate one coefficient via absolute-value two-tail bookkeeping" family before it starts. The proof mechanism is TotientTailCarryPeriod.lean:307 $c(n) \leq n$ sequence gets the same floor by the triangle inequality) and transfers verbatim to a #257 reformulation.

Route	Mechanism & site	Scope / salvage
Homogeneous Mersenne-primitive prime factor alone forces a contradiction	For the completely-multiplicative control $c(n) = n$ (zero totient content): if $K < q$, $q \mid 2^K - 1$, the corresponding tail shift is integral at every N yet $q \nmid K$. idCoeff_mersenneModulus_does_not_annihilate_tailShift [TotientTailCarryPeriod.lean:322] [Lean] scale:n/a coord:other:mersenne-modulus	A primitive prime factor of the homogeneous multiplier $2^K - 1$ alone supplies NO contradiction from tail integrality. A direct cross-problem warning: #257's denominators $2^n - 1$ are literally this $q \mid 2^K - 1$ object, so “a large primitive Mersenne prime factor alone forces a contradiction” should be checked against this lemma before being attempted on either problem.
Prime-power reduced-denominator unit-gap strengthening (rescuing extra Farey lattice points)	At a prime-power reduced denominator p^e , the unit-gap refinement can rescue at most ONE additional candidate lattice point beyond the ordinary gap certificate. reducedDenominatorCandidateCount_prime_pow_le_one [PrimitiveWeightCertificate.lean:111] (iff-characterisation PrimitiveWeightCertificate.lean [Lean] scale:uniform coord:other:farey-gap	A formal ceiling on how much this specific refinement can ever buy a known dead end for extending the $\approx 7.96 \times 10^{34}$ Farey rung (§10.4) unboundedly. Do not re-attempt this exact strengthening expecting more than +1 lattice point per prime power; the underlying reduced-denominator-implication PrimitiveWeightCertificate.lean [Lean] scale:uniform coord:other:farey-gap is reusable for other denominator-exclusion arguments.
Fixed rank-3 finite-difference kernel squeezing more than one dyadic bit of curvature information	$\text{fixedRankSecondDifference}(2^{n+1} \cdot 2, 2^{n+1} \cdot 3) = -2^{n+1}$ exactly, for all n : the two-adic valuation gained by the primitive kernel factor $2\varphi(j)$ is exactly tight. fixedRankSecondDifference_dyadic_fixture_exact [TotientFixedRankLcmAsymptotic.lean:162] [Lean] scale:n/a coord:binary-digit	Route-pruning ammunition: rules out extracting more than one dyadic bit from the bare 3-rank affine kernel $(1, -2, 1)$; any further progress on the fixed-rank curvature route needs new arithmetic input, not a squeeze from the same kernel. The affine rank-3 kernel classification itself (affine_rank_three_kernel_classification [TotientFixedRankLcmAsymptotic.lean:162] pure linear algebra over $\mathbb{Z}$, zero number-theoretic content) is fully reusable for any 3-point finite-difference argument on either problem.

11 Why the bounds are load-bearing

The near-miss interface between the corpus's proved results and the four open supply obligations #249-supply, #257-reset, #257-cofinal-rows, and #257-universal was swept for *promotion candidates*: proved theorems whose *stated* hypotheses looked, on a first read of the **yields** clause, close enough to the open obligation's shape that a routine strengthening (raising a scale bound, widening a hypothesis, or reindexing a quantifier) might close the gap without new mathemat-

ics. Fifty-six near-miss rows were catalogued across the four obligations; eighteen of them were flagged `promotable_claimed: True` on this first pass. **Every one of the eighteen proofs was then opened and read in full**, not merely re-inspected at the statement level, and the promotion claim was checked against the actual proof body. The audit table below reports the verdict for all eighteen.

Of the eighteen, **sixteen are conclusively NOT_PROMOTABLE**: reading the proof body exposes a documented arithmetic, logical, or coordinate obstruction that a routine strengthening cannot cross, and the row’s own record states exactly what new input would be required. **Two** (NM-02, NM-03 in the #257-universal obligation) are genuine exceptions: the auditor found the promotion is available with, in its own words, “no new mathematics” a pure widening of an already-general proof to a wider stated target. Crucially, *neither of these two closes anything*: they widen an equivalence or a corollary from one fixed target value to a family of targets, without touching the missing arithmetic content (a cofinal certificate supply, a rank upper bound, an anti-concentration estimate) that every genuine closure of #249 or #257 requires. So while the promotion audit is not literally eighteen-for-eighteen at the level of “can this exact statement be re-derived for a wider scope,” it is eighteen-for-eighteen at the level that matters for this paper’s obligations: **no promotion candidate in the audited set supplies missing arithmetic content toward closing #249 or #257**. The two mechanically-widenable rows are recorded honestly below rather than folded into the sixteen, because collapsing that distinction would itself be exactly the kind of quantifier-slippage error this paper’s accuracy rules forbid.

Obligation	Row / site	Gap kind	Verdict and exact blocker
249-supply	e1-companion, FirstHarmonicGap.lean	hypothesis strength	<code>NOT_PROMOTABLE</code> . The consumer already has the obligation’s exact quantifier shape; the missing in- [FirstHarmonicGap.lean:120] arithmetic fact a constant-saving first-harmonic (Weyl-sum) cancellation bound $\sum_{N \in [X, 2X)} \cos(2\pi \cdot (...)/2^L) \leq (9/10)X$ not proved at any X, h, L anywhere in the corpus.
249-supply	SGN-01, TotientActualStrength	hypothesis strength	<code>NOT_PROMOTABLE</code> . Positivity is exactly HALF of the need (to certificate the companion theorem in the same file proves integrality forces the residue to the TOP edge, not a kill the lower half of the band is discharged unconditionally and cofinally, the upper half is untouched).
249-supply	TE-04, TotientActualStrength	hypothesis strength	<code>NOT_PROMOTABLE</code> . The consumer side is finished TopEdgeAndCofinal (proved for every $LcmTopEdge$ is 1260) purely the one-sided residue-gap <i>producer</i> , unconditionally unsupplied at even one large a .
249-supply	TE-05-weakest, TotientActualLcmTopEdgeCarryOrbit.lean	hypothesis strength	<code>NOT_PROMOTABLE</code> . The exact identity pinning the target shows the dominance inequality is equivalent to CarryOrbit.lean and the signal in TopEdge (SGN-02) lean:1908 already proves the true carry orbit is strictly <i>positive</i> under integrality the corridor-escape branch the identity naturally supplies is exactly the branch already eliminated.

Obligation	Row / site	Gap kind	Verdict and exact blocker
249-supply	SEP-02, TotientActualOnlyOrbitSep	scale_- only	NOT_PROMOTABLE. The analytic tail is fully discharged with <code>TotientActualLcmOrbitSeparation.radius</code> only two exponents ($a = 4, 6$) have the resulting finite distance-to-integer question verified unconditionally (<code>actualLcmTailOrbit_four_and_six_notMem_int</code> [TotientActualLcmShortKill.lean:35]); nothing beyond $a = 6$ is proved.
249-supply	b7, MersenneShadowOnly	coordinate_ only	NOT_PROMOTABLE. The only proved unbounded-growth quantity <code>MersenneShadowMersenneCoordinate</code> is uniform in t , but there is no landed transport from a denominator lower bound to <code>certifiedKill</code> or to <code>totient-tail</code> non-integrality a different coordinate from the obligation.
249-supply	d4/d5, TotientMahlerDefect.lean TotientCarryKernelRigidity	coordinate_ only	NOT_PROMOTABLE. The scale side of a second, <code>defect.lean</code> (dependent on <code>defect.lean</code>) (unbounded carry-kernel <code>TotientCarryKernelRigidity</code> force <code>TotientCarryKernelRigidity</code> is fixed uniformly in e ; the opposite (rationality-side rank upper bound) is missing, and one natural route to it is proved dead (<code>false_of_compressedAdjointCertificate</code> [TotientMahlerDefect.lean:1051]).
257-reset	rc-2 / F4-adjacent, HalfCylinderMiddleCarry	multiple and upper bound	NOT_PROMOTABLE. The excess-side run-length law is only half-assembled: an exact iterate identity exists, <code>HalfCylinderMiddleCarry</code> [HalfCylinderMiddleCarry.lean:2309] but the intermediate carry does not exist separately, but no theorem in Lean composes the two into an excess-side run-length statement, even though every ingredient is present.
257-reset	E1, HalfCylinderOnlyLimit.lean	coordinate_ only	NOT_PROMOTABLE. Supplies an UPPER bound on the <code>HalfCylinderOnlyLimit.lean</code> (the picture); the obligation needs a LOWER bound on the deviation magnitude the two do not compose without an additional input.
257-reset	rc-8, <i>prose</i> erdos257_reset_crossing_unification_2026_07_24.md §6	multiple	NOT_PROMOTABLE. Two gaps at once: the sign law is <code>[Cert]/empirical</code> only (flagged <code>[Open]</code> as an unconditional lemma, not proved), and even if proved it delivers only the SIGN of the deviation, not its magnitude, which is what the obligation needs.
257-cofinal-rows	TH-sharp-capacity-progress, BooleanMobiusSkipRow.lean	hypothesis_ strength	NOT_PROMOTABLE. Unconditional and strictly more general than every actual consumer, but it needs a per- c input (<code>localBinarySuffix D 1 (2c-2) < 2^{c-2}) that <code>BooleanMobiusSkipRow.lean</code> [BooleanMobiusSkipRow.lean:286] through a route which then rigidly collapses onto one specific support family the deficit hypothesis is a self-imposed restriction of the consumers, and the general input itself is unsupplied.</code>
257-cofinal-rows	precriticalSuffixlt_of_-future_-skip_after_-takenBlock, BooleanMobiusCriticalCapacityCofinal.lean	hypothesis_ strength	NOT_PROMOTABLE. Unconditional and uniform in both parameters (c, t); the sole missing input is an orbit-level skip-gap bound on the rational half-greedy orbit, never proved and never previously isolated as a target anywhere in the banks.

Obligation	Row / site	Gap kind	Verdict and exact blocker
257-cofinal-rows	greedyHalfFrozenMargin_ nonneg,	only	NOT_PROMOTABLE. The margin-nonnegativity horizon is produced by a <i>non-effective</i> limit argument (an <i>existential horizon</i> from <code>made_converge</code>); the exact-row route needs the SPECIFIC effective horizon $J = k - 2$, and effectivising it is explicitly recorded as unfinished bookkeeping, not new mathematics but it is still unfinished.
257-cofinal-rows	C4a, <code>TerminalOnlyCofinal.lean</code>	multiple	NOT_PROMOTABLE. Trades off against the obligation in <code>TerminalOnlyCofinal.lean</code> on <i>support locality</i> : tolerates any support inside a depth window but demands a carry inside a tight $\sim 2\sqrt{M}$ strip, where the obligation tolerates an exponentially looser carry but demands the support be confined to a lower window the gap is support locality, not carry size, and neither socket implies the other.
257-universal	NM-02, <code>GreedyAchieveOnlySet.lean</code>	scale_ - only	PROMOTABLE (no new mathematics), but does not close anything. [248] The proof was opened and found already uniform in the target in every internal step; only the statement's own quantifier is artificially pinned to $t = 1/2$. Promoting it converts the whole half-greedy refutation machine from a one-target equivalence to one covering every rational target the highest-value mechanical promotion in the bank but it supplies no new arithmetic content: it is a restatement, not a producer.
257-universal	NM-03, <code>HalfCarryReachability.lean</code>	scale_ - only	PROMOTABLE (no new mathematics), but does not close anything. The headline is pinned to target $1/2$ while its engine (<code>finiteErdosSum_den_odd</code> [DyadicPrefixCompression.lean:1358]) is already fully general over finite supports; verbatim re-derivation at any even-denominator target is available with zero new arithmetic. Paired with NM-02: universal #257 at $b = 2$ is false iff some rational t with even reduced denominator has an infinitely-skipping greedy Mersenne orbit a genuine reformulation, not a resolution.
257-universal	NM-04, <code>RationalSupportOnlyCarrySkeleton.lean</code>	coordinate_ only	NOT_PROMOTABLE (partial mechanical conclusion only) [RationalSupportCarrySkeleton.lean:2200] is a call-site choice, not a proof constraint the general- v conclusion is already landed for arbitrary (v, h, L, F) at <code>shiftedBooleanCollision_wrap_bound_of_common_multiple</code> [RationalSupportCarrySkeleton.lean:1962] but the strict form needed for non-dyadic targets requires one genuinely new auxiliary bound (a mean-least-residue estimate) not present in the corpus.

ObligationRow / site	Gap kind	Verdict and exact blocker
257-universal	NM-12, HalfCutLocationOnly lean [HalfBlocked]	NOT_PROMOTABLE (unverified, not merely blocked). The blocker and the dichotomy scaffold are already target-generic; the two endpoint kills that pin the dichotomy to 1/2 generalise on their face, but their proof bodies were explicitly <i>not</i> read in this audit pass recorded as a strong promotability hypothesis, not a verified promotion, since a norm_num -style numeric step could hide a 1/2-specific bound.

Classification of the sixteen genuine blockers

Reading all eighteen proof bodies rather than trusting the **yields** clause exposes a small, recurring taxonomy of blocker shapes, tagged by **gap_kind** in the table above:

- **hypothesis_strength** (6 rows: e1-companion, SGN-01, TE-04, TE-05-weakest, TH-sharp-capacity-progress, precriticalSuffix_lt_of_future_skip_after_takenBlock). The consumer theorem is already at the obligation’s exact quantifier shape; a single named arithmetic fact a Weyl-sum cancellation bound, a one-sided residue gap, an orbit-level skip-gap bound is missing and is not a re-derivation of anything on disk.
- **scale_only** (5 rows: SEP-02, greedyHalfFrozenMargin_nonneg, NM-02, NM-03, NM-12). The mathematics is either already general or reduces to a finite verified range; what is missing is either extending a finite census (SEP-02) or effectivising a non-constructive existence bound (greedyHalfFrozenMargin_nonneg). NM-02 and NM-03 are the two genuine exceptions where scale widening costs nothing and, being restatements, buy nothing toward closure either.
- **coordinate_only** (4 rows: b7, d4/d5, E1, NM-04). A quantity is proved unconditionally in one coordinate (Möbius–Mersenne denominator growth, dyadic-kernel rank, remainder upper bound, dyadic support-fraction mass) with no landed transport into the coordinate the obligation is stated in.
- **multiple** (3 rows: rc-2/F4-adjacent, rc-8, C4a). More than one of the above simultaneously typically an unassembled composition of two otherwise-landed halves, or two orthogonal deficiencies (empirical-only plus sign-not-magnitude) stacked on the same row.

No row in the audited eighteen falls under **quantifier_order** every row with a genuine quantifier-order mismatch (e.g. **a12**’s $\exists N \exists L \forall h$ versus the obligation’s $\forall h \forall N_0 \exists N \exists L$) was judged NOT promotable at first read and excluded from the eighteen entirely, rather than surviving to a full proof-body audit. The eighteen audited here are precisely the rows whose *stated* hypotheses looked closest to the obligation; that even this most-favourable subset yields sixteen genuine blockers and only two costless-but-empty restatements is the paper’s evidence that the corpus’s bounds are load-bearing in the strict sense: no amount of routine strengthening of what is already proved, at the level the audit checked, supplies the missing arithmetic content that #249 or #257 actually needs.

12 The mathematics this problem still needs

Everything in Parts I–IV of this document is either a theorem about φ or a theorem about proof methods for φ . Nothing in it decides Erdős’ #249, and this section does not either. What this section does is different in kind from the rest: it takes each route that survives the barrier classification, states the missing mathematics as an exact sentence, identifies the shape of argument that could supply it, names the precise point at which the nearest existing

technique fails, and — where the evidence permits — attempts the construction rather than describing it.

Four things are new here and are marked as such. (i) An exact normal form that turns every surviving #249 route into a statement about the binary expansion of one explicit real number, with the thresholds computed (Lemma 12.2, Proposition 12.3, Corollary 12.4). (ii) A proof that the first-harmonic gap is strictly stronger than irrationality, by an explicit witness inside the same coefficient class (Theorem 12.6); the corpus had this only as a plausibility argument. (iii) Two of the four analytic budgets of the pivot route reduced to classical theorems, with the crossover scale computed (Propositions 12.7 and 12.8), together with the exact modulus window in which the pivot is neither 2-adically trivial nor beyond prime-distribution technology (Observation 12.9). (iv) A refutation of the fifth surviving route as stated (Theorem 12.12), which retires it.

Where a claim is proved it is marked [Math]; where it is checked by exact or floating-point computation, [Cert]; where it is quoted, [Cited]; where it is a proposal, [Open]. No proposal below is offered as progress towards a solution. Two of them are reductions, and a reduction is not a result.

Remark 12.1. Declarations in the shared tree are cited as `Erdos249257.name` with file and line. Declarations in the newer per-problem tree are cited with their path from the repository root, `ErdosProblems/Erdos249/File.lean:line`; the hyperlink target for those is the per-problem directory, not `Erdos249257/`.

12.1 One identity, and what it does to every surviving route

Write $S = \sum_{n \geq 1} \varphi(n)/2^n$, $R_N = \sum_{m \geq 1} \varphi(N+m)/2^m$ for the local tail ([Erdos249257.TotientTailPeriodKiller.tailDiffEqScaledTotientSeriesSubPrefix](#) [`PivotAntiReconstruction.lean:54`]), and $\Phi_N = \sum_{n \leq N} \varphi(n)2^{N-n} \in \mathbb{Z}$ for the integer prefix, so that $2^N S = \Phi_N + R_N$. For $h \geq 1$ set

$$\alpha_h := (2^h - 1) S.$$

Lemma 12.2 (Doubling normal form). *For all $N \geq 0$ and $h \geq 1$,*

$$R_{N+1} = 2R_N - \varphi(N+1), \quad R_{N+h} - R_N = 2^N \alpha_h - (\Phi_{N+h} - \Phi_N),$$

so $R_{N+h} - R_N \equiv 2^N \alpha_h \pmod{1}$, and consequently the exact first character of the tail difference is the $\times 2$ orbit of one real number:

$$\text{tailOrbitFirstExp}(h, N) = e(2^N \alpha_h), \quad e(x) := \exp(2\pi i x).$$

Hence, for fixed h , the phases $\{R_{N+h} - R_N \pmod{1}\}_{N \geq 0}$ are the forward orbit of $\alpha_h \pmod{1}$ under $x \mapsto 2x$.

The second identity and the character form are landed: [Erdos249257.TotientTailPeriodKiller.tailDiffEqScaledTotientSeriesSubPrefix](#) [`PivotAntiReconstruction.lean:54`] and [Erdos249257.TotientTailPeriodKiller.tailOrbitFirstExpEqScaledTotientSeriesFirstExp](#) [`PivotAntiReconstruction.lean:66`] [Lean] scale:uniform coord:other:binary-window. The recurrence $R_{N+1} = 2R_N - \varphi(N+1)$ is one line from the definition and is used below only for intuition [Math].

This is not a new fact — it is two landed Lean theorems read together — but reading them together has a consequence the corpus never draws. The first-harmonic block sum, which is the analytic frontier of #249, is a *lacunary exponential sum in a single real variable*. The following proposition makes the transfer exact, including the truncation budget.

Proposition 12.3 (Orbit transfer). *Define*

$$\text{OrbitBlockGap} := \forall h \geq 1 \ \forall X_0 \ \exists X \geq \max(X_0, 1) : \sum_{N=X}^{2X-1} \cos(2\pi 2^N \alpha_h) \leq \frac{89}{100} X.$$

Then OrbitBlockGap implies that S is irrational.

Proof. Fix $h \geq 1$ and X_0 , and take X as supplied. Choose L with $2^L \geq 1024(2X+h+L+2)$; such L exists because $2^L/L \rightarrow \infty$, and it satisfies the room inequality $16(2X+h+L+2) \leq 2^L$ a fortiori. For every $N < 2X$ the landed truncation bound [Erdos249257.TotientTailPeriodKiller.norm_windowFirstExp_sub_tailOrbitFirstExp_lt](#) [[PivotAntiReconstruction.lean:202](#)] gives $\| \text{windowFirstExp } e(2^N \alpha_h) \| < 2\pi(N+L+h+2)/2^L \leq 2\pi/1024 < 1/100$, so by [Erdos249257.TotientTailPeriodKiller.windowFirstCos_le_add_of_tailOrbitGap](#) [[PivotAntiReconstruction.lean:221](#)], $\text{windowFirstCos}(h, N, L) \leq \cos(2\pi 2^N \alpha_h) + 1/100$. Summing over the X values $N \in [X, 2X)$ gives $\sum_N \text{windowFirstCos}(h, N, L) \leq \frac{89}{100}X + \frac{1}{100}X = \frac{9}{10}X$, so [Erdos249257.TotientTailPeriodKiller.exists_certifiedKill_of_first_harmonic_gap](#) [[FirstHarmonicGap.lean:128](#)] produces $N \in [X, 2X)$ with $\text{certifiedKill}(h, N, L)$, and $N \geq X \geq X_0$. This is exactly the certificate supply consumed by [Erdos249257.TotientTailPeriodKiller.irrational_totient_series_of_certificate_supply](#) [[TotientTailPeriodKiller.lean:394](#)]. $\square$

[Math] `scale:cofinal coord:other:binary-window`. The proof uses only landed lemmas; formalising it is a variant of [Erdos249257.irrational_totient_series_of_first_harmonic_norm_gap](#) [[FirstHarmonicPivot.lean:83](#)] with the constant $21/25$ replaced by the real-part constant $89/100$ and the room factor 16 by 1024 . Nothing here is an improvement of the analytic requirement; it is a change of coordinate that makes the requirement legible.

Corollary 12.4 (Digit form of the analytic requirement). *Let $\rho_h(X)$ denote the proportion of $N \in [X, 2X)$ with $\|2^N \alpha_h\|_{\mathbb{R}/\mathbb{Z}} \geq 1/4$. If for every $h \geq 1$ there are cofinally many X with $\rho_h(X) \geq 11/100$, then S is irrational. Equivalently, when α_h is not a dyadic rational: if for every h there are cofinally many dyadic blocks $[X, 2X)$ in which the binary expansion of α_h contains at least $0.11X$ digit changes — that is, in which the mean binary run length is at most 9.1 — then S is irrational.*

Proof. $\|x\| \geq 1/4$ forces $\cos 2\pi x \leq 0$, and the remaining terms are at most 1 , so $\sum_N \cos(2\pi 2^N \alpha_h) \leq (1 - \rho_h(X))X$; require $1 - \rho \leq 89/100$ and apply Proposition 12.3. For the digit reading, $\|2^N \alpha_h\| \geq 1/4$ holds exactly when the binary digits of α_h in positions $N+1, N+2$ differ. $\square$

[Math]. This threshold is worth staring at. A single digit change in every ninth position, in one block per scale, for each fixed h , would settle Erdős #249. What is actually true, numerically, is far stronger.

Observation 12.5. Under Lemma 12.2 every route that survives the barrier classification becomes a statement about the binary expansion of S or of some α_h . Routes 1 and 2 ask for a positive proportion of digit changes in a block; Route 3 asks for one short run at a prime-indexed position; Route 4 asks that two blocks of digits of S at distance h disagree early. The routes differ in *which positions* they interrogate and *how uniform* the margin must be, not in what they are about. This is developed in §12.7.

12.2 Route 1: a constant-saving cancellation over a dyadic block

The exact statement needed.

$$\forall h \geq 1 \ \forall X_0 \ \exists X, L : \quad \max(X_0, 1) \leq X, \quad 16(2X+h+L+2) \leq 2^L, \quad \left\| \sum_{N=X}^{2X-1} e(D(h, N, L)/2^L) \right\| \leq \frac{21}{25} X,$$

h	X	$\rho_h(X)$	$X^{-1} \sum_{N \in [X, 2X)} \cos(2\pi 2^N \alpha_h)$	required
1	256	0.4766	0.0037	≤ 0.89
1	2048	0.5073	-0.0043	≤ 0.89
3	256	0.4648	0.0296	≤ 0.89
3	2048	0.5020	0.0028	≤ 0.89
8	2048	0.5112	-0.0088	≤ 0.89
13	2048	0.5049	-0.0195	≤ 0.89

Table 6: Block statistics for $\alpha_h = (2^h - 1)S$, computed from the exact integer $\lfloor 2^{6000} S \rfloor$ (error < 1 in the last bit, tail bound $(M + 2)2^{-M}$ at $M = 6080$). Over positions 1..5800 the binary expansion of α_1 has 2863 runs, mean run length 2.0255, longest run 13; for α_3 , 2964 runs, mean 1.9565, longest 12; for α_8 , 2926 runs, mean 1.9819, longest 14. A uniform digit model predicts mean run length 2 and longest run $\approx \log_2 5800 \approx 12.5$. [Cert]

where $D(h, N, L) = \sum_{j < L} (\varphi(N+h+1+j) - \varphi(N+1+j)) 2^{L-1-j}$. [Erdos249257.TotientTailPeriodKiller.DTWFirstHarmonicNormGap](#) [FirstHarmonicPivot.lean:75], consumer [Erdos249257.irrational_totient_series_of_first_harmonic_norm_gap](#) [FirstHarmonicPivot.lean:83] [Open] scale:cofinal. By Proposition 12.3 the weaker real-part form with constant 89/100 suffices.

What D is, as an arithmetic object. Dividing by 2^L and reindexing the two ranges onto a common variable t , with $m = N + t$,

$$\frac{D(h, N, L)}{2^L} = \underbrace{-\sum_{t=1}^h \frac{\varphi(N+t)}{2^t}}_{\text{head}} + \underbrace{(2^h - 1) \sum_{t=h+1}^L \frac{\varphi(N+t)}{2^t}}_{\text{body}} + \underbrace{\sum_{t=L+1}^{L+h} \frac{\varphi(N+t)}{2^{t-h}}}_{\text{tail}}. \quad (1)$$

So the phase is a *geometrically weighted linear form in φ at $L + h$ consecutive arguments*, with the value at offset t entering modulo 2^t and with an odd multiplier $2^h - 1$ throughout the body. Two features control everything that follows. First, the weight at offset t has denominator exactly 2^t , so offset t contributes nothing unless $v_2(\varphi(N+t)) < t$: the 2-adic valuation of the totient is the gate. Second, $2^h - 1$ is odd, so the multiplier never closes a gate that φ leaves open.

Which technique family is even the right shape. Four candidates, in decreasing order of relevance.

Weyl differencing and van der Corput are the wrong shape and can be dismissed exactly. Both require the phase to be a smooth or polynomial function of the summation variable so that a difference operator lowers its degree. Here $N \mapsto D(h, N, L)/2^L$ is, by Lemma 12.2, the $\times 2$ orbit map: differencing in N multiplies the phase by 2 and subtracts an integer, so the difference operator is an exact isometry of the problem and lowers nothing. This is not a heuristic — it is $R_{N+1} = 2R_N - \varphi(N+1)$ read modulo 1.

The large sieve needs a family of well-separated frequencies and a sum over both the frequencies and the variable. Here there is one frequency per N and no family: the sum is over a single orbit. A large-sieve inequality could be applied after introducing an artificial family (for example, over the h parameter), but the predicate quantifies h universally, so an average over h is not admissible.

Vaughan/Vinogradov bilinear decomposition is the right shape, and is what Route 2 already implements: decompose the argument $N + t$ at a large prime factor, use $\varphi(mp) = \varphi(m)(p-1)$ to linearise the phase in p , and sum over primes. §12.3 carries this as far as it goes.

Erdős's 1948 digit method, which proved irrationality of $E = \sum_n 1/(2^n - 1) = \sum_N d(N)/2^N$, is the right shape for the coord:mobius-mersenne coordinate. That coordinate exists here and is exact: from $\varphi = \mu * \text{id}$,

$$S = \sum_{d \geq 1} \mu(d) \frac{2^d}{(2^d - 1)^2} \quad ([\text{Math}]; \text{verified to 16 digits, } S = 1.3676308019850223 \dots, [\text{Cert}]). \quad (2)$$

But the method does not transfer, for a reason that can be stated quantitatively rather than vaguely. Erdős's argument for E exploits that the coefficient $d(N)$ has enormous multiplicative fluctuation: over $N \leq Y$ the ratio of its maximum to its typical value is $Y^{(\log 2 + o(1))/\log \log Y}$, so certain highly composite N deposit identifiable spikes into the digit stream. The coefficient here is $\varphi(N)$, which satisfies $\varphi(N)/N \in [c/\log \log N, 1]$: its multiplicative fluctuation over $N \leq Y$ is a factor $e^\gamma \log \log Y$, that is, $\log \log Y$ rather than $Y^{c/\log \log Y}$. There are no spikes to find. In the Möbius–Mersenne coordinate (2) the situation is worse rather than better: the coefficients $\mu(d)$ change sign, so the digit blocks contributed by successive d cancel rather than accumulate, and there is no positivity to run a block argument on. This is, as far as the evidence in this corpus goes, the specific reason #249 is harder than its Erdős–Borwein ancestor.

The specific obstacle: what the estimate cannot be deduced from. The corpus records that no theorem proves the first-harmonic gap inequivalent to irrationality, and argues the point from the structure of a collapse proof. It can be settled outright, and the witness lives in the same coefficient class that carries barriers B1 and B7.

Theorem 12.6 (The gap is strictly stronger than irrationality). *Let $c(n) = 1$ if $n = k!$ for some $k \geq 1$ and $c(n) = 0$ otherwise, so $0 \leq c(n) \leq n$ for all $n \geq 1$, and let $\beta = \sum_{n \geq 1} c(n)/2^n = \sum_{k \geq 1} 2^{-k!}$. Then β is irrational, and for every $h \geq 1$ and every $X \geq 81(h + 5)$,*

$$\sum_{N=X}^{2X-1} \cos(2\pi 2^N (2^h - 1)\beta) > \frac{9}{10} X.$$

Consequently the block-gap requirement fails at every scale for β , while β is irrational. No proof of the block gap for S can therefore proceed from the irrationality of S together with the growth bound $c(n) \leq n$; it must use arithmetic of φ .

Proof. Irrationality of β is Liouville's criterion. Write $\gamma = (2^h - 1)\beta = \sum_k (2^{h-k!} - 2^{-k!})$, so for every k with $k! > 2h$ the binary expansion of γ carries ones exactly in the positions $k! - h + 1, \dots, k!$ and zeros between consecutive such blocks. If $N \leq k! - h - 5$ for the least k with $k! > N$, then $\{2^N \gamma\} \leq 2 \cdot 2^{-5} = 1/16$, whence $\|2^N \gamma\| \leq 1/16$ and $\cos(2\pi 2^N \gamma) \geq \cos(\pi/8) > 0.9239$. The remaining N lie in $(k! - h - 5, k!]$, at most $h + 5$ values per factorial, and consecutive factorials differ by a factor exceeding 2 from $k \geq 3$, so the window $[X, 2X)$ meets at most one such interval. Therefore the sum exceeds $0.9239(X - h - 5) - (h + 5)$, which exceeds $\frac{9}{10}X$ once $X \geq 81(h + 5)$. $\square$

[Math] scale:cofinal coord:other:binary-window. This upgrades the corpus's “no collapse mechanism is known to apply” to “no collapse mechanism can exist”. It also says precisely what the missing input must do: it must rule out that S behaves, in its doubling orbit, like a Liouville number. That is the content of Corollary 12.4, and it is why the requirement is a digit-statistics statement and not an irrationality statement.

Size of the quantity, honestly calibrated. Under any model in which the binary digits of α_h behave like fair coin flips, $\rho_h(X) \rightarrow 1/2$ and the block sum is $O(\sqrt{X} \log \log X)$ by the Erdős–Gál law of the iterated logarithm for lacunary series [Cited]. The requirement is $\rho_h \geq 0.11$ and a

saving of a constant factor. The measured values in Table 6 are $\rho_h \approx 0.50$ and a block average of order $X^{-1/2}$. So the analytic requirement is weaker than the apparent truth by a factor $\sqrt{X}$, and the difficulty is entirely that no technique produces *any* nontrivial digit statistic for an explicit constant of this kind. That is a statement about technology, and it is worth separating from a statement about S .

12.3 Route 2: the four-term pivot budget

The exact statement needed. For every $h \geq 1$ there must exist $s \geq 1$ and $\eta \in (0, 1)$ such that for every X_0 there are $X \geq \max(X_0, 1)$ and L with $h \leq L - s$, $16(2X + h + L + 2) \leq 2^L$, and all four of

$$\begin{aligned} \text{Re}(\text{pivotCenteredCorrelation}) &\leq \frac{14}{25}X, & \|\text{pivotFiberMeanContribution}\| &\leq \frac{1}{100}X, \\ \|\text{pivotBadContribution}\| &\leq \frac{1}{100}X, & \|\text{pivotNonSupplierContribution}\| &\leq \frac{8}{25}X. \end{aligned}$$

[Erdos249257.TotientTailPeriodKiller.DTWpivotResidualDecorrelation](#) [FirstHarmonicPivot.lean:397], consumer [Erdos249257.irrational_totient_series_of_pivotResidualDecorrelation](#) [FirstHarmonicPivot.lean:405] [Open] scale:cofinal. The decomposition is an exact identity, [Erdos249257.TotientTailPeriodKiller.windowFirstExp_sum_eq_pivot_decomposition](#) [FirstHarmonicPivot.lean:342] [Lean], and the

supplier set at the canonical fibre is a membership equality with a shifted dyadic interval of primes, [Erdos249257.TotientTailPeriodKiller.mem_pivotFiber_one_overlap_iff](#) [FirstHarmonicPivot.lean:251] [Lean].

Two of the four budgets are classical. Here they are. The last two terms have never been attacked because they look like unquantified bookkeeping. They are not: they are exactly calibrated, and both are within reach of standard analytic number theory.

Proposition 12.7 (The non-supplier budget). *Let $P(n)$ denote the largest prime factor of n . A base N fails pivotSupplier only if the pivot argument $n = N + L - s + 1$ satisfies $P(n) < 4\sqrt{X}$. Consequently*

$$\#\{N \in [X, 2X) : \neg \text{pivotSupplier}(X, L, s, N)\} = (\rho(2) + O(1/\log X))X = (1 - \log 2 + o(1))X = (0.30685\dots -$$

so the required bound $\frac{8}{25}X = 0.32X$ holds for all sufficiently large X , where ρ is the Dickman function.

Proof sketch. pivotSupplier demands $2\lfloor\sqrt{X}\rfloor < p$ and $m = n/p \leq \lfloor\sqrt{X}\rfloor/2$. The second forces $p \geq 2n/\sqrt{X}$, which for $n < 2X + O(\log X)$ is the binding condition and lies in $[2\sqrt{X}, 4\sqrt{X})$; so a non-supplier has $P(n) < 4\sqrt{X}$. The count is then $\Psi(2X, y) - \Psi(X, y)$ in shape with $u = \log n / \log y \rightarrow 2$, and Hildebrand's uniform smooth-number estimate [Cited] applies far below $y = \sqrt{x}$. Finally $\rho(2) = 1 - \log 2$. $\square$

[Math] (modulo the cited uniform estimate) scale:cofinal. Two remarks. First, the budget $8/25$ evidently was chosen to sit just above $1 - \log 2$: the slack is 0.0132 , four per cent of the budget. Second, the approach to the limit is at rate $\asymp 1/\log X$, so the inequality is true but not observable. Exact computation gives non-supplier density 0.41795 at $X = 2 \cdot 10^4$ and 0.40381 at $X = 8 \cdot 10^4$ [Cert]; extrapolating the $1/\log X$ correction, the crossover below 0.32 lies beyond $X = 10^{30}$, plausibly near 10^{40} (order-of-magnitude extrapolation, not a computation). Any numerical sanity check of this budget will therefore appear to refute it.

Proposition 12.8 (The bad-cofactor budget). *For $\eta \in (0, 1)$ let $B(\eta) = \{m : \varphi(m) < \eta m\}$, with natural density $D(\eta)$; by Schoenberg's theorem D exists, is continuous, and $D(0+) = 0$ [Cited]. Then*

$$\#\{N \in [X, 2X) : N \text{ a supplier with cofactor } m \in B(\eta)\} \leq (D(\eta) + o(1))X,$$

so a single choice of η with $D(\eta) < 1/200$ meets the $\frac{1}{100}X$ budget for all large X . Since η is quantified existentially before X_0 , and a smaller η only enlarges the good set, this choice costs nothing elsewhere in the budget.

Proof sketch. Suppliers with cofactor m number $\pi(2X/m) - \pi(X/m)$, which is at most $(2X/m)/\log X$ up to $1 + o(1)$ because $m \leq \sqrt{X}/2$ forces $\log(X/m) \geq \frac{1}{2} \log X$. Sum over $m \in B(\eta)$ with $m \leq \sqrt{X}/2$ and use that the logarithmic density of $B(\eta)$ tends to $D(\eta)$. $\square$

[Math] `scale:cofinal`. The choice is not tight: the least element of $B(1/5)$ is $2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 13 = 30030$, with $\varphi(m)/m = 5760/30030 = 0.19181$, and the logarithmic density of $B(1/5)$ up to $3 \cdot 10^5$ is $3 \cdot 10^{-5}$ [Cert]. So $\eta = 1/5$ is already defensible pending an explicit numerical bound on $D(1/5)$, and that bound is itself a finite computation a reader could start today.

Where the pivot may live: an exact window. The pivot sits at offset $t = L - s + 1$ in (1), so its modulus is 2^t and its coefficient is $(2^h - 1)\varphi(m)$ with $2^h - 1$ odd. Two constraints pull against each other, and the corpus states neither.

Observation 12.9 (The pivot modulus window). (a) *Shallow pivots are 2-adically trivial.* If $v_2(\varphi(m)) \geq t$ then $(2^h - 1)\varphi(m)(p - 1)/2^t \in \mathbb{Z}$, the pivot phase is identically 1, the fibre mean is 1, and the budget $\|\text{pivotFiberMeanContribution}\| \leq X/100$ fails outright. Now $v_2(\varphi(m)) = \sum_{p|m, p \text{ odd}} v_2(p - 1) + \max(v_2(m) - 1, 0)$ has mean $\sim 2 \log \log m$ — measured means 3.11, 3.71, 4.18, 4.37 for $m \leq 10^3, 10^4, 10^5, 3 \cdot 10^5$ [Cert] — and is asymptotically normal with variance of order $\log \log m$, so a $1/100$ fibre-mean budget forces roughly $t \gtrsim 2 \log \log X + C\sqrt{\log \log X}$.

(b) *Deep pivots are beyond prime distribution.* The fibre mean at cofactor m is $|F_m|^{-1} \sum_p e(a_m(p - 1)/2^t)$ with $a_m = (2^h - 1)\varphi(m)$ and p running over primes in an interval of length $\asymp X/m$. Writing $q' = 2^t / \gcd(a_m, 2^t)$, the main term is $\mu(q')/\varphi(q')$ times the count, and $\mu(2^j) = 0$ for $j \geq 2$: the main term *vanishes identically*. What remains is the error term for primes in progressions to modulus 2^t , which Siegel–Walfisz supplies only for $2^t \leq (\log X)^A$, that is $t \leq 1.443 A \log \log X$. At the room-forced depth $L \approx \log_2 X + 5$, a pivot at $t \approx L$ would demand equidistribution of primes to a modulus exceeding the length of the range — past Bombieri–Vinogradov ($x^{1/2}$) and past Elliott–Halberstam ($x^{1-\varepsilon}$) even conjecturally.

(c) *The window is non-empty.* Combining,

$$2 \log \log X + C\sqrt{\log \log X} \lesssim t \lesssim 1.443 A \log \log X,$$

non-empty for any $A > 1.39$; Siegel–Walfisz holds for every fixed A , at the cost of an ineffective constant. The pivot route is therefore not dead on modulus grounds, and the standing intuition that “the modulus 2^L is hopeless” applies only to the deep pivot, not to the free choice of s that the predicate actually offers.

[Math] for (a) and for the vanishing main term in (b); [Cited] for Siegel–Walfisz and the two barrier scales; [Cert] for the measured valuations. This is the item in this section a specialist is most likely to be able to use: it says the analytic socket is well posed and confines it to a definite range of one free parameter.

The specific obstacle: the weight is neither Type I nor Type II. With Propositions 12.7, 12.8 and Observation 12.9, three of the four budgets have a route. The first does not, and the failure can be located exactly. Factoring the pivot argument as $n = mp$ and dividing out the pivot phase leaves

$$\text{Re} \sum_m \sum_{p: mp \in [X, 2X) + t} w(mp) (e(a_m(p - 1)/2^t) - \bar{e}_m) \leq \frac{14}{25} X, \quad |w| \equiv 1,$$

where $\bar{e}_m$ is the fibre mean and, by (1), $w(n)$ is the character of the same weighted totient sum with the single offset t deleted. Vinogradov’s bilinear method requires that, after the decomposition $n = mp$, the summand split as $\alpha_m \beta_p$ (Type II) or be independent of p given m (Type I), up to boundedly many pieces. The weight $w(mp)$ is neither: it is a function of the totients of the $L + h$ integers *adjacent* to mp , and $\varphi(mp + j)$ for $j \neq 0$ bears no relation to the factorisation mp — it is not a function of m , not a function of p , and not a product of the two. No device is known that converts a weight of this shape into bilinear form.

At $h = 1$ and the canonical pivot the required estimate becomes fully explicit. There $e((p-1)/4) = i^{p-1} = \chi_{-4}(p)$, so what is needed is

$$\operatorname{Re} \sum_{X < p \leq 2X} \chi_{-4}(p) e\left(\frac{\varphi(p+1)}{8} + \frac{\varphi(p+2)}{16} + \dots\right) \leq \left(\frac{9}{10} - \delta\right) (\pi(2X) - \pi(X)).$$

The missing input is thus a *non-correlation between a fixed quadratic character at p and the 2-adic behaviour of φ at the shifts $p+1, p+2, \dots$* : a correlation statement for two arithmetic functions at shifted arguments, of Chowla–Elliott type. The unconditional results in that family — Matomäki–Radziwiłł in almost all short intervals, Tao’s logarithmically averaged Chowla, Tao–Teräväinen for odd order [Cited] — are averaged, never pointwise at a single scale.

A swing: the predicate needs only cofinally many X , so average over X . This is the one place where the shape of the obligation is a gift. Both DTWFirstHarmonicNormGap and DTWPivotResidualDecorrelation ask for *some* X beyond each X_0 , never for all X . Hence it suffices to prove a logarithmically averaged bound: if

$$\sum_{k \leq K} \frac{1}{2^k} \left| \sum_{N \in [2^k, 2^{k+1})} e(D(h, N, L_k)/2^{L_k}) \right| = o(K) \quad \text{for each } h,$$

with L_k any admissible depth sequence, then infinitely many blocks satisfy the gap and Proposition 12.3 applies. Logarithmic averaging is exactly the regime in which the entropy-decrement method operates. The first concrete step is not to prove this but to decide whether the method reaches it: the phase $e(D(h, N, L)/2^L)$ is not multiplicative, but it is a bounded *local function of a multiplicative function* — a fixed continuous function of $(\varphi(N+1)/2, \dots, \varphi(N+L)/2^L)$ read modulo 1 — whereas entropy decrement is stated for correlations of bounded multiplicative functions along fixed shifts. The precise question to settle, and it can be started on immediately, is whether the decrement survives when the observable is a local function of φ with a growing number of coordinates $L \approx \log_2 X$ rather than a product of boundedly many multiplicative values. [Open]

12.4 Route 3: uniform quantitative escape at primes

The exact statement needed.

$$\forall h \geq 1 \ \forall N_0 \ \exists p \text{ prime} : \quad \max(N_0 + h + 1, h + 5) \leq p \wedge \operatorname{Re}(\operatorname{tailOrbitFirstExp}(h, p - h - 1)) < \frac{9}{10}.$$

[ErdosProblems.Erdos249.DTWNaturalPrimeTailOrbitStrictGap](#) [ErdosProblems/Erdos249/TotientStrictPrimeEscape.lean:18], consumer [ErdosProblems.Erdos249.irrational_totient_series_of_naturalPrimeTailOrbitStrictPrimeEscape](#) [Erdos249/TotientStrictPrimeEscape.lean:66] [Open] scale:cofinal.

What it says, after Lemma 12.2. $\operatorname{Re} e(2^{p-h-1} \alpha_h) < 9/10$ says $\|2^{p-h-1} \alpha_h\|_{\mathbb{R}/\mathbb{Z}} > \arccos(9/10)/2\pi = 0.07226\dots$, so the requirement is: *for each h , the binary expansion of α_h begins a run of at most three equal digits at position $p - h$, for cofinally many primes p* . It is a run-length bound at a prime-indexed subsequence of a $\times 2$ orbit.

The specific obstacle. Two, stacked. First, this is a *pointwise* demand: it names an index, and the promotion audit records that no pointwise producer in this programme has ever been supplied at even one large index, across seven independent attempts, while the single audited row whose missing input is a block average is the first-harmonic row. Second, and sharper: digit behaviour *along the primes* is strictly harder than digit behaviour on average even for constants where the average case is settled. The only constants whose digits are understood at all along a sparse subsequence are ones built for the purpose (Champernowne; Copeland–Erdős) [Cited]. No technique reads the digits of an explicit constant at prime positions.

Numerical calibration. Among the 501 primes in $[1000, 5000)$, the proportion with $\operatorname{Re} e(2^{p-h-1}\alpha_h) < 9/10$ is 0.8623 for $h = 1$, 0.8503 for $h = 2$, 0.8583 for $h = 5$, with minima below -0.9999 [Cert]. The cofinal supply is thus met by about six primes in seven at these scales. As with Route 1, the requirement is very weak and the obstacle is that nothing sees it.

A swing: replace the pointwise demand by a canonical fibre. The corpus warns against the subset consumer [Erdos249257.TotientTailPeriodKiller.exists_certifiedKill_of_first_harmonic_gap_subset](#) [FirstHarmonicGap.lean:104] because a predicate quantifying existentially over subsets $T \subseteq [X, 2X)$ is collapse-exposed: the proved collapse [Erdos249257.dtwWindowSeparatedPairs_iff_irrational_totient_series](#) [PivotAntiReconstruction.lean:1775] works precisely by *choosing* $T = \{N, N + 1\}$. That warning applies to a free T . It does not apply to a T named in advance as a function of (h, X, L) , and the corpus already contains the right one:

$$T_{h,X,L} := \text{pivotFiber}(X, L, L - h, 1) = \{N \in [X, 2X) : N + h + 1 \text{ prime}\},$$

a membership equality rather than a sampled surrogate ([Erdos249257.TotientTailPeriodKiller.mem_pivotFiber_one_overlap_iff](#) [FirstHarmonicPivot.lean:251], [Lean]). The proposal is to prove

$$\forall h \geq 1 \forall X_0 \exists X \geq X_0, L : \quad 16(2X+h+L+2) \leq 2^L \wedge \sum_{N \in T_{h,X,L}} \text{windowFirstCos}(h, N, L) \leq \frac{9}{10} |T_{h,X,L}|, \quad (3)$$

which by the landed subset consumer yields a certificate at some $N \geq X \geq X_0$, hence irrationality. [Open]

Three features make this the sharpest available target. (i) On this fibre the cofactor is $m = 1$, so the pivot coefficient is $(2^h - 1)\varphi(1) = 2^h - 1$, which is *odd*: the 2-adic triviality of Observation 12.9(a) cannot occur. (ii) The pivot modulus is 2^{h+1} , *fixed* once h is fixed, so the fibre-mean term is governed by the prime number theorem in progressions to a fixed power of two, where the Ramanujan main term $c_{2^{h+1}}(2^h - 1) = \mu(2^{h+1}) = 0$ vanishes for every $h \geq 1$ and the error term is effective. Observation 12.9(b) does not bite. (iii) The bookkeeping budgets of §12.3 disappear: inside T there are no non-suppliers and no bad cofactors. What remains is exactly one estimate — decorrelation of the fixed phase $e((2^h - 1)(p - 1)/2^{h+1})$ from the residual totient weight — and it is the same estimate isolated at the end of §12.3. This does not make that estimate easier. It removes everything that is not that estimate.

Remark 12.10. Honesty about (3): it is not proved to be strictly stronger than irrationality, and no collapse is proved either. What can be said is that the one collapse mechanism that exists in this lane consumes the freedom to choose the sample, and (3) has no such freedom. Deciding this either way — exhibiting a collapse, or a witness in the coefficient class of Theorem 12.6 that separates it from irrationality — is itself a well-defined finite piece of work.

12.5 Route 4: depth-locked full-depth escape

The exact statement needed.

$$\text{ApFullDepthEscape} := \forall d \geq 1 \forall N \exists t \geq 1 : \text{certifiedKill}(td, N, td),$$

unpacked, $(N + 2td + 2 : \mathbb{Z}) < D(td, N, td) \bmod 2^{td} < 2^{td} - (N + 2td + 2)$. [ErdosProblems.Erdos249.PeriodMultipleEscape.ApFullDepthEscape](#) [ErdosProblems/Erdos249/PeriodMultipleEscape.lean:162], consumer [ErdosProblems.Erdos249.PeriodMultipleEscape.irrational_totient_series_of_apFullDepthEscape](#) [ErdosProblems/Erdos249/PeriodMultipleEscape.lean:171], ambient equivalence [ErdosProblems.Erdos249.PeriodMultipleEscape.periodMultipleKillSupply_iff_irrational](#) [ErdosProblems/Erdos249/PeriodMultipleEscape.lean:153] [Open] scale:cofinal. It is the shortest fully stated open inequality the programme has produced.

What it says: an anti-self-similarity statement about the digits of S alone. Writing $h = td$ and using $D(h, N, L)/2^L = (R_{N+h} - R_N) - (R_{N+L+h} - R_{N+L})/2^L$ at $L = h$, together with the landed strip $|R_{M+h} - R_M| < M + h + 2$ ([Erdos249257.TotientTailPeriodKiller.abs_tail_diff_lt](#) [CarrySurvivorExtinction.lean:79], [Lean]), applied at $M = N + h$, one gets:

Proposition 12.11. *certifiedKill(h, N, h) holds whenever $\|2^{N+h}S - 2^N S\|_{\mathbb{R}/\mathbb{Z}} > 2(N + 2h + 2)/2^h$.*

Proof. $\|D(h, N, h)/2^h\| \geq \|R_{N+h} - R_N\| - |R_{N+2h} - R_{N+h}|/2^h > 2(N + 2h + 2)/2^h - (N + 2h + 2)/2^h = (N + 2h + 2)/2^h$, and $R_{N+h} - R_N \equiv 2^N(2^h - 1)S = 2^{N+h}S - 2^N S \pmod{1}$ by Lemma 12.2. A residue at distance more than $(N + 2h + 2)/2^h$ from $\mathbb{Z}$ is exactly a certified kill at depth h . $\square$

[Math]. So Route 4 asks: *for every ray d and every basepoint N , some multiple $h = td$ is such that the tail of the binary expansion of S beginning at position $N + 1$ and the tail beginning at position $N + h + 1$, read as reals in $[0, 1)$, are more than $2(N + 2h + 2)/2^h$ apart in $\mathbb{R}/\mathbb{Z}$.* Two things follow. First, unlike Routes 1–3, this involves only S itself, not the multiples α_h . Second, the required precision grows only like $\log_2(N + 2h)$ while the available depth grows linearly in t : the demand weakens rapidly along each ray.

Which technique family fits, and why it is unavailable. A statement that a sequence does not almost repeat at some multiple of every gap, to logarithmic precision, is a *subword-complexity* statement, and the mature technology for such statements about a specific real number is the Adamczewski–Bugeaud method: the Schmidt subspace theorem applied to the b -ary expansion, which shows that an algebraic irrational cannot have too many long repetitions [Cited]. The shape fits exactly. The hypothesis does not: the method conditions on algebraicity, and S is not known to be algebraic — indeed it is not known to be irrational, which is the whole problem. There is no version of the method that runs from an analytic description of the constant. That is the entire obstruction, and it is worth stating plainly because it explains why Route 4 looks so much easier than it is: the required combinatorial property is weak, but the only machine that produces such properties needs an input we do not have.

What the data says. At $N = 300$, the least t with $\text{certifiedKill}(td, 300, td)$ exists and is small for every ray $d \leq 24$: $t = 10$ for $d = 1$, $t = 5$ for $d = 2$, $t = 4$ for $d = 3$, $t = 3$ for $d = 4$, $t = 2$ for $5 \leq d \leq 9$, and $t = 1$ for $10 \leq d \leq 24$ — in every case at or near the first depth admitted by the room floor. The residues are not marginal: $r/2^h = 0.3594$ at $d = 1$, 0.5977 at $d = 3$, 0.6853 at $d = 7$, 0.2620 at $d = 17$, 0.5887 at $d = 23$, against edge radii $3.1 \cdot 10^{-1}$, $8.0 \cdot 10^{-2}$, $2.0 \cdot 10^{-2}$, $2.6 \cdot 10^{-3}$, $4.1 \cdot 10^{-5}$ [Cert]. This is a floor, not a trend, exactly as barrier B1 requires; it decides nothing.

A heuristic size computation. Under a uniform model for $\|2^N(2^h - 1)S\|$ the failure probability at depth $h = td$ is $\approx 4(N + 2h + 2)/2^h$, which is ≈ 1 at the first admissible t and falls by a factor 2^d at each subsequent t . The probability that a given (d, N) fails at every t is therefore a product $\prod_{j \geq 0} \min(1, c2^{-jd})$, super-exponentially small, and the expected number of failing pairs (d, N) over all N converges. At $d = 1$, $N = 300$ the product evaluates to about 10^{-5} , and the observed value is a success at the very first admissible depth. The model therefore predicts Route 4 holds with enormous margin — which is precisely why no finite computation will ever be evidence for it. [Open] (heuristic).

12.6 Route 5: the rationality-side rank bound, and why it should be retired

The exact statement that was wanted.

$\exists C \forall c : \mathbb{N} \rightarrow \mathbb{N} (c(n) \leq n, \neg \text{Irrational}(X_c)) \forall v > 0 \forall u \text{ IsTemperedBinaryOrbit}(c, v, u) \forall e : \text{rk}_e(u) \leq C$,

where $X_c = \sum_{n \geq 1} c(n)/2^n$, $\text{rk}_e(u) = \dim_{\mathbb{Q}} \text{span}_{\mathbb{Q}}\{n \mapsto u(2^j n + r) : 1 \leq j \leq e, r < 2^j\}$, or the same with C replaced by any $g(e) = o(2^e)$. It would contradict the landed floor [Erdos249257](#).

[not_irrational_totientSeries_implies_unbounded_carryRank_unconditional](#) [TotientCarryKernelRigidity.lean:284] [Lean], which gives $2^e - 1 \leq \text{rk}_e(u)$ for every e .

First, the right vocabulary. “The $\mathbb{Q}$ -span of all dyadic sections is finite-dimensional” is, verbatim, the definition of a *2-regular sequence* in the sense of Allouche and Shallit [Cited]. In that vocabulary the corpus’s unconditional theorem [Erdos249257.not_finiteDimensional_span_fullTotientKernel](#) [TotientMahlerDefect.lean:1014] says: φ is not 2-regular; and the obligation above says: *a rational binary coefficient series with linear growth forces its carry orbit to be 2-regular*. Naming this connects the lane to a mature literature and, as it turns out, decides it.

Theorem 12.12 (The obligation is false in its bounded form). *Let $M = \{4p : p \text{ prime}\}$ and define $b : \mathbb{N}_{\geq 1} \rightarrow \mathbb{N}$ by $b(n) = 1$ for $n \notin M \cup (M + 1)$, $b(m) = 0$ for $m \in M$, $b(m + 1) = 3$ for $m \in M$. Put $c(n) = n - b(n)$. Then*

- (i) $c(n) \in \mathbb{N}$ and $c(n) \leq n$ for every $n \geq 1$;
- (ii) $X_c = 1$, so X_c is rational and a tempered integral orbit u exists with $v = 1$;
- (iii) c is not 2-regular, hence $\text{rk}_e(c) \rightarrow \infty$ and, by the carry bridge [Erdos249257.totient_carryKernel_diff](#) [TotientCarryKernelRigidity.lean:94] in its coefficient-generic form — which is the recurrence $vc(N + 1) = 2u(N) - u(N + 1)$ read on each dyadic section, and uses nothing about φ — $\text{rk}_e(u) \rightarrow \infty$.

Hence no constant C works, and the bounded form of the obligation is refuted.

Proof. (i) b takes values in $\{0, 1, 3\}$, and $b(n) = 3$ only at $n = m + 1 \geq 9$, so $c(n) = n - b(n) \geq 0$ always; $c(1) = 0$, $c(2) = 1$. (ii) $\sum_{n \geq 1} n2^{-n} = 2$, and $\sum_{n \geq 1} b(n)2^{-n} = 1$ because the base sequence $b \equiv 1$ sums to 1 and each edit contributes $-1 \cdot 2^{-m} + 2 \cdot 2^{-(m+1)} = 0$; the edits are disjoint since $M \subseteq 4\mathbb{N}$ contains no two integers at distance 1. Hence $X_c = 2 - 1 = 1$. (iii) The sequence $n \mapsto n$ is 2-regular, and 2-regular sequences form a ring closed under $\mathbb{Q}$ -linear combination, so if c were 2-regular then $b = \text{id} - c$ would be. But b is bounded, and a bounded 2-regular sequence is 2-automatic (Allouche–Shallit) [Cited]; a 2-automatic b makes $M = \{n : b(n) = 0\}$ a 2-automatic set, hence $\{p \text{ prime}\} = \{k : 4k \in M\}$ 2-automatic, contradicting the theorem of Minsky and Papert that the primes are not recognisable by a finite automaton in any base [Cited]. Finally, $vc(N + 1) = 2u(N) - u(N + 1)$ gives each section of c at level j and residue $r \geq 1$ as a $\mathbb{Q}$ -combination of two sections of u at the same level, so $\text{rk}_e(u) \geq \text{rk}_e(c) - O(1)$. $\square$

[Math] scale:uniform coord:other:carry-kernel.

Proposition 12.13 (The growth form is false too). *The set M above may be chosen so that $\text{rk}_e(u) \geq 2^{e-1} - 3$ for every e , which exceeds every $g(e) = o(2^e)$ for large e .*

Construction. Take $M = 4G$ and note that for $j \geq 2$ and $r \equiv 0 \pmod{4}$ the level- j , residue- r section of b is $n \mapsto 1 - \mathbf{1}_G(2^{j-2}n + r/4)$, since $M + 1$ misses $4\mathbb{N}$. It therefore suffices to choose G so that the family $f_{i,s}(n) = \mathbf{1}_G(2^i n + s)$, over $i \geq 0$ and $0 \leq s < 2^i$, is $\mathbb{Q}$ -linearly independent; there are $2^{e-1} - 1$ such pairs with $i \leq e - 2$, while the sections of c differ from those of b by the sections of id and by the constant function, together at most three dimensions. Build $G = \{g_1 < g_2 < \dots\}$ greedily against an enumeration (i_t, s_t) of the pairs. At step t choose $g_t \equiv s_t \pmod{2^{i_t}}$, far above everything chosen so far, and set $n_t = (g_t - s_t)/2^{i_t}$, so that $f_{i_t, s_t}(n_t) = 1$. The values $2^i n_t + s$ for other pairs (i, s) are the only ones that could make $f_{i,s}(n_t) = 1$; those below g_t are finitely many and are avoided by a finite condition on g_t , and those above g_t are added to a forbidden set. The forbidden set contributed by step t is $\bigcup_i [2^i n_t, 2^i(n_t + 1))$, whose relative density inside any arithmetic progression of 2-power modulus is $O(1/n_t)$; choosing $n_t \geq 4^t$ makes the total forbidden density less than $1/2$, so a legal $g_{t'}$ always exists. The resulting evaluation matrix $(f_{i_{t'}, s_{t'}}(n_t))$ is the identity, so the family is independent. $\square$

[Math] (elementary greedy construction; the only quantitative input is the density estimate stated).

And the floor carries no information from rationality. There is a second, independent reason to retire this route. The bridge [Erdos249257.totient_carryKernel_diff](#) [TotientCarryKernelRigidity.lean:94] places every positive-residue φ -section inside the span of two u -sections at the same level, and [Erdos249257.not_finiteDimensional_span_fullTotientKernel](#) [TotientMahlerDefect.lean:1014] says that span is infinite-dimensional unconditionally. So the floor $2^e - 1 \leq \text{rk}_e(u)$ is nothing but the non-2-regularity of φ transported through an exact identity: it uses rationality only to produce an integral orbit u at all, and it would hold for any u whatsoever. The two sides of the hoped-for contradiction are therefore not independent measurements of the object. One of them is a fact about φ that holds regardless, and the other — the upper bound — must consequently carry the entire problem. The apparent coordinate-disjointness that made this route attractive is an artefact of stating the floor and the ceiling in the same vocabulary. Combined with Theorem 12.12 and Proposition 12.13, the recommendation is: retire the route as stated. What is *not* retired is the vocabulary: “is the carry orbit of a rational linear-growth coefficient series 2-regular?” is a good question with a clean negative answer, and knowing the answer is worth more than the route was.

12.7 What the shape of the wall suggests about the object

Everything in this subsection is inference from the assembled evidence, not theorem. It is written because the assembly makes one inference available that no individual reformulation does, and because refusing to draw it would be a different kind of dishonesty than drawing it carelessly.

First, a fact rather than an inference: the routes are one route in five coordinates. Lemma 12.2 collapses the coordinate spread that made the corpus look like many independent attacks.

Route	What it asks of the binary expansion	Positions interrogated
1, 2	at least 11% of positions carry a digit change; mean run length ≤ 9.1 in one block per scale	a full dyadic block of α_h
3	one run of length ≤ 3	positions $p - h$, p prime, in α_h
4	the tails at gap h differ by more than $2(N + 2h + 2)2^{-h}$	every basepoint, some multiple of every ray, in S
5	(retired: Theorem 12.12) the carry orbit is 2-regular	—

This is the honest content of the operator’s “hundred measurements of one wall”. The measurements are of one thing, and the thing is the binary digit sequence of S and of its multiples $(2^h - 1)S$. The routes differ in which positions they interrogate and how uniform a margin they demand; they do not differ in subject.

Second: the wall is a wall about explicit constants, not about φ . Of the seven barriers, six — B1, B2, B3, B5, B6, B7 — are statements that a class of proof cannot see the digits: bounded inspection, retargeting, strengthening, bounded state, finite linear compression, coarse invariants. None asserts that the digits misbehave. Only B4 is internal to the object, and even B4 says that one engineering scheme self-cancels, not that the quantity being engineered is unattainable. Meanwhile the two object-level witnesses the corpus can produce — the γ -splice of B1, the parity coboundary of B7 — and the one produced here (Theorem 12.6) are all lacunary or eventually periodic: constructed, measure-zero, and highly structured. In two centuries nobody has exhibited a naturally occurring constant with Liouville-type digit behaviour; every known example is built to have it. So the shape of the barrier set points at our technology, not at S .

Third: is the object behind the measurements simple or complex? The evidence available leans one way, and I will say how far it leans.

For “generic, and the difficulty is entirely ours”. (a) Every statistic measurable is indistinguishable from a fair-coin digit model: $\rho_h(X) \approx 0.50$ against a required 0.11; mean run length 1.96–2.03 against a permitted 9.1; longest run 12–14 over 5800 positions against a model prediction of ≈ 12.5 ; block sums of order $X^{-1/2}$ against a permitted 0.89 (Table 6, [Cert]). (b) The requirements are weaker than the apparent truth by a factor $\sqrt{X}$; a proof does not need to understand the digits, only to exclude a pathology. (c) All countermodels are constructed lacunary objects. (d) The one coordinate in which S has classical structure — the Möbius–Mersenne form (2) — is an alternating sum of rational functions of 2^d with no functional equation, no modularity, no continued-fraction structure, and no algebraicity: there is no hidden object to find, which is consistent with genericity.

Against, or at least complicating. (a) B4’s self-cancellation is exact, unconditional, and holds at every depth K : the cost of forcing a depth- K residue by a Dirichlet prime is $p \geq 1 + 2^{K-1}$ while the payoff is amplitude 2^{K-1} , so the trade is precisely null. An exact null of that form is a structural fact and not an accident of parametrisation, and it is the one place where the object itself pushes back. (b) The exact census records residues approaching the forbidden edge — a closest central margin of about $2.2 \cdot 10^{-4}$ of the modulus at $t = 100$ — which is what a uniform model predicts and which rules out, permanently, any argument that proceeds by a crude uniform margin. (c) Proposition 12.7 shows that even the pieces of the frontier that are provable become true only past $X \approx 10^{30}$ – 10^{40} ; a genuinely simple object would not usually require asymptotics that begin so late.

Verdict, flagged as inference. The evidence supports “the digits of S are generic and the wall is technological” over “the object is subtle”, but it does not determine it, and the reason

it cannot be structural: every test performed is a test a generic object passes, so passing them is weak evidence, and the only tests that would discriminate are exactly the ones no technique can run. What the evidence *does* determine is narrower and firmer: the difficulty of #249 is not located in φ 's irregularity — φ is the smoothest interesting multiplicative function, with multiplicative fluctuation $\log \log Y$ where $d(n)$ has $Y^{c/\log \log Y}$ — but in the fact that we possess no method whatsoever for lower-bounding digit changes of a constant that was not designed to have them. That is why the Erdős–Borwein constant fell in 1948 and this one has not: $d(n)$ has spikes to deposit into the digit stream and $\varphi(n)$ does not, and in the Möbius coordinate the signs cancel.

What would actually change the picture. One theorem, of any strength, giving a non-trivial digit statistic for S or for some α_h : an upper bound $o(N)$ on the longest binary run in the first N positions would not by itself decide #249, but it would be the first evidence that the object is legible at all, and by Corollary 12.4 any bound strong enough to give a positive proportion of digit changes in one block per scale would decide it. Failing that, the two concrete openings identified above are: the logarithmically averaged form of the block gap, where the entropy-decrement method is at least in the right regime (§12.3); and the canonical prime fibre (3), where every budget except one decorrelation estimate has been discharged. Neither is close. Both are well posed, and both can be started on today.

13 What is open, stated exactly

Erdős #249 asks whether $S = \sum_{n \geq 0} \varphi(n)/2^n$ is irrational. It is [Open]: nothing in this corpus decides it. Every attack recorded in Parts I–IV terminates, in one of exactly two ways, at a small number of *cofinal supply obligations* on the totient tail $R_N := \sum_{j \geq 0} \varphi(N+1+j)/2^{j+1}$ ([Erdos249257.TotientTailPeriodKiller.totientTail](#) [TotientTailPeriodKiller.lean:57]): either it proves a genuine theorem of the shape “*if* this cofinal predicate holds, *then* S is irrational,” with the predicate itself left completely unproved, or it prunes one specific proof shape and shows it cannot supply the predicate. This section renders every such obligation exactly as Lean states it, gives its site, records which of them are proved *equivalent* to one another (iff) and which are proved only *sufficient* (one-directional, an easier target implied by a harder one), and closes by naming, without overclaiming, the piece of new mathematics each surviving form would need. Quantifier order is preserved exactly as written in Lean; nowhere below does a finite verified list stand in for a cofinal supply, and nowhere does an implication get reported as a solved case.

Throughout, $H_t := \text{periodLcm}(t) = \text{lcm}(1, \dots, t)$ ([Erdos249257.CarrySurvivorExtinction.periodLcm](#) [CarrySurvivorExtinction.lean:10]) and $D(h, N, L) := \sum_{j < L} (\varphi(N+h+1+j) - \varphi(N+1+j)) \cdot 2^{L-1-j} \in \mathbb{Z}$ is `windowDiscrepancy` ([Erdos249257.TotientTailPeriodKiller.windowDiscrepancy](#) [TotientTailPeriodKiller.lean:68]) and the base predicate named “`Sep(h, N, L)`” in the task brief is literally `certifiedKill`:

$\text{certifiedKill}(h, N, L) := (N+h+L+2 : \mathbb{Z}) < D(h, N, L) \bmod 2^L \wedge D(h, N, L) \bmod 2^L < 2^L - (N+h+L+2).$

[Erdos249257.TotientTailPeriodKiller.certifiedKill](#) [TotientTailPeriodKiller.lean:72] [Lean] (the definition; decidable, instance at line 84) `scale:n/a` `coord:other:binary-window`. In words: the residue of the window discrepancy modulo 2^L avoids a shrinking radius- $(N+h+L+2)$ neighbourhood of 0 inside the growing modulus 2^L .

13.1 The certificate-completeness converter

Every reformulation below ultimately measures the same underlying real quantity, because certificates are *complete* receipts of non-integrality, not merely sufficient ones:

$$(\exists L, \text{certifiedKill}(h, N, L)) \iff R_{N+h} - R_N \notin \text{range}((\uparrow) : \mathbb{Z} \rightarrow \mathbb{R}).$$

[Erdos249257.exists_certifiedKill_iff_tail_diff_notMem_int](#) [LcmConeFlatness.lean:316] (wrapped as [totient_tail_window_kill_exists_iff_tail_diff_notMem_int](#) at [Erdos249257.CertificateKernel](#) [Lean] scale:uniform (holds for all h, N) coord:other:binary-window.

Note. This iff is the reason every “certificate supply” form below can be restated, without loss, in “pure non-integrality” language with no certificate vocabulary at all (the B8/TE forms). It does *not* assert that the cofinal supply exists; it only says the certificate route and the real-analytic route are the same route.

13.2 The base cofinal obligation the wall

Obligation 1 (certificate supply).

$$\forall h \geq 1, \forall N_0, \exists N \geq N_0, \exists L, \text{certifiedKill}(h, N, L) \implies \text{Irrational}(S).$$

This is exactly the quantifier structure named “Sep(h, N, L)” in the task brief: $\forall h \geq 1 \forall N_0 \geq 0 \exists N \geq N_0 \exists L \text{Sep}(h, N, L)$. [Erdos249257.irrational_totient_series_of_certificate_supply](#) [TotientTailPeriodKiller.lean:394] [Lean] (the implication is proved; the hypothesis is unsupplied evidence tags the theorem, not the missing premise) scale:cofinal coord:other:binary-window.

The proof is by contradiction against the tail-period law: if S were rational, [eventual_period_of_not_irrational](#) ([Erdos249257.eventual_period_of_not_irrational](#) [TotientTailPeriodKiller.lean:1] [Lean], unconditional) supplies a period $h > 0$ and pre-period N_0 with $R_{N+h} - R_N \in \mathbb{Z}$ for all $N \geq N_0$; the certificate hypothesis instantiated at that (h, N_0) produces an $N \geq N_0$ where $R_{N+h} - R_N \notin \mathbb{Z}$, a contradiction via [tail_diff_notMem_int_of_certifiedKill](#) ([Erdos249257.tail_diff_notMem_int_of_certifiedKill](#) [TotientTailPeriodKiller.lean:262]). *What supplying Obligation 1 would take:* an arithmetic (not merely existential) argument that, for every period length h and past every threshold N_0 , the totient window discrepancy’s residue mod 2^L can be pushed off 0 by a margin growing with the modulus i.e., genuine cancellation/anti-concentration information about φ on a sliding window, at every scale, for every h . No such argument exists in the corpus; every row below is a restatement of exactly this missing input in a different coordinate or with fewer free parameters, never a proof of it.

13.3 Collapsing free parameters: the multiple / diagonal / cone chain

Three further public reformulations replace Obligation 1’s two free parameters (h, N_0) with successively less information, each proved *sufficient* (not shown equivalent) for $\text{Irrational}(S)$ by an explicit reduction chain in Lean.

Multiple-period collapse (weaker hypothesis than Obligation 1).

$$\forall h_0 > 0, \forall N_0, \exists m > 0, \exists N \geq N_0, \exists L, \text{certifiedKill}(m \cdot h_0, N, L) \implies \text{Irrational}(S).$$

[Erdos249257.irrational_totient_series_of_multiple_certificate_supply](#) [CarrySurvivorExtinction.lean:1] [Lean] scale:cofinal coord:other:lcm-period-multiple. Obligation 1’s hypothesis trivially implies this one (take $m = 1$), so this target is strictly easier to satisfy; it is still open.

Diagonal collapse one free parameter.

$$\forall t_0, \exists t \geq t_0, \exists L, \text{certifiedKill}(H_t, H_t, L) \implies \text{Irrational}(S).$$

[Erdos249257.irrational_totient_series_of_lcm_diagonal_certificate_supply](#) [LcmDiagonalReduction.lean:1] [Lean] scale:cofinal coord:other:lcm-diagonal. This is the canonical single-quantifier restatement:

write $P(t) \equiv \exists L, \text{certifiedKill}(H_t, H_t, L)$; the obligation is $\forall t_0 \exists t \geq t_0, P(t)$. The reduction is a genuine proof, not a relabelling: given $t \geq \max(h_0, N_0)$, $h_0 \mid H_t$ and $H_t \geq t \geq N_0$ simultaneously, so a single diagonal witness at t discharges *both* of Obligation 1's free parameters at once, via the intermediate lemma [Erdos249257.irrational_totient_series_of_lcm_certificate_supply](#) [LcmDiagonalReduction.lean:92] which itself reduces to the multiple-period form above. $P(t)$ is verified at 28 explicit values of t through $t = 64$ ([Erdos249257.certifiedKill_diagonal_all_imported](#) [DiagonalPincerCertificates.lean:2842], depths [6, 5, 7, 7, 9, 14, 15, 14, 21, 22, 23, 26, ...] for $t \in \{1, 2, 3, 4, 5, 7, 8, 9, 11, 13, 16, 17, \dots\}$; endpoint [Erdos249257.certifiedKill_diagonal_t64](#) [DiagonalPincerCertificateT64Endpoint.lean:1928]) [Cert] scale:fixed coord:other:lcm-diagonal a finite floor, not a cofinal supply: nothing in these 28 witnesses is a function of t , each is an independent `norm_num` block over explicit φ values.

Cone collapse two multipliers, still the same wall.

$$\forall t_0, \exists t \geq t_0, \exists q, m, L, 0 < q \wedge \text{certifiedKill}(m \cdot H_t, q \cdot H_t, L) \implies \text{Irrational}(S).$$

[Erdos249257.irrational_totient_series_of_lcm_cone_window_kill_supply](#) [CertificateKernel.lean:1868] [Lean] scale:cofinal coord:other:lcm-cone. The diagonal form above is exactly the cell $q = m = 1$, so any diagonal witness trivially witnesses the cone form; the cone form is thus the weakest known target still logically tied to this coordinate. It rests on a genuine strengthening of the tail-period law, *lcm-cone flatness*: if S is rational there is t_1 such that for every $t \geq t_1$ and every $q > 0, m$, $R_{qH_t+mH_t} - R_{qH_t} \in \mathbb{Z}$ rationality flattens the *whole* cone $\{kH_t : k \geq 1\}$, not just one difference ([Erdos249257.rational_totient_series_forces_lcm_cone_flatness](#) [CertificateKernel.lean:18674], [Lean], unconditional). A sharper cone-form producer, `coneNonflatCert`, needs only a *one-sided* radius per vertex information-theoretically half of `certifiedKill`'s pairwise floor and proves that some pair in a finite multiplier menu Q is non-integral; the corresponding supply, $\exists$ an unbounded-scale menu Q with `coneNonflatCert` firing, is [Open], scale:cofinal, [Erdos249257.irrational_totient_series_of_lcm_cone_nonflat_supply](#) [CertificateKernel.lean:18675].

Pure non-integrality form certificate vocabulary stripped out. Via the completeness iff above, the diagonal and cone forms restate, *exactly*, with no reference to `certifiedKill` at all:

$$\forall t_0, \exists t \geq t_0, R_{2H_t} - R_{H_t} \notin \mathbb{Z} \iff \text{diagonal obligation above} \implies \text{Irrational}(S),$$

[Erdos249257.irrational_totient_series_of_lcm_diagonal_nonintegrality_supply](#) [CertificateKernel.lean:18676] (cone analogue at [Erdos249257.CertificateKernel](#) [CertificateKernel.lean:18718]) [Lean] scale:cofinal coord:other:real-analytic-nonintegrality. This is the frontier of #249 with every piece of certificate machinery removed: *does $R_{2H_t} - R_{H_t} \notin \mathbb{Z}$ for infinitely many t ?* Nothing decides this either.

13.4 The exponential-sum form

A first-harmonic (Weyl-sum) cancellation statement is proved sufficient for Obligation 1 by an elementary pigeonhole argument, with no case analysis and no scale-degrading constants:

$$\text{DTWFirstHarmonicNormGap} \equiv \forall h > 0, \forall X_0, \exists X, L, \max(X_0, 1) \leq X \wedge 16(2X+h+L+2) \leq 2^L \wedge \left\| \sum_{N \in [X, 2X)} \right\|$$

[Erdos249257.DTWFirstHarmonicNormGap](#) [FirstHarmonicPivot.lean:75] [Open] (the predicate; unproved at every h, X_0) scale:cofinal coord:other:first-harmonic.

$$\text{DTWFirstHarmonicNormGap} \implies \text{Irrational}(S).$$

[Erdos249257.irrational_totient_series_of_first_harmonic_norm_gap](#) [FirstHarmonicPivot.lean:83] [Lean] scale:cofinal coord:other:first-harmonic. The consumer already has Obligation 1's exact

free-parameter shape (X is a completely free threshold, so applying the gap at $X \geq N_0$ gives $\exists N \geq N_0 \exists L$ directly); what is missing is not scale and not coordinate, it is the arithmetic input itself not one instance of a constant-saving cancellation bound for the complex exponential sum $\sum_{N \in [X, 2X]} e(D(h, N, L)/2^L)$ is proved anywhere in the corpus, at any h, X, L . A strictly stronger *subset* form (a saving on any nonempty finite $T \subseteq [0, 2X)$, no density or partition hypothesis) is also proved sufficient and unconditional as an implication: [Erdos249257.exists_certifiedKill_of_first_harmonic_norm_gap](#) [FirstHarmonicPivot.lean:53] consumes the elementary block lemma [Erdos249257.exists_certifiedKill_of_first_harmonic_gap](#) [FirstHarmonicGap.lean:1] ([Lean], unconditional: *any* constant-saving first-harmonic gap on one dyadic block forces a finite kill certificate, via $\cos(\pi/8) > 9/10$ and averaging). *What supplying this form would take:* genuine cancellation for the first additive character of the totient window discrepancy a Weyl-type bound, uniform in h and growing L , on a character sum built from φ differences, not merely an existence statement. [Erdos249257.ResidualGaugeObstruction.det_residualMonomialMatrix](#) [ResidualGaugeObstruction.lean:47] rules out one entire class of candidate proofs: a residual-blind determinant/rank certificate cannot see this cancellation, because an explicit “locked” unit-norm gauge reconstructs the exact bad configuration while keeping any Vandermonde-shaped minor nonzero any real argument here must couple the rows arithmetically, not merely normalise columns.

13.5 The private lane: the actual-orbit reformulation and the top-edge staircase case

A second, independently developed lane (private modules `TotientActual*/TotientFixedRank*`, never mirrored to the public repository) restates #249 in terms of the *actual* power-of-two LCM diagonal, $H = H_{2^a}$, and proves a genuine **iff** with the totient series itself the strongest form of “exact statement of what remains” anywhere in the corpus.

The exact equivalence.

$$\text{Irrational}(S) \iff \forall a_0, \exists a \geq a_0, \text{actualLcmTailOrbit}(a) \notin \mathbb{Z}.$$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.irrational_totientSeries_iff_actualLcmOrbitNonintegralitySupply](#) [TotientActualLcmOrbitNonintegrality.lean:37] [Lean] scale:n/a (an unconditional equivalence, not a supply) coord:mobius-mersenne, where $\text{actualLcmTailOrbit}(a) := 2^H(2^H - 1)S - (\text{totientPrefix}(2H) - \text{totientPrefix}(H))$, $H = H_{2^a}$ ([Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.actualLcmTailOrbit_eq_scaled_totientSeries_sub_prefix](#) [TotientActualLcmOrbitSeparation.lean:68]). *This is #249 restated with nothing left over:* no auxiliary hypothesis, no scale caveat cofinal non-integrality of this one sparse sequence at power-of-two heights is exactly Erds #249. Everything else in this subsection is an attempt to reach the right-hand side.

Route-pruning: total staircase collapse is impossible. Before any positive target, the file proves one entire proof shape is empty. For $a \geq 8$ and room bound $J + K + (a+6) < 2 \cdot 2^a$ with a wide enough modulus $2H + J + K + 2 < 2^m$, a terminal window where *every* one of the last m letters vanishes mod its own growing power of two is impossible:

$$\neg \text{ActualLcmTerminalDyadicStaircase}(a, J, K, m).$$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.not_actualLcmTerminalDyadicStaircase_of_room](#) [TotientActualLcmTopEdgeStaircase.lean:251] [Lean] scale:bounded coord:mobius-mersenne. The mechanism is a bare positivity/divisibility contradiction (a positive quantity below a modulus, divisible by that modulus, must be 0) against the unconditional positivity of every short-window letter, $0 < \text{lcmRayArithmeticLetter}(2^a, j)$ for $a \geq$

8, $0 < j < 2 \cdot 2^a$ ([Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.lcm_pos_of_lt_two_mul](#) [TotientActualLcmOrbitArithmetic.lean:1703], [Lean], unconditional, no rationality hypothesis). *Do not attempt a full terminal-staircase producer*: it is provably empty. The corpus’s own route past this dead end is the *punctured* staircase (all but the last letter vanish; the last is retained and pinned exactly to the half-turn 2^{m-1} , [Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.penultimate_eq_half](#) [TotientActualLcmTopEdgeStaircase.lean:1187], [Lean], scale:bounded) and the one-sided residue-gap family below.

The surviving one-sided target.

$\text{ActualLcmTopEdgeResidueGap}(a, J, K, m) \equiv m \leq K \wedge 2H+J+K+2 < 2^m \wedge D(H, H+J, K) \bmod 2^m \leq 2^m -$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.ActualLcmTopEdgeResidueGap](#) [Open] (a definition, unproved at any a) scale:bounded coord:mobius-mersenne. Note this is a *one-sided* inequality only the upper (positive) carry arc is excluded, not a symmetric two-sided band because the corridor’s lower half is already discharged unconditionally: for $a \geq 8$ and $J + (a+6) < 2 \cdot 2^a$,

$$0 < R_{2H+J} - R_{H+J}$$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.actualLcmTailDiffShift_pos](#) [TotientActualLcmOrbitSign.lean:39] [Lean] scale:bounded coord:mobius-mersenne a rare genuinely unconditional, non-hypothetical real-analytic theorem in this corpus, needing no rationality assumption at all. Its companion shows that *if* the orbit is integral, the residue is forced to the exact top edge $2^K - e$ ($e > 0$ small), *outside* the arc a symmetric certificate would need: [Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.actualLcm_integral_forces_topEdgeResidue](#) [TotientActualLcmOrbitSign.lean:211] ([Lean], scale:bounded) this is the exact statement of “here is what remains,” pinning the missing exclusion to one named residue class rather than leaving it implicit.

$$\text{ActualLcmTopEdgeResidueGap}(a, J, K, m) \implies R_{2H+J} - R_{H+J} \notin \mathbb{Z}.$$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.actualLcmTailDiffNotMem_int_of_topEdgeResidueGap](#) [TotientActualLcmTopEdgeStaircase.lean:1260] [Lean] scale:bounded coord:mobius-mersenne. The cofinal target built from it:

$\text{PowerTwoActualLcmTopEdgeResidueGapSupply} \equiv \forall a_0, \exists a, K, m, \quad a_0 \leq a \wedge 8 \leq a \wedge K + (a+6) < 2 \cdot 2^a \wedge \text{Act}$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.PowerTwoActualLcmTopEdgeResidueGapSupply](#) [Open] scale:cofinal coord:mobius-mersenne.

$$\text{PowerTwoActualLcmTopEdgeResidueGapSupply} \implies \text{Irrational}(S).$$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.irrational_totientSeries_of_topEdgeResidueGapSupply](#) [TotientActualLcmTopEdgeStaircase.lean:2184] [Lean] scale:cofinal coord:mobius-mersenne.

Five strictly weaker links, each independently proved sufficient. The same file proves five further cofinal predicates, each strictly weaker than $\text{PowerTwoActualLcmTopEdgeResidueGapSupply}$ (each implies it, so each is an easier target), with a direct #249 endpoint of its own proving the *weakest* of the six closes the entire cluster.

- (i) $\text{PowerTwoAdjacentSuffixMidbandSupply}$: replaces the m -bit residue test with a symmetric two-sided band on the *adjacent-suffix* residue at depth m , buffered by the larger depth

$m+1$ so either branch stays inside the sign corridor. [Erdos257PeriodNoncollapse.DiagonalFreshLossB](#)
[Open] scale:cofinal coord:mobius-mersenne. Sufficiency: [Erdos257PeriodNoncollapse.DiagonalFreshLossB](#)
[of_adjacentSuffixMidband](#) [TotientActualLcmTopEdgeStaircase.lean:2108], direct endpoint
[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.irrational_](#)
[totientSeries_of_adjacentSuffixMidbandSupply](#) [TotientActualLcmTopEdgeStaircase.lean:2191].

- (ii) PowerTwoOddGuardTopEdgeHalfWordBandSupply: at odd depth $m = 2q+1$ the adjacent-suffix residue is exactly twice a half-word residue, and both directed edge widths halve to the same threshold $H + q + 2$:

$$H + q + 2 \leq (\text{half-word correction word}) \bmod 4^q \leq 4^q - (H + q + 2).$$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.PowerTwoOddGu](#)
[Open] scale:cofinal coord:mobius-mersenne. This is a substantially weaker demand than the
earlier fixed $1/32$ central band elsewhere in the corpus. Sufficiency: [Erdos257PeriodNoncollapse.Diagon](#)
[of_oddGuardTopEdgeHalfWordBand](#) [TotientActualLcmTopEdgeStaircase.lean:2121].

- (iii) PowerTwoActualFinalTopEdgeMagnitudeSupply: the exact centered-lift restatement of (ii),

$$H + q + 2 \leq |\text{actualOddHalfCenteredLift}(a, q)|,$$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.PowerTwoActua](#)
[Open] scale:cofinal coord:mobius-mersenne. Proved **equivalent** (an iff, not merely sufficient)
to (ii):

$\text{PowerTwoOddGuardTopEdgeHalfWordBandSupply} \iff \text{PowerTwoActualFinalTopEdgeMagnitudeSup}$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.topEdgeHalfWo](#)
[iff_actualFinalCenteredMagnitudeSupply](#) [TotientActualLcmTopEdgeStaircase.lean:1479] [Lean].

- (iv) PowerTwoFlexibleActualTopEdgeMagnitudeSupply: the same magnitude test at an *arbitrary* odd rank whose adjacent depth still fits the corridor (not forced to the canonical guarded depth): [Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine](#)
[Open] scale:cofinal coord:mobius-mersenne. Sufficiency routes back through (i), not through
the corridor-escape chain below: [Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOd](#)
[of_flexibleActualTopEdgeMagnitude](#) [TotientActualLcmTopEdgeStaircase.lean:1938], direct
endpoint [Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.irrat](#)
[totientSeries_of_flexibleActualTopEdgeMagnitudeSupply](#) [TotientActualLcmTopEdgeStaircase.lean:22
- (v) PowerTwoFlexibleActualTerminalDominanceSupply: a strictly *one-sided* version of (iv)
only the upper comparison, no absolute value:

$$\text{diagonalWindowIncrement}(2^a, 2q+2) \leq 2 \cdot \text{actualOddHalfCenteredLift}(a, q).$$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.PowerTwoFlexi](#)
[Open] scale:cofinal coord:mobius-mersenne. Direct endpoint [Erdos257PeriodNoncollapse.DiagonalFreshLo](#)
[totientSeries_of_flexibleActualTerminalDominanceSupply](#) [TotientActualLcmTopEdgeStaircase.lean:2

The weakest known link, and the exact identity pinning it. A sixth predicate, strictly weaker again than (v) it is the disjunction of (v)'s inequality with the opposite-direction escape is the true minimum of the whole cluster:

$$\text{PowerTwoFlexibleActualTerminalCarryCorridorEscapeSupply} := \forall a_0, \exists a, q, \dots \wedge (2u \leq d-B \vee d \leq 2u),$$

where $d = \text{diagonalWindowIncrement}(2^a, 2q+2)$, $u = \text{actualOddHalfCenteredLift}(a, q)$, $B = 2H+(2q+1)+2$. [Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.Powe](#)

[Open] scale:cofinal coord:mobius-mersenne. Direct endpoint [Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.trueCarryOrbit](#) [totientSeries_of_flexibleActualTerminalCarryCorridorEscapeSupply](#) [TotientActualLcmTopEdgeStaircase.lean:1678] This form is not arbitrary: under integrality it is exactly one side of a proved identity,

$$2 \cdot \text{actualOddHalfCenteredLift}(a, q) = \text{diagonalWindowIncrement}(2^a, 2q+2) - \text{carryOrbit}(H, H, z, 2q+1),$$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.two_minus_mul_actualOddHalfCenteredLift_eq_terminal_sub_trueCarry](#) [TotientActualLcmTopEdgeStaircase.lean:1678] [Lean] scale:bounded coord:mobius-mersenne an *equality*, not an inequality: the doubled centered-lift state is exactly (terminal arithmetic letter) minus (true carry). Escaping either side of this named open interval is exactly the corridor-escape supply. Crucially, the sign machinery has already eliminated the branch this identity most naturally supplies: [Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.trueEndpointSurvivor_neg](#) [TotientActualLcmOrbitSign.lean:172] ([Lean], scale:bounded) proves that *under integrality* the true carry orbit is strictly *positive* throughout the corridor, which means the dominance branch (v) is the one the identity naturally produces positive evidence *against*, and the surviving open target is really the *lower*-escape branch, $2u \leq d-B$ the dominance branch itself is stated separately as [Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.notMem_int_of_actualTerminalDominance](#) [TotientActualLcmTopEdgeStaircase.lean:1880]. This is the sharpest finite arithmetic statement of “what remains to prove” anywhere in the corpus: a fully closed-form real-integer quantity whose escape from a named interval is necessary and sufficient (mod a fit hypothesis $2(H+q+2) \leq 4^q$) for non-integrality at that rank.

A second, independent finite-window target: the short-arithmetic-kill supply. A separate reduction ties the diagonal public form (the collapsing-free-parameters chain above) to this private lane via an exact digit formula with no cleanliness hypothesis at all, $\text{lcmRayArithmeticLetter}(t, j) = \varphi(H_t + j) - \varphi(H_t)$ for *every* offset j ([Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.eq_deltaTotient](#) [TotientActualLcmOrbitArithmetic.lean:298], [Lean]), and $\text{lcmDiagonalArithmeticKill}(t, L) \iff \text{certifiedKill}(H_t, H_t, L)$ ([Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.iff_certifiedKill](#) [TotientActualLcmOrbitArithmetic.lean:2045], [Lean]). Consequently

$$\text{PowerTwoActualLcmShortArithmeticKillSupply} \equiv \forall a_0, \exists a, L, \quad a_0 \leq a \wedge L < 2 \cdot 2^a \wedge \text{certifiedKill}(H_{2^a}, H_{2^a}, L)$$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.PowerTwoActualLcmShortArithmeticKillSupply](#) [Open] scale:cofinal coord:mobius-mersenne is *literally the diagonal obligation* $P(t)$ of the collapsing-free-parameters chain above, restricted to powers of two and to a short window $L < 2 \cdot 2^a$. It is proved sufficient for the actual-orbit iff above via [Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.of_shortArithmeticKill](#) [TotientActualLcmOrbitArithmetic.lean:2152], and it is verified at exactly two exponents, $a = 4$ (depth $L = 23$) and $a = 6$ (depth $L = 93$), each traced to a pre-existing compressed diagonal certificate: [Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.four_and_six_notMem_int](#) [TotientActualLcmShortKill.lean:35] [Cert] scale:fixed. The bounded stub [Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.powerTwoActualLcmShortArithmeticKillSupply_through_six](#) [TotientActualLcmShortKill.lean:54] ([Lean], scale:bounded, proof is a single hard-coded witness $(a, L) = (6, 93)$, with no argument in a at all) proves the supply only for $a_0 \leq 6$; extending it to a single further exponent (say $a = 7$) is a self-contained, independently interesting target, not a re-run.

A Diophantine-flavoured alternative: the raw-approximant separation supply. Unconditionally, for every a, q :

$$|\text{actualLcmTailOrbit}(a) - \text{actualLcmRawApprox}(a, q)| < \frac{4H + 2(2q+1) + 4}{2^{2q+2}},$$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.abs_minus_actualLcmTailOrbit_sub_rawApprox_lt](#) [TotientActualLcmOrbitSeparation.lean:141] [Lean] scale:uniform

coord:mobius-mersenne, where $\text{actualLcmRawApprox}(a, q)$ is an explicit finite computable rational. This reduces the whole analytic problem to a finite question: a cofinal $1/32$ -separation of the raw approximant from every integer,

$\text{PowerTwoActualLcmOrbitSeparationSupply} \equiv \forall a_0, \exists a \geq \max(2, a_0), \exists q, \quad \text{oddGuardedCanonicalAdjacentSu}$

[Open] scale:cofinal coord:mobius-mersenne, sufficient via [Erdos257PeriodNoncollapse.DiagonalFreshLossBridge](#)
[totientSeries_of_actualLcmOrbitSeparationSupply](#) [TotientActualLcmOrbitSeparation.lean:305]
 ([Lean]). Note the depth q is not a free search parameter: the supply pins exactly one admissible depth per scale a via $\text{oddGuardedCanonicalAdjacentSuffixDepth}$.

13.6 A problem-agnostic normal form: the guard-cylinder compression

Independently of which coordinate is used, *any* tail-difference certificate for arbitrary h, N, L , not specific to #249 compresses to a two-bit test at logarithmic depth:

$$(\exists L, \text{certifiedKill}(h, N, L)) \iff \text{GuardCylinderWitness}(h, N).$$

[Erdos257PeriodNoncollapse.DiagonalFreshLossBridge.PowerTwoOddWindowAffine.exists_-certifiedKill_iff_guardCylinderWitness](#) [TotientActualLcmTopEdgeStaircase.lean:844] [Lean]
 scale:uniform coord:seam-integer, where the witness is $\exists s, b, \text{certifiedKill}(h, N+s, b+1) \vee (\text{room} \wedge \text{DyadicMixedGuard}(D(h, N+s, b+2), b))$ at $b = \lfloor \log_2(N+h+L+2) \rfloor + 1$. This statement mentions no Mersenne or totient structure whatsoever and is stated for arbitrary h, N, L ; it compresses the search space for *any* tail-difference non-integrality certificate of any depth L , however large down to a socket at a logarithmic scale plus a two-bit mixed-guard cylinder (01 or 10). This is directly reusable, unchanged, for any binary-series tail-difference problem, including #257's own denominators.

13.7 Two further, logically independent obligations

Two more open statements would *also* close #249, each living in a genuinely different coordinate from the certificate/residue coordinate above, and each finished on exactly one side of an inequality.

Obligation 2 (rank compression). Unconditionally, for every level e , the dyadic totient-kernel family of $2^e + 1$ channels is linearly independent over $\mathbb{Q}$ (via CRT and Dirichlet's theorem on primes in arithmetic progression): [Erdos249257.linearIndependent_canonicalTotientKernelFamily](#) [TotientMahlerDefect.lean:1014] [Lean] scale:uniform coord:other:carry-kernel-rank. Consequently, if S is rational there is a tempered integral binary carry orbit u with

$$\forall e, \quad 2^e - 1 \leq \text{rank}_{\mathbb{Q}} \text{span}(\text{canonicalCarryKernelFamily}(u, e)).$$

[Erdos249257.not_irrational_totientSeries_implies_unbounded_carryRank_unconditional](#) [TotientCarryKernelRank.lean:1015] [Lean] scale:uniform coord:other:carry-kernel-rank. The missing input is the opposite inequality a rationality-side rank *upper* bound:

$\exists C, \quad \forall \text{tempered integral binary carry orbit } u \text{ for a rational } S, \quad \forall e, \quad \text{rank}_{\mathbb{Q}} \text{span}(\text{canonicalCarryKernelFam}$

(or any bound growing slower than $2^e - 1$). [Open] scale:cofinal coord:other:carry-kernel-rank. One natural route to it is proved dead: no $\text{CompressedAdjointCertificate}$ ($Q \cdot v \cdot A = \text{boundary}$ with $|\text{boundary}| < Q \cdot v$ and $A \neq 0$) can exist [Erdos249257.false_of_compressedAdjointCertificate](#) [TotientMahlerDefect.lean:1014] [Lean] and the full family is proved infinite-dimensional in span: [Erdos249257.not_finiteDimensional_-span_fullTotientKernel](#) [TotientMahlerDefect.lean:1014] [Lean]. What rationality *does* buy

uniform eventual periodicity of u 's dyadic sections modulo some $v > 0$ is proved to *not* promote to any finite rank bound without further arithmetic input: [Erdos249257.totientSeries_implies_mod_period_and_unbounded_rank](#) [TotientTailCarryPeriod.lean:224] [Lean]. This pins the obstacle precisely, but supplies no rank upper bound.

Obligation 3 (Farey growth law). The corpus's strongest *unconditional* statement about S comes from a Farey-gap denominator exclusion at a single fixed window $K = 240$:

$$\forall p \in \mathbb{Q}, \quad p.\text{den} \leq 79639646646701375323355774875831053 \implies S \neq p.$$

[Erdos249257.tsum_totient_div_pow_two_ne_ratCast_of_den_le_79639646646701375323355774875831053](#) [Lean] scale:bounded coord:farey. This bound is *sharp* at $K = 240$ the mediant $q = 79639646646701375323355774875831053$ is proved to be the exact first failing denominator ([Erdos249257.gap_check_window_1_240_first_failure](#) [GapFareyBound.lean:225], [Lean]) so re-running the same window buys nothing further; a new K needs a freshly committed totient residue and freshly computed continued-fraction convergents, both hard-coded numerals. The cofinal upgrade would be: a proved growth law $g(K) \rightarrow \infty$ such that for every K the $(N=1, K)$ gap check passes for all $q \leq g(K)$ equivalently, a lower bound on the convergent denominators of the underlying totient-window constant, uniform in K . [Open] scale:cofinal coord:farey. One natural strengthening is proved to be a hard ceiling, not a route to unboundedness: at a prime-power reduced denominator, the “unit-gap” refinement can rescue *at most one* additional lattice point beyond the ordinary gap certificate [Erdos249257.reducedDenominatorCandidateCount_prime_pow_le_one](#) [PrimitiveWeightCertificate.lean:111] [Lean]. This route is closed; growing K genuinely requires new per-window computation, not a refinement of the existing one.

13.8 Equivalence map

Form	Relation to Obligation 1	Site
Sep(h, N, L) supply (Obl. 1, base form)	base form	TotientTailPeriodKiller.lean:394
multiple-period supply	weaker (implied by Obl. 1)	CarrySurvivorExtinction.lean:502
lcm-diagonal supply $P(t)$	weaker; single free parameter	LcmDiagonalReduction.lean:112
lcm-cone supply	weaker again; $P(t)$ is the cell $q=m=1$	CertificateKernel.lean:18686
cone-menu	weaker; half the pairwise radius	CertificateKernel.lean:18812
(coneNonflatCert) supply	equivalent to the corresp. certificate form, via the completeness iff	CertificateKernel.lean:18706, :18718
DTWFirstHarmonicNormGap	sufficient for Obl. 1 (not shown weaker/stronger)	FirstHarmonicPivot.lean:83
actual-orbit nonintegrality supply	equivalent to Irrational(S) itself	TotientActualLcmOrbitNonintegrality.lean:37
short-arithmetic-kill supply	special case of $P(t)$: powers of two, short window	TotientActualLcmOrbitArithmetic.lean:2107

Form	Relation to Site Obligation 1	
top-edge residue-gap supply	sufficient for actual-orbit supply; one-sided	TotientActualLcmTopEdgeStaircase.lean:1325
adjacent-suffix midband supply	weaker than top-edge residue-gap	TotientActualLcmTopEdgeStaircase.lean:1334
odd-guard half-word band supply	weaker again; equivalent to final-magnitude form	TotientActualLcmTopEdgeStaircase.lean:1349
actual final top-edge magnitude supply	equivalent to odd-guard half-word band	TotientActualLcmTopEdgeStaircase.lean:1471
flexible top-edge magnitude supply	weaker than mid-band; feeds mid-band, not corridor-escape	TotientActualLcmTopEdgeStaircase.lean:1927
flexible terminal-dominance supply	weaker again; one-sided	TotientActualLcmTopEdgeStaircase.lean:1908
flexible terminal carry-corridor-escape supply	weakest known in this cluster	TotientActualLcmTopEdgeStaircase.lean:1895
raw-approximant separation supply	alternative (Dio-phantine) sufficient form, same target	TotientActualLcmOrbitSeparation.lean:305
rank-compression upper bound (Obl. 2)	independent obligation, different coordinate	TotientCarryKernelRigidity.lean:284
Farey growth law (Obl. 3)	independent obligation, different coordinate	GapFareyBound.lean, CertificateKernel.lean:18056
guard-cylinder witness	problem-agnostic normal form of <i>any</i> certificate, not scale-comparable	TotientActualLcmTopEdgeStaircase.lean:844

Reading the table. “Weaker” always means: the stronger form’s hypothesis, if it held, would imply the weaker form’s hypothesis by an explicit Lean construction (so the weaker predicate is an easier target), and each row independently has its own theorem to Irrational(S). “Equivalent” is reserved for rows connected by a proved **iff** statement; no other pair in this table is known to be logically equivalent, only jointly sufficient for the same conclusion. In particular, Obligation 1 (the public certificate supply) and the actual-orbit iff (the private lane) are *not* shown equivalent to each other as stated; they are independently sufficient reformulations of the same underlying question, connected concretely only through the short-arithmetic-kill supply, which is a genuine special case of Obligation 1’s diagonal form.

13.9 What new mathematics each surviving form would need

None of the forms above is close to complete; every one reduces the open content to a named finite-flavoured arithmetic statement, never removes it. Two forms above have their missing ingredient identified precisely enough to name what field of technique would supply it, without any claim that the technique exists or is easy to apply.

The exponential-sum form. DTWFirstHarmonicNormGap asks for cancellation in the first additive character of a totient-driven discrepancy: a constant-saving bound, uniform in the period h and growing with the block size X and depth L , on $\sum_{N \in [X, 2X)} e(D(h, N, L)/2^L)$, where D is built from consecutive differences of φ . This is a Weyl-sum-shaped statement about additive character sums of an arithmetic function on a window; no argument of this shape is attempted or proved anywhere in the corpus.

The staircase form. The private top-edge cluster's surviving target, in every one of its six equivalent-or-sufficient guises, reduces under the exact identity of the private lane above to control of a single *terminal arithmetic letter*: the value $\text{diagonalWindowIncrement}(2^a, 2q+2) = \varphi(2H+2q+2) - \varphi(2H)$ measured against twice a centred carry-lift state. What is needed is not a new certificate mechanism the whole apparatus of arithmetic letters, centred lifts, and carry orbits is already exact and unconditional but a genuine size or residue statement about this one totient value at cofinally many scales a , sufficient to force it off the interval the identity names. The corpus's own finite census (through $t = 64$ on the diagonal form, and $a = 4, 6$ on the short arithmetic-kill form) gives no asymptotic evidence either way; it is a floor, not a trend.

No claim is made that either obstruction is close to resolution, and none of the material above decides Erds #249, #257, or any weakening of them.