

Cubic-Rate Irrationality and Reciprocal-Tail Rigidity

Erdős Problem #243

WILL COOK*

September 2026

ABSTRACT

We prove irrationality of the reciprocal sum when strictly increasing positive integers satisfy $a_n^2/a_{n+1} = 1 + 3/n + o(n^{-3})$. The proof extracts a cubic polynomial from an integer tail numerator, then uses a square in a cubic field to restrict its coefficients. Congruences modulo seven exclude the remaining possibilities. We also obtain Sylvester recurrence criteria from bounded numerator increases and weighted sums over new maxima. The unrestricted Erdős–Graham question remains open.

1 Introduction

The Sylvester recurrence $a_{n+1} = a_n^2 - a_n + 1$ gives a rational reciprocal sum by telescoping:

$$\frac{1}{a_n - 1} = \frac{1}{a_n} + \frac{1}{a_{n+1} - 1}.$$

We call an integer sequence which eventually satisfies this recurrence a *Sylvester tail*. Its tail sum at every sufficiently late index is $1/(a_n - 1)$. Erdős and Graham asked whether every rational reciprocal sum with $a_{n+1} \sim a_n^2$ arises in this way.

Problem 1.1 (Erdős Problem 243). Let $1 \leq a_1 < a_2 < \dots$ be a sequence of integers with

$$\lim_{n \rightarrow \infty} \frac{a_n}{a_{n-1}^2} = 1 \quad \text{and} \quad \sum \frac{1}{a_n} \in \mathbb{Q}.$$

Then $a_n = a_{n-1}^2 - a_{n-1} + 1$ for all sufficiently large n .

Erdős and Graham [5, p. 64] asked this question; see also Erdős [6, p. 105] and Bloom’s catalogue [11]. We prove the following irrationality result for a restricted class of quadratically growing sequences.

Theorem 1.2 (cubic-rate irrationality). *A strictly increasing sequence of positive integers with*

$$a_n^2/a_{n+1} = 1 + \frac{3}{n} + o(n^{-3})$$

has irrational reciprocal sum.

**Authorship and AI use.* Will Cook built and directed the research infrastructure and reviewed the claims when he could. AI agents did most of the research and drafting. Cook did not independently verify every claim.

For a Sylvester tail, $a_n^2/a_{n+1} - 1 = O(1/a_n)$, which is much smaller than $3/n$. Theorem 1.2 therefore treats a restricted case of the problem by excluding rationality altogether. The general quadratic limit $a_{n+1} \sim a_n^2$ remains insufficient for any argument given here.

For example, set $a_1 = 8$ and

$$a_{n+1} = \left\lceil \frac{na_n^2}{n+3} \right\rceil.$$

This gives 8, 16, 103, 5305, From $a_{n+1} \geq a_n^2/4$ we obtain $a_n \geq 4 \cdot 2^{2^{n-1}}$ by induction. The error introduced by the ceiling is less than one, and hence

$$0 \leq 1 + \frac{3}{n} - \frac{a_n^2}{a_{n+1}} < \frac{16}{a_n^2} = o(n^{-3}).$$

The theorem applies. This example also distinguishes the cubic argument from the recurrence criterion in Section 4. Indeed, putting $P_n = \prod_{j < n} a_j$ and $t_n = P_n/a_n$ gives $t_{n+1}/t_n = a_n^2/a_{n+1}$. We compare this ratio with that of $n(n+1)(n+2)$ and sum the rounding errors to obtain $t_n \sim Kn^3$ for some $K > 0$. Thus $t_{n+1} - t_n \sim 3Kn^2$, whereas the later criterion requires bounded upward increments.

Suppose the reciprocal sum were rational. Clearing its tails produces positive integers C_n whose ratios have the same cubic rate. We compare C_n with $F_n = n(n+1)(n+2)$, for which $F_{n+1}/F_n = 1 + 3/n$. The recurrence then makes $\Delta^4 C_n$ tend to zero. Since these differences are integers, they eventually vanish, and $C_n = AF_n + B$ on a tail. Dividing out a stable gcd gives a coprime recurrence whose cubic numerator must satisfy a square condition at every root modulo almost every prime. Chebotarev carries this condition to the cubic field itself. Traces there leave just two possible numerators, and four consecutive terms modulo seven exclude both. We give the details in Section 3.

For the recurrence criteria we use the integer remainders of Koizumi [10, Lemma 4]. In Section 4, a stable gcd and the Chinese remainder theorem exclude a rational tail whose P_n/a_n has bounded upward increments. An elementary infinite product argument gives the separate summability criterion of Section 5. We then clear the tails by a least common multiple and count crossings of new maxima (Section 6). The resulting weighted criterion isolates an arithmetic estimate still missing from the general problem. The companion paper also treats nonintegral rates and proves a stronger cubic disagreement theorem. Its detailed proof locations and the formal sources are listed in Appendix B.

2 Integer tails

Assume for this section that a_n are strictly increasing positive integers, $a_{n+1}/a_n^2 \rightarrow 1$, and $\sum_{n \geq 1} 1/a_n = p/q$ with positive integers p, q . Write

$$x_n = \sum_{k \geq n} \frac{1}{a_k}, \quad P_n = \prod_{j < n} a_j, \quad D_n = qP_n, \quad C_n = D_n x_n.$$

Rationality makes C_n an integer, because

$$C_n = pP_n - q \sum_{k < n} \frac{P_n}{a_k} \in \mathbb{N}_{>0}.$$

Removing the term $1/a_n$ from the tail gives the exact recurrences

$$C_{n+1} = a_n C_n - D_n, \quad D_{n+1} = a_n D_n.$$

These are Koizumi's integer-tail coordinates, up to a common rescaling and the indexing above [10, Lemma 4, pp. 11–12]. We initially keep the fraction C_n/D_n unreduced, so that its denominator has this simple multiplicative update.

For large n , $a_{n+1} \geq a_n^2/2 \geq 2a_n$. The terms after $1/a_{n+1}$ sum to at most $2/a_{n+2} \leq 4/a_{n+1}^2$, so

$$x_n = \frac{1}{a_n} + \frac{1}{a_{n+1}} + O(a_{n+1}^{-2}), \quad \frac{C_{n+1}}{C_n} = \frac{a_n^2}{a_{n+1}} + O(1/a_n). \quad (1)$$

For the second estimate, use $C_{n+1}/C_n = a_n x_{n+1}/x_n$ in the first. Quadratic growth makes $1/a_n = o(n^{-k})$ for every fixed k , so this error preserves the precision required in the cubic theorem. We keep the original index n throughout that proof. Although an eventual recurrence survives removal of a finite prefix, the expression $1 + 3/n + o(n^{-3})$ changes its lower-order terms under an index shift.

2.1 Coprimality after reduction

Both updates preserve common divisors, so $G_n = \gcd(C_n, D_n)$ satisfies $G_n \mid G_{n+1}$. Once this chain stabilises at a value g , we may divide by g without changing either recurrence:

$$u_{n+1} = a_n u_n - v_n, \quad v_{n+1} = a_n v_n, \quad \gcd(u_n, v_n) = 1.$$

Here $u_n > 0$ and $v_n \geq 0$; we call this a *reduced exact tail*.

Proposition 2.1 (persistent coprimality). *In a reduced exact tail, $\gcd(a_n, v_n) = 1$. Distinct multipliers are pairwise coprime, and every earlier multiplier is coprime to every later numerator.*

Lean
Comparator

Proof. A prime dividing both a_n and v_n would divide u_{n+1} and v_{n+1} , contrary to reducedness. For $i < t$, the factor a_i divides v_t . Reducedness gives $\gcd(a_i, u_t) = 1$, and $\gcd(a_t, v_t) = 1$ gives $\gcd(a_i, a_t) = 1$. $\square$

We shall obtain stabilisation in two ways: a fixed third difference bounds G_n in the cubic proof, and bounded negative errors bound it in the recurrence argument.

3 Proof of cubic-rate irrationality

Suppose, towards a contradiction, that the reciprocal sum in Theorem 1.2 is rational. Construct C_n, D_n as in Section 2. Equation (1) transfers the rate without changing its precision:

$$\frac{C_{n+1}}{C_n} = 1 + \frac{3}{n} + o(n^{-3}). \quad (2)$$

The ratio alone will give a polynomial for C_n . We then use the denominator recurrence to restrict that polynomial.

3.1 Polynomial extraction

Put $F_n = n(n+1)(n+2)$, so $F_{n+1}/F_n = 1 + 3/n$, and write the error in (2) as $\varepsilon_n = o(n^{-3})$. For $z_n = C_n/F_n$,

$$\frac{z_{n+1}}{z_n} = 1 + \frac{\varepsilon_n}{1 + 3/n}.$$

The logarithmic increments of z_n are absolutely summable, and their tails are $o(n^{-2})$. Thus z_n converges to some $A > 0$ with $z_n - A = o(n^{-2})$, giving $\delta_n = C_n - AF_n = o(n)$. This estimate alone gives no control of its differences. We obtain that control by subtracting the recurrence for AF_n from the recurrence for C_n :

$$\Delta\delta_n = \frac{3}{n}\delta_n + \varepsilon_n C_n = o(1), \quad \Delta h_n = h_{n+1} - h_n.$$

Taking three further differences of a sequence tending to zero gives $\Delta^4 C_n = \Delta^3(\Delta\delta_n) \rightarrow 0$. These are integers, so $\Delta^4 C_n = 0$ eventually. The Newton interpolation formula now expresses C_n on that tail as a rational polynomial of degree at most three. Comparing it with AF_n , their difference being $o(n)$, leaves

$$C_n = An(n+1)(n+2) + B, \quad A \in \mathbb{Q}_{>0}, \quad B \in \mathbb{Q}. \quad (3)$$

The alternating perturbation $F_n + (-1)^n$ has ratio $1 + 3/n + O(n^{-3})$ but never agrees eventually with a polynomial. It shows why this extraction uses little-oh precision. We make no claim that the perturbed sequence satisfies the denominator recurrence.

3.2 Normalisation and irreducibility

For a sufficiently late n , the integer $G_n = \gcd(C_n, D_n)$ divides $C_n, C_{n+1}, C_{n+2}, C_{n+3}$ and therefore divides $\Delta^3 C_n = 6A$. The positive divisibility chain G_n is bounded and stabilises at g . Proposition 2.1 applies to $u_n = C_n/g, v_n = D_n/g$. In addition, adjacent numerators are coprime: $\gcd(u_n, u_{n+1}) = \gcd(u_n, v_n) = 1$.

Write the polynomial for u_n as

$$Q(n) = \frac{m}{6}n(n+1)(n+2) + c. \quad (4)$$

The third difference $m = 6A/g$ is a positive integer. The constant c is also integral: choose a common denominator H for the coefficients of Q . For any integer t , the difference $Q(t + kH) - Q(t)$ is an integer. Taking k sufficiently large transfers integrality from the known tail values to $Q(t)$, and in particular to $Q(0) = c$. If a prime ℓ divides c , choose an arbitrarily late $n \equiv -1 \pmod{6\ell}$. Both $Q(n)$ and $Q(n+1)$ would be divisible by ℓ , contradicting adjacent coprimality. This also excludes $c = 0$. Thus

$$m \in \mathbb{N}_{>0}, \quad c \in \{1, -1\}.$$

The multipliers $a_n > 1$ on the reduced tail are pairwise coprime, so infinitely many distinct primes occur among them. Once a prime divides one such a_n , it divides every later v_t and no later u_t . If the cubic Q had a rational root, that root would reduce to a root modulo all but finitely many primes. Choose one of the persistent primes outside those

exceptions; a sufficiently late index in the root's residue class would have $\ell \mid Q(t) = u_t$, a contradiction. A cubic is reducible over $\mathbb{Q}$ only if it has a rational root. Hence Q is irreducible.

Set $\kappa = m/6$, $\eta = 6c/m$, and

$$f(T) = T^3 - T + \eta, \quad Q(n) = \kappa f(n+1).$$

Let α be a root of f and $K = \mathbb{Q}(\alpha)$, a cubic number field.

3.3 From finite fields to the cubic field

We first extract a condition at the roots of f modulo primes. Eliminating v_n from the reduced recurrence gives

$$u_{n+2} = (a_n + a_{n+1})u_{n+1} - a_n^2 u_n. \quad (5)$$

Let $\ell \nmid 6m$ be prime and let r be a root of f in $\mathbb{F}_\ell$. Since $\eta \neq 0$ modulo ℓ , we have $r \notin \{0, 1, -1\}$. Choose a sufficiently late index k with $k+1 \equiv r \pmod{\ell}$, so that $u_k \equiv 0$. Using $f(r) = 0$, we obtain

$$f(r-1) = -3r(r-1), \quad f(r+1) = 3r(r+1).$$

Multiplying (5) at index $k-1$ by u_{k-1} yields

$$-u_{k-1}u_{k+1} = (a_{k-1}u_{k-1})^2 = 9\kappa^2 r^2 (r^2 - 1) \quad \text{in } \mathbb{F}_\ell.$$

Dividing by the nonzero square $9\kappa^2 r^2$ shows that $r^2 - 1$ is a nonzero square. This holds for *every* root of f modulo every prime outside the finite exceptional set.

To pass to K , suppose that $\alpha^2 - 1$ is not a square there and choose β with $\beta^2 = \alpha^2 - 1$. Then $K(\beta)/K$ is quadratic. Its nontrivial automorphism fixes α and sends β to $-\beta$. We extend this automorphism to an element σ of $\text{Gal}(M/\mathbb{Q})$, where M is a Galois closure of $K(\beta)/\mathbb{Q}$.

Chebotarev's density theorem [9, §3, author-version p. 15] supplies infinitely many unramified rational primes with Frobenius conjugate to σ . By choosing the prime of M above each rational prime, we may take its Frobenius to be σ itself. Exclude residue characteristic two, primes dividing $6m$, and the finitely many primes at which α or β is nonintegral or β reduces to zero. At any remaining prime the reductions satisfy

$$\bar{\alpha}^\ell = \bar{\alpha}, \quad \bar{\beta}^\ell = -\bar{\beta} \neq \bar{\beta}.$$

The first equality puts $r = \bar{\alpha}$ in $\mathbb{F}_\ell$. The second keeps both roots $\pm\bar{\beta}$ of $X^2 - (r^2 - 1)$ outside $\mathbb{F}_\ell$. Thus $f(r) = 0$ and $r^2 - 1$ is a nonsquare, contrary to the condition just proved. We conclude that $\beta^2 = \alpha^2 - 1$ for some $\beta \in K$.

3.4 Traces in the cubic field

Set $z = \alpha + \beta$. The relation $(\alpha + \beta)(\alpha - \beta) = 1$ gives $z^{-1} = \alpha - \beta$ and $\alpha = (z + z^{-1})/2$, whence $\mathbb{Q}(z) = K$. We may therefore write the minimal polynomial of z as $z^3 + bz^2 + dz + w$, with rational b, d, w and $w \neq 0$. The polynomial for α gives

$$\text{Tr}(\alpha) = 0, \quad \text{Tr}(\alpha^2) = 2, \quad \text{Tr}(\alpha^3) = -3\eta.$$

We apply Newton's identities to z and z^{-1} and obtain

$$d = -bw, \quad b^2 + b(w - w^{-1}) = 1, \quad \eta = \frac{(b^2 + 1)(w + w^{-1})}{8}. \quad (6)$$

For the first equation, use $\text{Tr}(z) = -b$, $\text{Tr}(z^{-1}) = -d/w$. After substituting $d = -bw$, the second follows from $4\text{Tr}(a^2) = \text{Tr}(z^2) + 6 + \text{Tr}(z^{-2})$; the third follows from $8\text{Tr}(a^3) = \text{Tr}(z^3) + \text{Tr}(z^{-3})$, since $\text{Tr}(z + z^{-1}) = 0$.

The middle equation factors as $(bw - 1)(b + w) = 0$. Write $w = r/s$ in lowest terms, with $r \neq 0$, $s > 0$. Substituting $b = -w$ or $b = 1/w$ into the last equation and using $\eta = 6c/m$ gives respectively

$$m = \frac{48c rs^3}{(r^2 + s^2)^2} \quad \text{or} \quad m = \frac{48c r^3 s}{(r^2 + s^2)^2}.$$

Since $\gcd(r^2 + s^2, rs) = 1$ and $m \in \mathbb{Z}$, the square $(r^2 + s^2)^2$ divides 48. Hence $r^2 + s^2$ is 1, 2 or 4. Both r and s are nonzero, which leaves only 2: a sum of two nonzero integer squares cannot equal 1 or 4. Thus $|r| = s = 1$, and positivity of m gives $m = 12$. The only remaining numerators are

$$Q(n) = 2n(n+1)(n+2) + 1 \quad \text{and} \quad Q(n) = 2n(n+1)(n+2) - 1.$$

3.5 The contradiction modulo seven

In the plus case, choose a late block starting at $n \equiv 0 \pmod{7}$; in the minus case, start at $n \equiv 1 \pmod{7}$. The four successive numerators have residues

$$(u_n, u_{n+1}, u_{n+2}, u_{n+3}) \equiv \begin{cases} (1, 6, 0, 2) & (c = 1), \\ (4, 5, 0, 1) & (c = -1). \end{cases}$$

Call the first two denominator residues d_0, d_1 and the displayed numerator residues $c_0, c_1, 0, c_3$. From $0 = a_{n+1}c_1 - d_1$ and $c_3 = -a_{n+1}d_1$ we get $d_1^2 = -c_1c_3 = 2$. Thus $d_1 \in \{3, 4\}$. The first update, however, gives

$$d_1 = a_n d_0 = \frac{d_0(d_0 + c_1)}{c_0}.$$

As d_0 runs through $\mathbb{F}_7$, the plus case is $d_0(d_0 + 6)$ and the minus case is $2d_0(d_0 + 5)$. Both have image $\{0, 2, 5, 6\}$, which contains neither 3 nor 4. The zero numerator and its neighbours admit those two square roots, but the preceding update cannot produce either one. This is why we used four consecutive terms. Both cubics are excluded, proving Theorem 1.2.

Section 2 of the companion proves positive lower density of disagreement with each rational cubic of this form. That lower bound depends on the orbit and the cubic. For Theorem 1.2, exclusion of eventual equality suffices.

4 Bounded increases

We now return to the recurrence question. The numerator C_n need not have polynomial growth. Instead, we assume that its upward steps are bounded and prove that it eventually becomes constant. Decreases may be arbitrarily large. For a rational reciprocal tail, quadratic growth already supplies the relative limit in the following theorem.

Theorem 4.1 (bounded negative part). *Let $a, C, D : \mathbb{N} \rightarrow \mathbb{N}$ satisfy $a_n > 1$, $C_n > 0$, and*

Lean
Comparator

$$C_{n+1} + D_n = a_n C_n, \quad D_{n+1} = a_n D_n \quad (n \geq 0).$$

Put $E_n = D_n - (a_n - 1)C_n$. If

$$\frac{E_n}{C_n} \rightarrow 0 \quad \text{and} \quad E_n \geq -B \quad \text{for all sufficiently large } n$$

for some integer $B \geq 0$, then $E_n = 0$ for all sufficiently large n .

The limit implies $|E_n| < C_n$ eventually. Equivalently, for each $K \in \mathbb{N}$ we eventually have $K|E_n| < C_n$, the division-free form used in the linked theorem. We use the case $K = 1$ to propagate a zero error. If no zero occurs, the limit forces $C_n \rightarrow \infty$. The lower bound then stabilises the gcd and supplies bounded upward steps, contradicting a first-crossing argument below. For rational tails, this lower bound is the extra assumption.

4.1 Zero errors and descent

For brevity, an *exact state* consists of the sequences and recurrences in Theorem 4.1. In a reciprocal tail, $E_n = 0$ says $x_n = 1/(a_n - 1)$. We write $z_+ = \max(z, 0)$ and call $|E_n| < C_n$ *strict centring*.

Proposition 4.2 (error identities). *For an exact integer state,*

Lean
Comparator

$$C_{n+1} = C_n - E_n, \quad (a_{n+1} - a_n^2 + a_n - 1)C_{n+1} = a_n^2 E_n - E_{n+1}.$$

Proof. The first identity follows by substituting $D_n = E_n + (a_n - 1)C_n$ into the numerator recurrence. Substituting this expression at the next index and using $D_{n+1} = a_n D_n$ gives the second. $\square$

Equation (1) now gives $E_n/C_n = 1 - C_{n+1}/C_n \rightarrow 0$, hence strict centring eventually. Koizumi proves this relative-gap limit for an eventual pseudo-greedy tail [10, Corollary 3, p. 9]. His rounding convention also gives half-width centring, a stronger bound than is used here.

Theorem 4.3 (absorption and descent). *For a positive exact state with strict centring, $E_n = 0$ implies $E_{n+1} = 0$. For any positive integer state with $C_{n+1} = C_n - E_n$, eventual nonnegativity of E_n implies its eventual vanishing.*

Lean
Comparator

Proof. When $E_n = 0$, the second identity in Proposition 4.2 shows that C_{n+1} divides E_{n+1} . The inequality $|E_{n+1}| < C_{n+1}$ then forces $E_{n+1} = 0$. For the second assertion, $C_{n+1} = C_n - E_n$ makes the positive integer sequence C_n eventually nonincreasing, hence constant. $\square$

Corollary 4.4 (two zero errors). *If $E_n = E_{n+1} = 0$ and $C_{n+1} \neq 0$, then $a_{n+1} = a_n^2 - a_n + 1$. Thus eventual zero error in a positive exact state implies the eventual Sylvester recurrence.*

[Lean](#)
[Comparator](#)

Proof. Cancel the nonzero factor C_{n+1} in the second error identity. $\square$

After the centring threshold, one zero therefore makes every subsequent error zero. If there are no late zeros, integrality gives $|E_n| \geq 1$, and $|E_n|/C_n \rightarrow 0$ forces $C_n \rightarrow \infty$. The scalar update alone permits $C_n = n + 1$, $E_n = -1$ forever. To exclude such behaviour we shall use the denominator recurrence.

4.2 A stable gcd and a forbidden block

Proposition 4.5 (gcd stabilisation). *For a positive exact state, suppose that some fixed integer $B \geq 1$ satisfies $-B \leq E_n < 0$ at infinitely many indices. Then $G_n = \gcd(C_n, D_n)$ is eventually constant. Division by its stable value gives a reduced exact tail.*

[Lean](#)
[Comparator](#)

Proof. We have $G_n \mid G_{n+1}$ and $G_n \mid E_n$. Given n , choose $t \geq n$ with $-B \leq E_t < 0$. Then $G_n \leq G_t \leq -E_t \leq B$. Hence the positive divisibility chain stabilises. Division by its stable value preserves the recurrences and gives coprime states. $\square$

For this gcd argument, bounded negative errors need occur only arbitrarily late. To control the first crossing below, we shall need the upper bound on every sufficiently late upward step.

Lemma 4.6 (consecutive multiples). *For pairwise coprime integers $m_0, \dots, m_{B-1} \geq 2$ and every lower bound, there is a larger t such that $m_i \mid t + i$ for each $i < B$.*

[Lean](#)
[Comparator](#)

Proof. Solve $t \equiv -i \pmod{m_i}$ by the Chinese remainder theorem and add multiples of $\prod_i m_i$ to exceed the prescribed bound. $\square$

For instance, a sequence avoiding multiples of 2 and 3 cannot cross the pair $6k+2, 6k+3$ from below with upward steps at most two. Decreases do not affect this first-crossing argument.

Theorem 4.7 (Chinese remainder theorem and first crossing). *Let $u : \mathbb{N} \rightarrow \mathbb{N}$ tend to infinity and let $B \geq 1$ be an integer with $u_{n+1} \leq u_n + B$ for every n . There is no sequence of pairwise coprime integers $m_i \geq 2$ for which $\gcd(m_i, u_i) = 1$ whenever $i < t$.*

[Lean](#)
[Comparator](#)

Proof. Apply Lemma 4.6 to $m_0, \dots, m_{B-1}$, choosing $t > \max(u_0, \dots, u_B)$. There is a first $n > B$ with $u_n \geq t$. Then

$$t \leq u_n \leq u_{n-1} + B < t + B.$$

Thus $u_n = t + i$ for some $i < B < n$, and $m_i \mid u_n$, contrary to $\gcd(m_i, u_i) = 1$. $\square$

Bado proves a two-sided bounded-error criterion by this forbidden-block argument [7, Theorem 5.1, pp. 4–5]. The gcd stabilisation above allows us to use only the lower error bound.

Proof of Theorem 4.1. Suppose that the conclusion fails. Absorption excludes every sufficiently late zero. Since $|E_n| \geq 1$ there, the relative limit gives $C_n \rightarrow \infty$. There must also be infinitely many negative errors, for otherwise integer descent would make C_n constant. At those late negative indices, $1 \leq -E_n \leq B$, so $B \geq 1$ and Proposition 4.5 gives a stable gcd g . We divide by g and obtain $u_n = C_n/g \rightarrow \infty$ with

$$u_{n+1} - u_n = -E_n/g \leq B.$$

By Proposition 2.1, the multipliers $a_i \geq 2$ are pairwise coprime and each is coprime to every later numerator. They supply the moduli prohibited by Theorem 4.7, a contradiction. $\square$

Corollary 4.8. *Under Theorem 4.1, the multipliers satisfy $a_{n+1} = a_n^2 - a_n + 1$ eventually.*

Lean
Comparator

Proof. Theorem 4.1 gives eventual zero error; positivity of C_n and Corollary 4.4 give the recurrence. $\square$

4.3 The condition in the original sequence

Let $\gamma_n = a_n^2/a_{n+1} - 1$. For $t_n = P_n/a_n$,

$$t_{n+1} - t_n = t_n \gamma_n.$$

Thus the following hypothesis bounds upward increments of t_n . It allows t_n itself to be unbounded and allows decreases.

Corollary 4.9 (bounded increments of P_n/a_n). *Let $a_1 < a_2 < \dots$ be positive integers, $a_{n+1}/a_n^2 \rightarrow 1$, and $\sum_{n \geq 1} 1/a_n \in \mathbb{Q}$. Put $P_n = \prod_{j < n} a_j$. If*

Lean
Comparator

$$\limsup_{n \rightarrow \infty} \frac{P_n}{a_n} \left(\frac{a_n^2}{a_{n+1}} - 1 \right) < +\infty,$$

then $a_{n+1} = a_n^2 - a_n + 1$ for all sufficiently large n .

Proof. For the canonical integer tail, direct substitution gives

$$E_n + q \frac{P_n}{a_n} \gamma_n = q P_n \left(\frac{1}{a_{n+1}} - (a_n - 1) \sum_{k \geq n+2} \frac{1}{a_k} \right). \quad (7)$$

By the tail estimate, the subtracted term is at most $4(a_n - 1)/a_{n+1}^2 < 1/a_{n+1}$ eventually. The right side is positive, so an upper bound on $(P_n/a_n)\gamma_n$ gives a lower bound on E_n . The tail construction supplies positivity and the exact recurrences, while (1) gives $E_n/C_n \rightarrow 0$. Theorem 4.1 makes the error eventually zero, and Corollary 4.8 gives the Sylvester recurrence. $\square$

Koizumi's proof of Corollary 4(1) uses this product-weighted comparison [10, (11)–(12), p. 15]; that corollary assumes a nonpositive upper limit, whereas the argument here allows any finite upper limit. A corresponding LCM comparison appears below. For $a_n = b^{2^{n-1}}$, $b \geq 2$, the increments of P_n/a_n are zero. Since this is not a Sylvester tail, the corollary gives irrationality. For a Sylvester sequence starting at $a_1 > 1$, $a_n - 1 = (a_1 - 1)P_n$ and the increments tend to zero.

The inclusive rate $1/n$ gives another useful sufficient condition.

Corollary 4.10 (an inclusive one-sided $1/n$ bound). *Let $a_1 < a_2 < \dots$ be positive integers with $a_{n+1}/a_n^2 \rightarrow 1$ and $\sum_n 1/a_n \in \mathbb{Q}$. Suppose that for some $K \geq 0$ and $\varepsilon > 0$,*

Lean
Comparator

$$\gamma_n := a_n^2/a_{n+1} - 1 \leq \frac{1}{n} + \frac{K}{n^{1+\varepsilon}} \quad \text{for all large } n.$$

Then the sequence is eventually Sylvester. In particular, the conclusion holds under the pointwise eventual bound $\gamma_n \leq 1/n$.

Proof. The hypothesis bounds the positive part of γ_n eventually by $1/n + K/n^{1+\varepsilon}$. Since $t_{n+1}/t_n = 1 + \gamma_n$, comparison with n and the convergent product of $1 + K/n^{1+\varepsilon}$ gives $t_n = O(n)$. Hence $t_n \gamma_n \leq t_n(\gamma_n)_+ = O(1)$, and Corollary 4.9 applies. $\square$

The weaker condition $\limsup n(\gamma_n)_+ \leq 1$ does not suffice for the estimate $t_n = O(n)$. For example, the scalar ratios $1 + \gamma_n = 1 + (1 + 1/\log n)/n$ give a product of order $n \log n$. This example does not supply a reciprocal-tail counterexample.

For positive limiting increments, take $a_1 = 4$ and $a_{n+1} = \lceil na_n^2/(n+1) \rceil$, beginning 4, 8, 43, 1387. Here $a_n \geq 2 \cdot 2^{2^{n-1}}$ and

$$0 \leq 1 + \frac{1}{n} - \frac{a_n^2}{a_{n+1}} < \frac{4}{a_n^2}.$$

Consequently $t_n \sim Kn$ and $t_{n+1} - t_n \rightarrow K > 0$. The corollary proves irrationality because $\gamma_n \sim 1/n$ is incompatible with a Sylvester tail. The cubic example in the introduction has unbounded increments and requires the different proof in Section 3.

5 Finite total relative increase

An infinite product gives another way to bound a numerator. This argument needs only the scalar update $C_{n+1} = C_n - E_n$ and positivity on a discrete set. We assume summability of the relative upward increments. Mere convergence of those increments to zero would allow $C_n = n + 1$ and $C_n = (n + 1)^2$, both of which grow without bound.

Theorem 5.1 (a convergent sum of relative increases). *Let C_n be positive integers and E_n integers satisfying $C_{n+1} = C_n - E_n$. If*

Lean
Comparator

$$\sum_n \frac{(-E_n)_+}{C_n} < \infty,$$

then $E_n = 0$ eventually. Neither denominator dynamics nor vanishing relative error is required.

Proof. Set $\delta_n = (-E_n)_+/C_n$. Then

$$C_N \leq C_0 \prod_{n < N} (1 + \delta_n) \leq C_0 \exp\left(\sum_n \delta_n\right).$$

Choose an integer K above this bound. A strict rise of the integer sequence has size at least one, so it contributes at least $1/K$ to $\sum \delta_n$. Hence only finitely many rises occur. Thereafter C_n is a nonincreasing positive integer sequence, and its eventual constancy gives $E_n = 0$. $\square$

The proof also allows positive real C_n with integer E_n : the values lie in $C_0 + \mathbb{Z}$, whose bounded positive part is finite. Discrete increments are essential to this argument. With real errors, $C_n = 1 + 1/(n+1)$ decreases forever with zero relative-increase sum; without positivity, $C_n = -n-1$, $E_n = 1$ does so too. The companion's Section 13 gives the details.

On an exact reciprocal-tail orbit the conclusion gives the Sylvester recurrence. For the gap sequence of the pseudo-greedy expansion, the same criterion, with the same product bound and integer descent, appears in the Erdős Problem a Day working report on Problem #243, dated 12 August 2026 [8, "A global termination criterion"]; the statement above is for any positive integer sequence with $C_{n+1} = C_n - E_n$. This summability condition and the bound in Theorem 4.1 are different estimates. For exact reciprocal tails with $|E_n|/C_n \rightarrow 0$, each is equivalent to eventual vanishing of E_n ; neither estimate has been derived here from the unrestricted problem.

6 New maxima of an LCM numerator

Clearing the tails by a least common multiple removes repeated factors from their denominators. We shall control the resulting numerator by summing only at steps which exceed all preceding values. The argument counts first crossings of selected heights, so it also permits weights that decrease with the height. Put

$$L_n = \text{lcm}(q, a_1, \dots, a_{n-1}), \quad M_n = D_n/L_n, \quad U_n = C_n/M_n, \quad V_n = E_n/M_n.$$

The rational tail has denominator dividing L_n , so U_n and V_n are integers. This need not be a reduced fraction: U_n and L_n may share a factor. With $\rho_n = \gcd(L_n, a_n)$, the exact updates are

$$M_{n+1} = M_n \rho_n, \quad \rho_n U_{n+1} = U_n - V_n, \quad V_n = L_n - (a_n - 1)U_n.$$

These coordinates and updates also appear in Bado's September 2026 preprint. With his denominator parameter equal to q , his M_{n-1} , Δ_{n-1} , K_{n-1} , u_n and g_n are respectively our L_n , M_n , U_n , V_n and ρ_n ; his (19) is the middle update above [7, Proposition 7.1 and (16)–(20), pp. 6–7].

Since $L_n \mid qP_n$ and $V_n = (L_n/(qP_n))E_n$, multiplying (7) by $L_n/(qP_n)$ gives, eventually,

$$0 < V_n + \frac{L_n}{a_n} \left(\frac{a_n^2}{a_{n+1}} - 1 \right) \leq \frac{L_n}{a_{n+1}}. \quad (8)$$

The lower bound will transfer a growth condition to a bound on V_n . For the weighted version we will also need the upper bound, which makes the comparison error summable.

Write $R_n = \max_{j \leq n} U_j$ and $\mathcal{R} = \{n : U_{n+1} > R_n\}$. Strict centring already gives $U_{n+1} < U_n$ when $\rho_n \geq 2$, so every sufficiently late strict rise has $\rho_n = 1$. The stronger eventual bound $-U_n \leq 2V_n$, supplied by $V_n/U_n = E_n/C_n \rightarrow 0$, gives the quantitative estimate $U_{n+1} \leq 3U_n/4$ when $\rho_n \geq 2$. At a sufficiently late record step, where $\rho_n = 1$, the actual jump is $d_n = U_{n+1} - U_n = -V_n > 0$. This identity is not asserted at a contracting step with $\rho_n \geq 2$.

Theorem 6.1 (a convergent weighted sum over new maxima). *Assume the growth and rationality hypotheses of Problem 1.1. Let $f : [1, \infty) \rightarrow [0, \infty)$ be finite and nonincreasing, with $\int_1^\infty f(t) dt = \infty$. Then the sequence is eventually Sylvester if and only if, for some integer $B \geq 0$,*

$$\sum_{n \in \mathcal{R}} (-V_n - B)_+ f(U_n) < \infty.$$

The summand measures the actual increase $U_{n+1} - U_n$ beyond B , weighted by the starting height. Both $f(t) = 1/t$ and $f(t) = 1/\lceil t \log(et) \rceil$ are allowed, whereas $1/t^2$ has finite integral. We prove the theorem by showing that a non-Sylvester tail crosses enough heights to make the series diverge for every such f and B . The relative limit $V_n/U_n \rightarrow 0$ supplies no summability estimate.

Proof. Suppose the sequence is not eventually Sylvester. Absorption and integrality give

$$\frac{1}{U_n} \leq \frac{|V_n|}{U_n} = \frac{|E_n|}{C_n} \rightarrow 0.$$

Thus $U_n \rightarrow \infty$, so there are infinitely many record steps. At every sufficiently late one, $\rho_n = 1$, so a_n is coprime to L_n . The corresponding terms a_n are pairwise coprime: an earlier term divides the later L_n , while the term at that record is coprime to L_n . For any fixed $B \geq 1$, choose B such multipliers $m_0, \dots, m_{B-1} > B$ and take T after their indices and after the threshold beyond which records have $\rho_n = 1$. Their size follows from $a_n \rightarrow \infty$.

Put $P = \prod_i m_i$ and choose x with $m_i \mid x + i$ by the Chinese remainder theorem. We select the heights $\tau = x + B + kP > R_T, k \in \mathbb{Z}$. They are spaced P apart, and the first is at most $R_T + P$.

At the first crossing of a selected height, $U_n \leq R_n < \tau \leq U_n + d_n$. In particular this is a record step. We use its full jump $d_n = U_{n+1} - U_n$, including any recovery from an earlier decrease. If $d_n \leq B$, then $U_n \in [\tau - B, \tau)$, so some m_i divides U_n . It also divides L_n , hence divides $d_n = (a_n - 1)U_n - L_n$, contradicting $0 < d_n \leq B < m_i$.

It follows that every crossing has $d_n > B$. If one step first crosses $h \geq 1$ selected heights, their spacing gives $(h - 1)P < d_n$. Write $r = d_n - B \geq 1$. Since $P \geq B + 1$, $d_n = B + r \leq Pr$, and hence $h \leq r$. Each of those heights exceeds U_n , so monotonicity gives

$$\sum_{\substack{\tau \text{ first crossed} \\ \text{at step } n}} f(\tau) \leq (d_n - B)f(U_n).$$

Each selected height above R_T has exactly one first crossing. The first selected height is at most $R_T + P$; each later height is P further on. For $R_N \geq R_T + P$, comparison of each interval with its left endpoint gives

$$\sum_{\substack{T \leq n < N \\ n \in \mathcal{R}}} (-V_n - B)_+ f(U_n) \geq \frac{1}{P} \int_{R_T+P}^{R_N} f(t) dt. \quad (9)$$

Since $R_n \rightarrow \infty$, the right side diverges for every $B \geq 1$; $B = 0$ follows by domination. Conversely a Sylvester tail telescopes to $x_n = 1/(a_n - 1)$, so $V_n = 0$ eventually. $\square$

For example, $f(t) = 1/[t \log(et)]$ gives the sufficient condition

$$\sum_{n \in \mathcal{R}} \frac{(-V_n - B)_+}{U_n \log(eU_n)} < \infty.$$

Its finite lower bound in (9) is $P^{-1} \log(\log(eR_N)/\log(e(R_T + P)))$. Further fixed iterated logarithmic factors are allowed whenever the integral still diverges. These are specialisations of one crossing theorem.

The criterion also has an exact expression in the original growth defect. Put $\gamma_n = a_n^2/a_{n+1} - 1$ and $\theta_n = E_n/C_n$. The defect identity gives

$$\gamma_n + \theta_n = \frac{(1 - \theta_n)(a_n - 1 + \theta_{n+1})}{a_{n+1}}, \quad 0 < \gamma_n + \theta_n < 3/a_n$$

eventually. Thus the two nonnegative summands $U_n f(U_n)(\gamma_n - B/U_n)_+$ and $(-V_n - B)_+ f(U_n)$ differ by at most $3U_n f(U_n)/a_n$. We verify that this comparison error is summable. Since $a_n x_n \rightarrow 1$, we have $C_n/a_n \sim qP_n/a_n^2$, and

$$\frac{P_{n+1}/a_{n+1}^2}{P_n/a_n^2} = \frac{a_n^3}{a_{n+1}^2} \rightarrow 0.$$

The ratio test gives $\sum_n C_n/a_n < \infty$; now $U_n \leq C_n$ and $f(U_n) \leq f(1)$ make the comparison error summable. The errors in (7) and (8) are also absolutely summable, since each is eventually between zero and $qP_n/a_{n+1} = O(P_n/a_n^2)$. Consequently Theorem 6.1 is equivalent to finiteness of

$$\sum_{n \in \mathcal{R}} U_n f(U_n) \left(\frac{a_n^2}{a_{n+1}} - 1 - \frac{B}{U_n} \right)_+ \quad (10)$$

for some B . It remains to prove finiteness under the unrestricted hypotheses. The jump in either expression is $U_{n+1} - U_n$. Substituting $R_{n+1} - R_n$ loses the recovery from earlier decreases, which the crossing estimate uses. Nor would termwise convergence of the summands to zero establish finiteness.

Corollary 6.2 (a bound using the least common multiple). *Assume the hypotheses of Problem 1.1. Write $A_n = \text{lcm}(a_1, \dots, a_{n-1})$ with $A_1 = 1$. If*

[Lean](#)
[Comparator](#)

$$\limsup_{n \rightarrow \infty} \frac{A_n}{a_n} \left(\frac{a_n^2}{a_{n+1}} - 1 \right) < \infty,$$

then the sequence is eventually Sylvester.

Proof. Since $L_n/A_n = q/\gcd(q, A_n)$ lies in $[1, q]$, an upper bound on the expression in the corollary gives an upper bound on $(L_n/a_n)(a_n^2/a_{n+1} - 1)$; negative terms remain negative. Equation (8) therefore bounds the negative part of V_n . Choose an integer B above that eventual bound. The record series then vanishes after a finite prefix, so Theorem 6.1 applies. $\square$

Erdős and Straus assume a nonpositive upper limit in this LCM expression [1, Theorem 3, p. 132]: their N_k is A_{k+1} and their growth ratio has index $k+1$. Tijdeman and Yuan extend this type of criterion to positive numerators [4]. Here any finite upper bound suffices under the quadratic-limit assumption. Since $A_n \mid P_n$, the LCM hypothesis is no stronger than the product hypothesis in Corollary 4.9: if the product expression is at most B , the LCM expression is at most $\max(B, 0)$. When the earlier terms are pairwise coprime the two weights agree; repeated prime factors can make the LCM much smaller. This comparison does not assert the existence of a non-Sylvester rational example satisfying one bound but not the other.

The finite upper limit means an eventual upper bound; it does not require q to divide the LCM of the preceding denominators. For positive integer summand numerators b_n , the modified error is $V_n = b_n L_n - (a_n - 1)U_n$. The companion, Section 6 under “Integer coefficients”, separately proves boundedness from a lower bound on V_n and eventual constancy when $V_n/U_n \rightarrow 0$ is also assumed. The latter conclusion must not be read into the first hypothesis alone. The same section links finite examples separating them; compare Badea [3, p. 316] and Tijdeman–Yuan [4]. The companion’s Section 3 retains the signed-series comparison with Duverney [2, Corollary 3.2, p. 287] and the different growth hypotheses for irrationality exponents in [14, Theorem 1, author-version p. 2]. The crossing theorem uses neither of these further sets of hypotheses.

7 Further questions

Proposition 7.1 (necessary profile). *The integer tail of a sequence satisfying Problem 1.1’s hypotheses but not its conclusion has $E_n \neq 0$ eventually, $|E_n|/C_n \rightarrow 0$, unbounded negative magnitudes along negative indices, and*

Lean
Comparator

$$\sum_n \frac{(-E_n)_+}{C_n} = \infty.$$

Proof. Absorption excludes late zeros, descent excludes an eventually nonnegative error, and Theorems 4.1 and 5.1 exclude the two finiteness conditions. $\square$

For $B \geq 0$ put

$$F_B(X) = \sum_{\substack{n \in \mathcal{R} \\ U_n \leq X}} (-V_n - B)_+.$$

The weighted criterion reduces the problem to a statement about this cumulative sum. We would need to show, from growth and rationality, that for some fixed integer $B \geq 0$,

$$\liminf_{X \rightarrow \infty} \frac{F_B(X)}{X} = 0. \quad (11)$$

If the recurrence is not eventually Sylvester, first crossings instead give $F_B(X) \geq X/P_B - O_B(1)$, with an orbit-dependent CRT modulus P_B . The sum groups each record step by its starting numerator U_n , not by the running maximum R_n or the time index. On a non-Sylvester tail, $U_n \rightarrow \infty$, so only finitely many of these steps begin below any fixed X . To see the equivalence with an admissible weight, enumerate the record indices by

j and apply the next lemma with $u_j = U_n$ and $w_j = (-V_n - B)_+$. The lemma supplies a weight once (11) holds. Establishing that estimate would contradict the first-crossing lower bound and exclude every non-Sylvester rational tail.

Lemma 7.2 (weights and linear density). *Let u_j be positive integers, let $w_j \geq 0$, and put $F(X) = \sum_{u_j \leq X} w_j$, with the sum allowed a priori to be $+\infty$. Then $\liminf_{X \rightarrow \infty} F(X)/X = 0$ if and only if there is a finite nonincreasing $f : [1, \infty) \rightarrow [0, \infty)$ with $\int_1^\infty f(t) dt = \infty$ and $\sum_j w_j f(u_j) < \infty$.*

Lean
Comparator

Proof. Suppose the lower limit vanishes. Choose cutoffs at which the ratio $F(X)/X$ is small. Since F is constant on each $[m, m+1)$, there are integers $X_1 < X_2 < \dots$ with $X_k \geq 2^k$ and $F(X_k) \leq 2^{-k} X_k$. Set

$$f(t) = \sum_{k \geq 1} \frac{\mathbf{1}_{[1, X_k]}(t)}{X_k}.$$

This function is nonincreasing and bounded by $\sum_k 2^{-k}$. Each cutoff contributes $1 - 1/X_k$ to its integral, but at most 2^{-k} to the weighted sum. Interchanging nonnegative sums therefore gives

$$\int_1^\infty f(t) dt = \sum_k (1 - 1/X_k) = \infty, \quad \sum_j w_j f(u_j) = \sum_k \frac{F(X_k)}{X_k} \leq 1.$$

Conversely, let f be such a weight and suppose $F(m) \geq cm$ for some $c > 0$ and all integers $m \geq M$. A nonincreasing nonnegative function with divergent integral is positive everywhere, so $F(X) \leq f(X)^{-1} \sum_j w_j f(u_j)$ is finite. Partial summation gives, for $N > M$,

$$\sum_{u_j \leq N} w_j f(u_j) = F(N)f(N) + \sum_{m=1}^{N-1} F(m)(f(m) - f(m+1)) \geq c(Mf(M) + \sum_{m=M+1}^N f(m)),$$

every term being nonnegative. The right side diverges as $N \rightarrow \infty$, because $\sum_{m \geq 1} f(m) \geq \int_1^\infty f(t) dt = \infty$. $\square$

7.1 Arithmetic compatibility

The remaining estimate requires the denominator recurrence as well as control of the error. For example, $C_n = n^2 + 1$, $E_n = -(2n + 1)$ has vanishing relative error and the necessary scalar pattern. But at $n = 2, 3, 4$ its numerators are 5, 10, 17. The update $10 = 5a_2 - D_2$ forces $5 \mid D_2$, hence $5 \mid D_3 = a_2 D_2$; the next update would force $5 \mid 17$. It is not an exact tail. The companion's Section 14 also constructs sequences avoiding sparse fixed moduli and distinguishes that avoidance from compatibility with all later denominators. Neither property alone supplies (11).

Conversely, a full exact orbit already contains the analytic data. For a zero-indexed orbit with $a_n > 1$, $C_n > 0$, $D_n \geq 0$ and $E_n/C_n \rightarrow 0$, Section 4 of the companion proves $D_n > 0$, $C_n/D_n \rightarrow 0$, $\sum_n 1/a_n = C_0/D_0$ and quadratic growth. These are global assumptions on an infinite orbit. A scalar profile or a finite admissible prefix supplies only part of them.

Problem 7.3 (growth of repeated denominator factors). For every rational-tail orbit satisfying the hypotheses but not the conclusion of Problem 1.1, must

$$\limsup_{n \rightarrow \infty} \frac{\log M_n}{n} > 0?$$

Equivalently, must there be a $K \geq 1$ for which

$$2^n \leq M_n^K$$

at infinitely many indices?

Section 2 gives $C_{n+1}/C_n \rightarrow 1$; averaging logarithms therefore gives $\log C_n = o(n)$. Since $1 \leq M_n \leq C_n$, every such orbit already satisfies $\log M_n/n \rightarrow 0$. A positive answer therefore holds exactly when no such orbit exists: proving it from failure of Sylvester behaviour would exclude every counterexample.

Acknowledgements. I thank Wouter van Doorn for advice on the exposition, in particular for asking what the additional bound excludes and for pointing out unnecessary terminology and notation.

A A factorial residue reduction for forced orbits

A constant-negative error $E_n = -m$ with $C_0 = c$ produces $C_n = c + nm$ and the shape equation

$$D_n + m = (a_n - 1)C_n. \quad (12)$$

In the case $m = c = 1$, where (12) reads $D_n + 1 = (a_n - 1)(n + 1)$, each multiplier is determined by its predecessor; we call such an orbit *forced*. At index n the numerator of the next multiplier is

$$\text{num}(n, a) = (n + 1)a^2 - (n + 2)a + (n + 3),$$

the [forced numerator](#), and the divisor is $n + 2$. The orbit continues for another step exactly when this division is exact. Here survival counts exact divisions; it does not impose the positivity conditions of an infinite reciprocal-tail orbit. The [formal survival predicate](#) requires exact division at each step.

Example A.1 (the shape equation running until it fails). The forced orbit from $a = 3$ reads this way: $\text{num}(0, 3) = 6$ is divisible by 2 and gives $a_1 = 3$, while $\text{num}(1, 3) = 13$ is not divisible by 3, so the orbit stops there.

Direct iteration can produce very large intermediate integers. To decide whether the first h divisions are exact, however, it suffices to know the initial value modulo $(h + 1)!$. Computing a pseudo-greedy orbit through residues modulo a shrinking product modulus is Koizumi's method [10, Remark 2 and Algorithm 1, pp. 13–14]; for the forced numerator that product is a factorial.

Theorem A.2 (factorial residue reduction). *For all h and all integers $a \equiv b \pmod{(h + 1)!}$, the orbit from a survives h forced updates exactly when the orbit from b does.*

[Lean](#)
[Comparator](#)

Proof. Define $M(0, i) = 1$ and $M(h+1, i) = (i+2)M(h, i+1)$. Thus $M(h, 0) = (h+1)!$. We prove the stronger statement that, at index i , congruent inputs modulo $M(h, i)$ survive the same h updates. There is nothing to prove for $h = 0$.

For the inductive step, suppose $a \equiv b \pmod{(i+2)M(h, i+1)}$. Since $\text{num}(i, \cdot)$ is an integer polynomial, its values at a and b are congruent modulo that product. In particular, one is divisible by $i+2$ if and only if the other is. If neither is divisible, both orbits stop. Otherwise their quotients are congruent modulo $M(h, i+1)$, so the induction hypothesis applies to the remaining h updates at index $i+1$. Taking $i = 0$ proves the claim. $\square$

At $h = 1$, survival means $2 \mid a^2 - 2a + 3$, or equivalently that a is odd. This illustrates Theorem A.2 with modulus $2! = 2$.

The enumeration in Example 9.1 of the companion, over $2 \leq a_0 < 5000$, gives at most 17 successful updates (18 values including a_0). The seed $a_0 = 1$ is excluded because it is fixed and violates $a_n \geq 2$. This finite calculation establishes only the stated range. The infinite [constant-negative exclusion](#) requires the separate argument under the hypotheses given there.

B Sources and related work

The companion *Reciprocal-Tail Rigidity: Theorems, Proofs and Questions* gives full proofs and fixed-revision source links. Section 2 treats regular rates and the stronger cubic disagreement result; Sections 4–5 develop exact tails; Sections 11–12 give the gcd and bounded-increment arguments. The LCM results are in Section 6, scalar summability in Section 13, and open estimates and failed constructions in Section 14. A final map records the dependencies. The auxiliary constructions need not be rational reciprocal series; they satisfy only the hypotheses stated for them.

The older integer-remainder method is represented by Erdős–Straus [12, Theorem 2.1, pp. 85–86]. The polynomial Cantor-series hypotheses in Hančl–Tijdeman [13, Theorem 2.2, pp. 39–40] and the freely selectable series of Problem #270 [15, Theorems 1–2] concern different settings; the companion’s Section 3 explains the distinction. The Isabelle/HOL work [16] formalises older Erdős–Straus criteria, not Problem #243. The general regular-rate extraction in the companion also uses the standard Gamma-ratio asymptotics recorded in [17]; the cubic specialisation here uses only its explicit polynomial F_n .

The margin marks link statements to fixed-revision Lean declarations and recorded Comparator evidence. The printed cubic argument uses Chebotarev, while the supplied formal square-specialisation proof uses a Dedekind-zeta pole comparison. Agreement of their conclusions does not verify each step of the printed argument. No Lean build, Comparator replay or independent review was performed for this revision, and the evidence records retain source-pin and index gaps. The estimate in Section 7 remains open. The formal [factorial residue reduction](#) uses the [shrinking-modulus induction](#), [polynomial congruence](#), and [cancellation after exact division](#). Its modulus is identified by the [ascending-factorial formula](#) and its [factorial value at the initial index](#).

References

- [1] P. Erdős and E. G. Straus, *On the irrationality of certain Ahmes series*, J. Indian Math. Soc. (N.S.) **27** (1964), 129–133. MR 175848.
- [2] D. Duverney, *Irrationality of fast converging series of rational numbers*, J. Math. Sci. Univ. Tokyo **8** (2001), 275–316. MR 1837165.
- [3] C. Badea, *A theorem on irrationality of infinite series and applications*, Acta Arith. **63** (1993), no. 4, 313–323, doi:10.4064/aa-63-4-313-323.
- [4] R. Tijdeman and P. Yuan, *On the rationality of Cantor and Ahmes series*, Indag. Math. (N.S.) **13** (2002), no. 3, 407–418, doi:10.1016/S0019-3577(02)80018-0.
- [5] P. Erdős and R. L. Graham, *Old and New Problems and Results in Combinatorial Number Theory*, Monogr. Enseign. Math. **28**, Geneva, 1980, p. 64.
- [6] P. Erdős, *On the irrationality of certain series: problems and results*, in A. Baker (ed.), *New Advances in Transcendence Theory*, Cambridge UP, 1988, pp. 102–109, doi:10.1017/CBO9780511897184.009.
- [7] I. O. Bado, *Prime-Support Rigidity and Primitive Pseudo-Greedy Dynamics: Partial Progress on Erdős Problem #243*, preprint posted September 2026, doi:10.13140/RG.2.2.36612.08325.
- [8] P. White with Claude (Anthropic), *Erdős #243: working report*, Erdős Problem a Day, page dated 12 August 2026. AI-assisted, unrefereed working report.
- [9] P. Stevenhagen and H. W. Lenstra, Jr., *Chebotarëv and his density theorem*, Math. Intelligencer **18** (1996), no. 2, 26–37, doi:10.1007/BF03027290. Page references are to the linked author version dated 23 March 1995.
- [10] J. Koizumi, *Irrationality of the reciprocal sum of doubly exponential sequences*, Integers **26** (2026), Paper No. A28, 17 pp., doi:10.5281/zenodo.18714404. Locators refer to this published version.
- [11] T. F. Bloom, *Erdős Problem #243*, accessed 28 July 2026.
- [12] P. Erdős and E. G. Straus, *On the irrationality of certain series*, Pacific J. Math. **55** (1974), no. 1, 85–92.
- [13] J. Hančl and R. Tijdeman, *On the irrationality of polynomial Cantor series*, Acta Arith. **133** (2008), no. 1, 37–52, doi:10.4064/aa133-1-3. Locators refer to the published version.
- [14] D. Duverney, T. Kurosawa and I. Shiokawa, *Irrationality exponents of certain fast converging series of rational numbers*, Tsukuba J. Math. **44** (2020), no. 2, 235–250, doi:10.21099/tkbjm/20204402235. Theorem locators follow the linked 14-page author version.
- [15] T. Crmarić and V. Kovač, *On the irrationality of certain super-polynomially decaying series*, Colloq. Math. **179** (2025), 55–68, doi:10.4064/cm9628-5-2025. Theorem locators follow arXiv:2504.18712v1.
- [16] A. Koutsoukou-Argraki and W. Li, *Irrationality Criteria for Series by Erdős and Straus*, Archive of Formal Proofs, 12 May 2020. An Isabelle/HOL formalisation; the archive entry identifies the results formalised.
- [17] National Institute of Standards and Technology, *Digital Library of Mathematical Functions*, §5.11(iii), formula 5.11.12, accessed 16 September 2026.