

A Countermodel for Growth-and-Parity Arguments on the Prime-Gap Dyadic Series

*Complete-tail countermodels, sparse rationalisation, and integral-shift criteria for
Erdős Problem #251*

WILL COOK^{*}
September 2026

ABSTRACT

A sparse perturbation of the actual prime gaps can make their dyadic sum rational while preserving every fixed eventual congruence of the gaps and cumulative sums, and the empirical distributions of all unnormalised gap blocks of length $o(\log \log X)$. For every $0 < \varepsilon < 1$, its corrections can be bounded by $(\log n)^\varepsilon$ eventually. The construction uses pairs with fixed unweighted total and variable dyadic weight. Lean checks its pair identities and the proposition it proves, and the scheduling argument written here is ordinary mathematics. The simpler positive even word $a_n = 2(n^2 + 4n + 2)$ has complete dyadic tails $U_N = 2(N + 4)^2$ and rational sum $\sum_{j \geq 1} a_j 2^{-j} = 32$. We identify the actual complete prime-gap tails and classify rationality by integrality of their shifts. Two adjacent shifts in $(-1, 1)$ with unequal associated gaps cannot both be integral. Their joint occurrence cofinally for every positive shift would prove irrationality of the prime series; this prime-specific assertion remains unproved.

1 Introduction

The complete countermodel in one calculation. Set $a_n = 2(n^2 + 4n + 2)$ and $U_N = 2(N + 4)^2$. Since $2U_N - U_{N+1} = a_{N+1}$, finite telescoping gives

$$\sum_{j=1}^m \frac{a_{N+j}}{2^j} = U_N - \frac{U_{N+m}}{2^m} \rightarrow U_N.$$

Thus U_N is the actual infinite tail, not merely a solution of its recurrence, and the sum at $N = 0$ is 32. Moreover

$$a_{n+1} - a_n = 4n + 10, \quad U_{N+h} - U_N = 2h(2N + h + 8) \in \mathbb{Z}.$$

These positive even coefficients grow polynomially and are unbounded and nonperiodic, but every tail difference is integral. The unshifted normalisation $\sum_{n \geq 0} a_n 2^{-(n+1)}$ is 18. Proposition 2.4 states the resulting obstruction.

^{*}*Authorship and AI use.* Will Cook built and directed the research infrastructure and reviewed the claims when he could. AI agents did most of the research and drafting. Cook did not independently verify every claim.

Preserving more of the actual gaps. The quadratic example has the wrong cumulative growth to imitate the primes. A different construction addresses that limitation. If $A = \sum_{n \geq 0} a_n 2^{-(n+1)}$ converges, fix $M \geq 1$ and a cutoff K , and choose a rational $r \in (A, A + M2^{-K})$. Write

$$\frac{2^K(r - A)}{M} = \sum_{j \geq 0} \frac{\delta_j}{2^{j+1}}, \quad \delta_j \in \{0, 1\}.$$

Set $b_n = a_n$ for $n < K$ and $b_{K+j} = a_{K+j} + M\delta_j$. Then $\sum_{n \geq 0} b_n 2^{-(n+1)} = r$. Every coefficient retains its residue modulo M and changes by at most M ; each cumulative sum changes by at most Mn . Applied to prime gaps, this retains the cumulative $n \log n$ scale that the prime number theorem gives for p_n [13, Chapter 6]. Taking M even retains parity as well. Thus a bounded residue-preserving perturbation can have a rational value without sharing the polynomial example's growth. Neither construction asserts that its cumulative sums are prime.

Sparse corrections and local information. A single changed coefficient can alter every later cumulative residue. We therefore pair corrections: changing their allocation changes the dyadic value while leaving their unweighted total fixed. Sparsity has a separate job. Most short blocks then meet no correction at all, which preserves their empirical distributions without requiring the corrected positions to be prime.

Proposition 1.1 (sparse congruence-preserving rationalisation). *Let a_n be nonnegative integers with $\sum_{n \geq 0} a_n 2^{-(n+1)} = A < \infty$, let $K \in \mathbb{N}$, and let $f : \mathbb{N} \rightarrow \mathbb{R}$ tend to $+\infty$. There exist a set $S \subseteq [K, \infty)$ of upper Banach density zero and a nondegenerate interval $I \subset (A, \infty)$ such that every $r \in I$ is realised by a nonnegative correction e supported on S , with $0 \leq e_n \leq f(n)$ eventually, $\sum (a_n + e_n) 2^{-(n+1)} = r$, and every fixed modulus eventually dividing both e_n and $\sum_{i < n} e_i$. For the actual prime gaps and the specific choice $f(n) = (\log(n+3))^\varepsilon$, $0 < \varepsilon < 1$, one can in addition require $|S \cap [X, 2X]| = O_\varepsilon(X / \log \log X)$, so that all unnormalised blocks of length $o(\log \log X)$ are preserved in total variation.*

Construction. Use disjoint triples of coordinates $n_j - 1, n_j, n_j + 1$, with spacings $s_j = n_j - n_{j-1} \geq 4$, and moduli M_j forming a divisibility chain. Put $D_j = 2^{s_j} - 1$. Starting from $C_0 = 0$, define

$$c_j = (-C_j) \bmod M_j, \quad C_{j+1} = C_j + c_j + M_j D_j.$$

For a free digit $0 \leq d_j \leq D_j$, assign all three corrections:

$$e_{n_j-1} = c_j, \quad e_{n_j} = M_j d_j, \quad e_{n_j+1} = M_j(D_j - d_j),$$

and set $e_n = 0$ elsewhere. The complete triple has total $c_j + M_j D_j$, independent of d_j , so every buffer is fixed before the free digits are chosen. Immediately after the buffer, the cumulative correction is divisible by M_j ; both pair entries preserve it. Later buffers preserve every previously imposed divisor. Thus each fixed modulus eventually divides both individual and cumulative corrections if it eventually divides M_j .

Here is an explicit schedule for an arbitrary $f \rightarrow \infty$. Set

$$h(n) = \inf_{m \geq n} \min(f(m), m),$$

choose $n_{-1} \geq K$ so large that $h(n_{-1}) \geq 32$, and recursively let

$$k_j = \max\{k \geq 2 : k!2^{k+2} \leq h(n_{j-1})\}, \quad M_j = k_j!, \quad s_j = k_j + 2, \quad n_j = n_{j-1} + s_j.$$

The function h is nondecreasing, tends to infinity, and satisfies $h(n) \leq n$. Hence k_j is nondecreasing and tends to infinity. Each triple entry is at most $M_j 2^{s_j} \leq h(n_{j-1})$, which is at most $f(n)$ at every coordinate of that triple and at most n_{j-1} . This proves the size bound and summability of the correction. The spacings tend to infinity, so the union of triples has upper Banach density zero.

Put $w_j = M_j 2^{-n_j-2}$. The weighted correction separates as

$$\sum_n \frac{e_n}{2^{n+1}} = \beta + \sum_j d_j w_j, \quad \beta = \sum_j (c_j 2^{-n_j} + D_j w_j).$$

The buffer contribution is included in the fixed, positive number β . For $i > j$, monotonicity of the moduli gives

$$D_i w_i \geq M_j (2^{-n_{i-1}-2} - 2^{-n_i-2}), \quad \sum_{i>j} D_i w_i \geq w_j.$$

The digit intervals therefore overlap. Greedy selection fills the complete interval $[0, F]$, where $F = \sum_j D_j w_j > 0$: at each step the remainder is between zero and the remaining capacity, which tends to zero. This is the variable-digit covering of Fridy's lemma on generalised bases, whose condition $r_n \leq \sum_{i>n} k_i r_i$ is the overlap inequality above [6, Lemma, p. 194]; Fridy assumes nonincreasing weights, and the argument here uses only the overlap. Kovač and Tao use the same nested covering in [7, Lemma 5.1], where the choices at each stage are reciprocals of the integers in an interval; here they are the $D_j + 1$ equally spaced values $d w_j$. Take $I = (A + \beta, A + \beta + F)$.

For the polylogarithmic bound, instead set, after a sufficiently large initial cutoff,

$$s_j = \left\lfloor \frac{\varepsilon}{2} \log_2 \log(n_{j-1} + 3) \right\rfloor, \quad k_j = \max\{k \geq 2 : k! \leq (\log(n_{j-1} + 3))^{\varepsilon/4}\},$$

with $M_j = k_j!$, $n_j = n_{j-1} + s_j$ and the same D_j and buffers. Then $s_j \geq 4$ and $M_j 2^{s_j} \leq (\log(n_{j-1} + 3))^{3\varepsilon/4}$. The preceding congruence, convergence and interval arguments still apply. Spacing comparable to $\log \log n_j$ gives $|S \cap [X, 2X]| = O_\varepsilon(X / \log \log X)$. Couple the original and corrected length- m blocks at the same starting index in $[X, 2X]$. At most $m|S \cap [X, 2X + m]|$ starts meet a correction, so their empirical distributions differ in total variation by $O_\varepsilon(m / \log \log X)$ for $m = o(\log \log X)$. The complete argument and the all-index congruence check are expanded in the companion record, subsection "Paired corrections and cumulative congruences". The pair, buffer and conditional interval-filling identities are the Lean-checked core; the scheduling and density arguments are ordinary mathematics. $\square$

Every bounded test of these blocks has the same asymptotic empirical average. A density-zero witness family can lie inside the changed blocks, so its occurrence is not implied by this preservation statement. The reconstructed positions retain their congruences and cumulative growth; they are not asserted to be prime.

The prime series and its tails. Index the primes from zero: $p_0 = 2$, $p_1 = 3$, and $g_n = p_{n+1} - p_n$. In this notation Erdős Problem #251 asks whether

$$\Pi = \sum_{n \geq 0} \frac{p_n}{2^{n+1}} \quad \text{is irrational.}$$

This is the usual one-based series $2/2 + 3/4 + 5/8 + \dots$. The first gaps are 1, 2, 2, 4, 2, 4, 2, 4, 6, 2; only g_0 is odd. Let $G = \sum_{n \geq 0} g_n 2^{-(n+1)}$. The exact identities are

$$\Pi = 2 + G, \quad T_N = \sum_{j \geq 1} \frac{g_{N+j}}{2^j}, \quad T_0 = 2G - 1, \quad T_{N+1} = 2T_N - g_{N+1}.$$

Convergence and the identification of these actual tails are established below. They are not additional assumptions under a hypothetical rational value of Π .

What remains to be distinguished. For any integer-digit recurrence, iteration gives

$$T_N = 2^N T_0 - B_N, \quad T_{N+h} - T_N = 2^N (2^h - 1) T_0 - C_{N,h}, \quad B_N, C_{N,h} \in \mathbb{Z}.$$

A rational initial value forces an eventual integral shift; an irrational initial value permits no integral positive shift. This is the arithmetic reason for the lcm-diagonal criterion in Theorem 2.1. A more local sufficient condition uses two adjacent shift values in $(-1, 1)$ and a nonzero associated gap difference. If both values were integral they would be zero, contradicting their recurrence. The missing input is their joint occurrence at arbitrarily late indices for every positive shift. The countermodels show why several coarse hypotheses do not supply it.

Relation to prior work. Erdős stated the fixed-denominator question in his work on irrational series [1, p. 94][2, p. 62][4, p. 103]. His irrationality result for prime numerators with factorial denominators does not apply to this dyadic denominator sequence. Kovač's counterexample to a related variable-denominator expectation likewise does not decide the fixed-denominator problem [18]. An adjacent dyadic theorem of Erdős and Pomerance concerns bounded digits recording comparisons of the largest prime factors of consecutive integers [3, §7]; our prime numerators and gaps are unbounded. Assuming Kuperberg's uniform Hardy–Littlewood prime-tuples conjecture [8, Conjecture 1.3], Land posted a research draft dated 5 September 2026 that proves irrationality [9, Theorem 2], and Ringer a preprint dated 11 September 2026 that proves $\sum_n p_n B^{-n}$ normal to base B for each integer $B \geq 2$ [19, Corollary 1.2]; both arguments are formalised in Lean. Nothing in this note assumes a prime-tuples hypothesis, and Erdős Problem #251 remains open unconditionally. Interval filling of the kind used in Proposition 1.1 goes back to variable-digit expansions: for nonincreasing weights $r_i \rightarrow 0$ and digit bounds k_i , Fridy showed that every number in $(0, \sum_i k_i r_i)$ is a sum $\sum_i a_i r_i$ with integers $0 \leq a_i \leq k_i$ exactly when $r_n \leq \sum_{i > n} k_i r_i$ for every n [6, Lemma, p. 194]. It also has recent precedents for sums of unit fractions. Kovač and Tao use it to show that bounded perturbations of the denominators 2^n can give a rational sum, for instance $\sum_{n \geq 1} 1/(2^n + b_n) = 3/4$ with every $b_n \in \{1, \dots, 5\}$, answering a question of Erdős and Graham on irrationality sequences [7, Lemma 5.1, Theorem 2.5 and §5]. Van Doorn and Kovač combine dense

filling with a divisibility chain among distinguished terms that eventually absorbs every positive integer, so that finite reciprocal sums take every rational value in an interval [14, Lemma 7 and Proposition 8]. The corrections in Proposition 1.1 use a chain of factorial moduli with the same absorption property to preserve every fixed eventual congruence. The detailed historical comparisons and their hypotheses belong to the companion reasoning record. No priority claim is made for the elementary identities or constructions in this note.

Notation and organisation. Write $\mathbb{N} = \{0, 1, \dots\}$, $\mathbb{Z}$ for the integers, $\text{Irr}(x)$ for irrationality, $\text{den}(q)$ for the reduced denominator of $q \in \mathbb{Q}$, and φ for Euler's totient. Section 2 isolates the recurrence-level classifiers and the polynomial obstruction. Section 3 proves the actual prime-to-gap identity; Section 4 proves the shift classification and local criterion. Section 5 explains the remaining carry freedom, and Section 6 states the exact prime-specific obligation. The sources and verification boundary are recorded after the mathematics.

Keywords. irrationality; prime gaps; dyadic series; summation by parts; Lean 4.
MSC 2020. 11J72 (primary); 11N05, 68V20 (secondary).

2 Integral shifts and recurrence obstructions

We first record the recurrence-level conclusions. Modulo one the recurrence is multiplication by two, and the classifications below use only reduced denominators and Euler's congruence $d \mid 2^{\varphi(d)} - 1$ for odd d . They apply to arbitrary integer digits and therefore separate the finite algebra from the prime-specific input still missing in Problem #251.

For a real recurrence put

$$\Delta_h T(N) = T_{N+h} - T_N,$$

and let $L_0 = 1, L_j = \text{lcm}(1, \dots, j)$ for $j \geq 1$.

Theorem 2.1 (lcm-diagonal criterion). *Let $g : \mathbb{N} \rightarrow \mathbb{Z}$ and $T : \mathbb{N} \rightarrow \mathbb{R}$ satisfy $T_{N+1} = 2T_N - g_{N+1}$ for every N . Then*

$$\text{Irr}(T_0) \iff \Delta_{L_j} T(L_j) \notin \mathbb{Z} \text{ for every } j \geq 0.$$

Proof. Iteration gives $T_N = 2^N T_0 - B_N$ with $B_N \in \mathbb{Z}$, and hence

$$\Delta_h T(N) = 2^N (2^h - 1) T_0 - C_{N,h}, \quad C_{N,h} \in \mathbb{Z}.$$

For $h \geq 1$ the coefficient of T_0 is nonzero, so an integral shift forces $T_0 \in \mathbb{Q}$. Conversely, write $T_0 = a/(2^s d)$ in lowest terms with d odd. Euler's congruence gives $d \mid 2^{\varphi(d)} - 1$. Choose j with $L_j \geq s$ and $\varphi(d) \mid L_j$. Then $2^s d \mid 2^{L_j} (2^{L_j} - 1)$, making the displayed L_j -shift integral at $N = L_j$. This contradicts the right-hand condition. $\square$

The statement is **irrational initial iff all lcm diagonal nonintegral**.

The factorial schedule $j!$ has the same divisibility property, but the lcm schedule is pointwise no larger and is the canonical endpoint used here. The theorem is an exact

one-sequence reformulation; it gives no non-integrality statement for the actual prime-gap recurrence.

The adjacent-small-mismatch condition also admits an exact normal form. For a rational recurrence and a fixed h , write

$$D_N = \Delta_h T(N), \quad \delta_N = g_{N+h+1} - g_{N+1}.$$

Then $D_{N+1} = 2D_N - \delta_N$.

Theorem 2.2 (signed two-window normal form). *Assume that δ_N is even. The conjunction*

$$-1 < D_N < 1, \quad -1 < D_{N+1} < 1, \quad \delta_N \neq 0$$

is equivalent to

$$(\delta_N = 2 \text{ and } \frac{1}{2} < D_N < 1) \quad \text{or} \quad (\delta_N = -2 \text{ and } -1 < D_N < -\frac{1}{2}).$$

Proof. The recurrence and the two unit windows give $-3 < \delta_N < 3$. A nonzero even integer in that interval is 2 or -2 . Substitution into $D_{N+1} = 2D_N - \delta_N$ gives the stated half-window and, in the reverse direction, recovers the second unit window. $\square$

The rational normal form is [adjacent small mismatch](#) iff [signed two window](#). The same equivalence for real states is [real signed two window](#).

For the longer block, set

$$B_{h,N,r} = \sum_{i=0}^{r-1} 2^{r-1-i} \delta_{N+i}; \quad D_{N+r} = 2^r D_N - B_{h,N,r}.$$

Call the data-dependent affine condition

$$A_{N,r} : \quad D_{N+r} \in -B_{h,N,r} + 2^{r+1} \mathbb{Z}.$$

Theorem 2.3 (affine and fixed-lattice circularity). *For every rational dyadic tail recurrence and all $h, N, r \geq 0$,*

$$A_{N,r} \iff D_N \in 2\mathbb{Z}. \quad (2.1)$$

Consequently, if every δ_N is even, then

$$(\forall N_0 \exists N, r : N_0 < N \text{ and } \neg A_{N,r}) \iff D_N \notin \mathbb{Z} \text{ for arbitrarily large } N. \quad (2.2)$$

There is a second equivalence. Let $b : \mathbb{N} \rightarrow \mathbb{Q}$ satisfy $|D_N| \leq b(N)$ for every N , and suppose that for every N and every positive integer q there is an r with

$$2b(N+r)q < 2^r. \quad (2.3)$$

Then

$$\begin{aligned} \forall N_0 \exists N, r : N_0 < N \text{ and } \forall z \in \mathbb{Z}, \quad b(N+r) < |B_{h,N,r} - 2^r z| \\ \iff D_N \notin \mathbb{Z} \text{ for arbitrarily large } N. \end{aligned} \quad (2.4)$$

Proof. Substituting $D_{N+r} = 2^r D_N - B_{h,N,r}$ into $A_{N,r}$ cancels the observed block from both sides and leaves $D_N = 2z$. This proves (2.1). After one recurrence step, evenness of δ_N identifies even integrality at $N + 1$ with ordinary integrality at N . The forward implication in (2.2) follows, while the reverse implication chooses a nonintegral D_N and the legal depth $r = 0$.

For (2.4), eventual integrality and the block identity give some $z \in \mathbb{Z}$ with

$$B_{h,N,r} - 2^r z = -D_{N+r},$$

so the displayed strict separation contradicts $|D_{N+r}| \leq b(N+r)$. Conversely, if D_N is nonintegral and has reduced denominator q , then its distance from every integer is at least $1/q$. Choose r from (2.3), scale this separation by 2^r , and use the block identity together with $|D_{N+r}| \leq b(N+r)$. The triangle inequality gives $|B_{h,N,r} - 2^r z| > b(N+r)$ for every integer z . $\square$

The three displayed equivalences are assembled in one declaration, [affine circularity bundle](#).

The first equivalence shows that the apparent affine hierarchy contains no depth-dependent information: it is the pullback of even integrality through the recurrence identity. The fixed lattice removes that data dependence, but under the stated growth condition rational denominator separation already forces every required escape. Hence neither cofinal condition is an independent source of information about consecutive primes.

Finally, the coarse coefficient profile itself has an exact infinite counterexample.

Proposition 2.4 (quadratic polynomial-shift countermodel). *Put*

$$a_n = 2(n^2 + 4n + 2), \quad U_n = 2(n + 4)^2.$$

Then a_n is positive, even and strictly increasing, $U_{n+1} = 2U_n - a_{n+1}$, every shift $U_{N+h} - U_N$ is integral,

$$a_{N+2} - a_{N+1} \neq 2, -2 \quad \text{for every } N,$$

and

$$\sum_{j \geq 1} \frac{a_j}{2^j} = 32$$

as the kernel-checked identity for the countermodel series value.

Proof. Expand $a_{n+1} - a_n = 4n + 10$ and $2U_n - U_{n+1} = a_{n+1}$. The h -shift is $U_{N+h} - U_N = 2h(2N + h + 8)$, an even integer. The finite telescope $\sum_{j=1}^N a_j/2^j = 32 - U_N/2^N$ is exact rational algebra from the recurrence; $U_N/2^N \rightarrow 0$ because a quadratic is dominated by 2^N . The nonnegative terms therefore have infinite sum 32. Strict growth makes the word unbounded and nonperiodic. $\square$

The eight displayed clauses are assembled as one statement in [polynomial countermodel](#).

3 Summation by parts, with the endpoint retained

Summation by parts trades a sequence for its consecutive differences. The form recorded here is exact: it carries no error term, it assumes nothing about the sequence, and it retains the endpoint term rather than absorbing it into an estimate. For a sequence P of rational numbers and $n \geq 0$ put

$$D(P, n) = \sum_{i=0}^{n-1} \frac{P(i)}{2^{i+1}}, \quad \Delta(P, n) = \sum_{i=0}^{n-1} \frac{P(i+1) - P(i)}{2^{i+1}},$$

both empty, hence zero, at $n = 0$. We call these the [dyadic partial sum](#) and the [dyadic difference sum](#) of P .

Proposition 3.1 (finite summation by parts). *For every rational sequence P and every $n \geq 0$,*

$$D(P, n+1) = P(0) + \Delta(P, n) - \frac{P(n)}{2^{n+1}}.$$

Proof. Expand the difference sum and shift its first index. All interior coefficients of $P(i)$ become 2^{-i-1} ; the remaining terms are $P(0)$ and the displayed endpoint $-P(n)2^{-n-1}$. This also covers $n = 0$. $\square$

This is the [finite summation by parts](#). No positivity, monotonicity or convergence is required.

Specialising to $P(i) = p_i$, whose first value is $p_0 = 2$, and writing $g_i = p_{i+1} - p_i$ for the zero-based gaps, gives the reformulation. The displayed partial sums with denominators 2^i are twice these, by the [factor-of-two identity](#).

Theorem 3.2 (prime-gap reformulation). *Let $p_0 = 2, p_1 = 3, \dots$ be the primes in increasing order and $g_i = p_{i+1} - p_i$. For every $n \geq 0$,*

$$\sum_{i=0}^n \frac{p_i}{2^{i+1}} = 2 + \sum_{i=0}^{n-1} \frac{g_i}{2^{i+1}} - \frac{p_n}{2^{n+1}}.$$

This is the [prime-gap summation by parts](#). The leading 2 is p_0 . At $n = 2$ both sides equal $19/8$. The identity is summation by parts specialised to the primes; Tao posted this reduction on the problem's forum thread on 7 October 2025 [17, comment of 7 October 2025].

3.1 The infinite identity and the irrationality equivalence

Write

$$u_n = \frac{p_n}{2^{n+1}}, \quad v_n = \frac{g_n}{2^{n+1}}$$

for the [prime dyadic terms](#) and the [prime-gap dyadic terms](#), so that $\sum_{n \geq 0} u_n = \Pi$. The termwise identity $v_n = 2u_{n+1} - u_n$ is the [dyadic discrete derivative](#). It expresses each gap term as an integer combination of two consecutive prime terms, so once (u_n) is summable the gap series can be summed by rearranging two copies of the prime series, which is what the following proof does.

Theorem 3.3 (infinite prime-gap identity). *Let $u_n = p_n/2^{n+1}$ and $v_n = g_n/2^{n+1}$. If (u_n) is summable, then (v_n) is summable and*

$$\sum_{n \geq 0} u_n = 2 + \sum_{n \geq 0} v_n.$$

Proof. The shifted sequence (u_{n+1}) is summable. Sum $v_n = 2u_{n+1} - u_n$ and use $u_0 = 1$:

$$\sum_{n \geq 0} v_n = 2 \left(\sum_{n \geq 0} u_n - u_0 \right) - \sum_{n \geq 0} u_n = \sum_{n \geq 0} u_n - 2.$$

□

The summability transfer is the [gap-series summability theorem](#), and the displayed identity is the [infinite prime-gap identity](#). Summability of both series is part of the conclusion in [infinite prime gap identity](#) rather than a hypothesis carried by the reader.

Corollary 3.4 (exact irrationality reformulation). *With u_n and v_n as in Theorem 3.3, and (u_n) summable,*

$$\text{Irr} \left(\sum_{n \geq 0} u_n \right) \iff \text{Irr} \left(\sum_{n \geq 0} v_n \right).$$

The corresponding [zero-based series with denominator \$2^n\$](#) is $4 + 2 \sum_{n \geq 0} v_n$ and has the same irrationality status. Termwise it is twice the normalised series, by the [infinite factor-of-two identity](#).

These are the [normalised irrationality equivalence](#), the [displayed-series identity](#), and the [displayed-series irrationality equivalence](#). The three clauses are also available as the single declaration [irrationality reformulation](#).

The summability hypothesis. The formal source now proves the elementary polynomial bound $p_n \leq 1250(n+1)^4$, using prime counting and central-binomial growth, and deduces summability directly ([polynomial bound](#), [summability](#)). Thus Theorem 3.3 and Corollary 3.4 are now unconditional. The [unconditional infinite prime-gap identity](#) removes the summability hypothesis; the prime number theorem is therefore needed neither for convergence nor for this identity. It enters this note only through the $n \log n$ scale of the primes [13, Chapter 6]. The open content is exactly irrationality of the prime-gap series. Concretely, $\Pi = 3.674643966 \dots$ is irrational if and only if $\sum_{n \geq 0} g_n 2^{-(n+1)} = 1.674643966 \dots$ is, and neither is known.

4 The tail recurrence and integral shifts

The series is not attacked directly. Suppose $\sum_{i \geq 0} a_i 2^{-(i+1)}$ converges, with every a_i an integer, and rescale its tails by putting

$$T_N = 2^{N+1} \sum_{i \geq N} \frac{a_i}{2^{i+1}} = \sum_{j \geq 1} \frac{a_{N+j}}{2^j}.$$

Two facts follow immediately. First $T_{N+1} = 2T_N - a_{N+1}$, so moving one level along doubles the rescaled tail and subtracts a single coefficient. Second $T_0 = 2 \sum_{i \geq 0} a_i 2^{-(i+1)} - a_0$, so the sum is rational exactly when T_0 is. The whole of Section 4 therefore studies that recurrence in isolation, assuming nothing about the coefficients beyond the fact that they are integers; the prime-gap instance is resumed at the end of the section. The following definition is the object so obtained, stripped of its origin.

Definition 4.1. Let $g : \mathbb{N} \rightarrow \mathbb{Z}$ and $T : \mathbb{N} \rightarrow \mathbb{Q}$. Say T satisfies the *dyadic tail recurrence* with *digits* g when

$$T_{N+1} = 2T_N - g_{N+1} \quad \text{for every } N.$$

We call the sequence $(T_N)_{N \geq 0}$ the *orbit* of T_0 under g , write $\sigma_h(N) = T_{N+h} - T_N$ for the *shift* of length h at N , and call a rational number *integral* when it is the image of an integer.

These are the [tail recurrence](#), the [shift](#), and [integrality](#). The digits are arbitrary integers. In the intended instance they are the prime gaps and T_N is the complete gap tail $\sum_{j \geq 1} g_{N+j} 2^{-j}$. That identification, and the recurrence over $\mathbb{R}$ with no rationality hypothesis, already sit in `RealPrimeGapTail.lean`. Every statement below is a theorem about Definition 4.1.

One small orbit, referred to again below, is worth having in view. Take every digit $g_N = 0$ and $T_0 = 1/12$. Then $T_{N+1} = 2T_N$, so the orbit is $\frac{1}{12}, \frac{1}{6}, \frac{1}{3}, \frac{2}{3}, \frac{4}{3}, \dots$, and $\sigma_h(N) = (2^h - 1)2^N/12$. The shift of length 2 is not integral at $N = 0$, where $\sigma_2(0) = \frac{1}{3} - \frac{1}{12} = \frac{1}{4}$, and is integral at $N = 2$, where $\sigma_2(2) = \frac{4}{3} - \frac{1}{3} = 1$; the shift of length 1 is $\sigma_1(N) = 2^N/12$ and is never integral. The change at $N = 2$ is accounted for by Theorem 4.6, and the failure at $h = 1$ by Theorem 4.3.

Iterating the recurrence h times multiplies T_N by 2^h and accumulates an explicit integer, which we now name. Define $B_{0,N} = 0$ and $B_{h+1,N} = 2B_{h,N} + g_{N+h+1}$, so that $B_{h,N} = g_{N+1}2^{h-1} + \dots + g_{N+h}$: the [tail block](#). Thus $B_{1,N} = g_{N+1}$, $B_{2,N} = 2g_{N+1} + g_{N+2}$ and $B_{3,N} = 4g_{N+1} + 2g_{N+2} + g_{N+3}$: the block puts the weights $2^{h-1}, \dots, 2^0$ on the h digits following index N .

Theorem 4.2 (block identity). *Let $T : \mathbb{N} \rightarrow \mathbb{Q}$ satisfy the dyadic tail recurrence with integer digits g , and let $B_{h,N}$ be as above. For every N and h ,*

$$T_{N+h} = 2^h T_N - B_{h,N}, \quad \text{hence} \quad \sigma_h(N) = (2^h - 1) T_N - B_{h,N}.$$

Proof. Induct on h using the recursions for T and B , then subtract T_N to obtain the shift identity. $\square$

The first identity is the [iterated block identity](#); the second is the [scaled shift identity](#). Both are also assembled in [block identity](#), and the real-state form is [real block identity](#).

The shifts themselves satisfy the [shift step identity](#)

$$\sigma_h(N+1) = 2\sigma_h(N) - (g_{N+h+1} - g_{N+1}).$$

Since $B_{h,N}$ is an integer, Theorem 4.2 converts a question about the shift into a question about the single scaled term $(2^h - 1)T_N$.

Theorem 4.3 (integral-shift criterion). *Let $T : \mathbb{N} \rightarrow \mathbb{Q}$ satisfy the dyadic tail recurrence with integer digits g . For every N and h , the shift $\sigma_h(N)$ is integral if and only if $(2^h - 1)T_N$ is integral.*

Proof. By Theorem 4.2 the two differ by the integer $B_{h,N}$, and subtracting an integer does not change integrality, by the [integer-shift invariance](#). $\square$

This is the [integral-shift criterion](#). Thus integrality depends only on the reduced denominator of T_N .

The denominator criterion is exact:

$$\sigma_h(N) \in \mathbb{Z} \iff \text{den}(T_N) \mid 2^h - 1. \quad (3.4)$$

This is the [denominator classification](#). Euler's totient supplies one admissible shift length when the denominator is odd; it is a witness, not the classification itself.

Theorem 4.4 (a shift of totient length). *Let $T : \mathbb{N} \rightarrow \mathbb{Q}$ satisfy the dyadic tail recurrence with integer digits g , and let φ be Euler's totient function. If the reduced denominator d of T_N is odd, then $\sigma_{\varphi(d)}(N)$ is integral.*

Proof. Since d is odd, 2 and d are coprime, so Euler's congruence gives $2^{\varphi(d)} \equiv 1 \pmod{d}$, that is $d \mid 2^{\varphi(d)} - 1$. Writing $2^{\varphi(d)} - 1 = dk$ and $T_N = u/d$ in lowest terms, $(2^{\varphi(d)} - 1)T_N = ku$ is an integer, and Theorem 4.3 transfers this to the shift. The Euler multiplier itself is the [totient multiplier at odd denominator](#). $\square$

This is the [totient shift](#).

For $\text{den } T_N = 3$, the shift of length 2 is integral while the shift of length 1 is not. For denominator 5, length 4 is admissible.

The hypothesis is a genuine restriction: the argument uses coprimality of 2 with the denominator, and the even part of a denominator is exactly what the doubling in the recurrence acts on. It cannot be dropped. If $T_N = 1/2$ then $2^h - 1$ is odd for every $h \geq 1$, so $(2^h - 1)T_N$ is never an integer and, by Theorem 4.3, no shift at N is integral.

Theorem 4.5 (propagation). *Let $T : \mathbb{N} \rightarrow \mathbb{Q}$ satisfy the dyadic tail recurrence with integer digits g , and fix h and N . If $\sigma_h(N)$ is integral, then $\sigma_h(N + k)$ is integral for every $k \geq 0$.*

Proof. By the shift step identity, $\sigma_h(N + 1) = 2\sigma_h(N) - (g_{N+h+1} - g_{N+1})$ is an integer combination of an integer and two digits; this is the [one-step propagation](#), and the claim follows by induction on k . $\square$

This is the [propagation to every later index](#).

The three preceding theorems combine as follows, and this is the statement the rest of the note rests on. The special case is immediate: if $\text{den } T_0$ is already odd, then Theorem 4.4 at $N = 0$ makes the shift of length $\varphi(\text{den } T_0)$ integral and Theorem 4.5 keeps it integral at every later index, so one may take $N_0 = 0$. In general a denominator carries a power of two as well, and the key point is that the doubling in the recurrence annihilates exactly the 2-adic part of a denominator, and nothing else: after finitely many steps the orbit therefore reaches a term with odd reduced denominator, which is precisely the situation Theorem 4.4 handles. No control of the digits is needed anywhere.

Theorem 4.6 (exact denominator dynamics and eventual integrality). *Let $T : \mathbb{N} \rightarrow \mathbb{Q}$ satisfy the dyadic tail recurrence with integer digits g (Definition 4.1). Then some fixed positive shift is integral from some point onwards:*

$$\exists h \geq 1 \exists N_0 \forall N \geq N_0, \quad \sigma_h(N) \in \mathbb{Z}.$$

Proof. At every step the reduced denominator obeys the exact recurrence

$$\text{den}(T_{N+1}) = \frac{\text{den}(T_N)}{\gcd(2, \text{den}(T_N))}.$$

Thus each even denominator loses exactly one factor of 2, while an odd denominator is unchanged. After finitely many steps the denominator is odd. Theorem 4.4, applied at that index s , supplies the positive shift $h = \varphi(\text{den } T_s)$, and Theorem 4.5 keeps that shift integral at every later index. $\square$

The one-step formula is the [denominator recurrence](#), with its [odd case](#) and [even case](#).

In the orbit displayed after Definition 4.1 the proof runs as follows: $\text{den } T_0 = 12 = 2^2 \cdot 3$, so $s = 2$, the orbit reaches $T_2 = 1/3$ with odd denominator, and $h = \varphi(3) = 2$. That is exactly the shift length seen to be integral there from index 2 onwards, and no shorter one works.

Three features of the argument are used later. The proof may take as its number of preparatory steps the 2-adic valuation of $\text{den } T_0$. The resulting shift length h is the totient of the odd denominator reached from a hypothetical rational initial value, and is not known in advance for the prime-gap orbit. This is why this argument requires Problem 6.2 for every h , rather than for one preassigned shift length. Finally, the digits enter only through the integer $B_{s,0}$, so the conclusion holds for an arbitrary integer digit sequence.

Denominator factorisation and cancellation are the [power-of-two/odd denominator scaling](#) and the [multiplier at the two-adic exponent](#), packaged in the [fixed-denominator theorem](#); the quantified conclusion is the [eventual integral-shift theorem](#) for the actual prime-gap tail state. The displayed indexed form is [eventual integral positive shift](#).

The useful contrapositive is stated for a real recurrence. Call its shifts *cofinally non-integral* when, for every fixed $h \geq 1$ and every threshold N_0 , some $N \geq N_0$ has $\sigma_h(N) \notin \mathbb{Z}$. This is precisely the negation of the conclusion of Theorem 4.6: no shift length whatever becomes integral and stays integral.

Theorem 4.7 (exact rationality classification). *Let $T : \mathbb{N} \rightarrow \mathbb{R}$ satisfy $T_{N+1} = 2T_N - g_{N+1}$ with integer g . If its shifts are defined by $\sigma_h(N) = T_{N+h} - T_N$, then the following are equivalent:*

- (i) T_0 is rational;
- (ii) $\sigma_h(N)$ is integral for some $h \geq 1$ and some N ;
- (iii) for some fixed $h \geq 1$, $\sigma_h(N)$ is integral at every sufficiently large N .

Consequently T_0 is irrational if and only if every positive-length shift is non-integral at every index, equivalently if and only if the shifts are cofinally non-integral.

Proof. For (i) $\Rightarrow$ (iii), choose $q \in \mathbb{Q}$ whose real cast is T_0 . The real block identity identifies the whole orbit with the cast of the rational recurrence starting at q ; Theorem 4.6 applied to that rational orbit then gives (iii). The implication (iii) $\Rightarrow$ (ii) is immediate. For (ii) $\Rightarrow$ (i), the real block identity gives

$$\sigma_h(N) = (2^h - 1)T_N - B_{h,N}.$$

Here $B_{h,N}$ and $\sigma_h(N)$ are integers and $2^h - 1 \neq 0$, so T_N is rational. Iterating the recurrence backwards through the block identity then makes T_0 rational. Negating the pointwise and eventual forms gives the two irrationality formulations. $\square$

The [rational actual-tail state](#) obeys the [tail recurrence](#), and a hypothetical rational value of the gap series supplies the [rational-tail representation](#). The exact real classifiers are [one integral positive shift](#), [eventual integrality](#), [pointwise non-integrality](#), and [cofinal non-integrality](#). All four equivalences are collected in [rationality classification](#).

The actual prime-gap orbit. The concrete tails

$$T_N = \sum_{j \geq 1} \frac{g_{N+j}}{2^j}$$

are already identified, with no rationality hypothesis. The polynomial prime bound gives convergence, and $T_{N+1} = 2T_N - g_{N+1}$ is the same identification. If $G = \sum_{n \geq 0} g_n/2^{n+1}$, then $T_0 = 2G - 1$. Together with Theorem 3.3, this shows that Π , G , and T_0 have the same rationality status. Thus Theorem 4.7 identifies Problem #251 exactly with cofinal shift escape for T . The older rational-candidate representation remains in `PrimeGapDyadicTail.lean`. What is not proved is the cofinal non-integrality needed by Theorem 4.7.

4.1 Two adjacent small shifts cannot both be integral

The exact classifier reduces irrationality to cofinal non-integrality, a condition of infinite precision imposed on a complete tail. The key point is that inside the open interval $(-1, 1)$ integrality is equality with zero, so on that range the one-step shift recurrence

$$\sigma_h(N + 1) = 2\sigma_h(N) - (g_{N+h+1} - g_{N+1})$$

turns simultaneous integrality of two adjacent shifts into a single comparison of digits. What this buys is a *certificate*: a condition attached to a single pair of adjacent indices, whose verification already contradicts integrality at that pair. The point of arranging the argument this way is that the distance of a shift from the integers never has to be estimated; it suffices to know that both shifts lie in $(-1, 1)$ and that two digits differ.

Theorem 4.8 (adjacent small-shift obstruction). *Let $T : \mathbb{N} \rightarrow \mathbb{Q}$ satisfy the dyadic tail recurrence with integer digits g . Fix h and N . If*

$$-1 < \sigma_h(N) < 1, \quad -1 < \sigma_h(N + 1) < 1, \quad g_{N+h+1} \neq g_{N+1},$$

then $\sigma_h(N)$ and $\sigma_h(N + 1)$ cannot both be integral. Consequently, if such a pair occurs beyond every threshold, the h -shift is not eventually integral.

Proof. An integral rational strictly between -1 and 1 is zero. If both shifts were integral, both would therefore vanish, and substitution in the displayed shift step identity would give $g_{N+h+1} = g_{N+1}$, a contradiction. The cofinal statement chooses one such adjacent pair after the alleged onset of integrality. $\square$

The finite contradiction is the [adjacent small-shift obstruction](#); its quantified form is the [cofinal small-mismatch theorem](#); and the actual-prime-gap specialisation is the [prime-gap specialisation](#). Theorem 4.8 is conditional on its two tail inequalities; no theorem asserting that such pairs occur is claimed here. The local obstruction and its cofinal consequence are stated as one declaration in [rational small pair bundle](#).

All three are stated for a rational orbit, while the prime-gap tail T of Section 4 is real, so the rational statement does not on its own discharge the instances that arise downstream. The same argument gives the real form directly.

Corollary 4.9 (adjacent small-shift obstruction, real form). *Let $T : \mathbb{N} \rightarrow \mathbb{R}$ satisfy $T_{N+1} = 2T_N - g_{N+1}$ with integer digits g , and write $\sigma_h(N) = T_{N+h} - T_N$. Fix h and N . If*

$$-1 < \sigma_h(N) < 1, \quad -1 < \sigma_h(N+1) < 1, \quad g_{N+h+1} \neq g_{N+1},$$

then $\sigma_h(N)$ and $\sigma_h(N+1)$ do not both lie in $\mathbb{Z}$. Consequently, if such a pair occurs beyond every threshold, the h -shift is not eventually integral.

Proof. The shift identity holds over $\mathbb{R}$, and the only integer in $(-1, 1)$ is still zero. Thus the proof of Theorem 4.8 applies without change, including its cofinal conclusion. $\square$

Corollary 4.9 is [real small pair bundle](#), and the resulting implication for the actual prime series is [real small pair prime endpoint](#). These live declarations can postdate the historical public snapshot used by the source links.

The third hypothesis, on its own, is available. For the actual gaps tabulated in Section 1 it reads $g_4 = 2 \neq 4 = g_3$ at $h = 1, N = 2$; and Proposition 4.10 below says precisely that for each fixed $h \geq 1$ the inequality $g_{N+h+1} \neq g_{N+1}$ holds for arbitrarily large N , since its failure from some index onwards is eventual periodicity with period h . What is missing is their joint occurrence with the digit mismatch at the same indices. Each inequality constrains a complete infinite tail.

Proposition 4.10 (prime gaps do not become periodic). *For every positive h , the actual consecutive-prime-gap sequence is not eventually periodic with period h .*

Proof. The gaps are unbounded, by the standard construction: the interval from $n! + 2$ to $n! + n$ contains no prime. Far stronger lower bounds for large gaps are known [5, Theorem 1, p. 66], but unboundedness is all that is needed. An eventually periodic natural-valued sequence has finite range after its preperiod, while its finite initial segment is bounded as well; hence it is bounded, a contradiction. $\square$

This is the [unboundedness of the actual gaps](#) and the [non-eventual periodicity](#) of the gap sequence. The statement at the index convention displayed above is [prime gaps not eventually periodic](#). Combining nonperiodicity with eventual strict smallness of one positive shift would also exclude eventual integrality, but eventual smallness at every sufficiently large index is stronger than the cofinal adjacent-pair hypothesis in Theorem 4.8 and is not asserted here.

5 Nonperiodic coefficients with a rational sum

There is an exact finite obstruction recorded in the pinned source: a second example has finite sum $-n/2^n$ and non-eventually-periodic coefficients ([finite endpoint; coefficient nonperiodicity](#)). The actual prime gaps are also not eventually periodic ([prime-gap nonperiodicity](#)), but that fact alone is insufficient.

For a bounded canonical base-two digit expansion, rationality forces eventual periodicity. Prime gaps are unbounded coefficients, so that criterion does not apply. The following telescoping construction shows explicitly how nonperiodic coefficients can have a rational sum.

The construction runs the emission of coefficients backwards. Let $K : \mathbb{N} \rightarrow \mathbb{Q}$ be arbitrary, read K_n as the value carried into level n , and put $\kappa_n = 2K_n - K_{n+1}$: the [carry coefficient](#). That definition is exactly the statement that

$$\frac{K_n}{2^n} = \frac{\kappa_n}{2^{n+1}} + \frac{K_{n+1}}{2^{n+1}},$$

so at each level the carried value splits into one emitted coefficient and a new carry. Nothing at all is assumed about K : not integrality, not positivity, not any bound. That is the sense in which the carry is free, and it is what the counterexample below exploits.

Proposition 5.1 (exact telescoping). *Let $K : \mathbb{N} \rightarrow \mathbb{Q}$ be arbitrary and $\kappa_n = 2K_n - K_{n+1}$. For every $n \geq 0$,*

$$\sum_{i=0}^{n-1} \frac{\kappa_i}{2^{i+1}} = K_0 - \frac{K_n}{2^n}.$$

Proof. A routine induction on n ; the added term is $(2K_n - K_{n+1})/2^{n+1} = K_n/2^n - K_{n+1}/2^{n+1}$. $\square$

This is the [carry telescoping identity](#).

Consequence for the coefficients. If $K_n 2^{-n} \rightarrow 0$, the emitted partial sums converge to K_0 . A parity-compatible positive example is

$$K_0 = \frac{5}{2}, \quad K_n = 2n + 2 \ (n \geq 1), \quad \kappa_0 = 1, \quad \kappa_n = 2n \ (n \geq 1),$$

for which

$$\sum_{i=0}^{n-1} \frac{\kappa_i}{2^{i+1}} = \frac{5}{2} - \frac{K_n}{2^n} \rightarrow \frac{5}{2}.$$

Thus the emitted coefficients $1, 2, 4, 6, 8, 10, \dots$ are positive, even after the first term, unbounded and not eventually periodic, although their dyadic sum is rational. Rationality alone therefore cannot imply eventual periodicity even for a positive, parity-correct integer coefficient sequence. The example does not claim that these coefficients are prime gaps; it isolates the additional arithmetic information any successful argument must use.

6 Complements and further questions

Problem #251 is open. Unconditional convergence of both dyadic series, the [unconditional infinite prime-gap identity](#), and the [real-to-rational scaled-tail representation](#) are proved. Every rational candidate supplies one positive fixed shift which is integral at every sufficiently late tail index: the [eventual integral-shift theorem](#). The denominator construction selects that same shift so that prime-gap nonperiodicity also rules out its eventual confinement to the open unit interval ([not eventually small](#)). This conclusion is compatible with rationality: it says that the rationally forced shift cannot support an eventual-smallness contradiction. It supplies neither eventual smallness nor cofinally many adjacent small mismatches for that shift. The remaining issue is recurrence or anti-concentration of the *actual* prime-gap shifts. Throughout, $g_i = p_{i+1} - p_i$ are the zero-based prime gaps of Section 1. Write $E(h)$ for the cofinal escape statement in (5.1) at shift h .

Problem 6.1 (divisor-hitting shift escape). For every $r \geq 1$, does some positive multiple of r escape cofinally?

$$\forall r \geq 1 \exists m \geq 1 : \quad E(mr).$$

Divisor-hitting escape changes the organisation of the quantifiers. Within an integer-digit recurrence it is equivalent to irrationality, just as universal shift escape is: a rational state has a positive eventual order, and every positive multiple of that order is an integral shift. A quantitatively stated sufficient estimate can still be easier to attack, but that requires a separately proved analytic bound. Forward propagation is Theorem 4.5.

Problem 6.2 (universal prime-gap shift escape). For every $h \geq 1$ and every N_0 , prove that some $N \geq N_0$ satisfies

$$\sum_{j \geq 1} \frac{g_{N+h+j} - g_{N+j}}{2^j} \notin \mathbb{Z}. \quad (5.1)$$

The series in (5.1) is exactly $T_{N+h} - T_N$. Theorem 4.7 makes Problem 6.2 equivalent to irrationality of Π . Within the integer-digit recurrence, divisor-hitting and universal escape are equivalent: a hypothetical rational value chooses a shift length from the odd part of its reduced denominator, and the cocycle then makes every positive multiple eventually integral. The two formulations organise the quantifiers differently; they do not give a strict logical weakening.

Problem 6.3 (cofinal adjacent small mismatch). For every fixed $h \geq 1$ and every N_0 , prove that some $N \geq N_0$ satisfies

$$\begin{aligned} \left| \sum_{j \geq 1} \frac{g_{N+h+j} - g_{N+j}}{2^j} \right| &< 1, \\ \left| \sum_{j \geq 1} \frac{g_{N+h+1+j} - g_{N+1+j}}{2^j} \right| &< 1, \\ g_{N+h+1} &\neq g_{N+1}. \end{aligned} \quad (5.2)$$

The two sums are adjacent h -shifts. Corollary 4.9 turns each instance of (5.2) into a finite contradiction to simultaneous integrality; Theorem 4.8 is the rational case. A cofinal family of such pairs therefore proves Problem 6.2; Theorem 4.7 then gives irrationality of the gap series, and Corollary 3.4 transfers it to Π . The endpoint takes every positive h : a proof only at $h = 1$ eliminates one eventual-integrality possibility and does not discharge the classifier. Digit mismatches and small shifts must occur at the same indices; separate cofinal occurrence statements do not establish that intersection. This formulation deliberately asks only for sporadic adjacent pairs; the stronger assertion that a fixed shift is eventually always smaller than one is unnecessary. The conjunction may have density zero: that excludes a positive-proportion lower bound, but an averaging argument may still produce an unbounded sparse count.

Several natural prime-distribution inputs are insufficient. Isolated small gaps, isolated large gaps, average gap estimates, and the occurrence of any one fixed finite pattern do not suffice: both inequalities in (5.2) contain the complete infinite continuation. Nor does parity: after the first gap all g_n are even. Unboundedness and non-eventual-periodicity of the actual gaps, though now checked, do not imply the required small-tail recurrence. In particular, Section 5 shows that no argument can deduce eventual periodicity from rationality alone.

A finite truncation criterion. Both problems above are stated in terms of infinite tails, but a dominated truncation reduces the first of them to a finite quantity. For $L \geq 1$ put

$$S_{h,N,L} = \sum_{j=1}^L \frac{g_{N+h+j} - g_{N+j}}{2^j},$$

the truncation of the series in (5.1) after L terms. This is a finite sum of gap differences and can be computed; the point of the following proposition is to say how far from an integer it must be before the discarded tail is irrelevant.

Equivalently, introduce the integral dyadic block

$$D_{h,N,L} = \sum_{j=1}^L 2^{L-j} (g_{N+h+j} - g_{N+j}), \quad S_{h,N,L} = \frac{D_{h,N,L}}{2^L}.$$

Then

$$\text{dist}(S_{h,N,L}, \mathbb{Z}) = 2^{-L} \min\{D_{h,N,L} \bmod 2^L, 2^L - (D_{h,N,L} \bmod 2^L)\}.$$

Here $D_{h,N,L} \bmod 2^L$ denotes the least nonnegative residue, including when $D_{h,N,L} < 0$. Thus the finite criterion below is an exact modular small-arc problem: the residue of $D_{h,N,L}$ must avoid the two arcs of radius $2^L R_{h,N,L}(M)$ around 0 modulo 2^L . A one-block certificate would be a prime-gap theorem producing such an avoided arc on a logarithmic block.

Proposition 6.4 (finite truncation). *Let $M(n) \geq g_n$ for every n , assume that the series below converges, and put*

$$R_{h,N,L}(M) = \sum_{j>L} \frac{M(N+h+j) + M(N+j)}{2^j}.$$

Suppose that for every fixed $h \geq 1$ and every N_0 there exist $N \geq N_0$ and $L \geq 1$ with

$$\text{dist}(S_{h,N,L}, \mathbb{Z}) > R_{h,N,L}(M). \quad (5.3)$$

Then Problem 6.2 holds.

Proof. The part of $\sum_{j \geq 1} (g_{N+h+j} - g_{N+j})2^{-j}$ omitted from $S_{h,N,L}$ has absolute value at most $R_{h,N,L}(M)$, so under (5.3) the full sum lies at positive distance from every integer. $\square$

The last distance-to-integers inference is the elementary paper argument in the displayed proof: an error at most R cannot reach an integer when the approximation is farther than R from every integer. It is [not real integral of error bound](#), and the proposition itself is [cofinal escape of finite truncation](#). For instance, if $S_{h,N,L} = 3/8$ and $R_{h,N,L}(M) = 1/32$, the full sum lies within $1/32$ of $3/8$ and so at distance at least $11/32$ from every integer, which settles (5.1) at that N . The prime-gap tail bound, the convergence used above, and the existence of blocks satisfying (5.3) are not consequences of that inference. The proposition's full sum is real-valued, and the formal declaration is stated for the real tail throughout. Its conditional endpoint for the actual prime series is [irrational prime series of finite truncation](#).

Taking the classical bound $M(n) \ll n \log n$, a choice $L = \lceil A \log_2(N + h + 2) \rceil$ with any fixed $A > 1$ makes the right side a negative power of N up to logarithms. Thus (5.3) asks for a finite dyadic anti-concentration estimate on a logarithmic-length block, not control of an infinite tail and not eventual periodicity of the full gap sequence. For Problem 6.3, the same truncation must certify two adjacent full-tail values inside the open unit interval, together with the displayed gap mismatch. A finite prefix is useful only when its omitted tail is rigorously dominated.

What (5.3) requires is joint control of the finite block of weighted differences $(g_{N+h+1} - g_{N+1}, \dots, g_{N+h+L} - g_{N+L})$ modulo powers of two, along a block of logarithmic length. We do not know how to obtain such control and we make no progress on it here. The following results on prime gaps address a different shape of question. Zhang's bounded-gap theorem [15, Theorem 1, p. 1122] produces infinitely many bounded consecutive-prime gaps. Maynard proves substantially more than an individual-gap statement: [12, Theorem 1.1, p. 384] bounds $\liminf_n (p_{n+m} - p_n)$ for every fixed m , and hence gives bounded clusters of every fixed size; [12, Theorem 1.3, p. 385] gives the explicit unconditional bound $\liminf_n (p_{n+1} - p_n) \leq 600$. The large-gap theorem of Ford, Green, Konyagin, Maynard and Tao [5, Theorem 1] gives an effective lower bound for the largest single consecutive-prime gap below X . None of these results supplies the joint dyadic distribution of a logarithmic block of consecutive gap differences required by (5.3). The same introduction notes a separate sequel on chains of large gaps; the cited theorem itself supplies no residue-sensitive block estimate of the kind needed here.

The remaining estimate. The actual tails satisfy convergence, the prime-to-gap identity, and the recurrence used above. The unresolved input is the joint occurrence in (5.2) for every positive shift. A separate first-moment approach in the companion record uses

the growth $p_n \asymp n \log n$ [13, Chapter 6] to obtain

$$\sum_{X \leq N < 2X} T_{N+L} \ll X \log X \quad (L \leq X).$$

It controls the number of excessive omitted tails. A lower bound for the corresponding finite prime-gap configurations is still required. Sparse witness counts can suffice when they exceed that error budget. The independent finite continued-fraction enclosure gives $q \geq 2^{39997} > 10^{12040}$ for a rational representation of Π ; it makes no assertion about an irrationality exponent.

Statements and declarations

Artefact and data availability. The [source links](#) identify a fixed public snapshot. More recent live declarations are identified explicitly where they are used; a local declaration locator does not establish its presence at a historical public pin. The repository’s versioned source and verification records, rather than this manuscript’s navigation links, identify the checked object. The three finite-computation receipts and their generating programs are public in the [computation guide](#) at an immutable revision. That guide records dependencies, replay commands, source provenance, and the exact versus floating-point evidence boundary.

Lean 4 [10] checks each proof term against the fixed version of the mathlib library [11], and the sources linked here contain no proof placeholders and no project-defined axioms; Lean does not authorise the exposition, the citation choices, or the interpretation, for which the author remains responsible.

For Proposition 1.1, the pair identities, congruence buffer and abstract variable-alphabet filling are checked in `SparseRationalisationCore.lean`. Lean checks the proposition itself. [Arbitrary word sparse rationalisation](#) gives, for every $f \rightarrow \infty$, a support in $[K, \infty)$ of upper Banach density zero, an interval above A , and a correction realising each point of that interval; [arbitrary word sparse rational target](#) realises a rational value above A ; and [polylogarithmic word interval](#) gives the polylogarithmic clause, with block limits [growing block TV](#) and [growing block tests uniform](#). The explicit three-coordinate schedule in the construction is the ordinary proof, expanded in the companion record, subsection “Paired corrections and cumulative congruences”. The proposition has no Comparator entry.

Funding and competing interests. This work received no external funding. The author declares no competing interests.

Acknowledgements. The problem numbering and status follow the Erdős Problems catalogue maintained by Thomas Bloom [16].

A Guide to the formal sources

Each linked phrase opens its Lean declaration at the pinned source revision 99f4bf47422a. The note’s declarations live in the #251 modules, including the actual real-tail bridge. A local locator is not evidence that the historical public pin contains the file. The summation-by-parts declarations are prime-specific; most of

Section 4 is stated for arbitrary integer digits and arbitrary rational or real orbits, while Section 4.1 records the actual-gap specialisation.

References

- [1] P. Erdős, *Sur certaines séries à valeur irrationnelle*, Enseign. Math. (2) **4** (1958), 93–100, doi:10.5169/seals-34629.
- [2] P. Erdős and R. L. Graham, *Old and New Problems and Results in Combinatorial Number Theory*, Monogr. Enseign. Math. 28, Geneva, 1980.
- [3] P. Erdős and C. Pomerance, *On the largest prime factors of n and $n + 1$* , Aequationes Math. **17** (1978), 311–321, doi:10.1007/BF01818569.
- [4] P. Erdős, *On the irrationality of certain series: problems and results*, in A. Baker (ed.), *New Advances in Transcendence Theory*, Cambridge UP, 1988, pp. 102–109, doi:10.1017/CBO9780511897184.009.
- [5] K. Ford, B. Green, S. Konyagin, J. Maynard and T. Tao, *Long gaps between primes*, J. Amer. Math. Soc. **31** (2018), 65–105, doi:10.1090/jams/876; preprint arXiv:1412.5029v3.
- [6] J. A. Fridy, *Generalized bases for the real numbers*, Fibonacci Quart. **4** (1966), no. 3, 193–201.
- [7] V. Kovač and T. Tao, *On several irrationality problems for Ahmes series*, Acta Math. Hungar. **175** (2025), 572–608, arXiv:2406.17593v4. Statement and section numbers cited here follow arXiv version 4.
- [8] V. Kuperberg, *Sums of singular series with large sets and the tail of the distribution of primes*, arXiv:2210.09775v2, 15 June 2023.
- [9] J. Land, *A conditional proof of the irrationality of $\sum_{n \geq 1} p_n 2^{-n}$ under a uniform Hardy–Littlewood prime-tuples conjecture*, research draft, 5 September 2026, with an accompanying Lean formalisation of the conditional argument.
- [10] L. de Moura and S. Ullrich, *The Lean 4 theorem prover and programming language*, in A. Platzer and G. Sutcliffe (eds.), CADE 28, Lecture Notes in Comput. Sci. 12699, Springer, 2021, pp. 625–635, doi:10.1007/978-3-030-79876-5_37.
- [11] The mathlib Community, *The Lean mathematical library*, in CPP 2020, ACM, 2020, pp. 367–381, doi:10.1145/3372885.3373824. The article describes a December 2019 Lean 3-era snapshot; the repository lock owns the current revision.
- [12] J. Maynard, *Small gaps between primes*, Ann. of Math. (2) **181** (2015), 383–413, doi:10.4007/annals.2015.181.1.7.
- [13] H. L. Montgomery and R. C. Vaughan, *Multiplicative Number Theory I: Classical Theory*, Cambridge Stud. Adv. Math. 97, Cambridge UP, 2007, doi:10.1017/CBO9780511618314.
- [14] W. van Doorn and V. Kovač, *Lacunary sequences whose reciprocal sums represent all rational numbers in an interval*, Acta Arith. **223** (2026), 275–295, arXiv:2509.24971v3. Statement numbers cited here follow arXiv version 3.
- [15] Y. Zhang, *Bounded gaps between primes*, Ann. of Math. (2) **179** (2014), 1121–1174, doi:10.4007/annals.2014.179.3.7.
- [16] T. F. Bloom, *Erdős Problem #251*, erdosproblems.com/251, accessed 6 September 2026 (page displays “last edited 28 September 2025”).
- [17] *Erdős Problem #251 discussion thread*, erdosproblems.com/forum/thread/251, accessed 6 September 2026: comment of T. Tao, 7 October 2025, and comments of J. Land, 6 September 2026.

- [18] ChatGPT 5.4 Pro (orchestrated by Vjeko Kovač), *On the Erdős problem #251*, unpublished note, 2026, hosted by the Department of Mathematics, University of Zagreb, web.math.pmf.unizg.hr, accessed 6 September 2026; posted in the problem’s forum thread on 15 April 2026 [17].
- [19] S. Ringer, *Local gap statistics, telescoping, and normality: a local-pattern approach to Erdős problem 251*, preprint dated 11 September 2026, with a Lean formalisation; submitted as a partial proof claim at erdosproblems.com/251 on 13 September 2026.

Companion system context. The [claim and trust boundary](#), [cold-clone route to proof authority](#), and [public contribution protocol](#) are described in sibling papers. Those descriptions do not change the mathematical status of this note.