

Distinct running least common multiples

Affine tails and Erdős Problem #269

WILL COOK*

July 2026; revised 29 September 2026

ABSTRACT

We give a five-map proof that the reciprocals of the distinct running least common multiples of the 5-smooth integers have an irrational sum. The spacing of powers of 5 separates the tail images, so rationality would force a periodic jump sequence. We also derive a residue criterion for the sum counted with multiplicity, whose irrationality remains open here.

1 Two sums from running least common multiples

The 5-smooth integers are the positive integers whose prime factors belong to $\{2, 3, 5\}$. Their running least common multiples begin $1, 2, 6, 12, 60, 60, 120, 360, 360, \dots$. More generally, for a finite set of primes P , let $\mathcal{S}_P$ be the positive integers supported on P , including 1, and put

$$L(x) = \text{lcm}\{u \in \mathcal{S}_P : u \leq x\}, \quad H_P(x) = \prod_{p \in P} p^{\lfloor \log_p x \rfloor} \quad (x \geq 1).$$

Every integer in the prefix divides $H_P(x)$, and each maximal prime power occurs there. Hence $L(x) = H_P(x)$. The running value changes only at a positive prime power, where it is multiplied by that prime. Counting each distinct height once gives

$$\mathcal{D}_P = 1 + \sum_{\substack{t=p^n \\ p \in P, n \geq 1}} \frac{1}{H_P(t)}.$$

Positive powers of distinct primes never coincide, and geometric growth of the successive heights gives convergence. We use $\mathbb{N} = \{0, 1, 2, \dots\}$.

Theorem 1.1 (the distinct-height sum for $\{2, 3, 5\}$). *The number $\mathcal{D}_{\{2,3,5\}}$ is irrational.*

[Lean](#)

We group the jumps into intervals between successive powers of 2 and express their contributions by five affine maps. To recover the jumps from a tail value, we use the spacing of powers of 5 to find a bound on which the five images are disjoint. Under

**Authorship and AI use.* Will Cook built and directed the research infrastructure and reviewed the claims when he could. AI agents did most of the research and drafting. Cook did not independently verify every claim.

rationality the normalised tails take only finitely many values, so a repeated value would force periodicity of the later jumps, contradicting the irrational frequency of the powers of 3.

Erdős asserted irrationality of $\mathcal{D}_P$ for every finite P with $|P| \geq 2$ in a letter dated 1 January 1973, without printing a proof [3, p. 335]. Thus the theorem above is a special case of that earlier assertion. We claim no priority for its conclusion. The [companion's general argument](#) uses an isolated interchange of jumps on a torus. That argument is retained there as an ordinary proof awaiting formalisation. The proof below is independent of that argument.

The question in Erdős Problem #269 concerns the sum with multiplicity,

$$\mathcal{R}_P = \sum_{u \in \mathcal{S}_P} \frac{1}{L(u)},$$

as recorded in [1, p. 65] and [2, p. 106]. For example, $L(5) = L(6) = 60$ for $P = \{2, 3, 5\}$, giving $2/60$ in $\mathcal{R}_P$ and $1/60$ in $\mathcal{D}_P$. The theorem treats a related, distinct target. Our arguments leave the repeated sum unresolved for finite P with $|P| \geq 3$. For $P = \{p\}$ both sums equal $p/(p-1)$.

Section 2 gives the five-map proof. For two primes, Fan's factorisation [7] reduces both sums to a Hecke–Mahler value and yields transcendence (Section 3). We then derive the recurrence for the repeated three-prime sum and its residue criterion in Sections 4 and 5. Appendix A explains why the two-prime factorisation has no finite separated analogue for the three-prime kernel.

2 Recovering the prime-power jumps

We group the prime-power jumps into the right-closed intervals $(2^a, 2^{a+1}]$. Each contains its terminal power of 2, at most one power of 3 and at most one power of 5. We write τ_a for the sequence of interior prime labels, so that $\tau_a \in \{\emptyset, 3, 5, 35, 53\}$. Both orders occur: the block $(16, 32]$ contains $25 < 27$, whereas $(64, 128]$ contains $81 < 125$. We omit the terminal 2 from τ_a .

Proof of Theorem 1.1. To remove the denominators of an initial segment, put $P_a = H_{\{2,3,5\}}(2^a)$ and define

$$Y_a = P_a \sum_{\substack{t > 2^a \\ t=2^n, 3^n \text{ or } 5^n, n \geq 1}} \frac{1}{H_{\{2,3,5\}}(t)}. \quad (1)$$

Every subsequent jump multiplies the height by at least 2, and a jump by 3 eventually occurs. Comparison with $\sum_{j \geq 1} 2^{-j}$ therefore gives $0 < Y_a < 1$. Splitting off one block gives

$$Y_a = G_{\tau_a}(Y_{a+1}), \quad G_{\tau}(y) = \frac{\mu_{\tau} + y}{b_{\tau}}, \quad \begin{array}{c|ccccc} \tau & \emptyset & 3 & 5 & 35 & 53 \\ \hline b_{\tau} & 2 & 6 & 10 & 30 & 30 \\ \mu_{\tau} & 1 & 3 & 3 & 13 & 9 \end{array} \quad (2)$$

For example, a block of type 35 contributes $1/3 + 1/15 + 1/30 = 13/30$ to the normalised tail. The same calculation for type 53 gives $1/5 + 1/15 + 1/30 = 9/30$. Thus the order of the two interior jumps changes the constant term of the map.

We need a narrower interval than $(0, 1)$ to recover that order from a tail value. The table gives $Y_a \geq 3/10$. A power 5^f belongs to block $\lfloor f \log_2 5 \rfloor$; these block indices start at 2, and their successive differences are 2 or 3. Every three consecutive blocks therefore include a power of 5. At such a block, the bound $Y_{a+1} < 1$ gives $Y_a \leq G_{35}(1) = 7/15$, which also bounds $G_5(1)$ and $G_{53}(1)$. Before the next such block there are at most two maps, each equal to $G_\emptyset$ or G_3 . Since $G_3(y) \leq G_\emptyset(y) = (1+y)/2$ for $y \geq 0$, we obtain

$$Y_a \in I = \left[\frac{3}{10}, \frac{13}{15} \right], \quad G_\emptyset(G_\emptyset(7/15)) = \frac{13}{15}. \quad (3)$$

The spacing condition was needed to bound the arithmetic tails. The interval I is not invariant under arbitrary compositions of the five maps, since $G_\emptyset(13/15) = 14/15$.

On I the five images are disjoint. Their endpoints, listed in increasing order, are

τ	$G_\tau(I)$
53	$[31/100, 74/225]$
5	$[33/100, 29/75]$
35	$[133/300, 104/225]$
3	$[11/20, 29/45]$
$\emptyset$	$[13/20, 14/15]$

The consecutive gaps are $1/900, 17/300, 79/900, 1/180$. The two smallest gaps depend on $3(3/10) - 13/15 = 1/30 > 0$, so the preliminary upper bound 1 would not suffice. Since $Y_{a+1} \in I$, the value Y_a identifies τ_a uniquely. We can then calculate $Y_{a+1} = b_{\tau_a} Y_a - \mu_{\tau_a}$ and repeat the argument to recover every later block.

Suppose that $\mathcal{D}_{\{2,3,5\}} = N/K$, with integers N and $K \geq 1$. Every height at or before 2^a divides P_a . Consequently

$$KY_a = P_a N - K \left(P_a + \sum_{\substack{t \leq 2^a \\ t=2^n, 3^n \text{ or } 5^n, n \geq 1}} \frac{P_a}{H_{\{2,3,5\}}(t)} \right) \in \mathbb{Z}.$$

There are only finitely many integers in $[3K/10, 13K/15]$, so two tail values agree. Recovery of all subsequent blocks then makes (τ_a) eventually periodic.

To exclude this possibility, let $\alpha = \log 2 / \log 3$. The number of powers of 3 in block a is $\lfloor (a+1)\alpha \rfloor - \lfloor a\alpha \rfloor$, and is determined by τ_a . If the block word has eventual period $\ell \geq 1$, let c be the number of 3-powers in one period. For every sufficiently large fixed a and every $n \geq 1$ we then have

$$\lfloor (a+n\ell)\alpha \rfloor = \lfloor a\alpha \rfloor + nc.$$

Dividing by n and taking the limit gives $\ell\alpha = c$, contrary to unique factorisation, since it would imply $2^\ell = 3^c$. $\square$

The [companion's account of this proof](#) also compares one-jump and dyadic bounds. Its finite-state extension uses intervals depending on earlier blocks. The saved exact computations satisfy that criterion for every set of three primes at most 31 and for 292 of the 330 four-prime sets in that range. The general criterion and those computations are recorded as remarks awaiting formalisation. For $\{2, 3, 5, 7\}$ two admissible automaton words give the same affine map at every subsequent refinement, preventing that automaton from recovering its first block. This identity does not establish a collision between two words realised by the arithmetic sequence.

3 Two-prime transcendence

For two primes we can retain the multiplicities and still evaluate the sum through one Hecke–Mahler value. Fan gave the factorisation of the repeated sum and its transcendence consequence in his post of 26 June 2026 [7]. The following calculation also expresses the distinct-height sum in that value.

Theorem 3.1 (both two-prime sums). *Let $p < q$ be distinct primes. Put $\theta = \log p / \log q$ and $A = \sum_{n \geq 0} p^{-n} q^{-\lfloor n\theta \rfloor}$. Let $\mathcal{R}_{p,q}$ sum the reciprocal running LCM at every positive $\{p, q\}$ -smooth integer, and let $\mathcal{D}_{p,q}$ count each distinct running LCM once. Then* Lean[†]

$$\mathcal{D}_{p,q} = \frac{(q-p)A + p}{q-1}, \quad \mathcal{R}_{p,q} = \frac{(p+q-1)A - (p-1)A^2}{q-1}. \quad (4)$$

Both numbers are transcendental.

The Lean proof assumes the transcendence theorem of Bugeaud and Laurent; the two identities in (4) are proved in Lean without it.

Proof. Write $x = 1/p$, $y = 1/q$, $m_n = \lfloor n\theta \rfloor$ and $\delta_n = m_{n+1} - m_n$. Unique factorisation makes θ irrational, and $0 < \theta < 1$ gives $\delta_n \in \{0, 1\}$. The initial value and the positive powers of p contribute $A = \sum_{n \geq 0} x^n y^{m_n}$. There is one q -power strictly between p^n and p^{n+1} exactly when $\delta_n = 1$, so the positive powers of q contribute

$$B = \sum_{n \geq 0} \delta_n x^n y^{m_n+1}, \quad \mathcal{D}_{p,q} = A + B.$$

Since $0 < x, y < 1$, these series converge absolutely. The identity $y^{m_{n+1}} - y^{m_n} = \delta_n y^{m_n} (y - 1)$ and an index shift in A give $A - 1 - xA = x(y - 1)B/y$. Therefore

$$B = \frac{p - (p-1)A}{q-1}.$$

We compute the repeated sum from the height $p^{i+\lfloor j/\theta \rfloor} q^{j+m_i}$ at $p^i q^j$, obtaining

$$\mathcal{R}_{p,q} = A \sum_{j \geq 0} y^j x^{\lfloor j/\theta \rfloor} = A(1 + B).$$

For the last equality, each $j \geq 1$ corresponds to $n = \lfloor j/\theta \rfloor$ with $m_n = j - 1$ and $\delta_n = 1$. Substitution proves (4).

To apply the transcendence theorem, we express A in terms of the Hecke–Mahler series $F_\theta(x, y) = \sum_{n \geq 1} \sum_{k=1}^{m_n} x^n y^k$. Geometric summation gives

$$A = \frac{1}{1-x} - \frac{1-y}{y} F_\theta(x, y). \quad (5)$$

Bugeaud and Laurent’s theorem [5, Theorem 1.1] applies with intercept $\rho = 0$, $\beta = x$ and $\alpha = y$: the slope θ is irrational and lies in $(0, 1)$, x and y are nonzero algebraic numbers, $|x| < 1$ and $|xy^\theta| = p^{-2} < 1$. Thus $F_\theta(x, y)$ is transcendental; this case $\rho = 0$ is due to Loxton and van der Poorten [6, Theorem 8, p. 40]. Thus A is transcendental. The displayed affine and quadratic polynomials are nonconstant, so an algebraic value of either would force A to be algebraic. $\square$

The [companion’s two-prime calculation](#) records an ordinary extension to the monoid generated by coprime integers $1 < p < q$, such as 4, 9. That monoid need not contain all integers supported on the prime divisors of pq ; multiplicatively dependent generators such as 4, 8 require a different calculation.

For larger prime sets, the sums over pure powers $E_p = \sum_{\alpha \geq 0} L(p^\alpha)^{-1}$ satisfy $\mathcal{D}_P = \sum_{p \in P} E_p - (|P| - 1)$. The [companion’s prime-power subseries argument](#) treats their irrationality as a remark with an ordinary proof awaiting formalisation. The source record contains no separate priority search for this claim. Irrationality of the individual subseries would not imply irrationality of their sum. The repeated series also contains mixed products of primes, whose multiplicities change the tail estimates.

4 The repeated three-prime sum

We now count the smooth integers contributing each height. For three pairwise distinct primes p, q, r , write $H = H_{\{p, q, r\}}$ and $K(i, j, k) = H(p^i q^j r^k)^{-1}$. The next identities identify the heights and their multiplicities before we group the repeated series into intervals between powers of 2.

Proposition 4.1 (the running least common multiple). *Let p, q, r be pairwise distinct primes and $x \geq 1$. Then the running LCM equals the three-prime height: $L(x) = H(x)$.*

[Lean](#)
[Comparator](#)

Proof. We have $u \mid H(x)$ for every supported $u \leq x$. For the reverse divisibility, take the maximal powers of p, q, r below x : they belong to the prefix and are pairwise coprime, so their product divides its LCM. $\square$

For each prime, $x/p < p^{\lfloor \log_p x \rfloor} \leq x$. Multiplying gives

$$x^3 / (pqr) < H(x) \leq x^3. \quad (6)$$

Proposition 4.2 (cells and jumps). *The running LCM is constant when the three integer logarithms are constant. A jump in exactly one logarithm multiplies it by the corresponding prime. The first n positive powers of each prime, together with 1, form $3n + 1$ distinct points.*

[Lean](#)
[Comparator](#)

Proof. The height is a product of the three maximal pure powers, which proves the constancy and jump assertions. Unique factorisation separates the $3n$ positive pure powers from one another and from 1. $\square$

Proposition 4.3 (grouping terms with the same height). *For a finite exponent box $\mathcal{B}$, set $F(H) = \{(i, j, k) \in \mathcal{B} : H(p^i q^j r^k) = H\}$. Then*

Lean
Comparator

$$\sum_{(i,j,k) \in \mathcal{B}} K(i, j, k) = \sum_H \frac{\#F(H)}{H}. \quad (7)$$

Proof. Partition the finite exponent box by the value of the height. The terms in the part $F(H)$ all equal $1/H$. $\square$

Henceforth $P = \{2, 3, 5\}$ and $S = \mathcal{R}_P$. To clear the finite prefix of this series, we use half-open intervals $[2^a, 2^{a+1})$ and normalise by half the height at their left endpoint. These endpoint conventions differ from those in Section 2. Put

$$s_a = \sum_{\substack{x \in \mathcal{S}_P \\ 2^a \leq x < 2^{a+1}}} \frac{1}{H(x)}, \quad T_a = \sum_{j \geq 0} s_{a+j}, \quad h_a = \frac{H(2^a)}{2}, \quad X_a = h_a T_a.$$

The factor $1/2$ makes h_a a common multiple of the heights strictly before 2^a . Indeed, each such height has exponent of 2 at most $a - 1$. Here $h_0 = 1/2$, and h_a is an integer for $a \geq 1$. Put

$$b_a = \frac{H(2^{a+1})}{H(2^a)}, \quad m_a = \sum_{\substack{x \in \mathcal{S}_P \\ 2^a \leq x < 2^{a+1}}} \frac{H(2^{a+1})}{2H(x)}. \quad (8)$$

Lemma 4.4 (integer coefficients and four possible bases). *For every $a \geq 0$, m_a is a positive integer and $b_a \in \{2, 6, 10, 30\}$. The word “numerator” does not impose the positional-digit restriction $m_a < b_a$; that restriction need not hold.*

Lean
Comparator

Proof. For $x < 2^{a+1}$ we have $2H(x) \mid H(2^{a+1})$, so every summand in m_a is integral. The interval contains 2^a , giving positivity. Between successive powers of 2 there is at most one power of 3 and at most one power of 5. Along with the final factor 2, these give $b_a \in \{2, 6, 10, 30\}$. $\square$

For example, $[2, 4)$ contains 2, 3 and gives $(b_1, m_1) = (6, 4)$. At $[16, 32)$ we obtain $(b_4, m_4) = (30, 65)$, so these coefficients cannot be treated as positional digits.

Proposition 4.5 (the tail recurrence and a quadratic bound). *The series defining S, T_a converge. For every $a \geq 0$,*

Lean
Comparator

$$X_{a+1} = b_a X_a - m_a, \quad 0 < X_a \leq \frac{8640}{343}(a+1)^2 < 90(a+1)^2.$$

For every integer $B \geq 1$, either some BX_a is integral and all later states are integral, or $\text{dist}(BX_a, \mathbb{Z}) \geq 1/31$ at arbitrarily large indices.

Proof. We count at most $(a+1)^2$ smooth integers in $[2^a, 2^{a+1})$: each pair of exponents of 3, 5, both at most a , allows at most one exponent of 2. By (6), $s_a \leq 30(a+1)^2/8^a$. As $h_a \leq 8^a/2$ and $a+j+1 \leq (a+1)(j+1)$, we have

$$X_a \leq 15(a+1)^2 \sum_{j \geq 0} \frac{(j+1)^2}{8^j} = \frac{8640}{343}(a+1)^2.$$

This also proves convergence. Splitting off s_a gives the recurrence, since $h_{a+1} = b_a h_a$ and $m_a = h_{a+1} s_a$.

For the final assertion, suppose that every sufficiently late distance is strictly less than $1/31$, and write $BX_a = z_a + e_a$, where z_a is integral and $|e_a| < 1/31$. The recurrence makes $e_{a+1} - b_a e_a$ an integer of absolute value less than $(1 + b_a)/31 \leq 1$. Thus $e_{a+1} = b_a e_a$. Since $b_a \geq 2$, boundedness forces every such error to vanish. Once integral, the states remain integral by the recurrence. $\square$

We can also write the repeated series as a Cantor series: $S/2 = \sum_{a \geq 0} m_a / (b_0 \cdots b_a)$. The criteria in [9, Theorem 2.1] and [10, Theorem 3.1] require a small-numerator hypothesis $m_a / (b_{a-1} b_a) \rightarrow 0$. Here the b_a are bounded and m_a grows quadratically. The [companion's tail estimates](#) prove this lower bound as well as sharper upper bounds. In particular, $X_a > m_a/30$ is unbounded, so the finite-value argument in Section 2 no longer applies to these normalised tails.

Theorem 4.6 (rationality gives positive integer tails). *If $S = A/D$ in lowest terms, where $D = 2^u 3^v 5^w B$ and $\gcd(B, 30) = 1$, then for every $a \geq a_0 = u + 1 + 2v + 3w$,*

Lean
Comparator

$$d_a = BX_a \in \mathbb{Z}_{>0}, \quad d_{a+1} = b_a d_a - B m_a, \quad d_a \leq 90B(a+1)^2.$$

Proof. For $a \geq 1$, clearing the finite prefix gives

$$X_a = h_a S - \sum_{\substack{x \in \mathcal{S}_p \\ x < 2^a}} \frac{h_a}{H(x)}, \quad \sum_{\substack{x \in \mathcal{S}_p \\ x < 2^a}} \frac{h_a}{H(x)} \in \mathbb{Z}. \quad (9)$$

If $a \geq u + 1 + 2v + 3w$, the exponent $a - 1$ of 2 in h_a is at least u , and $2^a \geq 3^v, 5^w$. Therefore $2^u 3^v 5^w \mid h_a$, and BX_a differs from $h_a A / (2^u 3^v 5^w)$ by an integer. Positivity, recurrence and the bound follow from Proposition 4.5. $\square$

Corollary 4.7 (the first index at which the denominator clears). *Under the same lowest-terms hypothesis, put $M = 2^u 3^v 5^w$ and let den denote the positive reduced denominator. For $a \geq 1$,*

Lean
Comparator

$$\text{den}(BX_a) = \frac{M}{\gcd(M, h_a)}.$$

Consequently BX_a is integral exactly when $2^a \geq \max(2^{u+1}, 3^v, 5^w)$. The first such a can be found by integer comparisons, without logarithmic rounding.

Proof. Equation (9) shows that BX_a and $h_a A / M$ have the same reduced denominator. Since $\gcd(A, M) = 1$, this denominator is $M / \gcd(M, h_a)$. Divisibility by each of $2^u, 3^v, 5^w$ gives the three stated inequalities. $\square$

For a hypothetical reduced denominator $2^3 3^2 5 \cdot 7$, the first integral $7X_a$ would occur at $a = 4$. The sufficient onset in the theorem is 11. The [companion's denominator calculation](#) also determines the reduced denominator of X_a itself.

5 A residue criterion

Iterating the recurrence gives $X_{\ell+2} = b_{\ell+1}b_\ell X_\ell - (b_{\ell+1}m_\ell + m_{\ell+1})$. For a window of h steps starting at ℓ , define

$$W_{\ell,0} = 1, \quad F_{\ell,0} = 0, \quad W_{\ell,h+1} = b_{\ell+h}W_{\ell,h}, \quad F_{\ell,h+1} = b_{\ell+h}F_{\ell,h} + m_{\ell+h}.$$

Induction yields

$$X_{\ell+h} = W_{\ell,h}X_\ell - F_{\ell,h}. \quad (10)$$

Under rationality, this determines the residue of a positive integer tail modulo $W_{\ell,h}$. We compare that residue with its quadratic bound. For integers $W \geq 1, t$, let $\text{lpr}_W(t) = 1 + ((t-1) \bmod W)$, taking the remainder in $\{0, \dots, W-1\}$. Thus $\text{lpr}_W(0) = W$. Throughout this section set

$$K(B, a) = 90B(a+1)^2. \quad (11)$$

Lemma 5.1 (least positive residues). *If d is a positive integer with $d \leq K$ and $d \equiv -BF \pmod{W}$, where $W \geq 1$, then $\text{lpr}_W(-BF) \leq K$.*

Lean
Comparator

Proof. The least positive representative is at most every positive representative of its class, and in particular at most d . $\square$

For $\ell = 1, h = 6$, direct iteration gives $W_{1,6} = 648000$ and $F_{1,6} = 524431$. Hence $\text{lpr}_{W_{1,6}}(-F_{1,6}) = 123569 > K(1,7) = 5760$. This excludes an integral X_1 . A rational denominator might clear only later, so a criterion for irrationality must allow every late start and every denominator part coprime to 30.

Theorem 5.2 (a residue criterion for irrationality). *The number S is irrational if and only if*

Lean
Comparator

$$\begin{aligned} &\text{for every } B \geq 1 \text{ with } \gcd(B, 30) = 1 \text{ and every } a_0 \geq 1, \\ &\text{there are } \ell \geq a_0, h \geq 1 \text{ such that } \text{lpr}_{W_{\ell,h}}(-BF_{\ell,h}) > K(B, \ell + h). \end{aligned} \quad (12)$$

Proof. Suppose that (12) holds and S is rational. By Theorem 4.6, $d_a = BX_a$ is a positive integer for every sufficiently large a . Choose a window beyond that onset. Equation (10) gives $d_{\ell+h} \equiv -BF_{\ell,h} \pmod{W_{\ell,h}}$, contrary to Lemma 5.1 and (12).

Conversely, suppose that S is irrational and fix $B, \ell \geq 1$. By (9), BX_ℓ is nonintegral. Write $\delta = \lceil BX_\ell \rceil - BX_\ell \in (0, 1)$. Since $W_{\ell,h} \geq 2^h$ and $X_{\ell+h} = O((\ell+h+1)^2)$, $\delta + BX_{\ell+h}/W_{\ell,h}$ lies in $(0, 1)$ for sufficiently large h . The window identity then shows that

$$\text{lpr}_{W_{\ell,h}}(-BF_{\ell,h}) = \lceil BX_\ell \rceil W_{\ell,h} - BF_{\ell,h} = \delta W_{\ell,h} + BX_{\ell+h}.$$

The first term eventually exceeds $K(B, \ell + h)$. This proves the required inequality from every fixed start. $\square$

The upper-bound property of K is essential: choosing the zero function would make every residue inequality automatic. The [companion's residue analysis](#) treats smaller bounds and identifies when more general bounds give the same criterion. Finite searches cover only finitely many multipliers and starts.

The repeated three-prime question can therefore be written in three equivalent forms.

Problem 5.3 (the repeated three-prime target). Prove that $S = \mathcal{R}_{\{2,3,5\}}$ is irrational.

Problem 5.4 (no reduced tail is an integer). Prove that for every $a \geq 1$ and every integer $B \geq 1$ coprime to 30,

$$BX_a \notin \mathbb{Z}. \quad (13)$$

Problem 5.5 (residue inequalities after every starting index). Prove (12).

An integral BX_a makes S rational by (9); Theorem 4.6 proves the converse implication. Theorem 5.2 gives the third form. All three remain unproved for S . The [companion's discussion of the repeated sum](#) records the finite calculations and the hypotheses missing from proposed finite-difference and function-theoretic arguments.

A The three-prime kernel

We ask whether the two-prime factorisation can be replaced, for three primes, by a finite sum of products separating one exponent from the other two. The kernel $K(i, j, k) = H(p^i q^j r^k)^{-1}$ rules this out. Diagonal rescaling reduces the question to a matrix with entries 1 and $1/r$; separate choices of row and column indices then give nonzero minors of every order.

Theorem A.1 (no finite separation of the kernel). *Let p, q, r be primes with $p \neq q$, $p \neq r$ and $q \neq r$. For every $n \geq 0$ there are injective maps $I, J : \{0, \dots, n-1\} \rightarrow \mathbb{N}$ such that, for every $k \geq 0$,*

[Lean](#)
[Comparator](#)

$$\det(K(I(a), J(b), k))_{0 \leq a, b < n} \neq 0.$$

Consequently, for no finite d do there exist rational-valued functions $f_\ell(i)$ and $G_\ell(j, k)$, $0 \leq \ell < d$, satisfying

$$K(i, j, k) = \sum_{\ell < d} f_\ell(i) G_\ell(j, k) \quad \text{for all } i, j, k.$$

Proof. For fixed k , we rescale rows and columns to obtain a matrix independent of k . Put $c = r^{-1}$, $x_i = \{i \log_r p\}$ and $y_j = \{j \log_r q\}$, where braces denote fractional parts. Splitting the floor exponents gives

$$K(i, j, k) = U_i(k)^{-1} C_{ij} V_j(k)^{-1}, \quad C_{ij} = c^{\mathbf{1}_{\{x_i + y_j \geq 1\}}},$$

where

$$U_i(k) = p^i q^{\lfloor \log_q(p^i r^k) \rfloor} r^{k + \lfloor \log_r p^i \rfloor},$$

$$V_j(k) = q^j p^{\lfloor \log_p(q^j r^k) \rfloor} r^{\lfloor \log_r q^j \rfloor}.$$

These positive factors preserve nonvanishing of minors. The remaining entry records whether $x_i + y_j \geq 1$. Distinct primes make $\log_r p$ and $\log_r q$ irrational, so each fractional-part orbit is dense in $[0, 1]$. We choose row and column indices independently, using only these two one-dimensional density statements. For $n \geq 1$, choose distinct rows with $0 < x_{I(0)} < \dots < x_{I(n-1)} < 1$. We choose each column so that its entries change from 1 to c at a different selected row. Writing $s_b = 1 - y_{J(b)}$, density of (y_j) lets us choose

$$s_0 \in (0, x_{I(0)}), \quad s_b \in (x_{I(b-1)}, x_{I(b)}) \quad (1 \leq b < n).$$

The intervals for the s_b are disjoint, so the column indices are distinct. The strict inequalities also avoid the threshold itself, and give $x_{I(a)} + y_{J(b)} \geq 1$ exactly when $b \leq a$. Thus multiplying row a by $U_{I(a)}(k)$ and column b by $V_{J(b)}(k)$ reduces the selected kernel minor to

$$T_n(c) = \begin{pmatrix} c & 1 & \cdots & 1 \\ c & c & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ c & c & \cdots & c \end{pmatrix}, \quad \det T_n(c) = c(c-1)^{n-1}.$$

Indeed, subtracting each preceding row from the next, working upwards from the last row, leaves diagonal entries $c, c-1, \dots, c-1$. The original determinant is therefore

$$c(c-1)^{n-1} \prod_{a \leq n} U_{I(a)}(k)^{-1} \prod_{b \leq n} V_{J(b)}(k)^{-1} \neq 0.$$

The indices were chosen from C , independently of k . For $n = 0$ the empty determinant equals 1.

For the final assertion, suppose a separation with d summands existed and fix any k . On the rows $I(0), \dots, I(d)$ and columns $J(0), \dots, J(d)$ its matrix would be the product of the $(d+1) \times d$ matrix $(f_\ell(I(a)))_{a,\ell}$ and the $d \times (d+1)$ matrix $(G_\ell(J(b), k))_{\ell,b}$. Its rank would be at most d , so its determinant would vanish, contrary to the minor just constructed. $\square$

No continuity or boundedness is assumed for the separated factors.

Corollary A.2 (the same minors modulo integers coprime to 30). *For $(p, q, r) = (2, 3, 5)$ and every $n \geq 1$, there are injective maps $I, J : \{0, \dots, n-1\} \rightarrow \mathbb{N}$, chosen independently of B and k , such that for every $B \geq 2$ coprime to 30 and every $k \geq 0$, the selected $n \times n$ kernel matrix has unit determinant over $\mathbb{Z}/B\mathbb{Z}$, with each reciprocal prime power interpreted by its modular inverse.*

[Lean](#)
[Comparator](#)

Proof. Choose the maps from Theorem A.1. Every row and column factor is a unit modulo B . The normalised determinant is $5^{-1}(-4/5)^{n-1}$, also a unit. $\square$

The modulus may be composite, for example 49 or 77. Coprimality with 30 makes both the reciprocal entries and the determinant units: the latter introduces only a power of 4 in its numerator.

Example A.3. For $(p, q, r) = (2, 3, 5)$, the leading two-by-two determinant is $-1/15$: its four entries are $1, 1/6, 1/2, 1/60$, so it equals $1/60 - 1/12$.

Leading minors. The leading 4×4 block at $\{2, 3, 5\}$ is singular:

$$K(3, j, 0) = \frac{1}{120} K(0, j, 0) \quad (0 \leq j < 4).$$

These equalities follow by evaluating the height at 3^j and $8 \cdot 3^j$. But at $j = 4$, $K(3, 4, 0) - K(0, 4, 0)/120 = -1/19440000$. Thus the leading minors do not supply the arbitrary-order theorem.

Proposition A.4 (rank of a matrix of threshold columns). *Let $m \geq 1$ and let c lie in a field with $c \neq 0, 1$. For $0 \leq k \leq m$ write v_k for the length- m column with a 1 in each of the first k coordinates and c thereafter. A matrix whose distinct columns are v_k for k in a nonempty set E has rank $|E| - \mathbf{1}_{\{0, m\} \subseteq E}$.*

Lean
Comparator

The correction accounts for the two constant columns: $v_0 = cv_m$. There is no other dependence among distinct threshold columns.

Proof. List $E = \{k_1 < \dots < k_t\}$. The $t - 1$ differences $v_{k_{j+1}} - v_{k_j} = (1 - c)\mathbf{1}_{\{k_j, \dots, k_{j+1}-1\}}$ have disjoint nonempty supports, hence are linearly independent. If $k_1 > 0$ or $k_t < m$, their union misses a coordinate where v_{k_1} is nonzero, so the rank is t . If $k_1 = 0$ and $k_t = m$, then $v_0 = c\mathbf{1}$ lies in the span of the differences (their sum is $(1 - c)\mathbf{1}$), so the rank is $t - 1$. $\square$

The restrictions $c \neq 0, 1$ exclude a zero column or identical columns; they hold for $c = 1/r$ over $\mathbb{Q}$. In one fixed layer k , ordering the sampled row phases puts every column in the stated form. The formula therefore determines the rank of each nonempty rectangular sample. For fixed row and column indices this rank is independent of k . The [companion's kernel analysis](#) gives the exact rational comparisons.

Approximation. Truncating the original kernel to $i < N$ gives a sum of N separated terms. The bound $H(x) > x^3/(pqr)$ makes the truncation error tend to zero both uniformly and in $\ell^1(\mathbb{N}^3)$. The rescaled matrix in the rank proof has uniform distance $(1 - 1/r)/2$ from the matrices of finite rank. Rescaling preserves exact rank but removes the decay responsible for the first approximation. The [companion's approximation theorem](#) proves both assertions for their stated norms. Neither the rank theorem nor the approximation bound implies irrationality of the scalar sum.

B Proof sources and further comparisons

The margin marks link to the recorded Lean declarations and, where present, Comparator comparisons. For the five-map proof the declarations are in [the distinct-height development](#) and [its block-series criterion](#); no comparison is recorded. The [evidence record](#) gives the versions. Neither checker was rerun for this revision, and the public links may precede this manuscript's local revision. The named input for two-prime transcendence stays beside that theorem.

The [distinct-height programs](#) calculate the interval endpoints and finite-state examples using exact arithmetic; the proof in Section 2 derives its constants directly. The [companion's source inventory](#) also covers the ordinary general and subseries arguments. Its historical note reports a second AI agent's check on 26 September 2026 and the incorporation of its corrections. No human mathematical review is reported.

The classical criteria of Erdős–Straus [9], Hančl–Tijdeman [10] and Diananda–Oppenheim [8] are compared with the bounded-base series in the companion. It also compares the bounded-ratio argument with Erdős–Taylor [11, Theorem 1, p. 600] and Fan [12, Lemma 3.1, p. 7], and the carry restrictions with the polynomial Cantor-series results of Hančl–Tijdeman [13, Theorems 2.2, 3.1 and 4.2]. Koutsoukou–Argyrazi and

Li’s Isabelle development [19] formalises classical irrationality criteria. It does not treat the present repeated series.

For transcendence, the relevant comparisons are the finite-difference conditions of Luca–Ouaknine–Worrell [14], the echoing conditions in [15], the fixed-base complexity theorem of Adamczewski–Bugeaud [20], and multivariate Mahler theory [21]. Quadratic growth of m_a supplies none of the missing cancellation or recurrence hypotheses. The companion exhibits nonzero boundary terms in a proposed third-difference calculation and distinguishes variable-base digits from fixed-base expansions. Work on fixed-prime semigroups [16, 17] and other Ahmes-series problems [18] provides context; no estimate from those papers enters our dyadic counting argument.

Statements and declarations

Authorship and availability. AI agents carried out most of the research and drafting. The companion contains the longer proofs and exact computations, together with the hypotheses and limitations of the unsuccessful approaches.

Funding and competing interests. This work received no external funding. The author declares no competing interests.

Acknowledgements. The problem number and its status as of 28 July 2026 are taken from Thomas Bloom’s Erdős Problems catalogue [4]. We thank Wouter van Doorn for advice on exposition, particularly on unexplained terminology, unnecessary notation and restrictive hypotheses. His advice concerned a note on Problem #243; he has not reviewed this paper’s mathematics.

References

- [1] Paul Erdős and Ronald L. Graham, *Old and New Problems and Results in Combinatorial Number Theory*, Monographies de L’Enseignement Mathématique **28**, L’Enseignement Mathématique (1980), [source](#).
- [2] Paul Erdős, *On the irrationality of certain series: problems and results*, in *New Advances in Transcendence Theory*, Cambridge University Press (1988), 102–109, [doi:10.1017/CB09780511897184.009](#).
- [3] Paul Erdős, *Letter to the Editor*, *Fibonacci Quarterly* **12**, no. 4 (1974), 335, [source](#).
- [4] Thomas F. Bloom, *Erdős Problem #269* (2026), [source](#). Accessed 28 July 2026.
- [5] Yann Bugeaud and Michel Laurent, *Transcendence and continued fraction expansion of values of Hecke–Mahler series*, *Acta Arithmetica* **209** (2023), 59–90, [doi:10.4064/aa220323-18-1](#); [arXiv:2203.12901](#).
- [6] John H. Loxton and Alfred J. van der Poorten, *Arithmetic properties of certain functions in several variables III*, *Bulletin of the Australian Mathematical Society* **16** (1977), 15–47, [doi:10.1017/S0004972700022978](#).

- [7] Steve Fan, *Comment on Erdős Problem #269, thread 269, post 7218* (2026), [source](#). Public forum post, 26 June 2026, thread 269, post 7218.
- [8] P. H. Diananda and A. Oppenheim, *Criteria for irrationality of certain classes of numbers II*, *American Mathematical Monthly* **62**, no. 4 (1955), 222–225. Cited in the form stated by Serbenyuk, arXiv:[1706.03124](#), Theorem 1.
- [9] Paul Erdős and Ernst G. Straus, *On the irrationality of certain series*, *Pacific Journal of Mathematics* **55**, no. 1 (1974), 85–92, doi:[10.2140/pjm.1974.55.85](#).
- [10] Jaroslav Hančl and Robert Tijdeman, *On the irrationality of Cantor and Ahmes series*, *Publicationes Mathematicae Debrecen* **65**, no. 3–4 (2004), 371–380, doi:[10.5486/PMD.2004.3254](#).
- [11] Paul Erdős and S. James Taylor, *On the set of points of convergence of a lacunary trigonometric series and the equidistribution properties of related sequences*, *Proceedings of the London Mathematical Society* **s3-7**, no. 1 (1957), 598–615, doi:[10.1112/plms/s3-7.1.598](#).
- [12] Steve Fan, *Strongly complete sets and a conjecture of Erdős* (2026), [source](#); arXiv:[2607.14071](#). The cited Lemma 3.1 is in arXiv v1, 15 July 2026.
- [13] Jaroslav Hančl and Robert Tijdeman, *On the irrationality of polynomial Cantor series*, *Acta Arithmetica* **133**, no. 1 (2008), 37–52, doi:[10.4064/aa133-1-3](#).
- [14] Florian Luca, Joël Ouaknine and James Worrell, *Transcendence of Hecke–Mahler Series*, *Bulletin of the London Mathematical Society* **57**, no. 5 (2025), 1360–1368, doi:[10.1112/blms.70033](#); arXiv:[2412.07908](#). Numbered references use the published article.
- [15] Pavol Kebis, Florian Luca, Joël Ouaknine, Andrew Scoones and James Worrell, *On Transcendence of Numbers Related to Sturmian and Arnoux–Rauzy Words*, in *51st International Colloquium on Automata, Languages, and Programming (ICALP 2024)*, *Leibniz International Proceedings in Informatics* **297**, Schloss Dagstuhl – Leibniz-Zentrum für Informatik (2024), 144:1–144:15, doi:[10.4230/LIPIcs.ICALP.2024.144](#).
- [16] Robert Tijdeman and H. G. Meijer, *On integers generated by a finite number of fixed primes*, *Compositio Mathematica* **29**, no. 3 (1974), 273–286, [source](#).
- [17] Alessandro Languasco, Florian Luca, Pieter Moree and Alain Togbé, *Sequences of integers generated by two fixed primes*, *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg* **95** (2025), 123–148, doi:[10.1007/s12188-025-00293-9](#); arXiv:[2309.12806](#).
- [18] Vjekoslav Kovač and Terence Tao, *On several irrationality problems for Ahmes series*, *Acta Mathematica Hungarica* **175** (2025), 572–608, doi:[10.1007/s10474-025-01528-0](#); arXiv:[2406.17593](#).

- [19] Angeliki Koutsoukou-Argyraki and Wenda Li, *Irrationality Criteria for Series by Erdős and Straus*, Archive of Formal Proofs (2020), [source](#). Entry dated 12 May 2020; proof-document version consulted: 6 February 2026.
- [20] Boris Adamczewski and Yann Bugeaud, *On the complexity of algebraic numbers I. Expansions in integer bases*, Annals of Mathematics **165**, no. 2 (2007), 547–565, [doi:10.4007/annals.2007.165.547](#).
- [21] Boris Adamczewski and Colin Faverjon, *Mahler’s method in several variables and finite automata*, Annals of Mathematics **204**, no. 2 (2026), 455–533, [doi:10.4007/annals.2026.204.2.1](#). Online 13 September 2026; locators here refer to the 68-page author manuscript.