

# Prime-Gap Dyadic Series: Perturbations, Exact Criteria and Certificates

*Proofs, counterexamples and computations for Erdős Problem #251*

WILL COOK\*

September 2026

---

## ABSTRACT

We give a sparse perturbation construction for convergent dyadic series with nonnegative integer coefficients, together with exact criteria for the prime-gap series. Varying adjacent corrections with a fixed sum gives an interval of dyadic sums while preserving eventual congruences. One sparse set contains all correction supports, and the congruence cutoffs are independent of the value attained. Applied to prime gaps with eventual correction bound  $\log(n+3)$ , the construction gives vanishing total variation error for blocks of length  $o(\log \log X)$  sampled on  $[X, 2X)$  and preserves the cumulative asymptotic  $n \log n$ , without ensuring primality. The actual tails satisfy a recurrence whose integral shifts characterise rationality. We give counterexamples to weaker coefficient conditions and integer certificates excluding rational denominators below  $2^{40062} > 10^{12059}$ . The irrationality question remains unresolved.

## 1 Introduction

The primes  $p_0 = 2, p_1 = 3, \dots$  define the dyadic series

$$\Pi = \sum_{n \geq 0} \frac{p_n}{2^{n+1}} = 2/2 + 3/4 + 5/8 + 7/16 + \dots$$

Its irrationality is Erdős Problem #251 [22], recorded in [1, p. 94], [2, p. 62] and [4, p. 103]. Writing  $g_n = p_{n+1} - p_n$ , summation by parts gives  $\Pi = 2 + \sum_n g_n 2^{-(n+1)}$ . We use this identity to compare what can be deduced from properties of the coefficients with what would be needed from the actual prime-gap tails.

Our main construction is Theorem 2.1. For any convergent dyadic series with nonnegative integer coefficients, we choose one sparse set containing all correction supports. By varying the corrections on this set, we attain every value in an interval. Given any function tending to infinity, we can choose corrections bounded by it for all sufficiently large indices. Every fixed modulus eventually divides both their terms and their partial sums, with cutoffs fixed before the target. The construction varies adjacent entries while

---

\**Authorship and AI use.* Will Cook built and directed the research infrastructure and reviewed the claims when he could. AI agents did most of the research and drafting. Cook did not independently verify every claim.

keeping their ordinary sum fixed. This separates the choice of weighted sum from the subsequent congruence calculations.

With a polylogarithmic bound on the corrections, the original and altered empirical block distributions have total variation distance tending to zero for lengths  $o(\log \log X)$  sampled in  $[X, 2X)$ . Applied to prime gaps, the theorem gives rational sums and cumulative positions asymptotic to  $n \log n$ . These positions are comparison integers and are not shown to be prime. These congruences and statistics are therefore compatible with a rational sum. The irrationality of the actual prime series remains unresolved.

The proofs about the actual primes begin in Section 4 and are independent of the sparse construction. Finite summation by parts identifies the endpoint that must vanish. The elementary prime bound in Appendix A gives this limit, and a boundary condition distinguishes the complete tails from other recurrence solutions. Their integral differences then characterise rationality. We use finite enclosures of these differences for the tests in Section 6, and enclosures of the series itself for the denominator bounds in Section 7. Section 8 records the counterexamples and what each coefficient condition fails to control. The appendices give the integer calculations, further equivalent criteria and source concordance.

For reference, the first primes and gaps are

| $i$   | 0 | 1 | 2 | 3 | 4  | 5  | 6  | 7  | 8  | 9  |
|-------|---|---|---|---|----|----|----|----|----|----|
| $p_i$ | 2 | 3 | 5 | 7 | 11 | 13 | 17 | 19 | 23 | 29 |
| $g_i$ | 1 | 2 | 2 | 4 | 2  | 4  | 2  | 4  | 6  | 2  |

Thus  $g_0 = 1$  and all later gaps are even. This parity restriction will be used to characterise two consecutive tail differences of absolute value less than one. The alternative normalisation with denominator  $2^i$  satisfies

$$\sum_{i=0}^{n-1} \frac{p_i}{2^i} = 2 \sum_{i=0}^{n-1} \frac{p_i}{2^{i+1}},$$

so it has the same rationality status. The [formal factor-of-two identity](#) uses these same finite sums.

**Notation.** We write  $\mathbb{N} = \{0, 1, \dots\}$ , use  $\varphi$  for Euler's totient function, and let  $\text{den } q$  be the positive denominator of  $q \in \mathbb{Q}$  in lowest terms. A real number is *integral* if it belongs to  $\mathbb{Z}$ , and  $\text{dist}(x, \mathbb{Z})$  is its distance to the nearest integer. A sequence is eventually periodic with period  $h \geq 1$  if  $a_{n+h} = a_n$  for every sufficiently large  $n$ . A set of indices has density zero if its counting function is  $o(x)$ , and a property holds for almost all indices if its exceptional set has density zero. A property may hold at arbitrarily large indices even when those indices form a set of density zero. An empty sum is zero.

## 2 Sparse congruence-preserving perturbations

At indices  $n, n+1$ , the pairs  $(0, 6), (2, 4), (4, 2), (6, 0)$  all have ordinary sum 6, while their dyadic contributions are 6, 8, 10, 12 divided by  $2^{n+2}$ . If the cumulative correction before the pair is odd, we first add 1 at  $n-1$ . Choosing the pair then leaves the cumulative

residue after the triple unchanged. We repeat this operation with nested moduli eventually divisible by every fixed integer.

Finite changes alone preserve rationality, so we use infinitely many triples. Making their separations tend to infinity gives upper Banach density zero, but also makes each weighted contribution smaller. We compensate by increasing the range of choices inside a pair. The estimate (5) shows that all later ranges together cover the gap between adjacent choices at the current stage. The set containing the supports and the interval of sums are chosen before the value to be attained.

**Density and block conventions.** All intervals of indices contain integers, and log is natural unless a base is displayed. Upper Banach density zero means

$$\lim_{H \rightarrow \infty} \sup_{u \in \mathbb{N}} \frac{|S \cap [u, u + H]|}{H} = 0.$$

For integers  $X \geq 1$  and  $m \geq 1$ , let  $\mu_{a,X,m}$  be the empirical probability measure obtained by choosing an integer  $n \in [X, 2X)$  uniformly and observing  $(a_n, \dots, a_{n+m-1})$ . Equal blocks are counted with multiplicity. We use  $d_{TV}(\mu, \nu) = \sup_B |\mu(B) - \nu(B)|$ . For the same function of the block, bounded in absolute value by  $B_0$ , its means under the two measures differ by at most  $2B_0 d_{TV}(\mu, \nu)$ . No rescaling of the coefficients is implicit.

**Theorem 2.1** (sparse changes preserving congruences). *Let  $a_n \in \mathbb{N}$  and  $A = \sum_{n \geq 0} a_n 2^{-n-1} < \infty$ . For every cutoff  $K$  and every  $f : \mathbb{N} \rightarrow \mathbb{R}$  tending to infinity, there are a set  $S \subseteq [K, \infty)$  of upper Banach density zero and a nondegenerate interval  $I \subset (A, \infty)$  with the following property. Every  $r \in I$  has the form*

$$r = \sum_{n \geq 0} (a_n + e_n) 2^{-n-1},$$

where  $e_n \in \mathbb{N}$  is supported on  $S$ ,  $e_n \leq f(n)$  eventually, and, for every integer  $q \geq 1$ , there is an  $N_q$ , chosen independently of  $r$ , such that

$$q \mid e_n \quad \text{and} \quad q \mid \sum_{i < n} e_i \quad (n \geq N_q).$$

For every  $\varepsilon > 0$ , the construction can instead be chosen with  $e_n \leq (\log(n+3))^\varepsilon$  eventually and  $|S \cap [X, 2X)| = O_\varepsilon(X / \log \log X)$ . In that case the empirical distributions of original and corrected unnormalised blocks of length  $m(X) = o(\log \log X)$ , sampled at the same integer starts in  $[X, 2X)$ , have total variation distance tending to zero, uniformly over  $r$ . Precisely the coupling error is at most  $m|S \cap [X, 2X + m)|/X$ ; with  $\|\Phi\|_\infty \leq B_0$  the bounded-test error is at most twice this quantity times  $B_0$ , also for tests depending on the starting index.

The hypotheses allow any nonnegative integer sequence of polynomial growth, but not  $a_n = 2^n$ , whose dyadic series diverges. The bound on the corrections can grow as slowly as  $\log \log(n+3)$ . A fixed bound is not possible here: eventual boundedness together with eventual divisibility by every fixed modulus would make the corrections eventually zero; their fixed total would then have to be divisible by every modulus and hence be zero. Nonnegative corrections could not change the sum. The set  $S$ , the interval

$I$  and every congruence cutoff are fixed before the target value is chosen. Each correction depends on that value, and its support may be a proper subset of  $S$ . Requiring upper Banach density zero is stronger than requiring ordinary density zero. For example, the union of the integer intervals  $[k!, k! + k)$ ,  $k \geq 3$ , has ordinary density zero but upper Banach density one. If  $k! \leq x < (k+1)!$ , at most  $k(k+1)/2$  indices have occurred, so their proportion is at most  $k(k+1)/(2k!) \rightarrow 0$ . Nevertheless each interval is filled and their lengths are unbounded. Upper Banach density zero rules out this concentration in long intervals, however far apart those intervals lie.

*Proof. Fixed totals.* At stage  $j$  we put a correction at  $n_j - 1$  and a variable pair at  $n_j, n_j + 1$ . We keep the pair's ordinary sum fixed. Consequently, the residue to be cancelled at a later stage is independent of all earlier pair choices. Let  $n_{-1} \geq K$ , write  $s_j = n_j - n_{j-1} \geq 4$ , and take positive integers  $M_j$  with  $M_j \mid M_{j+1}$ . For now let  $D_j$  be a positive integer bounding the digit at stage  $j$ . We first describe the corrections for these parameters, then choose them to obtain both sparsity and an interval of sums.

Starting from  $C_0 = 0$ , define

$$c_j = (-C_j) \bmod M_j \quad (0 \leq c_j < M_j), \quad C_{j+1} = C_j + c_j + M_j D_j.$$

For integers  $0 \leq d_j \leq D_j$ , set

$$e_{n_j-1} = c_j, \quad e_{n_j} = M_j d_j, \quad e_{n_j+1} = M_j(D_j - d_j), \quad (1)$$

and put  $e_n = 0$  elsewhere. The two sums of a triple are

$$\begin{aligned} \sum_{r=n_j-1}^{n_j+1} e_r &= c_j + M_j D_j, \\ \sum_{r=n_j-1}^{n_j+1} e_r 2^{-r-1} &= c_j 2^{-n_j} + M_j D_j 2^{-n_j-2} + d_j M_j 2^{-n_j-2}. \end{aligned}$$

The first identity verifies that  $C_j$  is the cumulative correction before  $n_j - 1$ , whatever the digits are. Thus all  $C_j$  and  $c_j$  can be computed before choosing a single  $d_j$ . The second identity shows that increasing  $d_j$  by one transfers  $M_j$  from  $n_j + 1$  to  $n_j$  and increases the weighted sum by

$$w_j = M_j 2^{-n_j-2}.$$

After  $c_j$  has been inserted, the cumulative correction is divisible by  $M_j$ ; the two entries of the pair preserve this divisibility. Fix  $q \mid M_j$ . At index  $n_j$  both the coefficient correction and the cumulative correction are divisible by  $q$ . At every later triple,  $q$  divides  $C_j$  and  $M_j$ , so it also divides  $c_j$  and both pair entries. The intervening corrections are zero. Hence

$$q \mid e_n, \quad q \mid \sum_{i < n} e_i \quad (n \geq n_j).$$

The entry  $c_j$  need not be divisible by  $q$ , which is why it is placed at  $n_j - 1$ , before the cutoff. The index  $n_j$  depends on  $q$  and the chosen moduli, but not on the digits or the value to be attained.

*Spacing and size.* The variable part at stage  $j$  takes the values  $0, w_j, \dots, D_j w_j$ . To fill the gaps between these choices, we need the later variable parts to have total range at least  $w_j$ . This suggests making each later range a difference of successive dyadic scales. We therefore choose  $D_i = 2^{s_i} - 1$ , for which

$$D_i w_i = (2^{n_i - n_{i-1}} - 1) M_i 2^{-n_i - 2} = M_i (2^{-n_{i-1} - 2} - 2^{-n_i - 2}).$$

Because  $M_i \geq M_j$  for  $i > j$ , replacing every later modulus by  $M_j$  gives the finite estimate

$$\begin{aligned} \sum_{i=j+1}^J D_i w_i &\geq M_j \sum_{i=j+1}^J (2^{-n_{i-1} - 2} - 2^{-n_i - 2}) \\ &= w_j - M_j 2^{-n_J - 2} \quad (J > j). \end{aligned} \tag{2}$$

The digit range  $D_j = 2^{s_j} - 1$  therefore gives the required overlap estimate even when the separations increase. Every entry of (1) is at most  $M_j 2^{s_j}$ , so we must keep this quantity below the prescribed bound at the three indices. We must also have  $s_j \rightarrow \infty$  and ensure that every fixed positive integer divides all sufficiently late  $M_j$ .

To allow a nonmonotone bound  $f$ , put

$$h(n) = \inf_{m \geq n} \min(f(m), m).$$

This nondecreasing lower envelope tends to infinity. A bound chosen using  $h(n_{j-1})$  is therefore valid at each of the later coordinates of the triple. The extra cap by  $m$  serves a separate purpose:  $e_n \leq n$  will make the weighted corrections summable. Choose  $n_{-1}$  with  $h(n_{-1}) \geq 32$ , and set

$$\begin{aligned} k_j &= \max\{k \geq 2 : k! 2^{k+2} \leq h(n_{j-1})\}, \\ M_j &= k_j!, \quad s_j = k_j + 2, \quad n_j = n_{j-1} + s_j. \end{aligned} \tag{3}$$

The maximum exists:  $k = 2$  is admissible because  $2! 2^4 = 32$ , and  $k! 2^{k+2}$  tends to infinity. Since  $h$  is nondecreasing, so is  $k_j$ . Moreover  $n_j \rightarrow \infty$ , hence  $h(n_j) \rightarrow \infty$  and  $k_j \rightarrow \infty$ . It follows that  $M_j \mid M_{j+1}$ , every fixed  $q$  divides all sufficiently late  $M_j = k_j!$ , and the separations  $s_j = k_j + 2$  tend to infinity. The same choice gives the size estimate: at each supported coordinate  $n$  of the  $j$ th triple,

$$0 \leq e_n \leq M_j 2^{s_j} \leq h(n_{j-1}) \leq \min(f(n), n).$$

This proves the size bound and convergence for every choice of digits.

Let  $S = \bigcup_{j \geq 0} \{n_j - 1, n_j, n_j + 1\}$ . For any  $R$ , all but finitely many successive centres are at least  $R$  apart. An interval of length  $H$  therefore contains at most  $3(H/R + 2)$  points of  $S$ , in addition to a fixed finite set. The bound is uniform in the position of the interval. Divide by  $H$ , take the supremum over positions and let  $H \rightarrow \infty$ ; then let  $R \rightarrow \infty$ . This proves upper Banach density zero.

*Attaining the interval.* We have now fixed the centres, moduli, digit ranges and residue corrections, but have left every digit free. Define

$$\beta = \sum_{j \geq 0} (c_j 2^{-n_j} + D_j w_j), \quad F_j = \sum_{i \geq j} D_i w_i.$$

The first quantity is the contribution with all digits zero. The second is the maximum possible variable contribution from stage  $j$  onwards. Both are finite by the bounds just proved, with  $\beta > 0$ ,  $F_0 > 0$  and  $F_j \rightarrow 0$ . The total correction is

$$\sum_n e_n 2^{-n-1} = \beta + \sum_j d_j w_j. \quad (4)$$

Taking  $J \rightarrow \infty$  in (2) now gives

$$F_{j+1} \geq M_j 2^{-n_j-2} = w_j. \quad (5)$$

Every variable sum from stage  $j$  onwards lies in  $[0, F_j]$ ; we must still show that every point of this interval is attained. A target in  $[0, F_j]$  leaves a remainder in  $[0, F_{j+1}]$  after choosing digit  $d$  exactly when it belongs to

$$[dw_j, dw_j + F_{j+1}], \quad d = 0, \dots, D_j.$$

These intervals meet or overlap by (5), and their union is  $[0, F_j]$  because  $F_j = D_j w_j + F_{j+1}$ .

Given  $x \in [0, F_0]$ , start with  $\rho_0 = x$ . Having chosen the first  $j$  digits, choose  $d_j$  so that

$$\rho_{j+1} = \rho_j - d_j w_j \in [0, F_{j+1}].$$

The covering just proved makes this possible at every stage. Since  $0 \leq \rho_j \leq F_j$  and  $F_j \rightarrow 0$ , we obtain

$$x - \sum_{i=0}^j d_i w_i = \rho_{j+1} \rightarrow 0.$$

Thus every  $x \in [0, F_0]$  is attained, and we may take  $I = (A + \beta, A + \beta + F_0)$ . The target  $r \in I$  determines only the digits, through  $x = r - A - \beta$ ; it changes neither  $S$ ,  $I$  nor the cutoff  $N_q = n_j$  chosen using  $q \mid M_j$ . This is the finite-choice covering argument of [27, Lemma 4]: the  $j$ th choice set has largest gap  $w_j$  and diameter  $D_j w_j$ , its maxima are summable, and the sum of subsequent diameters is at least  $w_j$ . Equality is allowed. Fridy's generalised-base lemma [9, p. 194] and the reciprocal-choice argument in [11, Lemma 5.1] are antecedents. Increasing the centres alone need not make  $w_j$  decrease, since  $M_j$  also increases. The finite-choice argument uses (5), without a monotonicity assumption on the weights.

*The logarithmic scale.* For  $f(n) = (\log(n+3))^\varepsilon$ , we can choose the separations explicitly at the scale  $\log \log n$ . Keep the same triples and  $D_j = 2^{s_j} - 1$ , and now choose

$$\begin{aligned} s_j &= \left\lfloor \frac{\varepsilon}{2} \log_2 \log(n_{j-1} + 3) \right\rfloor, \\ k_j &= \max\{k \geq 2 : k! \leq (\log(n_{j-1} + 3))^{\varepsilon/4}\}, \end{aligned} \quad (6)$$

with  $M_j = k_j!$  and  $n_j = n_{j-1} + s_j$ . Start far enough out that  $s_j \geq 4$  and all choices exist. The exponents  $\varepsilon/4$  and  $\varepsilon/2$  add to  $3\varepsilon/4 < \varepsilon$ , so the corrections satisfy the prescribed bound:

$$M_j 2^{s_j} \leq (\log(n_{j-1} + 3))^{3\varepsilon/4} \leq (\log(n + 3))^\varepsilon$$

at each coordinate  $n$  of the triple. After enlarging the initial index, this is also at most  $n$ . The moduli remain nested, and every fixed positive integer divides all sufficiently late moduli. Summability, the congruence induction and (2) therefore apply without change. Here  $s_j \asymp_\varepsilon \log \log n_j$ , so

$$|S \cap [X, 2X]| = O_\varepsilon(X / \log \log X).$$

Choose the same uniform starting index  $n \in [X, 2X)$  for the two blocks. A changed coordinate  $t$  can affect only the starts  $n \in [t - m + 1, t]$ , of which there are at most  $m$ . Only  $t \in S \cap [X, 2X + m)$  can occur in a sampled block. The union bound consequently gives

$$\begin{aligned} d_{\text{TV}}(\mu_{a,X,m}, \mu_{a+e,X,m}) &\leq \mathbb{P}(e_{n+i} \neq 0 \text{ for some } 0 \leq i < m) \\ &\leq \frac{m|S \cap [X, 2X + m)|}{X}. \end{aligned}$$

For  $m \leq X$ , two dyadic bands cover  $[X, 2X + m)$  and give the bound  $O_\varepsilon(m / \log \log X)$ . In particular it tends to zero when  $m = o(\log \log X)$ , uniformly over the values attained, since the same set  $S$  contains every correction support. The block length enters solely through the number of starts that one altered coordinate can affect. For a test bounded in absolute value by  $B_0$  that also depends on the starting index, the coupled values agree at unchanged blocks and differ by at most  $2B_0$  elsewhere. Their mean difference is consequently at most  $2B_0 m |S \cap [X, 2X + m)| / X$ . This uses the common starting index, not merely the marginal distributions. No limiting distribution, randomness or independence of the original coefficients is assumed.  $\square$

Identical marginal distributions alone would not control an index-dependent test. For example, let  $a_n = n \bmod 2$  and  $b_n = 1 - a_n$ . For every even  $X$ , their one-coordinate empirical distributions on  $[X, 2X)$  are equal, but the test  $\mathbf{1}_{\{x=n \bmod 2\}}$  has mean 1 for  $(n, a_n)$  and 0 for  $(n, b_n)$ . This example is not a sparse perturbation; it isolates why the proof couples blocks at the same index.

The comparison concerns absolute, not relative, error in event frequencies. For example, it applies to fixed block lengths and to  $m(X) = \lfloor \sqrt{\log \log X} \rfloor$  for large  $X$ , but makes no vanishing-error assertion for lengths comparable to  $\log \log X$ . An  $o(1)$  change may erase an event whose probability tends to zero; relative preservation requires an error small compared with that probability. The comparison is of finite coefficient blocks, not complete weighted tails: coefficients beyond an unchanged block can still change its tail.

**Corollary 2.2** (target intervals arbitrarily close to the original sum). *For the sparse theorem for an arbitrary sequence, the target interval can additionally be required to lie in  $(A, A + \eta)$  for any prescribed  $\eta > 0$ .*

*Proof.* In the construction for a general bound  $f$ , every nonzero correction satisfies  $e_n \leq n$ . Choose an integer  $K' \geq K$  sufficiently large that  $\sum_{n \geq K'} n 2^{-(n+1)} = (K' + 1)2^{-K'} < \eta$ , and start the construction beyond  $K'$ . This still leaves the originally prescribed prefix unchanged. The complete interval of corrections has positive lower endpoint and upper endpoint at most this sum. Thus the interval may be chosen as close to  $A$  as desired; this does not say that one fixed bound on the corrections permits every target above  $A$ .  $\square$

### 3 Context and mathematical dependencies

Tao's forum comment of 7 October 2025 gives the prime-to-gap reduction and proposes sufficiently uniform quantitative control of about  $\log \log n$  consecutive gaps as an input to irrationality [23, comment of 7 October 2025]. Theorem 4.3 records the reduction with its endpoint and convergence proof. An elementary prime bound suffices for it. We use the prime number theorem separately, to deduce  $P_n \sim n \log n$  in Corollary 8.6.

**Conditional prime-pattern results.** Land's draft gives conditional irrationality [13, Conjecture 1, Theorem 2], and Ringer's draft gives conditional normality [26]. For each fixed integer base  $b \geq 2$ , Ringer's number is  $\sum_{n \geq 0} p_n b^{-(n+1)}$  and its normality is to base  $b$ . Varying  $b$  varies the number, so the result makes no assertion of absolute normality for one constant. The authors supply formalisation material, which we have not rebuilt.

The uniform Hardy–Littlewood hypothesis in [12, Conjecture 1.3] counts prime translates of each nonempty admissible tuple of  $k \leq (\log \log x)^3$  distinct integer shifts in  $[0, (\log x)^2]$ . Admissibility means that the shifts omit a residue class modulo each prime. The main term is the singular series of the tuple multiplied by  $\int_2^x (\log t)^{-k} dt$ , with absolute error at most  $Cx^{1-\delta}$ . The constants  $C, \delta > 0$  and the starting threshold precede the choice of  $x$  and of the tuple. Replacing the integral by its leading asymptotic can exceed this error, and qualitative asymptotics for individual fixed tuples do not give the stated uniformity.

In the version rechecked on 18 September 2026, Ringer also states weaker hypotheses for the joint positions of the first  $L$  primes after a prime basepoint, with  $L$  of order  $\log \log X$ . The basepoints are uniform among primes in  $(X, 2X]$ , where  $X$  measures prime values. One hypothesis controls a weighted sum of adverse tuple-count errors. The adverse sign alternates with the order of inclusion–exclusion, so both undercounts and overcounts can matter. Another hypothesis allows bounded domination by a comparison law in place of total variation approximation. Neither is a single-gap marginal estimate or a consequence of an upper sieve bound alone. We assume none of these hypotheses here.

Our index band  $[X, 2X)$  corresponds to  $[p_X, p_{2X})$ . The prime number theorem gives  $\pi(2p_X) = 2X + o(X)$ , so its symmetric difference with the prime basepoints in  $(p_X, 2p_X]$  has size  $o(X)$ . Each set has  $X + o(X)$  elements. The two uniform measures have common mass equal to the intersection size divided by the larger cardinality, and hence have total variation distance  $o(1)$ . This compares the same uniformly bounded observable with the same normalisation at common basepoints. It supplies no relative or quantitative control of the growing patterns required by the preceding conditional results.

**Related irrationality problems.** Erdős's 1958 article states that  $\sum_{n \geq 1} p_{n-1}^k / n!$  is irrational for each  $k \geq 1$ , and prints a proof only for  $k = 1$ , explaining that the higher-power proof is more complicated [1, pp. 93–95]. Schlage-Puchta later proved the stronger linear independence over  $\mathbb{Q}$  of  $1, S_0, S_1, \dots$ , where  $S_k = \sum_{n \geq 1} p_{n-1}^k / n!$  [18, Theorem 3]. The fixed dyadic denominator question is stated separately in [4, p. 103], including the difficulty of the case  $k = 1$ .

The same page proposes irrationality of  $\sum_{n \geq 1} p_{n-1} / (d_1 \cdots d_n)$  whenever  $d_n \geq 2$  and  $d_n = o(p_{n-1})$ . Kovač's note of 15 April 2026 disproves this variable-denominator asser-

tion with a sum equal to 1 [24, Theorem 1 and proof, pp. 1–2]. Its printed attribution is ChatGPT 5.4 Pro, orchestrated by Vjeko Kovač. Section 3 of the 1958 article also concerns variable product denominators [1, pp. 96–97]. We make no quantified reinterpretation of its nonstandard asymptotic notation in condition (5), which is unused here. The end-point example can be read directly: with  $G_0 = 1$  and  $G_n = \prod_{j=1}^n (p_{j-1} + 1)$ ,

$$\frac{p_{n-1}}{G_n} = \frac{1}{G_{n-1}} - \frac{1}{G_n}, \quad \sum_{n \geq 1} \frac{p_{n-1}}{G_n} = 1.$$

These examples leave fixed dyadic denominators unchanged as a separate question.

A different dyadic series with bounded coefficients is due to Erdős and Pomerance. If  $P(m)$  is the largest prime factor of  $m$ , they proved irrationality of

$$\sum_{m \geq 2} \mathbf{1}_{\{P(m) > P(m+1)\}} 2^{-m} \quad [3, §7, p. 320].$$

Erdős and Graham record the complementary indicator [2, p. 62]. Since consecutive integers cannot have the same largest prime factor, its series is  $1/2$  minus the displayed one and is also irrational. These binary coefficients differ from the unbounded numerators of  $\Pi$ . Schlage-Puchta's concatenation theorem [18, Theorem 2, p. 1] concerns a base  $b \geq 2$  that is not a proper power, with its stated monotonicity and growth assumptions. Concatenation places blocks according to their lengths, whereas our coefficients occupy fixed dyadic positions. We do not use that theorem. The analytic input used below is his Lemma 4, proved by Selberg's sieve: for every fixed nonzero  $F \in \mathbb{Z}[x_0, \dots, x_k]$ ,  $F(g_n, \dots, g_{n+k}) \neq 0$  for almost all  $n$  [18, Lemma 4, pp. 5–6].

**Interval covering and divisibility.** The bounded perturbation in Theorem 8.1 is a binary expansion. The sparse construction uses the finite-choice lemma of Crmarić and Kovač [27, Lemma 4]. For finite nonnegative choice sets with at least two elements and summable maxima, their lemma fills a single interval if each largest successive gap is at most the sum of the subsequent diameters. Here the sets are  $\{0, w_j, \dots, D_j w_j\}$ , and the condition is  $w_j \leq \sum_{i>j} D_i w_i$ . The proof checks this inequality at every stage. Fridy's lemma [9, Lemma, p. 194] assumes nonincreasing weights, a restriction avoided by the finite-choice form. Kovač and Tao use reciprocal choices [11, Lemma 5.1, Theorem 2.5].

Van Doorn and Kovač combine finite subsum covering with a distinguished subsequence of denominators [20, Lemma 7, Proposition 8]. Each distinguished denominator is divisible by all earlier denominators, and every positive integer divides some denominator. Thus every fixed integer divides all sufficiently late distinguished denominators, as it does our factorial moduli. Their condition is imposed on the distinguished subsequence, not on every denominator. At a distinguished cutoff the finite sums lie in the reciprocal grid of the last denominator. Adding the covering inequalities between successive cutoffs gives Lemma 7 for the full finite prefix, leaving no gaps in that grid. Once the denominator of a rational target divides the last denominator and the target lies in the available range, it has a finite representation. Their proposition fills a half-open interval of rational targets by finite subsums. Our construction fills a fixed interval of real targets by infinite choices.

Van Doorn's exchange  $\{21d, 28d\} \leftrightarrow \{20d, 30d\}$  preserves the reciprocal sum  $1/(12d)$  while changing the ordinary sum from  $49d$  to  $50d$  [29, proof of Theorem 3]. The pair in (1) preserves the ordinary sum and changes the weighted one. The analogy concerns the conserved quantity, and does not invoke his partition theorem. The present argument adds no new interval-covering principle.

**Binary subsums.** The *achievement set* of a positive summable sequence consists of sums obtained by retaining or omitting each term. The sequence  $(2^{-n})_{n \geq 1}$  fills  $[0, 1]$ . The sequence  $(3^{-n})_{n \geq 1}$  omits  $(1/6, 1/3)$  because its terms after the first sum to  $1/6$ . Nevertheless a failure of this simple overlap test at some indices does not exclude interior elsewhere in the achievement set. Bartoszewicz, Filipczak and Szymonik obtain interior from a central run of integer block sums in a multigeometric family [32, Theorem 2]. Prus-Wiśniowski and Ptak construct Cantorvals with density-zero overlap indices  $\{n : a_n \leq \sum_{i>n} a_i\}$  [31, Theorem]. These compact sets are the closures of their interiors, and the endpoints of every nondegenerate interval component are accumulation points of one-point components.

The surrounding classification is discussed in [30, §1] and in Nitecki's exposition [36, Theorem 14], which credits Guthrie and Nymann. The cited preprint's title and theorem numbers differ from those in the published Monthly article. Even prescribing the exact inequality pattern leaves the presence of interior undetermined. Marchwicki and Miska's construction [38, Theorem 2.1], with the proof repair in [37], realises any infinite strict term-dominating index set with a Cantor achievement set. The repair preserves uniqueness, and the latter paper gives a simpler proof of a weaker conclusion without uniqueness. Nowakowski's sufficient condition [39, Definition 2, Theorem 3.1] instead requires every stage of the Star Procedure's positivity recursion to succeed. Knowing that neither term-tail inequality holds eventually therefore cannot replace the overlap calculation in our variable-digit proof.

**Uniformity and finite-state comparisons.** Adamczewski, Drmota and Müllner compute logarithmic densities of each fixed automatic sequence along primes [28, Theorems 1.2, 1.4]. Their Theorem 1.2 gives a sufficient condition for natural densities to exist and be rational, while Theorem 1.4 reduces logarithmic-density calculation to integers coprime to a fixed modulus. The general logarithmic densities need not be rational. Applying these results to our tails would require a specified finite-alphabet automaton computing the observable, together with uniform error bounds if its state space grows. The theorems themselves concern neither unbounded consecutive gaps nor a growing family of carry automata.

Kuperberg's singular-series estimate permits a specified growing cardinality [12, Theorem 1.1]. Her arithmetic-progression and smooth-weight estimates [33] provide a fixed-parameter comparison, without giving a law for ordered consecutive-gap blocks. Jha's revised preprint [34] studies growing short-interval prime counts under a strong Hardy–Littlewood hypothesis. Such counts alone do not control the weighted consecutive-gap differences in our criteria. A fixed-dimensional limit theorem likewise leaves the required growing-block errors to be proved. At the cited revision, the *Formal*

*Conjectures* repository leaves Problem #251 unproved [25]. Its zero-based sum starts with the zeroth prime divided by  $2^0$ , and is twice our normalisation.

**The tail argument.** An ordinary binary expansion is rational exactly when its digits are eventually periodic. Applying carries to integer coefficients changes the sequence to which this criterion refers. Even bounded nonperiodic coefficients can give a rational sum, as Proposition D.6 shows. A classical alternative is the growth criterion for Ahmes series  $\sum_k 1/a_k$  with strictly increasing positive integer denominators:  $a_k^{1/2^k} \rightarrow \infty$  suffices for irrationality [6][11, §1]. Replacing divergence by any sufficiently large fixed threshold fails, since shifted Sylvester sequences grow like  $C^{2^k}$  for arbitrarily large  $C$  and have rational reciprocal sum. The criterion is sufficient and is not necessary. Splitting  $p_i/2^{i+1}$  into  $p_i$  equal unit fractions introduces repetitions, and even the distinct denominators  $2^n$  satisfy  $(2^n)^{1/2^n} \rightarrow 1$ .

We instead classify reduced denominators of the complete tails under rationality. Euler's congruence turns rationality into eventual integrality of a fixed positive-length tail difference. This argument needs only integer coefficients and does not force them to repeat: Proposition 8.10 with  $K_n = n$  gives unbounded coefficients  $n - 1$  and sum zero. The criterion involving two small differences then uses parity and a finite enclosure of an actual tail difference. Establishing its simultaneous inequalities arbitrarily late for every positive shift would prove irrationality. We prove neither that prime estimate nor its necessity. The appendix calculations and counterexamples separate these remaining questions from the algebraic equivalences.

## 4 Summation by parts, with the endpoint retained

We return to the actual prime gaps  $g_n = p_{n+1} - p_n$ , with  $p_0 = 2$  and  $g_0 = 1$ . We write

$$\Pi = \sum_{n \geq 0} p_n 2^{-n-1}, \quad S = \sum_{n \geq 0} g_n 2^{-n-1}.$$

Here  $S$  denotes the gap sum. The set of indices used in Section 2 will not be needed below. The two calculations below have different jobs: summation by parts gives  $\Pi = 2 + S$ , and deleting the first gap gives the initial tail  $T_0 = 2S - 1$ .

We begin with a finite identity, so no convergence hypothesis is needed. Its endpoint identifies the estimate required before passing to the infinite series. The linked definitions are the [dyadic partial sum](#) and the [weighted sum of consecutive differences](#). The formula below writes both sums explicitly.

**Proposition 4.1** (finite summation by parts). *For every rational sequence  $P$  and every  $n \geq 0$ ,*

$$\sum_{i=0}^n \frac{P(i)}{2^{i+1}} = P(0) + \sum_{i=0}^{n-1} \frac{P(i+1) - P(i)}{2^{i+1}} - \frac{P(n)}{2^{n+1}}.$$

*Proof.* At  $n = 0$  both sides equal  $P(0)/2$ . When  $n$  is replaced by  $n + 1$ , the change on the right is

$$\frac{P(n+1) - P(n)}{2^{n+1}} - \frac{P(n+1)}{2^{n+2}} + \frac{P(n)}{2^{n+1}} = \frac{P(n+1)}{2^{n+2}}.$$

This is the next term on the left, which proves the identity by induction.  $\square$

Specialising to  $P(i) = p_i$ , whose first value is  $p_0 = 2$ , and writing  $g_i = p_{i+1} - p_i$  for the zero-based gaps, gives the reformulation.

**Theorem 4.2** (prime-gap reformulation). *Let  $p_0 = 2, p_1 = 3, \dots$  be the primes in increasing order and  $g_i = p_{i+1} - p_i$ . For every  $n \geq 0$ ,*

$$\sum_{i=0}^n \frac{p_i}{2^{i+1}} = 2 + \sum_{i=0}^{n-1} \frac{g_i}{2^{i+1}} - \frac{p_n}{2^{n+1}}.$$

The leading 2 is the first prime, not a normalising constant. At  $n = 2$ , for instance, the left side is  $2/2 + 3/4 + 5/8 = 19/8$  and the right side is  $2 + (1/2 + 2/4) - 5/8 = 19/8$ .

#### 4.1 The infinite identity and the irrationality equivalence

To pass to the infinite series, let

$$u_n = \frac{p_n}{2^{n+1}}, \quad v_n = \frac{g_n}{2^{n+1}}.$$

Thus  $\sum_n u_n = \Pi$ . The termwise identity  $v_n = 2u_{n+1} - u_n$  is the [dyadic discrete derivative](#). It expresses each gap term as an integer combination of two consecutive prime terms, so once  $(u_n)$  is summable the gap series can be summed by rearranging two copies of the prime series.

**Theorem 4.3** (infinite prime-gap identity). *Both series converge and*

$$\sum_{n \geq 0} \frac{p_n}{2^{n+1}} = 2 + \sum_{n \geq 0} \frac{g_n}{2^{n+1}}.$$

Writing  $S = \sum_{n \geq 0} g_n 2^{-(n+1)}$ , the complete tails  $T_N = \sum_{j \geq 1} g_{N+j} 2^{-j}$  satisfy  $T_0 = 2S - 1$  and  $T_{N+1} = 2T_N - g_{N+1}$ . Thus  $\Pi$ ,  $S$  and  $T_0$  have the same rationality status.

*Proof.* The bound  $p_n \leq 1250(n+1)^4$  of Appendix A makes  $(u_n)$  summable, and  $0 < g_n \leq p_{n+1}$  then makes  $(v_n)$  summable. The shifted sequence  $(u_{n+1})$  is summable, so summing  $v_n = 2u_{n+1} - u_n$  and using  $u_0 = 1$  gives

$$\sum_{n \geq 0} v_n = 2 \left( \sum_{n \geq 0} u_n - u_0 \right) - \sum_{n \geq 0} u_n = \sum_{n \geq 0} u_n - 2.$$

□

The tail assertions follow by splitting off the first term and using  $g_0 = 1$ . Affine transformations with rational coefficients and nonzero slope preserve rationality.

The prime number theorem is needed neither for this convergence nor for the identity. Its asymptotic  $p_n \sim n \log n$  is used in Corollary 8.4 and in the argument counting repeated tail values [17, Chapter 6].

**Corollary 4.4** (exact irrationality reformulation).  *$\Pi$  is irrational if and only if  $S = \sum_{n \geq 0} g_n 2^{-(n+1)}$  is irrational. The corresponding zero-based series with denominator  $2^n$  equals  $4 + 2S$  and has the same irrationality status.*

*Proof.* The identity just proved gives  $\Pi = 2 + S$ . Adding a rational constant or multiplying by a nonzero rational number preserves rationality and irrationality.  $\square$

Concretely,  $\Pi = 3.674643966 \dots$  is irrational if and only if  $S = 1.674643966 \dots$  is. Neither irrationality statement is established here.

## 5 The tail recurrence and the exact criteria

We rescale each tail so that its first denominator is again 2. For a convergent series  $\sum_{i \geq 0} a_i 2^{-i-1}$  with  $a_i \in \mathbb{Z}$ , put

$$T_N = 2^{N+1} \sum_{i \geq N} a_i 2^{-i-1} = \frac{a_{N+1}}{2} + \frac{a_{N+2}}{4} + \dots.$$

Thus  $T_N$  starts with coefficient  $a_{N+1}$ , not  $a_N$ . Doubling and removing that first coefficient gives  $T_{N+1} = 2T_N - a_{N+1}$ . Modulo integers, a step is simply doubling; the detailed values of the integer coefficients disappear. This is why the recurrence gives a rationality criterion for arbitrary integer coefficients.

At  $N = 0$  we have  $T_0 = 2 \sum_{i \geq 0} a_i 2^{-i-1} - a_0$ . In particular, for the prime gaps  $T_0 = 2S - 1 = 2\Pi - 5$ , so  $T_0$ ,  $S$  and  $\Pi$  have the same rationality status. We first analyse the recurrence abstractly. We then impose the vanishing boundary condition that singles out complete tails and apply the criterion to these prime-gap tails.

**Definition 5.1.** Let  $g : \mathbb{N} \rightarrow \mathbb{Z}$  and  $T : \mathbb{N} \rightarrow \mathbb{Q}$  or  $T : \mathbb{N} \rightarrow \mathbb{R}$ . Say  $T$  satisfies the *dyadic tail recurrence* with coefficients  $g$  when  $T_{N+1} = 2T_N - g_{N+1}$  for every  $N$ . Write  $\sigma_h(N) = T_{N+h} - T_N$  for the *shift* of length  $h$  at  $N$ , and call a real number *integral* when it belongs to  $\mathbb{Z}$ .

For a complete-tail example, take  $g_n = 2$  for odd  $n \geq 1$  and  $g_n = 4$  for even  $n \geq 1$ . Its complete tail at index 0 is  $T_0 = (2/2 + 4/4)/(1 - 1/4) = 8/3$ , and the tails alternate between  $8/3$  and  $10/3$ . Thus  $\sigma_1(N)$  is always  $2/3$  or  $-2/3$ , whereas  $\sigma_2(N) = 0$  for every  $N$ . Nonintegrality at one prescribed shift length does not imply irrationality, even for positive even coefficients and genuine tails.

Modulo integers, the recurrence is repeated doubling: the coefficient term does not affect the fractional part. An integral shift means that the fractional part has returned to an earlier value. Iterating the recurrence  $h$  times makes this observation explicit: it multiplies  $T_N$  by  $2^h$  and accumulates an integer. Define  $B_{0,N} = 0$  and  $B_{h+1,N} = 2B_{h,N} + g_{N+h+1}$ , so that  $B_{h,N} = g_{N+1}2^{h-1} + \dots + g_{N+h}$  is the **weighted integer sum** accumulated in those  $h$  steps.

**Theorem 5.2** (block identity). *For every  $N$  and  $h$ ,*

$$T_{N+h} = 2^h T_N - B_{h,N}, \quad \text{hence} \quad \sigma_h(N) = (2^h - 1) T_N - B_{h,N}.$$

*Proof.* The case  $h = 0$  follows from  $B_{0,N} = 0$ . If the first identity holds at  $h$ , then

$$T_{N+h+1} = 2(2^h T_N - B_{h,N}) - g_{N+h+1} = 2^{h+1} T_N - B_{h+1,N}.$$

Subtracting  $T_N$  gives the second identity.  $\square$

Subtracting the tail recurrences at  $N + h$  and  $N$  gives

$$\sigma_h(N + 1) = 2\sigma_h(N) - (g_{N+h+1} - g_{N+1}), \quad (7)$$

which is the [recurrence for the tail differences](#). Since  $B_{h,N}$  is an integer, Theorem 5.2 also gives the [integrality criterion](#):  $\sigma_h(N)$  is integral if and only if  $(2^h - 1)T_N$  is integral. For fixed  $h$  in this general recurrence,  $T_N$  therefore determines whether the shift is integral, regardless of the later integer coefficients. For a complete tail, however, changing those coefficients also changes  $T_N$ ; the observation does not permit arbitrary changes to prime gaps.

For rational  $T_N = u/v$  in lowest terms, the recurrence gives  $T_{N+1} = (2u - g_{N+1}v)/v$ . The numerator has greatest common divisor  $\gcd(2, v)$  with  $v$ , because  $\gcd(u, v) = 1$ . Thus

$$\text{den}(T_{N+1}) = \frac{\text{den}(T_N)}{\gcd(2, \text{den}(T_N))}, \quad \sigma_h(N) \in \mathbb{Z} \iff \text{den}(T_N) \mid 2^h - 1. \quad (8)$$

Each even denominator loses exactly one factor of 2 and an odd denominator is unchanged, so after finitely many steps the denominator is an odd integer  $d$ ; Euler's congruence then gives  $d \mid 2^{\varphi(d)} - 1$ , and the multiplicative order of 2 modulo  $d$  is the least positive such exponent, with the convention that this order is 1 when  $d = 1$ . The hypothesis that  $d$  is odd cannot be dropped: if  $T_N = 1/2$  then  $2^h - 1$  is odd for every  $h \geq 1$ , so no shift at  $N$  is integral. The resulting periodicity is modulo  $\mathbb{Z}$ : the fractional parts eventually repeat, not necessarily the full tail values or the integer coefficients.

**Theorem 5.3** (exact rationality classification). *Let  $T : \mathbb{N} \rightarrow \mathbb{R}$  satisfy  $T_{N+1} = 2T_N - g_{N+1}$  with integer coefficients  $g$ . The following are equivalent:*

- (i)  $T_0$  is rational;
  - (ii)  $\sigma_h(N)$  is an integer for some  $h \geq 1$  and some  $N$ ;
  - (iii) for some fixed  $h \geq 1$ ,  $\sigma_h(N)$  is an integer at every sufficiently large  $N$ .
- More precisely, if  $T_0 = u/(2^s d)$  is in lowest terms, with  $d$  odd, then

$$\text{den}(T_N) = 2^{\max(s-N, 0)} d, \quad \sigma_h(N) \in \mathbb{Z} \iff N \geq s \text{ and } d \mid 2^h - 1 \quad (h \geq 1).$$

Consequently  $T_0$  is irrational if and only if every positive-length shift is nonintegral at every index, equivalently if and only if for every  $h \geq 1$  and every cutoff some later  $N$  has  $\sigma_h(N) \notin \mathbb{Z}$ .

*Proof.* For (i)  $\Rightarrow$  (iii), the block identity makes every  $T_N$  rational. By (8), its denominator is a fixed odd integer  $d$  for all sufficiently large  $N$ . Choose  $h = \varphi(d)$ ; Euler's congruence and the same formula show that every subsequent  $\sigma_h(N)$  is integral. The implication (iii)  $\Rightarrow$  (ii) is immediate. For (ii)  $\Rightarrow$  (i), the block identity gives  $\sigma_h(N) = (2^h - 1)T_N - B_{h,N}$  with  $B_{h,N}$  and  $\sigma_h(N)$  integers and  $2^h - 1 \neq 0$ , so  $T_N$  is rational, and  $T_N = 2^N T_0 - B_{N,0}$  then makes  $T_0$  rational. Negating the three conditions gives the two irrationality formulations.  $\square$

**Lemma 5.4** (the boundary condition for the tail recurrence). *Let  $a_1, a_2, \dots$  be real numbers with  $\sum_{j \geq 1} |a_j| 2^{-j} < \infty$ , and let  $U_{N+1} = 2U_N - a_{N+1}$ . Then*

$$U_N = \sum_{j \geq 1} a_{N+j} 2^{-j} \quad \text{for every } N \quad \Longleftrightarrow \quad 2^{-N} U_N \rightarrow 0.$$

*Proof.* Iteration gives  $2^{-N} U_N = U_0 - \sum_{j=1}^N a_j 2^{-j}$ . The limit is zero exactly when  $U_0$  equals the full series; subtracting its first  $N$  terms then gives the claimed tail formula. Without the boundary condition, one may add  $C2^N$  to every  $U_N$  without changing the recurrence.  $\square$

For instance, zero coefficients and initial value  $U_0 = 1/12$  give  $\frac{1}{12}, \frac{1}{6}, \frac{1}{3}, \frac{2}{3}, \frac{4}{3}, \dots$ , with  $U_{N+h} - U_N = (2^h - 1)2^N/12$ . Shifts of length 2 become integral at index 2; shifts of length 1 never do. These are not complete tails of the zero sequence, since  $2^{-N} U_N = 1/12$  does not tend to zero. The denominator calculation remains valid, but the boundary condition excludes this extra homogeneous term.

**The actual prime-gap tails.** Put

$$T_N = \sum_{j \geq 1} \frac{g_{N+j}}{2^j},$$

which converges by Theorem 4.3. Reindexing identifies this sum with the rescaled complete tail of  $S$ , as in the [shifted-gap series identity](#). Splitting off the first term gives  $T_{N+1} = 2T_N - g_{N+1}$  without a rationality hypothesis, as formalised in the [real tail recurrence](#). Since  $T_0 = 2S - 1 = 2.349287932\dots$ , the numbers  $\Pi$ ,  $S$  and  $T_0$  have the same rationality status, and Theorem 5.3 specialises to the actual series: irrationality of  $\Pi$  is equivalent to nonintegrality arbitrarily far out for each positive shift of  $T$ . The formal statement is the [irrationality equivalence for the actual prime tails](#). Thus the passage from the series to its complete tails is proved. What remains is to establish the required nonintegrality for these prime-gap tails.

### 5.1 Pairs of congruent indices and least-common-multiple shifts

We now allow the distance between the indices to vary and require congruence modulo a prescribed positive integer. Applying this test for every modulus gives another characterisation of irrationality. Thus the variable offset changes the form of a possible witness while leaving the logical strength of the condition unchanged.

**Theorem 5.5** (pairs of congruent indices).  *$S$  is irrational if and only if for every  $t \geq 1$  and every  $N_0$  there are  $N, M \geq N_0$  with  $M \equiv N \pmod{t}$  and  $T_M - T_N \notin \mathbb{Z}$ .*

*Proof.* If  $S$  is irrational, take  $M = N + t$  for the  $N$  supplied by Theorem 5.3 at shift length  $t$ . Conversely, suppose  $S$  is rational. By (8) the orbit reaches an index  $N_0$  beyond which the reduced denominator is a fixed odd  $d$ ; let  $t$  be the multiplicative order of 2 modulo  $d$ , taking  $t = 1$  when  $d = 1$ . For  $N, M \geq N_0$  with  $M \equiv N \pmod{t}$ , the difference  $T_M - T_N$  is then an integer, contradicting the stated property at this  $t$  and this cutoff.  $\square$

Modulo one, the tail recurrence is multiplication by two. Once the reduced denominator is the fixed odd integer  $d$ , a difference  $T_M - T_N$  is integral precisely when  $M - N$  is divisible by the order of 2 modulo  $d$ . Theorem 5.5 therefore allows the distance between the indices to vary, but requires a nonintegral pair beyond every cutoff for every prescribed modulus.

Equal tail values do not help this criterion, since their difference is zero. Section D.4 shows that rationality and the prime number theorem would in fact force many equal tail values. It also explains why, beyond the rational denominator cutoff, an interval condition excluding integers already contradicts rationality at indices congruent modulo the corresponding multiplicative order. These are ordinary deductions, not additional formalised statements.

A second reformulation tests one prescribed sequence of pairs of indices. Let  $L_0 = 1$  and  $L_j = \text{lcm}(1, \dots, j)$ .

**Theorem 5.6** (criterion using least common multiples). *Let  $g : \mathbb{N} \rightarrow \mathbb{Z}$  and  $T : \mathbb{N} \rightarrow \mathbb{R}$  satisfy  $T_{N+1} = 2T_N - g_{N+1}$ . Then  $T_0$  is irrational if and only if  $T_{2L_j} - T_{L_j} \notin \mathbb{Z}$  for every  $j \geq 0$ .*

*Proof.* Theorem 5.3 gives the forward implication. Conversely, if  $T_0$  is rational then some positive shift length  $h$  is integral at every basepoint beyond an index  $N_0$ . Choose  $j$  so large that  $N_0 \leq L_j$  and  $h \mid L_j$ . The telescoping identity  $\sigma_{a+b}(N) = \sigma_a(N) + \sigma_b(N + a)$  shows by induction that every positive multiple of  $h$  is integral at every such basepoint, so  $T_{2L_j} - T_{L_j}$  is an integer.  $\square$

One could use factorials instead of least common multiples, since they have the same eventual divisibility property. The least common multiples are no larger at each index. Theorems 5.5 and 5.6 change which tail differences one must test, but remain exact equivalences for integer-coefficient recurrences. Neither proves the needed nonintegrality for the actual prime gaps.

## 6 A local certificate, and one actual pair

To rule out an integral shift, we can bound two consecutive complete differences in  $(-1, 1)$  and check that the corresponding gaps are unequal. Indeed, simultaneous integrality would force both differences to vanish, and their recurrence would then give  $g_{N+h+1} = g_{N+1}$ . For even gaps we can state these conditions as a single signed interval for the first difference.

**Theorem 6.1** (two consecutive small tail differences). *Let  $T$  satisfy the dyadic tail recurrence with integer coefficients  $g$ , and fix  $h$  and  $N$ . If*

$$-1 < \sigma_h(N) < 1, \quad -1 < \sigma_h(N + 1) < 1, \quad g_{N+h+1} \neq g_{N+1},$$

*then  $\sigma_h(N)$  and  $\sigma_h(N + 1)$  cannot both be integers. Consequently, if such a pair occurs beyond every threshold, the  $h$ -shift is not eventually integral.*

*Proof.* An integer strictly between  $-1$  and  $1$  is zero. If both shifts were integers, both would vanish, and (7) would give  $g_{N+h+1} = g_{N+1}$ . The final assertion chooses one such adjacent pair after the alleged onset of integrality.  $\square$

**Corollary 6.2** (real form and the sufficient condition). *The same statement holds for a real orbit, with the same proof. If for every  $h \geq 1$  and every cutoff some later  $N$  satisfies the three displayed conditions for the actual prime gaps, then  $\Pi$  is irrational.*

The unequal-gap condition alone is available: at  $h = 1, N = 2$  it reads  $g_4 = 2 \neq 4 = g_3$ , and Proposition 6.3 gives such inequalities at arbitrarily large indices for each fixed  $h$ . The missing assertion is that both small-tail inequalities and the unequal-gap condition hold at the same arbitrarily late indices. Separate infinite sets of witnesses for the three requirements need not intersect.

**Proposition 6.3** (prime gaps do not become periodic). *For every positive  $h$ , the actual consecutive-prime-gap sequence is not eventually periodic with period  $h$ .*

*Proof.* For  $m \geq 2$  the integers  $(m+1)! + 2, \dots, (m+1)! + m + 1$  are composite, so the gaps are unbounded. An eventually periodic sequence of natural numbers has finite range after its preperiod and a bounded initial segment, hence is bounded.  $\square$

Far stronger lower bounds for large gaps are known [8, Theorem 1]; the **unboundedness of the gaps** is all that is needed.

With even gap differences, the three conditions in Theorem 6.1 reduce to two possible signed intervals, each accompanied by a prescribed gap difference. For fixed  $h$  write  $D_N = \sigma_h(N)$  and  $\delta_N = g_{N+h+1} - g_{N+1}$ , so that  $D_{N+1} = 2D_N - \delta_N$ .

**Theorem 6.4** (an equivalent signed interval test). *Assume  $\delta_N$  is even. The conjunction  $-1 < D_N < 1, -1 < D_{N+1} < 1, \delta_N \neq 0$  is equivalent to*

$$(\delta_N = 2 \text{ and } \frac{1}{2} < D_N < 1) \quad \text{or} \quad (\delta_N = -2 \text{ and } -1 < D_N < -\frac{1}{2}).$$

*In either case, with  $s = \delta_N/2 \in \{-1, 1\}$ ,  $sD_{N+1} \in (-1, 0)$  and both differences are nonintegral. The same statement holds for any real triple with  $D' = 2D - \delta$  and even  $\delta$ , without a hypothesis on prime gaps.*

*Proof.* Since  $\delta_N = 2D_N - D_{N+1}$ , the bounds  $|D_N| < 1$  and  $|D_{N+1}| < 1$  give  $-3 < \delta_N < 3$ . A nonzero even integer in this interval is 2 or  $-2$ . If  $\delta_N = 2$ , the inequality  $-1 < 2D_N - 2 < 1$ , together with  $|D_N| < 1$ , is equivalent to  $1/2 < D_N < 1$ . Negating both differences gives the case  $\delta_N = -2$ . Each substitution is reversible.  $\square$

For the actual gaps, both indices  $N + 1$  and  $N + h + 1$  are positive, so  $\delta_N$  is even for every  $N \geq 0$ . Theorem 6.4 therefore applies to the actual tails. Its proof uses order, the recurrence and the evenness of  $\delta_N$ ; it does not require rational tail values. This is the real version used in Theorem 8.7.

## 6.1 An explicit remainder and a certified actual pair

We make the signed interval test finite by bounding the omitted part of each complete tail. The polynomial prime bound gives a rational remainder, so after clearing denominators we need only strict integer inequalities.

**Proposition 6.5** (explicit remainder). *For integers  $h, N \geq 0$  and  $L \geq 1$  put*

$$F_{h,N,L} = \sum_{j=1}^L \frac{g_{N+h+j} - g_{N+j}}{2^j}, \quad P(x) = x^4 + 8x^3 + 36x^2 + 104x + 150,$$

$$E_{h,N,L} = \frac{1250}{2^L} (P(N+h+L+2) + P(N+L+2)).$$

*Then  $|\sigma_h(N) - F_{h,N,L}| \leq E_{h,N,L}$ . In particular  $|F_{h,N,L}| + E_{h,N,L} < 1$  certifies  $|\sigma_h(N)| < 1$ , and  $\text{dist}(F_{h,N,L}, \mathbb{Z}) > E_{h,N,L}$  certifies  $\sigma_h(N) \notin \mathbb{Z}$ .*

*Proof.* Splitting the complete tails after  $L$  terms gives

$$\sigma_h(N) = F_{h,N,L} + 2^{-L} \sigma_h(N+L).$$

The last term is the part not determined by the finite gap blocks. The checked bound  $p_n \leq 1250(n+1)^4$  gives  $0 \leq g_m \leq p_{m+1} \leq 1250(m+2)^4$ , so its absolute value is at most

$$1250 \sum_{j>L} \frac{(N+h+j+2)^4 + (N+j+2)^4}{2^j}.$$

The polynomial identity  $2P(x) = (x+1)^4 + P(x+1)$  telescopes to  $\sum_{k=1}^J (m+k)^4 2^{-k} = P(m) - P(m+J)2^{-J}$ , whose last term tends to zero. Apply it with  $m = N+h+L+2$  and  $m = N+L+2$  after writing  $j = L+k$ . The two tests follow from the triangle inequality and the definition of distance to  $\mathbb{Z}$ .  $\square$

**Proposition 6.6** (a certified adjacent pair). *For the actual prime gaps,  $h = 1$  and  $N = 2$  satisfy the three hypotheses of Theorem 6.1: both  $\sigma_1(2)$  and  $\sigma_1(3)$  lie in  $(-1, 1)$  and are nonintegral, and  $g_4 = 2 \neq 4 = g_3$ .*

*Proof.* Take  $L = 40$  and  $Q = 2^{40} = 1099511627776$ . Exact integer arithmetic over the first 46 primes gives

| $N$ | $QF_{1,N,40}$ | $QE_{1,N,40}$ |
|-----|---------------|---------------|
| 2   | -662838684750 | 11764181250   |
| 3   | 873345886050  | 12805761250   |

In each row  $|QF| + QE < Q$ , which puts the whole certified interval inside  $(-1, 1)$ , and  $QE$  is smaller than the distance from  $QF$  to the nearest multiple of  $Q$ , which excludes every integer. The margins are 424908761776 and 213359980476 respectively. These are strict integer comparisons rather than inferences from rounded numerical tails. Appendix B replays them.  $\square$

This computation proves the local condition at one pair of indices. The quantifiers over every  $h$  and every cutoff remain.

**Proposition 6.7** (certifying two small differences from one enclosure). *Let  $D_{N+1} = 2D_N - \delta_N$  with real  $D_N$ , and suppose  $\delta_N = 2s$  for  $s \in \{-1, 1\}$ . If integers  $A, B, Q$  satisfy  $Q > 0$ ,  $B \geq 0$ ,  $|QD_N - A| \leq B$ , and*

$$2sA - Q > 2B, \quad Q - sA > B,$$

then  $1/2 < sD_N < 1$ ,  $|D_{N+1}| < 1$ , and both  $D_N, D_{N+1}$  are nonintegral. In particular, the two small-difference conditions are certified from just one tail enclosure.

*Proof.* Since  $|s| = 1$ , the enclosure gives

$$\frac{sA - B}{Q} \leq sD_N \leq \frac{sA + B}{Q}.$$

The first strict inequality says that the lower endpoint exceeds  $1/2$ ; the second says that the upper endpoint is less than 1. Thus  $sD_N \in (1/2, 1)$ . The recurrence and  $\delta_N = 2s$  now give  $sD_{N+1} = 2sD_N - 2 \in (-1, 0)$ . Both intervals exclude integers, and the second also gives  $|D_{N+1}| < 1$ .  $\square$

For  $h = 1, N = 2$ , the exact data already used above give  $Q = 2^{40}$ ,  $A = -662838684750$ ,  $B = 11764181250$  and  $s = -1$ . The two positive integer margins are respectively 202637379224 and 424908761776. Proposition 6.6 encloses each difference separately. Here the first enclosure and the exact gap difference give the second interval through the recurrence. The second enclosure is therefore an independent cross-check, not a necessary input.

## 7 Two lower bounds on a possible rational denominator

We can also use a finite enclosure of  $\Pi$  itself. A rational number strictly between two fractions of determinant one has denominator at least the sum of their denominators. We apply this observation to two enclosures obtained from the polynomial prime bound. Appendix B prints the smaller certificate; the linked Lean development checks the integer comparisons for both. The conclusions exclude finite ranges of possible denominators and do not establish irrationality.

**Theorem 7.1** (a denominator bound checked by exact integer comparisons). *Let  $a \in \mathbb{Z}$  and let  $b$  be a positive integer. If  $\Pi = a/b$  then  $b \geq 2^{589} > 10^{177}$ , and the same floor holds for every rational equal to  $S$ .*

*Proof.* Put  $c = 1229$ , the number of primes below  $10^4$ , and  $A = \sum_{i=0}^{c-1} p_i 2^{c-i-1}$ , so that  $A/2^c$  is the  $c$ th partial sum of  $\Pi$ . All terms are positive, so  $A/2^c \leq \Pi$ . For the upper bound,  $p_{c+j} \leq 1250(c+j+1)^4$ . Since  $c \geq 9$ , each successive ratio of  $(c+1+j)^4$  is at most  $(11/10)^4 < 3/2$ . Hence  $(c+1+j)^4 \leq (c+1)^4 (3/2)^j$  for all  $j \geq 0$ . The omitted dyadic terms are therefore bounded by a geometric series with ratio  $3/4$ , whose sum gives

$$\frac{A}{2^c} \leq \Pi \leq \frac{2A + 5000(c+1)^4}{2^{c+1}}.$$

The four positive integers  $u, v, u', v'$  printed in Appendix B satisfy

$$u'v - uv' = 1, \quad \frac{u}{v} < \frac{A}{2^c}, \quad \frac{2A + 5000(c+1)^4}{2^{c+1}} < \frac{u'}{v'}, \quad v + v' \geq 2^{589},$$

all four being integer comparisons after clearing the positive denominators. If  $\Pi = a/b$ , the strict lower and upper bounds make  $av - bu$  and  $bu' - av'$  positive integers. Each

is therefore at least 1. The determinant identity expresses  $b$  as their positive integer combination of  $v'$  and  $v$ :

$$b = b(u'v - uv') = (av - bu)v' + (bu' - av')v \geq v + v' \geq 2^{589}.$$

The only properties needed of the bracketing fractions are their order, positive denominators and determinant one. Their decimal expansions and the way they were found play no part in this denominator bound. Since  $589 \log_{10} 2 > 177$ , the floor exceeds  $10^{177}$ . If  $S = a/b$ , apply the same argument to  $\Pi = (a + 2b)/b$ .  $\square$

The Lean kernel evaluates the four inequalities itself, on the same integers, after recomputing the primes below  $10^4$  by trial division; no compiled code is trusted. Its analytic input is the same enclosure, derived from the [polynomial bound](#)  $p_n \leq 1250(n+1)^4$ . The four integers  $u, v, u', v'$  have 177 and 178 digits.

**Theorem 7.2** (certified continued-fraction exclusion). *Every rational equal to  $\Pi$ , and hence every rational equal to  $S$ , has reduced denominator  $q \geq 2^{39997}$ , and therefore  $q > 10^{12040}$ .*

*Proof.* Put  $c = 80200$ , the number of primes below 1 023 068, and  $A = \sum_{i=0}^{c-1} p_i 2^{c-i-1}$ . The estimate in the proof of Theorem 7.1 holds for every  $c \geq 9$  and gives

$$\frac{A}{2^c} \leq \Pi \leq \frac{2A + 5000(c+1)^4}{2^{c+1}}.$$

Four positive integers  $u, v, u', v'$ , described after this proof, satisfy

$$u'v - uv' = 1, \quad \frac{u}{v} < \frac{A}{2^c}, \quad \frac{2A + 5000(c+1)^4}{2^{c+1}} < \frac{u'}{v'}, \quad v + v' \geq 2^{40062},$$

all four being integer comparisons after clearing the positive denominators. If  $\Pi = a/b$  with  $a \in \mathbb{Z}$  and  $b$  a positive integer, the determinant calculation in the proof of Theorem 7.1 gives

$$b \geq v + v' \geq 2^{40062} > 10^{12059}.$$

The last inequality holds because  $10^{12059} = (10^{59})^{204} 10^{23}$  and  $2^{40062} = (2^{196})^{204} 2^{78}$ , with  $10^{59} < 2^{196}$  and  $10^{23} < 2^{78}$ . The reduced denominator  $q$  is one such  $b$ , so  $q \geq 2^{40062} > 2^{39997}$  and  $q > 10^{12059} > 10^{12040}$ . If  $S = a/b$ , apply the same argument to  $\Pi = (a + 2b)/b$ .  $\square$

The integers  $u, v, u', v'$  come from the continued fractions of the two endpoints  $A/2^c$  and  $(2A + 5000(c+1)^4)/2^{c+1}$ , which agree in their first 23407 partial quotients. The fraction  $u/v$  is the convergent formed by those partial quotients, and  $u'/v'$  is the mediant of  $u/v$  and the convergent before it. Only the four comparisons enter the proof. The numerators  $u, u'$  have 12061 decimal digits and the denominators  $v, v'$  have 12060.

The Lean kernel checks every step of this proof and evaluates every computation itself; no compiled code is trusted. It recomputes  $c$  and  $A$  from the definition of primality, scanning the integers below 1 023 068 in 250 consecutive blocks and checking the prime count and partial numerator at the end of each block. It decides primality by trial division below 1012. On  $[1012, 1013^2)$ , which contains the rest of the scan because

$1013^2 = 1\,026\,169$ , it uses the [fact](#) that such an integer is prime exactly when it is coprime to the product of the primes below 1012, and it recomputes that product by trial division. Lean proves that the two primality tests agree at every natural number. The kernel then decides the four comparisons, together with  $10^{59} < 2^{196}$  and  $10^{23} < 2^{78}$ ; the enclosure and the determinant step are proved in general form. Lean proves the [bound](#)  $b \geq 2^{40062}$  and  $b > 10^{12059}$  for both series and derives the theorem as stated from it, using only Lean's three standard axioms.

An independent [program](#) outside Lean recomputes  $A$  by a prime sieve and uses exact integer arithmetic. Its [recorded output](#) has the same 23407 common partial quotients. It brackets the interval with the last two consecutive convergents, whose denominator sum has bit length 40062, and checks the advertised bound  $2^{39997} > 10^{12040}$ . This pair differs from the convergent–mediant pair in the Lean certificate above, whose denominator sum has bit length 40063 and is at least  $2^{40062}$ . The two bit lengths refer to different valid pairs.

The constants  $2^{39997}$  and  $10^{12040}$  in the statement come from a separate computation with a bracket of width  $2^{-80000}$ , which used the best-approximation property of convergents [10, §6, Theorems 16 and 17]; Short gives an accessible statement and proof [19, Theorem 1.1]. Its [recorded output](#) lists the generating program and its digest and a last convergent whose denominator has bit length 39998, hence is at least  $2^{39997}$ ; the program also checked its arithmetic against the known expansions of  $e$  and  $\pi$ . That bit length describes one convergent only; the bound for every rational in the enclosure comes from the certified sum  $v + v'$  above, and  $2^{39997} > 10^{12040}$  is a separate integer comparison.

A rational number may have a denominator beyond either bound. Extending the computation can exclude a larger finite range, but a longer prefix need not improve the bound at every step and cannot by itself exclude all denominators. These statements are denominator exclusions, not estimates for an irrationality exponent.

## 8 Coefficient conditions and counterexamples

We examine which coefficient conditions leave room for a rational dyadic sum. Bounded residue-preserving changes give the first examples, and finite unions of translated polynomial zero sets explain why they preserve nonconcentration. Sparse changes then give the simultaneous comparison with prime gaps. For the actual gaps, the indices satisfying the two small-difference conditions and the unequal-gap condition have density zero. They may nevertheless include arbitrarily large indices.

### 8.1 Bounded residue-preserving perturbations

For one fixed positive modulus  $M$ , we may add either 0 or  $M$  after a prescribed prefix. The weighted choices form a binary expansion, so we can choose a rational sum while preserving every coefficient modulo  $M$ . This construction permits changes at every index after the prefix; it imposes neither density-zero support nor eventual preservation of all moduli.

**Theorem 8.1** (bounded perturbations preserving congruences). *Let  $a_n$  be natural numbers with  $\sum_{n \geq 0} a_n 2^{-(n+1)}$  convergent. For every integer  $M \geq 1$  and every cutoff  $K$  there are digits  $\varepsilon_n \in \{0, 1\}$ , zero for  $n < K$ , such that  $\sum_{n \geq 0} (a_n + M\varepsilon_n) 2^{-(n+1)}$  is rational.*

*Proof.* Write  $A = \sum_{n \geq 0} a_n 2^{-(n+1)}$  and choose a rational  $r \in (A, A + M2^{-K})$ . A binary expansion of  $(r - A)/M$  has the form  $\sum_{n \geq K} \varepsilon_n 2^{-(n+1)}$  with  $\varepsilon_n \in \{0, 1\}$ ; set  $\varepsilon_n = 0$  for  $n < K$ . The perturbed series converges and equals  $r$ .  $\square$

Hence a property shared by every allowed perturbation cannot by itself force irrationality, since one such perturbation has a rational sum; this consequence is not stated separately in Lean. Lean also checks the construction at the actual consecutive prime gaps, for every  $M \geq 1$  and every  $K$ : the [perturbed prime-gap series is rational](#), and each perturbed gap lies in  $[g_n, g_n + M]$  and is [congruent to  \$g\_n\$  modulo  \$M\$](#) . For each fixed  $k \geq 1$ , the sum of  $k$  consecutive gaps increases by at most  $kM$ . Thus infinitely many bounded gaps and bounded clusters of each fixed size survive, but their original numerical bounds need not. Nonnegative corrections also preserve lower bounds for large gaps. Taking  $M = 2q$  preserves parity and the cumulative positions modulo a prescribed  $q$ , hence any limiting residue frequencies of those positions. These conclusions concern the size and residue information in the results discussed in Section 10; they establish neither primality of the new positions nor preservation of every quantitative assertion of those theorems.

## 8.2 Algebraic nonconcentration survives the rationalising perturbation

Schlage-Puchta's lemma states that  $F(g_n, \dots, g_{n+k}) \neq 0$  for almost all  $n$ , for every fixed nonzero  $F \in \mathbb{Z}[x_0, \dots, x_k]$  [18, Lemma 4, pp. 5–6]. We say that an integer sequence has *fixed-block nonconcentration* if it satisfies this conclusion for every  $k \geq 0$  and every such  $F$ . A bounded integer perturbation preserves the property: there are only finitely many translated polynomials to consider for each fixed block. No randomness or independence assumption is required.

This nonconcentration condition excludes every bounded integer sequence: if its values lie in a finite set  $E$ , the nonzero polynomial  $\prod_{c \in E} (x_0 - c)$  vanishes at every index. It also excludes a sequence satisfying a fixed nonzero polynomial relation on every short block, such as  $a_n = n$ , for which  $x_1 - x_0 - 1$  vanishes. The density-zero exceptional set depends on the fixed polynomial. There cannot be one such set for all polynomials:  $x_0 - a_n$  vanishes on the block beginning at  $n$ , so a common exceptional set would contain every index. No uniform estimate for growing families of polynomials or block lengths is asserted.

In the next proposition,  $N, H$  are nonnegative integers,  $h$  is an integer, and all counts are over integer indices and integer triples.

**Proposition 8.2** (finite counting for shifted gap differences). *Write  $p_n$  for the primes indexed from  $p_0 = 2$  and  $g_n = p_{n+1} - p_n$ . For  $h \geq 2$  and  $r \in \mathbb{Z}$ , let  $M_{h,r}(N)$  count  $n < N$  with  $g_{n+h} - g_n = r$ . Let  $Q_{N,H,r}$  count triples  $(x, d, s)$  with  $x < p_N$ ,  $0 < d < s \leq H$ ,  $d + r > 0$ , and all four integers  $x, x + d, x + s, x + s + d + r$  prime. Then, for every  $N, H \geq 0$ ,*

$$(H + 1)M_{h,r}(N) \leq (h + 1)p_{N+h+1} + (H + 1)Q_{N,H,r}.$$

*Proof.* For a counted index  $n$ , put  $x = p_n$ ,  $d = g_n$  and  $s = p_{n+h} - p_n$ . If the total span  $p_{n+h+1} - p_n$  is at most  $H$ , then  $0 < d < s \leq H$ ,  $d + r = g_{n+h} > 0$ , and the four primes are precisely  $x, x + d, x + s, x + s + d + r$ . The map is injective because  $x = p_n$  determines  $n$ . For the remaining indices the integer span is at least  $H + 1$ . Each gap appears in at most  $h + 1$  of the spans, so their total is at most  $(h + 1) \sum_{i < N+h} g_i < (h + 1)p_{N+h+1}$ . Multiply the resulting bound on the exceptional count by  $H + 1$  and add the small-span contribution.  $\square$

The displayed finite shifted-count bound is unconditional by the elementary proof above. To obtain zero density from this particular finite bound, one would need, for every  $\varepsilon > 0$  and all large  $N$ , a choice of  $H$  making its right side less than  $\varepsilon N(H + 1)$ . For  $h = 1$ , the same definitions give  $s = d$ , so  $x + d = x + s$ . The strict condition  $d < s$  in the four-prime count fails; that case requires a three-prime count instead. Zero density itself is already known for every fixed  $h \geq 1$  and  $r \in \mathbb{Z}$ : apply Schläge-Puchta's lemma above to the nonzero polynomial  $x_h - x_0 - r$ . Thus the finite reduction should not be read as leaving that upper bound open. Neither upper bound supplies the late occurrences or weighted-tail separation required for irrationality.

For the perturbation claim, first suppose that  $a$  has fixed-block nonconcentration and every correction is either 0 or a fixed integer  $M \geq 1$ . For a fixed  $r \in \mathbb{Z}$ , an equality  $b_{n+1} - b_n = r$  then requires  $a_{n+1} - a_n \in \{r - M, r, r + M\}$ . Each of these three equalities holds on a set of density zero. The general argument uses the same finite-union step for translated polynomials.

**Theorem 8.3** (nonconcentration is perturbation-stable). *Let  $a : \mathbb{N} \rightarrow \mathbb{Z}$  have fixed-block nonconcentration, let  $E \subset \mathbb{Z}$  be finite, and let  $b_n = a_n + e_n$  with  $e_n \in E$  for every  $n$ . Then  $b$  has fixed-block nonconcentration.*

*Proof.* Fix  $k \geq 0$  and a nonzero  $F \in \mathbb{Z}[x_0, \dots, x_k]$ . For a tuple  $\mathbf{e} = (e^{(0)}, \dots, e^{(k)}) \in E^{k+1}$  put  $F_{\mathbf{e}}(x_0, \dots, x_k) = F(x_0 + e^{(0)}, \dots, x_k + e^{(k)})$ . The substitution  $x_i \mapsto x_i + e^{(i)}$  is a ring automorphism of  $\mathbb{Z}[x_0, \dots, x_k]$  with inverse  $x_i \mapsto x_i - e^{(i)}$ , so  $F_{\mathbf{e}} \neq 0$ . If  $F(b_n, \dots, b_{n+k}) = 0$  then  $F_{\mathbf{e}}(a_n, \dots, a_{n+k}) = 0$  for the tuple  $\mathbf{e} = (e_n, \dots, e_{n+k})$ , whence

$$\{n : F(b_n, \dots, b_{n+k}) = 0\} \subseteq \bigcup_{\mathbf{e} \in E^{k+1}} \{n : F_{\mathbf{e}}(a_n, \dots, a_{n+k}) = 0\}.$$

Each set on the right has density zero by hypothesis, and there are only finitely many such sets because  $E^{k+1}$  is finite. Their union therefore has density zero.  $\square$

Finiteness of the correction set is essential to this argument; no independence between the corrections and the sequence is needed. Without a restriction on the corrections, the choice  $e_n = -a_n$  would give  $b_n = 0$ , which fails nonconcentration even for the polynomial  $x_0$ .

**Corollary 8.4** (nonconcentration does not force irrationality). *Fix  $M \geq 1$  and  $K \geq 0$ , and let  $b$  be the perturbed sequence supplied by Theorem 8.1 at the actual prime gaps. Then  $\sum_{n \geq 0} b_n 2^{-(n+1)}$  is rational,  $b_n = g_n$  for  $n < K$ ,  $b_n - g_n \in \{0, M\}$  and  $b_n \equiv g_n \pmod{M}$  for every  $n$ ,  $b$  has fixed-block nonconcentration, and the cumulative sequence  $P_n = 2 + \sum_{i < n} b_i$  satisfies  $p_n \leq P_n \leq p_n + Mn$  and hence  $P_n \sim n \log n$ .*

The Lean proof assumes Schlage-Puchta's Lemma 4 [18] and the prime number theorem.

*Proof.* Theorem 8.1 gives the rational sum, prefix and congruence assertions. Summing  $0 \leq b_i - g_i \leq M$  and using  $\sum_{i < n} g_i = p_n - 2$  gives  $p_n \leq P_n \leq p_n + Mn$ . Now  $p_n \sim n \log n$  [17, Chapter 6] implies  $P_n \sim n \log n$ . The perturbation takes values in the fixed two-element set  $E = \{0, M\}$ , so Theorem 8.3 applies to the actual gaps, which have fixed-block nonconcentration by Schlage-Puchta's lemma [18, Lemma 4].  $\square$

Fixed-block polynomial nonconcentration, taken together with a prescribed finite prefix of actual gaps, a pointwise bound  $b_n \leq g_n + M$ , every residue modulo  $M$ , positivity and cumulative growth at the prime-number-theorem scale, is therefore compatible with a rational dyadic value. Taking  $M$  even also preserves eventual evenness. To preserve a prescribed modulus  $q$  together with parity, take  $M = 2q$ , with the corresponding pointwise bound  $b_n \leq g_n + 2q$ . The terms  $P_n$  are not asserted to be prime. The nonconcentration conclusion is for each fixed polynomial in a fixed block; it gives no uniform bound for polynomial families or block lengths growing with the index.

**Proposition 8.5** (nonconcentration under sparse changes). *Let  $a, b : \mathbb{N} \rightarrow \mathbb{Z}$  agree off a set  $S$  of ordinary density zero. If  $a$  has fixed-block nonconcentration, then so does  $b$ . No boundedness assumption on  $a - b$  is needed.*

*Proof.* Fix  $k$  and a nonzero  $F \in \mathbb{Z}[x_0, \dots, x_k]$ . The two blocks agree unless  $n + i \in S$  for some  $0 \leq i \leq k$ . Consequently

$$\begin{aligned} & |\{n < N : F(b_n, \dots, b_{n+k}) = 0\}| \\ & \leq |\{n < N : F(a_n, \dots, a_{n+k}) = 0\}| + \sum_{i=0}^k |S \cap [i, N + i]|. \end{aligned}$$

The first count is  $o(N)$  by nonconcentration for  $a$ ; the remaining finitely many counts are  $o(N)$  because  $S$  has density zero. Only the number of changed blocks enters, not the sizes of the corrections or of the polynomial values. This proves the assertion for the already fixed sequence  $b$  and every fixed  $F, k$ , without a uniform estimate for growing polynomial families or block lengths.  $\square$

**Corollary 8.6** (a rational sum with the stated prime-gap statistics). *For every prescribed finite prime-gap prefix and  $0 < \varepsilon \leq 1$ , there is an altered sequence  $b = g + e$ , with  $P_n = 2 + \sum_{i < n} b_i$ , for which one can simultaneously impose a rational dyadic value, nonnegative integer corrections eventually at most  $(\log(n+3))^\varepsilon$ , all fixed eventual coefficient and cumulative congruences, fixed-block polynomial nonconcentration, and vanishing total variation distance between unnormalised block distributions for lengths  $o(\log \log X)$ . The cumulative positions satisfy*

$$0 \leq P_n - p_n = O_\varepsilon \left( \frac{n(\log(n+3))^\varepsilon}{\log \log n} \right), \quad P_n \sim n \log n.$$

Lean proves this for every  $\varepsilon > 0$ , assuming Schlage-Puchta's Lemma 4 [18] only for the nonconcentration clause and the prime number theorem only for  $P_n \sim n \log n$ .

*Proof.* Apply Theorem 2.1 at a rational target. Proposition 8.5 transfers the nonconcentration supplied by Schlage-Puchta's Lemma 4 [18] to the corrected sequence. To obtain the cumulative support bound, split  $[\sqrt{n}, n)$  into dyadic intervals, use  $\log \log x \asymp \log \log n$  there, and bound the first  $\sqrt{n}$  indices trivially. Thus  $|S \cap [0, n)| = O(n / \log \log n)$ . The pointwise correction bound gives the displayed estimate, including a fixed constant for the exceptional prefix. Since  $(\log n)^{\varepsilon-1} / \log \log n \rightarrow 0$  for  $0 < \varepsilon \leq 1$ , the prime number theorem [17, Chapter 6] gives the last conclusion. The position series is rational as well: reversing the order of summation of nonnegative terms gives

$$\sum_{n \geq 0} \frac{P_n}{2^{n+1}} = 2 + \sum_{i \geq 0} b_i \sum_{n > i} 2^{-n-1} = 2 + \sum_{i \geq 0} \frac{b_i}{2^{i+1}} \in \mathbb{Q}.$$

The finite value on the right also proves convergence on the left.  $\square$

These conclusions follow from the sparse theorem, Schlage-Puchta's Lemma 4 and the prime number theorem. In fact, all conclusions also hold for every  $\varepsilon > 1$ : apply the corollary with exponent 1. Its correction is smaller than the prescribed bound, and its cumulative estimate  $O(n \log(n+3) / \log \log n)$  is stronger than the displayed bound and still gives  $P_n \sim n \log n$ . Thus the stated range  $0 < \varepsilon \leq 1$  suffices to obtain every positive exponent in the correction bound; no larger correction is required.

The congruences imply that for every fixed integer  $y \geq 2$ , all sufficiently late  $P_n$  have no prime divisor at most  $y$ : use modulus  $y!$  and  $p_n > y$  to get  $\gcd(P_n, y!) = 1$ . This is stronger than eventual oddness, but it does not establish primality. Excluding divisors up to  $\sqrt{P_n}$  would require a bound growing with  $n$ , whereas the congruence cutoff depends on the fixed  $y$ . Even primality of every  $P_n$  would not identify the sequence with the consecutive primes, since a prime-valued sequence may skip primes.

The two preservation arguments are different. The bounded construction uses a finite set of translated polynomials to preserve nonconcentration; it need not preserve empirical block frequencies. The sparse construction instead compares blocks at the same indices, most of which are unchanged. Its total variation bound is absolute, so it need not preserve relative frequencies of rare events. Moreover, its vanishing-error range is  $m = o(\log \log X)$ , not  $m$  comparable to  $\log \log X$ . Neither comparison establishes preservation of the quantitative prime-pattern hypotheses of the conditional irrationality and normality results discussed in Section 3.

### 8.3 Density of indices with two small tail differences

When the two tail differences have absolute value less than one and the gaps are unequal, the calculation above restricts the gap difference to 2 or  $-2$ . For each fixed shift, Schlage-Puchta's lemma gives density zero for both events. We use this to bound the set of possible witnesses, without excluding witnesses at arbitrarily large indices.

**Theorem 8.7** (density zero for the small-difference condition). *Fix  $h \geq 1$ . The set of  $N \geq 1$  at which the three hypotheses of Theorem 6.1 hold for the actual prime gaps has density zero. For the same  $h$ , the set of  $N$  with  $g_{N+h+1} = g_{N+1}$  also has density zero.*

The Lean proof assumes Schlage-Puchta's Lemma 4 [18].

*Proof.* For  $N \geq 1$  every gap involved is even, so  $\delta_N = g_{N+h+1} - g_{N+1}$  is even. If  $|\sigma_h(N)| < 1$ ,  $|\sigma_h(N+1)| < 1$  and  $\delta_N \neq 0$ , then  $\delta_N = 2\sigma_h(N) - \sigma_h(N+1)$  lies in  $(-3, 3)$ , so  $\delta_N = \pm 2$ . Apply Schlage-Puchta's lemma [18, Lemma 4] at index  $n = N + 1$  with  $k = h$  to the two polynomials  $x_h - x_0 - 2$  and  $x_h - x_0 + 2$ , neither of which vanishes identically: each of  $\{N : \delta_N = 2\}$  and  $\{N : \delta_N = -2\}$  has density zero, and the event is contained in their union. The second assertion is the same lemma applied to  $x_h - x_0$ .  $\square$

Unequal gaps occur at almost every index, a stronger conclusion than Proposition 6.3. But combining unequal gaps with the two small tail differences forces their difference to be exactly 2 or  $-2$ , which occurs only on a density-zero set. Thus a proof of Problem 10.2 must find arbitrarily large successful indices in that sparse set. A positive-density lower bound for the same event is impossible. An averaging argument that detects a sparse set is not ruled out.

The proof of Schlage-Puchta's lemma gives a quantitative upper bound as well. For fixed  $h$ , it bounds the number of eligible indices in  $[X, 2X]$  by  $O_h(X / \log \log X)$ . To see the two contributions, discard starts whose block of  $h + 1$  gaps contains a gap exceeding  $\log X \log \log X$ . The sum of these block sums is  $O_h(X \log X)$  by the prime number theorem, so there are at most  $O_h(X / \log \log X)$  discarded starts. For the remaining starts, the sieve calculation in the cited proof bounds the zeros of each of  $x_h - x_0 - 2$  and  $x_h - x_0 + 2$  by

$$O_h\left(\frac{X(\log \log X)^{2h+2}}{\log X}\right) = o_h\left(\frac{X}{\log \log X}\right).$$

The shift from  $n = N + 1$  changes only the band endpoints. Thus the same upper bound applies to the indices satisfying the three conditions of Theorem 6.1. This is an ordinary consequence of the cited sieve argument, not a new lower bound or an asymptotic formula for the number of successful indices.

The observed proportions in Section 9 decrease across the sampled bands. These finite observations establish neither an asymptotic rate nor the required existence beyond every cutoff. Counting by prime size and counting by gap index also give different cut-offs and must not be compared without converting between them.

## 8.4 Recurring gap values differing by two do not suffice

Theorem 6.4 needs values differing by 2 at a prescribed distance  $h$  and a complete tail difference in a signed interval. Recurrence of two values in a residue class supplies neither requirement. We demonstrate this with an integer sequence whose dyadic sum is rational and whose complete tail differences are integral, while retaining the recurrence and growth properties stated below. This is not a sequence of actual prime gaps.

The mechanism is to choose integer tails first, then recover the coefficients from  $a_n = 2U_{n-1} - U_n$ . Constant tails  $U_n = 4$  give constant coefficients  $a_n = 4$ . Raising one isolated tail to 6 changes the two affected coefficients from 4, 4 to 2, 8, with the same combined dyadic contribution. A slowly growing even background will supply cumulative growth  $n \log n$ ; factorial indices will impose the recurrence in residue classes while keeping these changes sparse.

**Theorem 8.8** (recurring values are not enough). *There is a sequence  $(a_n)_{n \geq 1}$  of positive even integers with the following properties. The values 2 and 4 each occur infinitely often at indices divisible by every fixed  $t \geq 1$ . The sequence is unbounded, not eventually periodic, and satisfies  $a_n = O(\log n)$ . The series  $\sum_{n \geq 1} a_n 2^{-n}$  equals 6, and every scaled tail  $\sum_{j \geq 1} a_{N+j} 2^{-j}$  is an integer, so every tail shift is integral. The increasing odd sequence  $P_n = 3 + \sum_{j \leq n} a_j$  satisfies  $P_n \sim n \log n$ .*

*Proof.* Put  $b_n = 2\lceil \frac{1}{2} \log(n+64) \rceil$ , so that  $b_n$  is even,  $b_n \geq 6$ , and  $b_n = \log(n+64) + O(1)$ . Since  $\frac{1}{2} \log(n+65) - \frac{1}{2} \log(n+63) < 1$  for every  $n \geq 1$ , consecutive increments of  $b$  lie in  $\{0, 2\}$  and  $b_{n+1} - b_{n-1} \leq 2$ . Call an index  $n$  *special* when  $n = k!$  or  $n = 2k!$  for some  $k \geq 5$ , and define

$$U_n = \begin{cases} 2b_{n-1} - 2, & n = k!, k \geq 5, \\ 2b_{n-1} - 4, & n = 2k!, k \geq 5, \\ b_n, & \text{otherwise,} \end{cases} \quad a_n = 2U_{n-1} - U_n \quad (n \geq 1).$$

For  $k \geq 5$  one has  $k! < 2k! < (k+1)!$ , so all special indices are distinct. They are even and therefore never adjacent. Each  $U_n$  is an even integer by construction, so each  $a_n$  is an even integer.

At an ordinary index whose predecessor is also ordinary,  $a_n = 2b_{n-1} - b_n = b_{n-1} - (b_n - b_{n-1}) \geq 6 - 2 = 4$ . At a special index the predecessor is ordinary and  $a_n = 2b_{n-1} - (2b_{n-1} - r) = r$ , which is 2 at  $n = k!$  and 4 at  $n = 2k!$ . Immediately after a special index carrying  $r$ ,

$$a_{n+1} = 2(2b_{n-1} - r) - b_{n+1} \geq 4b_{n-1} - 2r - (b_{n-1} + 2) = 3b_{n-1} - 2r - 2 \geq 8.$$

Every  $a_n$  is therefore a positive even integer, and  $U_n = O(\log n)$  gives  $a_n = O(\log n)$ . For  $k \geq 5$ , the index  $n = k! + 2$  and its predecessor are ordinary: both lie strictly between  $k!$  and  $2k!$ , where there are no special indices. Along these indices  $a_n \geq b_{n-1} - 2 \rightarrow \infty$ . Thus  $(a_n)$  is unbounded and hence not eventually periodic.

The definition  $a_n = 2U_{n-1} - U_n$  is exactly  $a_n 2^{-n} = U_{n-1} 2^{-(n-1)} - U_n 2^{-n}$ , so for every  $N \geq 0$  and  $m \geq 1$

$$\sum_{j=1}^m \frac{a_{N+j}}{2^j} = U_N - \frac{U_{N+m}}{2^m}.$$

Since  $U_n = O(\log n)$  the endpoint tends to zero, so the tail at  $N$  equals the integer  $U_N$ ; at  $N = 0$  the series equals  $U_0 = b_0 = 6$ . Every difference  $U_{N+h} - U_N$  is an integer, so every tail shift is integral.

For each  $t$  and every sufficiently large  $k$  we have  $t \mid k!$ , hence  $t \mid 2k!$ , and  $a_{k!} = 2$  while  $a_{2k!} = 4$ : both values recur infinitely often at indices congruent to 0 modulo  $t$ .

Finally, summing  $a_j = 2U_{j-1} - U_j$  gives  $\sum_{j=1}^n a_j = 2U_0 + \sum_{j=1}^{n-1} U_j - U_n$ . The baseline  $\sum_{j \leq n} b_j = n \log n + O(n)$ . The special indices up to  $n$  number  $O(\log n / \log \log n)$  and each changes the summand by  $O(\log n)$ , so their total contribution is  $O((\log n)^2) = o(n)$ . Hence  $P_n = 3 + \sum_{j \leq n} a_j \sim n \log n$ , and  $P_n$  is odd and increasing.  $\square$

The factorial indices used above give recurrence in the zero residue class. The following separate construction strengthens this to every residue class.

### 8.5 A rational series with recurring values in every residue class

There is a synthetic integer sequence  $(a_n)_{n \geq 1}$  with all of the following properties. Each  $a_n$  is positive and divisible by 2, and

$$a_n \leq 4 \log(n+1) + 24.$$

For every  $t \geq 1$ , every  $0 \leq r < t$ , and every cutoff  $N$ , there are  $i, j \geq N$  such that

$$i \equiv j \equiv r \pmod{t}, \quad a_i = 2, \quad a_j = 4.$$

For every integer  $B$  and cutoff  $N$ , some  $n \geq N$  satisfies  $a_n > B$ . For every  $h \geq 1$  there is no cutoff after which  $a_{n+h} = a_n$  for all  $n$ . Nevertheless, all the complete tails

$$U_N = \sum_{j \geq 1} \frac{a_{N+j}}{2^j} \quad (N \geq 0)$$

are integers,  $U_0 = 6$ , and  $U_{N+h} - U_N \in \mathbb{Z}$  for every  $N, h \geq 0$ . The sequence

$$P_N = 3 + \sum_{j=1}^N a_j$$

is strictly increasing and odd, with  $P_N/(N \log N) \rightarrow 1$ . These are synthetic positions; no  $P_N$  is asserted to be prime.

To obtain both recurring values in every residue class, enumerate the triples  $(t, r, m)$  with  $t \geq 1$ ,  $0 \leq r < t$  and  $m \geq 0$ , each once. At the  $k$ th step we choose a centre  $c_k \equiv r \pmod{t}$ , subject to  $c_0 \geq 100$ ,  $c_{k+1} \geq c_k + 3$  and  $c_k \geq 2^{k^2}$  for  $k \geq 1$ . Each arithmetic progression is unbounded, so these lower bounds can be met recursively. Set  $v_k = 2$  for even  $m$  and  $v_k = 4$  for odd  $m$ . For each  $(t, r)$  both choices then occur infinitely often. Use the even baseline

$$b_n = 6 + 2 \left\lfloor \frac{\log(n+1)}{2} \right\rfloor, \quad U_n = \begin{cases} 2b_{n-1} - v_k, & n = c_k, \\ b_n, & n \notin \{c_k : k \geq 0\}, \end{cases}$$

and define  $a_n = 2U_{n-1} - U_n$ . The separation of the centres will ensure that  $a_{c_k} = v_k$ .

The baseline is even and at least 6. For  $n \geq 1$ , the increase of  $\log(n+1)$  over two consecutive steps is less than 2, so the floor in the definition gives  $b_n - b_{n-1} \in \{0, 2\}$  and  $b_{n+1} - b_{n-1} \leq 2$ . Away from a centre and its successor,  $a_n = 2b_{n-1} - b_n \geq 4$ . At a centre,  $a_{c_k} = v_k$ . Immediately afterwards,

$$a_{c_k+1} = 4b_{c_k-1} - 2v_k - b_{c_k+1} \geq 3b_{c_k-1} - 2v_k - 2 \geq 8.$$

Since the centres are at least three apart, these cases exhaust all indices. Also  $0 < U_n \leq 2b_n$ , so  $a_n \leq 4b_{n-1} \leq 4 \log(n+1) + 24$ . The indices  $n = c_k + 2$  lie outside the centres and their successors, because consecutive centres are at least three apart. Along these indices  $a_n \geq b_{n-1} - 2 \rightarrow \infty$ . Thus the sequence is unbounded and cannot be eventually periodic.

Finite telescoping gives

$$\sum_{j=1}^m a_{N+j} 2^{-j} = U_N - U_{N+m} 2^{-m}.$$

The bound  $U_n = O(\log(n+1))$  makes the last term tend to zero. Hence the complete tail equals the integer  $U_N$ , and its initial value is 6. For the cumulative growth,  $b_n = \log(n+1) + O(1)$  and there are at most  $1 + \sqrt{\log_2 N}$  centres up to  $N$ . Replacing  $b_n$  by  $U_n$  changes its partial sum by  $O((\log N)^{3/2})$ . Therefore

$$\sum_{j=1}^N a_j = 2U_0 + \sum_{j=1}^{N-1} U_j - U_N = N \log N + O(N),$$

which gives the stated asymptotic for  $P_N$ .

The full statement is [kernel-checked in Lean](#).

Thus positivity, evenness, unboundedness, nonperiodicity, logarithmic size, cumulative growth  $N \log N$ , and recurrence of the values 2 and 4 in every residue class are jointly compatible with integral tails. These properties alone cannot prove that a tail difference is nonintegral. The sequence is not asserted to consist of prime gaps: its logarithmic bound in particular excludes the extreme large gaps of the primes. The construction therefore does not address hypotheses that use those extreme gaps.

## 8.6 Polynomial coefficients and telescoping sums

We can construct further rational sums by starting with a polynomial  $U_n$  and defining  $c_{n+1} = 2U_n - U_{n+1}$ . Polynomial growth gives the vanishing dyadic boundary condition, so the recurrence identifies  $U_n$  as the complete tail. A quadratic choice already makes the coefficients positive and increasing.

**Proposition 8.9** (a quadratic sequence with integral tails). *Put  $c_n = 2(n^2 + 4n + 2)$  and  $U_n = 2(n+4)^2$ . Then  $c_n$  is positive, even and strictly increasing,  $U_{n+1} = 2U_n - c_{n+1}$ , every shift  $U_{N+h} - U_N$  is integral,  $c_{n+1} - c_n = 4n + 10$  is never  $\pm 2$ , and*

$$\sum_{j \geq 1} \frac{c_j}{2^j} = 32.$$

*Proof.* Direct expansion gives the recurrence, and the finite telescope is  $\sum_{j=1}^n c_j 2^{-j} = 32 - 2(n+4)^2 2^{-n}$ , whose last term tends to zero. The remaining assertions follow from the integer values of  $U_n$  and from  $c_{n+1} - c_n = 4n + 10 \geq 10$ .  $\square$

The series starts at  $j = 1$  and equals the initial rescaled tail  $U_0 = 32$ . Under the opening convention its sum is  $\sum_{n \geq 0} c_n 2^{-(n+1)} = (c_0 + 32)/2 = 18$ . Positivity, parity, strict growth, unboundedness and nonperiodicity are therefore jointly compatible with rationality, and the adjacent-gap condition of Theorem 6.4 fails at every index. The telescoping mechanism is the one used in the note of ChatGPT 5.4 Pro (orchestrated by Vjeko Kovač) against the variable-denominator expectation of Erdős [24, Theorem 1 and proof, pp. 1–2]; the sequence is not a result about Problem #251.

Rationality also fails to force the coefficients to repeat. Let  $K : \mathbb{N} \rightarrow \mathbb{Q}$  be arbitrary and put  $\kappa_n = 2K_n - K_{n+1}$ : the [coefficient of the telescoping series](#).

**Proposition 8.10** (exact telescoping). *For every  $n \geq 0$ ,  $\sum_{i=0}^{n-1} \kappa_i 2^{-(i+1)} = K_0 - K_n 2^{-n}$ .*

*Proof.* The sum is empty when  $n = 0$ . Passing from  $n$  to  $n+1$  adds  $(2K_n - K_{n+1})2^{-(n+1)} = K_n 2^{-n} - K_{n+1} 2^{-(n+1)}$ , which replaces the terminal term by the required one.  $\square$

The identity  $\kappa_n = 2K_n - K_{n+1}$  has the algebraic form used in the rationality criterion of Erdős and Straus when  $a_n = 2$ . That criterion has additional hypotheses: for integers  $b_n$  and positive integers  $a_n$  with  $a_n > 1$  for all large  $n$  and  $|b_n|/(a_{n-1}a_n) \rightarrow 0$ , the series  $\sum_n b_n/(a_1 \cdots a_n)$  is rational exactly when some positive integer  $B$  and integers  $c_n$  satisfy  $Bb_n = c_n a_n - c_{n+1}$  and  $|c_{n+1}| < a_n/2$  for all large  $n$  [7, Theorem 2.1, pp. 85–86]. With  $a_n = 2$ , the smallness hypothesis becomes  $|b_n|/4 \rightarrow 0$ . Since the  $b_n$  are integers, this forces  $b_n = 0$  eventually. Thus the criterion in this specialisation does not apply even to a bounded integer sequence that is nonzero infinitely often, let alone to the positive prime gaps. Proposition 8.10, by contrast, imposes no growth condition on  $K$ . The infinite series converges precisely when  $2^{-n}K_n$  has a finite limit, and then its sum is  $K_0 - \lim_{n \rightarrow \infty} 2^{-n}K_n$ . The sum equals  $K_0$  precisely when that limit is zero; polynomial growth suffices. For  $K_n = 2^n$ , every  $\kappa_n$  is zero, so the series converges to 0, not to  $K_0 = 1$ .

Taking  $K_0 = \frac{5}{2}$  and  $K_n = 2n + 2$  for  $n \geq 1$  gives  $\kappa_0 = 1$  and  $\kappa_n = 2n$ , so the coefficients  $1, 2, 4, 6, 8, \dots$  are positive, even after the first term, unbounded and not eventually periodic, while their dyadic sum is  $\frac{5}{2}$ . Rationality alone therefore cannot imply eventual periodicity even for a positive, parity-correct integer coefficient sequence, and rationality cannot be contradicted merely by Proposition 6.3.

The arithmetic-progression formulations of nonintegrality do not create new prime-distribution information. Their full definitions and exact hypotheses are preserved in Appendix E.

## 9 Finite numerical searches

The scans below evaluate prime-gap tails in double precision over the reported finite ranges. No bound for accumulated rounding error was supplied, so their hits and counts remain numerical observations even when the omitted infinite tail has a rigorous bound. For exact comparisons, Proposition 6.6 and the certificates in Section 7 use integer arithmetic and have linked Lean checks.

Over the 6 841 648 primes below  $1.2 \times 10^8$  and offsets  $h = 1, \dots, 16$ , the scan records indices satisfying the equivalent signed inequalities in Theorem 6.4 at every tested offset, with observed proportions between 0.00418 and 0.008248 and nearly balanced signs. At  $h = 1$  it records 56 427 hits among 6 841 564 candidate indices, the last at the prime 119 995 753. Multiplying the observed proportion in each band by that band's mean of  $\log p$  gives 0.17671 in the first band and 0.13148 in the last. This finite observation establishes no asymptotic counting law and no occurrence beyond every cutoff. The [search output](#) is public.

A second scan uses the 1 270 607 primes below  $2 \times 10^7$  and pairs of indices in the half-open range  $1\,143\,545 \leq n, m < 1\,270\,540$ . It searches each residue class modulo each  $1 \leq t \leq 20$  for the condition of Theorem 5.5, selecting pairs with a next-gap difference of 2 and a tail difference in the specified interval. At  $t = 20$ , every residue class has at least 90 150 counted pairs; pairs may share indices. For each class, the program also records

the larger index of a selected late witness. The minimum of those recorded indices is 1 270 520; it need not come from the class with the fewest pairs. For the saved sample pairs, floating-point arithmetic gives both small-difference inequalities. The scan supplies neither arbitrary moduli nor witnesses beyond every prescribed cutoff. Its [search output](#) is public. The [computation guide](#) gives the programs, dependencies and commands to rerun both searches and the 80000-bit continued-fraction calculation of Section 7. Appendix D.7 corrects the stated justification of the tail bound in the free-pair program.

## 10 Remaining prime-gap estimates

For the actual primes, irrationality is equivalent to the nonintegrality condition in Theorem 5.3. We record its quantifiers together with the stronger simultaneous small-pair condition. A finite certificate supplies one witness; these questions require witnesses for every prescribed shift and cutoff.

**Problem 10.1** (nonintegral tail differences of each positive length). For every  $h \geq 1$  and every  $N_0$ , prove that some  $N \geq N_0$  satisfies

$$\sum_{j \geq 1} \frac{g_{N+h+j} - g_{N+j}}{2^j} \notin \mathbb{Z}. \quad (9)$$

**Problem 10.2** (two small tail differences at arbitrarily large indices). For every  $h \geq 1$  and every  $N_0$ , prove that some  $N \geq N_0$  satisfies

$$|\sigma_h(N)| < 1, \quad |\sigma_h(N+1)| < 1, \quad g_{N+h+1} \neq g_{N+1}. \quad (10)$$

Problem 10.1 and the variable-offset criterion of Theorem 5.5 are each equivalent to irrationality. By Corollary 6.2, Problem 10.2 is sufficient; no converse is asserted. These logical reformulations do not supply the required prime-gap estimate.

The counterexamples retain a fixed prefix and modulus (Theorem 8.1), polynomial nonconcentration and cumulative growth  $n \log n$  (Corollary 8.4), or all the congruences and block statistics of the sparse construction. The distinct construction in Section 8.5 makes two values differing by 2 recur in every residue class. Each has a rational sum; none guarantees prime cumulative positions.

For the actual gaps, Theorem 8.7 shows that the event in (10) has density zero for each fixed  $h$ . A proof of its occurrence must therefore find arbitrarily late witnesses inside a sparse set, rather than prove that they occupy a positive proportion. A finite prefix alone cannot do this: it can be retained while the dyadic sum is made rational. A finite block together with a proved bound on the omitted tail can still certify one pair, as in Proposition 6.6. The distinction is between a finite certified instance and an occurrence theorem beyond every cutoff.

Proposition 6.5 gives sufficient integer inequalities ensuring  $|\sigma_h(N)| < 1$  at a chosen truncation length  $L$ . Failure of such a test need not mean that the complete tail lies outside the required interval: its error interval may still cross an endpoint. For fixed  $h$  and  $\varepsilon > 0$ , the choice  $L = \lceil (4 + \varepsilon) \log_2(N + 2) \rceil$  gives  $E_{h,N,L} = O_{h,\varepsilon}(N^{-\varepsilon})$ . Indeed,

$L = O_\varepsilon(\log N)$ , both arguments of the quartic  $P$  are  $O_{h,\varepsilon}(N)$ , and  $2^{-L} \leq (N+2)^{-4-\varepsilon}$ . Requiring a fixed positive margin from both endpoints is stronger than the strict inequalities in (10). At this depth, the test bounds the signed integer  $2^L F_{h,N,L}$ , not just its residue modulo  $2^L$ , and also requires  $g_{N+h+1} - g_{N+1} = \pm 2$ . For fixed  $h$  and  $\varepsilon$ , checking one certificate uses  $O_\varepsilon(\log N)$  gap values and the explicit remainder bound. This counts data, not bit operations. The unproved input is the existence of certified witnesses beyond every cutoff for each fixed  $h$ ; constants uniform in  $h$  are not required.

The following published results control gap sizes and clusters, not the signed weighted sums required by this criterion. Zhang’s bounded-gap theorem [21, Theorem 1, p. 1122] produces infinitely many bounded consecutive-prime gaps. Maynard bounds  $\liminf_n (p_{n+m} - p_n)$  for every fixed  $m$ , giving bounded clusters of every fixed size [16, Theorem 1.1, p. 384], with the explicit unconditional bound  $\liminf_n (p_{n+1} - p_n) \leq 600$  [16, Theorem 1.3, p. 385]. Polymath subsequently improved this to 246 [35, Theorem 1.4(i) of arXiv v4]; neither numerical bound controls the signed weighted tails required here. The large-gap theorem of Ford, Green, Konyagin, Maynard and Tao gives an effective lower bound for the largest single consecutive-prime gap below  $X$  [8, Theorem 1]. A theorem giving bounded clusters at each fixed cluster size does not by itself supply a weighted condition on blocks whose length grows with the basepoint, and none of these results is used as a proof input here.

Conditionally the picture is different. Tao’s comment names uniform quantitative prime-tuples control of about  $\log \log n$  consecutive gaps as a possible approach [23, comment of 7 October 2025], and Land’s draft carries that out under Kuperberg’s uniform prime-tuples conjecture [13]. The counterexamples in Section 8 show that the listed size, congruence and block-statistical properties alone do not force irrationality. These constructions are not shown to enumerate the consecutive primes, so they do not rule out arguments that combine those properties with further information about primes.

**Funding and competing interests.** This work received no external funding. The author declares no competing interests.

**Acknowledgements.** I thank Wouter van Doorn for advice on explaining unfamiliar hypotheses, removing unnecessary terminology, and using notation only when it helps the reader. His comments concerned an earlier note on Problem #243; this acknowledgement does not imply that he reviewed or endorsed the mathematics of the present paper. The problem numbering follows the Erdős Problems catalogue maintained by Thomas Bloom [22]. The logarithmic-scale construction and the transfer of nonconcentration come from earlier working materials of this project, which were not an independent mathematical review.

## A An elementary polynomial bound for the primes

We derive the bound  $p_n \leq 1250(n+1)^4$  by comparing the size and prime factorisation of a central binomial coefficient, following Erdős’s proof of Chebyshev’s theorem [5, §1, pp. 194–196]. With  $\pi(x)$  denoting the number of primes at most  $x$ , we first prove, for

$m \geq 4$ ,

$$4^m < m \binom{2m}{m} \leq m(2m)^{\pi(2m)}. \quad (11)$$

For the first inequality,  $m \binom{2m}{m} / 4^m$  equals  $70/64$  at  $m = 4$  and its ratio at successive indices is  $(2m+1)/(2m) > 1$ . For the second, the exponent of a prime  $\ell$  in  $\binom{2m}{m}$  is  $\sum_{k \geq 1} (\lfloor 2m/\ell^k \rfloor - 2\lfloor m/\ell^k \rfloor)$ , each summand is 0 or 1, and every summand with  $\ell^k > 2m$  vanishes; the exponent is therefore at most  $\lfloor \log_\ell(2m) \rfloor$ . Thus the full power of each prime appearing in  $\binom{2m}{m}$  is at most  $2m$ . Multiplying over at most  $\pi(2m)$  distinct primes gives the second inequality.

Now fix  $n \geq 0$ , put  $x = n + 5$  and  $m = x^4 \geq 625$ , and suppose  $\pi(2m) \leq n$ . Since  $x \leq 2^x$  and  $n + 4(n+1)x = 4x^2 - 15x - 5 \leq 2x^4$ ,

$$m(2m)^n = 2^n x^{4(n+1)} \leq 2^{n+4(n+1)x} \leq 2^{2x^4} = 4^m,$$

contradicting (11). Hence  $\pi(2m) > n$ , so at least  $n+1$  primes lie below  $2m$  and therefore

$$p_n \leq 2(n+5)^4 \leq 1250(n+1)^4,$$

the last step because  $(n+5) \leq 5(n+1)$  for  $n \geq 0$ . The coarser bound  $p_n \leq 1250(n+1)^4$  is checked by the kernel at [line 360](#).

## B Integer certificates

The Python calculation below recomputes the primes by trial division, then checks the two small-pair rows of Proposition 6.6 and every integer inequality in Theorem 7.1. It uses the four certificate integers from the linked Lean proof and performs only integer arithmetic. The adjacent quoted strings are concatenated by Python; they split long integers across printed lines without changing their values.

```
from math import isqrt

primes = [n for n in range(2, 10000)
           if all(n % d for d in range(2, isqrt(n) + 1))]
gaps = [q - p for p, q in zip(primes, primes[1:])]
P = lambda x: x**4 + 8*x**3 + 36*x**2 + 104*x + 150
Q = 2**40
expected = [(-662838684750, 11764181250),
            (873345886050, 12805761250)]
for N, target in zip((2, 3), expected):
    D = sum((gaps[N+1+j] - gaps[N+j])*2**(40-j)
            for j in range(1, 41))
    B = 1250*(P(N+43) + P(N+42))
    distance = min(D % Q, Q - D % Q)
    if (D, B) != target or not (abs(D)+B < Q and B < distance):
        raise ArithmeticError("tail certificate failed")
if gaps[4] == gaps[3]:
    raise ArithmeticError("gap mismatch failed")

u = int(
    "8065641857152652932176019632186898003271162829171466334827"
    "3083607794415278717445033509407855988903369988525550746159"
```

```

"7355889792250084202344821020139160956663658789718168152662"
"0217"
)

v = int(
"2194945124413663232143970924541263312422069524635615360518"
"4247077351958221816830720189289904831662955084392690248683"
"1216291723988537733235173040607254496838530213867781442335"
"1745"
)

up = int(
"6539437101498162626882411892470905222108260008568555445975"
"3026163315521784668689909712759876562484620059098138417469"
"5232839888185316374272333277389611483117334000493867584923"
"912"
)

vp = int(
"1779611075789866551198427241621709630120136323281598176637"
"8490605249229735501968764904036152970729491656650873938580"
"6203025466313389810291243786005499164537499383612081805160"
"767"
)

c = len(primes)
A = sum(p*2**(c-1-i) for i, p in enumerate(primes))
checks = (c == 1229,
          u > 0 and v > 0 and up > 0 and vp > 0,
          up*v - u*vp == 1,
          u*2**c < A*v,
          (2*A + 5000*(c+1)**4)*vp < up*2**(c+1),
          v+vp >= 2**589,
          2**589 > 10**177)
if not all(checks):
    raise ArithmeticError("denominator certificate failed")
print("Both tail rows and the denominator floor verified.")

```

## C Verification and reproducibility

The verification concordance lists the formal proofs by statement. *Lean* links to the supporting declarations; a dagger identifies a proof that assumes a named input. *Comparator* links to a recorded kernel check against a separately written statement; *pending* means that this comparison has not been recorded. The [verification record](#) gives the precise correspondence, dependencies and reproducible checks. A row with only a record link has no complete formal proof recorded. The source links identify definitions and proof steps at fixed revisions; the short paper's source index retains its different revision in [Appendix G](#). The formal sparse construction changes one coordinate per stage, whereas the printed proof uses triples. Both fix a set containing all correction supports, the interval of sums and the modulus cutoffs before choosing a value. The formal proofs use the published inputs stated below; the linked declarations do not check the different printed arguments step by step.

Lean 4 [14] and mathlib [15] supply the proof-checking environment. The [evidence record](#) lists the declarations and checks. The programs for the searches in Section 9 are in the [computation guide](#). The program in Appendix B checks the adjacent pair and the bound  $2^{589}$ ; the independent program of Section 7 checks the certificate of Theorem 7.2. Formal checking does not establish novelty or attribution.

The summation-by-parts identity and the polynomial prime bound concern the actual primes. The recurrence results apply to arbitrary integer coefficients and, as stated, to rational or real sequences; substituting the actual prime-gap tail requires its convergence and boundary condition. The finite certificates use the same polynomial bound to control the omitted terms. The counterexamples concern other coefficient sequences and construct no new primes.

The Lean proofs of three results assume published inputs that the Lean development does not prove. The Lean proof of Theorem 8.7 assumes Schlage-Puchta's Lemma 4 [18, Lemma 4, pp. 5–6]; for a fixed nonzero integer polynomial in a fixed block of consecutive prime gaps, it states that the set of indices where the polynomial vanishes has density zero. Those of Corollaries 8.4 and 8.6 assume that lemma and the prime number theorem in the form  $p_n \sim n \log n$  [17, Chapter 6]. What each statement adds to these inputs is checked, including the estimate that the corrected sequence of Corollary 8.6 differs from the prime gaps at  $O_\varepsilon(n / \log \log n)$  indices below  $n$ , which uses no information about the distribution of primes.

## D Further criteria, examples and computational details

This appendix gives further forms of the recurrence criterion and their proofs. The finite truncation argument includes an example separating refinement of one fixed tail difference from estimates at changing indices. A bounded counterexample and the complete numerical table follow. None of these arguments uses the sparse construction.

### D.1 Totient-length shifts and persistence of integrality

Once the denominator is odd, Euler's congruence specifies an integral shift of totient length. We record this choice and then prove that an integral shift persists under each subsequent recurrence step.

**Proposition D.1** (a shift of totient length). *Let  $T : \mathbb{N} \rightarrow \mathbb{Q}$  satisfy the dyadic tail recurrence with integer coefficients. If the reduced denominator  $d$  of  $T_N$  is odd, then  $\sigma_{\varphi(d)}(N)$  is an integer.*

*Proof.* Since  $d$  is odd, 2 and  $d$  are coprime, so Euler's congruence gives  $d \mid 2^{\varphi(d)} - 1$ . Writing  $2^{\varphi(d)} - 1 = dk$  and  $T_N = u/d$  in lowest terms,  $(2^{\varphi(d)} - 1)T_N = ku$  is an integer, and the integral-shift criterion transfers this to the shift.  $\square$

**Proposition D.2** (propagation). *Let  $T : \mathbb{N} \rightarrow \mathbb{R}$  satisfy  $T_{n+1} = 2T_n - a_{n+1}$  with integer coefficients, and define  $\sigma_h(N) = T_{N+h} - T_N$ . For fixed  $h, N \geq 0$ , if  $\sigma_h(N)$  is an integer, then  $\sigma_h(N+k)$  is an integer for every  $k \geq 0$ .*

*Proof.* The shift step identity gives  $\sigma_h(N+1) = 2\sigma_h(N) - (a_{N+h+1} - a_{N+1})$ , an integer combination of an integer and two coefficients. Induct on  $k$ .  $\square$

These are the totient shift and the propagation theorems cited in the note. For example, if  $\text{den } T_N = 3$  then  $\varphi(3) = 2$  and  $3T_N$  is an integer, so  $\sigma_2(N)$  is integral while  $\sigma_1(N)$  is not; if  $\text{den } T_N = 5$  then  $\sigma_4(N)$  is integral. In the orbit with every coefficient zero and  $T_0 = 1/12$ , the denominator is  $12 = 2^2 \cdot 3$ , the orbit reaches  $T_2 = 1/3$  with odd denominator at  $s = 2$ , and  $h = \varphi(3) = 2$  is exactly the shift length seen to be integral from index 2 onwards. The totient supplies one length, while the exact criterion is  $\text{den}(T_N) \mid 2^h - 1$ . At an index with odd denominator  $d > 1$ , the least positive length is the multiplicative order of 2 modulo  $d$ . When  $d = 1$ , every positive length works; before the denominator becomes odd, no positive length works.

## D.2 The finite truncation criterion

For the actual gaps, we enclose each complete weighted difference by a finite sum and a bound on the omitted terms. A nonnegative summable majorant gives such an enclosure, but need not provide a computable remainder. Proposition 6.5 supplies the rational bound used in the finite tests.

**Proposition D.3** (finite truncation). *Let  $M : \mathbb{N} \rightarrow \mathbb{R}$  satisfy  $M(n) \geq g_n$  for every  $n$  and  $\sum_{n \geq 0} M(n)2^{-n} < \infty$ , and put*

$$S_{h,N,L} = \sum_{j=1}^L \frac{g_{N+h+j} - g_{N+j}}{2^j}, \quad R_{h,N,L}(M) = \sum_{j>L} \frac{M(N+h+j) + M(N+j)}{2^j}.$$

*If for every  $h \geq 1$  and every  $N_0$  there are  $N \geq N_0$  and  $L \geq 1$  with  $\text{dist}(S_{h,N,L}, \mathbb{Z}) > R_{h,N,L}(M)$ , then the nonintegrality condition in Problem 10.1 holds. Consequently  $\Pi$  is irrational.*

*Proof.* The part of  $\sum_{j \geq 1} (g_{N+h+j} - g_{N+j})2^{-j}$  omitted from  $S_{h,N,L}$  has absolute value at most  $R_{h,N,L}(M)$ , so under the displayed inequality the full sum lies at positive distance from every integer.  $\square$

With  $M(n) = 1250(n+2)^4$ , the quantities  $S_{h,N,L}$  and  $R_{h,N,L}(M)$  are exactly  $F_{h,N,L}$  and  $E_{h,N,L}$  from Proposition 6.5. The same enclosure therefore tests separation from the integers, without requiring either of the two consecutive differences to be small. To compute the distance from the truncated sum to  $\mathbb{Z}$ , write the weighted block sum as

$$D_{h,N,L} = \sum_{j=1}^L 2^{L-j} (g_{N+h+j} - g_{N+j}), \quad S_{h,N,L} = \frac{D_{h,N,L}}{2^L}.$$

Then

$$\text{dist}(S_{h,N,L}, \mathbb{Z}) = 2^{-L} \min\{D_{h,N,L} \bmod 2^L, 2^L - (D_{h,N,L} \bmod 2^L)\},$$

where  $D_{h,N,L} \bmod 2^L$  is the least nonnegative residue, including when  $D_{h,N,L} < 0$ . The criterion requires this residue to be more than  $2^L R_{h,N,L}(M)$  from both 0 and  $2^L$ . One successful comparison certifies nonintegrality at that index; a failed comparison is inconclusive. Irrationality requires such certificates arbitrarily late for every positive  $h$ . At prescribed logarithmic depth, the needed residue must be farther from both endpoints than the error bound. With the polynomial majorant in Proposition 6.5, the remainder bound is rational and explicit, so the whole test reduces to integer comparisons. The modular rewriting is an ordinary deduction, not a separately checked Lean declaration.

**Proposition D.4** (completeness of finite separation). *Suppose  $D \in \mathbb{R}$ ,  $S_L \in \mathbb{R}$  and  $R_L \geq 0$  satisfy  $|D - S_L| \leq R_L$  and  $R_L \rightarrow 0$ . Then*

$$D \notin \mathbb{Z} \iff \text{there exists } L \text{ with } \text{dist}(S_L, \mathbb{Z}) > R_L.$$

*Proof.* The reverse implication follows from the triangle inequality. For the forward implication put  $\delta = \text{dist}(D, \mathbb{Z}) > 0$ . For all sufficiently large  $L$ , one has  $R_L < \delta/2$ , and the 1-Lipschitz property of distance gives  $\text{dist}(S_L, \mathbb{Z}) \geq \delta - R_L > R_L$ . No monotonicity of the errors is required.  $\square$

For fixed  $h, N$  and the explicit polynomial remainder, testing  $L = 1, 2, \dots$  therefore stops with a certificate whenever  $\sigma_h(N)$  is nonintegral. An unsuccessful comparison is inconclusive. A proof that every depth fails, unlike a finite unsuccessful run, would imply integrality by the same equivalence. None of this shows that suitable indices occur arbitrarily late. Quantitative estimates are still needed to guarantee separation at a prescribed truncation length as  $N$  varies. For a numerical example unrelated to primes,  $[1 - 2^{-N}, 1 + 2^{-N}]$  contains both the noninteger  $1 + 2^{-2N}$  and the integer 1 for every  $N \geq 1$ . Its radius tends to zero, yet it never certifies the changing value as nonintegral. This does not contradict completeness, which keeps the value fixed while refining its enclosure.

### D.3 Nonintegral differences at multiples of each shift

We may ask for a nonintegral multiple of each prescribed shift length. The following formulation still has the full strength of irrationality, as the argument after the question shows.

**Problem D.5** (multiples of each shift). For every  $r \geq 1$ , is there an  $m \geq 1$  such that the tail differences of length  $mr$  are nonintegral at arbitrarily large indices, as in (9)?

This is an exact reformulation, not a weaker mathematical target. If  $T_0$  is irrational, every positive shift is nonintegral by the block identity, so the stated condition follows with  $m = 1$ . Conversely, rationality gives a shift  $h \geq 1$  that is integral at every sufficiently late index. For each positive integer  $m$ , the identity

$$T_{N+mh} - T_N = \sum_{j=0}^{m-1} (T_{N+(j+1)h} - T_{N+jh})$$

makes every multiple  $mh$  eventually integral. Taking  $r = h$  contradicts the stated condition. This ordinary deduction uses the checked rationality classification and a finite telescope; rearranging the quantifiers supplies no new estimate for the actual prime gaps.

### D.4 Repeated tail values and congruent indices

We take  $T_N = \sum_{j \geq 1} g_{N+j} 2^{-j}$  to be the complete actual prime-gap tail. Under rationality, the prime number theorem and a counting argument give many repeated tail values. We then compare these equalities with the nonintegral differences required at congruent indices. These are ordinary deductions in the text. The linked formal results establish the recurrence and the congruent-index criterion.

**Repeated tail values.** For an integer  $X \geq 0$ , summing  $g_{N+1} = 2T_N - T_{N+1}$  over  $0 \leq N < X$  gives the exact identity

$$\sum_{N=0}^X T_N = p_{X+1} - p_1 - T_0 + 2T_X \leq (p_{X+1} - p_0) + 2T_X.$$

The inequality uses  $T_0 \geq 0$  and  $p_1 \geq p_0$ . The prime number theorem [17, Ch. 6] also gives

$$\frac{T_X}{p_X} = \sum_{j \geq 1} \frac{g_{X+j}}{2^j p_X} \rightarrow 0. \quad (12)$$

Indeed, for each fixed  $j$ , both  $p_{X+j+1}/p_X$  and  $p_{X+j}/p_X$  tend to 1, so their difference  $g_{X+j}/p_X$  tends to zero. For the uniform bound needed to pass the limit through the sum, the two-sided estimates  $p_n \asymp n \log n$  give one constant  $K$  such that, for  $X \geq 2$  and  $j \geq 1$ ,

$$\begin{aligned} \frac{g_{X+j}}{p_X} &\leq K \frac{(X+j+1) \log(X+j+1)}{X \log X} \\ &\leq K(1+j) \left( 1 + \frac{\log(1+j)}{\log X} \right) \leq K(1+j)^2. \end{aligned}$$

The second line uses  $X+j+1 \leq X(1+j)$  and  $\log(1+j) \leq j \log 2 \leq j \log X$ . Since  $\sum_{j \geq 1} (1+j)^2 2^{-j} < \infty$ , dominated convergence applies. Subtracting the exact summation identities at  $2X$  and  $X$  now gives

$$\sum_{X < N \leq 2X} T_N = p_{2X+1} - p_{X+1} + 2(T_{2X} - T_X) \sim X \log X.$$

Indeed, the prime difference is asymptotic to  $X \log X$ , and (12) makes both tail terms  $o(X \log X)$ . Thus the mean complete tail in this band is asymptotic to  $\log X$ , without a rationality assumption. This is a mean, not a pointwise estimate. For the counting argument we only need the upper bound  $\sum_{X < N \leq 2X} T_N \leq K_0 X \log X$ , valid for any fixed  $K_0 > 1$  and all sufficiently large  $X$ . For each fixed  $C > 1$ , Markov's inequality then leaves at least  $(1 - 1/C)X$  indices with  $T_N \leq K_0 C \log X$ .

Under rationality, take  $X$  beyond the point where the reduced denominator is the fixed odd integer  $d$ . Among the indices just selected by Markov's inequality, the values lie in  $d^{-1} \mathbb{Z}_{\geq 0} \cap [0, K_0 C \log X]$ . There are at most  $dK_0 C \log X + 1$  such values. Pigeonhole therefore gives one value occurring at least

$$\frac{(1 - 1/C)X}{dK_0 C \log X + 1}$$

times. Taking, for example,  $K_0 = 2$  gives a lower bound of order  $\gg_{C,d} X / \log X$ . The value may depend on  $X$ ; no single value recurring in every band is established.

This count uses the growth of the actual primes. It is false for a general rational integer-coefficient recurrence: the quadratic example in Proposition 8.9 has strictly increasing tails  $T_N = 2(N+4)^2$ . Even for the primes, equal-tail pairs have difference zero and hence do not supply the nonintegral difference required by the criterion using congruent indices.

The terminal term in the exact identity cannot simply be omitted. The factorial-gap argument in the proof of Proposition 6.3 shows that  $T_X \geq g_{X+1}/2$  is unbounded, whereas  $p_1 + T_0$  is fixed. The exact identity therefore gives  $\sum_{N=0}^X T_N > p_{X+1}$  for arbitrarily large  $X$ . It does not establish that this strict inequality holds eventually.

**The interval condition at congruent indices.** Suppose the sum is rational, let  $d$  be the eventual odd reduced denominator, and let  $t$  be the order of 2 modulo  $d$ , with  $t = 1$  when  $d = 1$ . Beyond the denominator cutoff,  $M \equiv N \pmod{t}$  forces  $T_M - T_N \in \mathbb{Z}$ . Neither this difference nor its negative can then lie in  $(\frac{1}{2}, 1)$ . Thus a witness in that interval, at two such congruent indices, would contradict rationality without a further condition on the intervening gaps. This uses only the interval occurring in Theorem 6.4; it does not assert that its other hypotheses hold for arbitrary congruent pairs.

## D.5 A bounded sequence with recurring values and a rational sum

The following bounded example isolates the recurring-values argument without a growth estimate. Logarithmic growth was needed in Theorem 8.8 to obtain cumulative size  $n \log n$ , but is unnecessary for the recurrence itself.

**Proposition D.6** (recurring values with integral tails). *Put  $U_0 = 4$  and, for  $n \geq 1$ ,  $U_n = 6$  when  $n = k!$  for some  $k \geq 3$  and  $U_n = 4$  otherwise, and set  $a_n = 2U_{n-1} - U_n$  for  $n \geq 1$ . Then  $a_n \in \{2, 4, 8\}$ . For every  $k \geq 3$ , the value 2 occurs at index  $k!$  and the value 4 at index  $2k!$ , so both recur infinitely often at indices divisible by any fixed  $t \geq 1$ . The series  $\sum_{n \geq 1} a_n 2^{-n}$  equals 4 and every tail  $\sum_{j \geq 1} a_{N+j} 2^{-j}$  equals the integer  $U_N$ .*

*Proof.* For  $k \geq 3$  the index  $k!$  is even and at least 6, and  $k! - 1$  is odd, so the preceding tail is 4; hence  $a_{k!} = 8 - 6 = 2$ ,  $a_{k!+1} = 12 - 4 = 8$ , and  $a_n = 8 - 4 = 4$  elsewhere. For  $k \geq 2$  one has  $k! < 2k! < (k+1)!$ , so  $2k!$  is not a factorial. Its odd predecessor  $2k! - 1 \geq 3$  is not a factorial either; therefore  $a_{2k!} = 4$ . Since  $t \mid k!$  for every large  $k$ , both values recur in the residue class 0 modulo  $t$ . The identity  $a_n 2^{-n} = U_{n-1} 2^{-(n-1)} - U_n 2^{-n}$  telescopes to  $\sum_{j=1}^m a_{N+j} 2^{-j} = U_N - U_{N+m} 2^{-m}$ , and  $U$  is bounded, so the endpoint vanishes.  $\square$

The indices with coefficient 2 are precisely the factorials  $k!$ ,  $k \geq 3$ . They occur infinitely often with unbounded gaps, which is impossible for a value recurring in an eventually periodic sequence. Thus even this bounded coefficient sequence is nonperiodic, despite its rational dyadic sum. Its cumulative sum has linear growth. The logarithmic construction in Theorem 8.8 adds unboundedness and cumulative growth  $N \log N$  without changing the telescoping mechanism. Neither construction guarantees prime cumulative positions.

## D.6 Numerical counts and proportions

The scan in Section 9 used the 6 841 648 primes below  $1.2 \times 10^8$  and computed double-precision tails for offsets  $h \leq 16$ . We reproduce its reported counts for  $h = 1, \dots, 4$ . The denominator of each proportion is the number of tested basepoints, 6 841 565  $- h$ , rather than the prime count.

| $h$ | events | proportion | $\delta = +2$ | $\delta = -2$ | last event at prime |
|-----|--------|------------|---------------|---------------|---------------------|
| 1   | 56 427 | 0.008248   | 28 022        | 28 405        | 119 995 753         |
| 2   | 31 979 | 0.004674   | 15 742        | 16 237        | 119 993 807         |
| 3   | 30 233 | 0.004419   | 15 093        | 15 140        | 119 998 321         |
| 4   | 29 262 | 0.004277   | 14 606        | 14 656        | 119 993 473         |

The scan records hits at every tested offset  $h \leq 16$ . The observed proportions range from 0.00418 to 0.008248, with nearly balanced signs at each offset. For  $h = 1$ , the first row below gives the proportions in the eight bands; the second multiplies each by its band's mean of  $\log p$ :

0.011285, 0.008951, 0.008303, 0.007936, 0.007651, 0.007507, 0.007253, 0.007094;  
0.17671, 0.15058, 0.14420, 0.14067, 0.13766, 0.13667, 0.13333, 0.13148.

The ratio of the last rescaled proportion to the first is 0.744 at  $h = 1$  and lies between 0.744 and 0.8121 across the measured offsets. Theorem 8.7 gives limiting density zero for each fixed offset. The bound  $O_h(X/\log \log X)$  derived after its proof is only an upper bound; it predicts neither a monotone decline nor an asymptotic proportion. The finite-band ratios therefore do not establish an asymptotic rate.

## D.7 Two corrections

We distinguish a bound for the actual tail from the polynomial series used to compute it, and then correct an insufficient offset condition.

**The tail bound in the free-pair program.** The 4096-bit free-pair program linked through Section 9 bounds  $T_s$  by summing the first 400 terms of  $1250(s + j + 2)^{42-j}$  after rounding each down, then adding 1. This does not necessarily bound that polynomial series: at the recorded  $s = 1\,270\,306$ , the returned value is 40 below the exact sum  $1250P(s + 2)$ , with  $P$  as in Proposition 6.5.

The returned bound nevertheless has a valid justification. The stronger estimate  $p_n \leq 2(n + 5)^4$  proved in Appendix A gives

$$T_s \leq 2P(s + 6) < 625(s + 3)^4 \quad (s \geq 0).$$

The last expression is already the first summand of the program, and all its remaining summands are nonnegative. The strict inequality follows from

$$625(s + 3)^4 - 2P(s + 6) = 623s^4 + 7436s^3 + 32958s^2 + 62972s + 40437 > 0.$$

Thus the rounding issue is in the stated majorant argument, not a failure of the returned enclosure. This argument justifies the analytic bound; it is not a rerun of the denominator-exclusion search.

**An insufficient condition.** Theorem 5.5 allows the offset to vary. For each modulus  $t$ , one must still find pairs of congruent indices beyond every prescribed cutoff whose tail difference is nonintegral. Two even coefficient values differing by 2, each occurring infinitely often in the same residue class of indices, do not give the required tail difference for an integer-coefficient recurrence: the sequence of Theorem 8.8 has both properties and only integral tail shifts.

## E Arithmetic-progression reformulations of integrality

We compare an iterated tail difference with an arithmetic progression determined by the intervening coefficients. The recurrence reduces membership in this progression to integrality of the original difference. Under the stated growth bound, the fixed progression  $2^r \mathbb{Z}$  gives the same criterion. The two reductions are recorded separately below.

Fix  $h$  and abbreviate  $D_N = \sigma_h(N)$  and  $\delta_N = g_{N+h+1} - g_{N+1}$ . Iterating over a longer block gives

$$B_{h,N,r} = \sum_{i=0}^{r-1} 2^{r-1-i} \delta_{N+i}; \quad D_{N+r} = 2^r D_N - B_{h,N,r}.$$

Both the progression  $-B_{h,N,r} + 2^{r+1} \mathbb{Z}$  and the value  $D_{N+r}$  contain the same integer offset. Substituting the recurrence cancels it, leaving a condition on  $D_N$  alone. Thus increasing the block length in this particular test adds no arithmetic information.

**Theorem E.1** (equivalent arithmetic-progression tests). *For every rational dyadic tail recurrence and all  $h, N, r \geq 0$ ,*

$$D_{N+r} \in -B_{h,N,r} + 2^{r+1} \mathbb{Z} \iff D_N \in 2\mathbb{Z}. \quad (13)$$

Consequently, if every  $\delta_N$  is even, then

$$\begin{aligned} & (\forall N_0 \exists N, r : N_0 < N \text{ and } D_{N+r} \notin -B_{h,N,r} + 2^{r+1} \mathbb{Z}) \\ & \iff D_N \notin \mathbb{Z} \text{ for arbitrarily large } N. \end{aligned} \quad (14)$$

There is a second equivalence. Let  $b : \mathbb{N} \rightarrow \mathbb{Q}$  satisfy  $|D_N| \leq b(N)$  for every  $N$ , and suppose that for every  $N$  and every positive integer  $q$  there is an  $r$  with

$$2b(N+r)q < 2^r. \quad (15)$$

Then

$$\begin{aligned} & \forall N_0 \exists N, r : N_0 < N \text{ and } \forall z \in \mathbb{Z}, \quad b(N+r) < |B_{h,N,r} - 2^r z| \\ & \iff D_N \notin \mathbb{Z} \text{ for arbitrarily large } N. \end{aligned} \quad (16)$$

*Proof.* Substituting  $D_{N+r} = 2^r D_N - B_{h,N,r}$  into the first membership condition cancels the observed block from both sides and leaves  $D_N = 2z$ . This proves (13). When every  $\delta_N$  is even, the recurrence also gives  $D_{N+1} \in 2\mathbb{Z}$  if and only if  $D_N \in \mathbb{Z}$ . Thus eventual integrality would force eventual even integrality and contradict the left side of (14). This proves the forward implication. Conversely, given a cutoff, choose a nonintegral  $D_N$  beyond it and take  $r = 0$  in (13).

For (16), eventual integrality and the block identity give some  $z \in \mathbb{Z}$  with

$$B_{h,N,r} - 2^r z = -D_{N+r},$$

so the displayed strict separation contradicts  $|D_{N+r}| \leq b(N+r)$ . Conversely, if  $D_N$  is nonintegral, choose a positive integer  $q$  with  $1/q \leq \text{dist}(D_N, \mathbb{Z})$ . Such a  $q$  exists because  $\mathbb{Z}$  is closed and  $D_N \notin \mathbb{Z}$ . Choose  $r$  from (15). The block identity and the triangle inequality give, for every  $z \in \mathbb{Z}$ ,

$$|B_{h,N,r} - 2^r z| \geq 2^r |D_N - z| - |D_{N+r}| \geq \frac{2^r}{q} - b(N+r) > b(N+r).$$

The same  $r$  works for all integers  $z$ , as required.  $\square$

Evenness is essential in (14). Without it, the recurrence  $T_N = N+1$ ,  $g_n = n-1$  for  $n \geq 1$ , has  $D_N = \sigma_1(N) = 1$  and  $\delta_N = 1$  at every index. Thus  $D_N$  is never an even integer but is always an integer. By (13), the left side of (14) holds while the right side fails.

The growth hypothesis (15) is equivalent to

$$\liminf_{n \rightarrow \infty} 2^{-n} b(n) = 0.$$

Indeed, writing  $n = N+r$ , its inequality becomes  $2^{-n} b(n) < 1/(2^{N+1}q)$ . Since  $b(n) \geq 0$ , finding such an  $n \geq N$  for every  $N$  and every positive integer  $q$  is exactly the stated liminf condition. In particular it holds for polynomial bounds and for  $Cc^n$  with fixed  $C > 0$  and  $1 < c < 2$ , but fails for  $b(n) = 2^n$ . A limit of zero is not necessary: the bound  $b(n) = 1$  at odd indices and  $b(n) = 2^n$  at even indices also satisfies the hypothesis. This example concerns the growth condition alone, not a bound for the prime tails. Only one suitable scale is needed to magnify the positive distance to  $\mathbb{Z}$  beyond the error; no comparison at every sufficiently large scale is required.

The proof in fact uses no rationality of  $D_N$ : the integer  $q$  is chosen from its positive distance to  $\mathbb{Z}$ , not from a reduced denominator. Thus (16) holds for real integer-coefficient recurrences under the same bound and growth hypothesis. The same is true of (13), and of (14) when the coefficient differences are even. These extensions follow from the printed argument; the Lean proof covers the rational case stated in the theorem.

The first test is exactly even integrality of  $D_N$ , regardless of  $r$ . The fixed-progression test compares  $2^r \text{dist}(D_N, \mathbb{Z})$  with the bound on the remaining tail. When  $D_N$  is nonintegral, the growth hypothesis makes the former larger than twice the latter at a suitable scale. Rewriting the test does not provide an estimate about the distribution of consecutive primes.

## E.1 A signed comparison with more analytic data

A signed variant appears in the totient companion, *The Binary Totient Series*, under “Signed interpolation with entire Dirichlet defects”. We recall its different hypotheses to delimit the comparison. Around the totient sequence, signed integer changes bounded

by any positive  $f(n) \rightarrow \infty$  fill an interval at every reciprocal integer base. They can preserve the odd coefficients, every fixed eventual congruence, all arithmetic-progression Dirichlet singularities, and prescribed finite root-of-unity jets. The proof takes finite differences of increasing order on separated finite sets of indices. The support of the changes has upper Banach density zero. The union of the intervals spanned by those sets has ordinary density zero and upper Banach density one.

This theorem allows the bound on the corrections to grow arbitrarily slowly and retains more analytic observations. It uses both signs, however, and does not subsume the one-sided prime-gap perturbation theorem proved here. In particular, it constructs comparison integer sequences, not consecutive prime gaps, and neither theorem supplies an irrationality proof for the actual prime series. The companion contains a full ordinary proof and the precise block estimate in terms of the prescribed bound; no infinite formal theorem is claimed.

## F Conclusions

The sparse construction preserves the specified congruences and block statistics while attaining an interval containing rational and irrational values. These properties therefore leave rationality undetermined for the comparison integer sequences. Ensuring that the cumulative positions are consecutive primes remains outside the construction.

For the actual primes, the tail recurrence gives exact criteria, and the finite certificates exclude particular possibilities. A proof using the small-difference condition would still need simultaneous witnesses arbitrarily late for every positive shift. Neither the computations nor the counterexamples establish that prime-gap estimate.

## G Source index for the short paper

The links below identify further formal definitions and statements used in the short paper, at its original revision 99f4bf47422a and line numbers. This local source pin applies only to the index below; Appendix C retains the long record's separate pin. Each link is labelled by the mathematics it states.

**Finite series.** The first prime gap is one; The second prime gap is two; Finite dyadic sums; Finite dyadic sums of consecutive differences; Summation by parts for a rational sequence; The two prime-series normalisations differ by a factor of two; Summation by parts for the primes; Terms of the prime series; Terms of the prime-gap series; The identity relating prime and gap terms.

**Convergent series.** Convergence of the gap series from convergence of the prime series; The infinite prime-to-gap identity under summability; Prime terms with denominator powers starting at zero; The factor-of-two identity for individual terms; Equivalent irrationality of the prime and gap series; The infinite identity in the second normalisation; Equivalent irrationality in the second normalisation; An elementary polynomial upper bound for the primes; Convergence of the prime series; The prime-to-gap identity with convergence proved.

**Recurrences.** The recurrence satisfied by rescaled dyadic tails; The difference between two terms of a recurrence; The weighted integer sum accumulated by iteration; Iteration of the recurrence; A shift as a multiple of a tail minus an integer; The recurrence for consecutive tail differences; Subtracting an integer preserves integrality; The criterion for a shift to be integral.

**Denominators and integral shifts.** The denominator divisibility criterion for an integral shift; Euler's congruence for an odd denominator; An integral shift of totient length; An integral shift remains integral at the next index; An integral shift remains integral at every later index; The denominator after one recurrence step; An odd denominator is unchanged by a step; An even denominator is halved by a step; Removing the power of two from a rational denominator; An integral multiple after the power of two is removed; Eventual integral shifts from the denominator factorisation; Eventual integral shifts under rationality.

**Rationality criteria.** Rational representation of a rescaled prime-gap tail; The recurrence for that rational representation; Existence of the rational representation under rationality; Rationality is equivalent to one integral positive-length shift; Rationality is equivalent to eventual integrality of a fixed shift; Irrationality is equivalent to nonintegrality of every positive-length shift; Irrationality is equivalent to arbitrarily late nonintegral shifts of each length.

**Small differences and counterexamples.** Two small differences with unequal coefficients cannot both be integral; Arbitrarily late small unequal pairs exclude eventual integrality; The small-pair implication for prime gaps; Prime gaps exceed every prescribed bound; Prime gaps are not eventually periodic; Coefficients obtained by telescoping; The finite telescoping identity; The telescoping example with coefficients equal to the index minus one; Nonperiodicity of the telescoping example; Rationality forces some fixed shift to fail eventual smallness.

**Further finite identities.** Finite prime sums in the second normalisation; Adding one term to a dyadic partial sum; Adding one term to a difference partial sum; Consecutive primes are in increasing order; The natural-number gap agrees with subtraction in the rationals; Finite sums of weighted prime gaps; Finite sums for the telescoping coefficients; A nonnegative version of the telescoping coefficients; Agreement of the nonnegative and integer formulas.

## References

- [1] Paul Erdős, *Sur certaines séries à valeur irrationnelle*. L'Enseignement Mathématique **4** (1958), 93–100, doi:10.5169/seals-34629.
- [2] Paul Erdős and Ronald L. Graham, *Old and New Problems and Results in Combinatorial Number Theory*. Monographies de L'Enseignement Mathématique, L'Enseignement Mathématique, 1980.
- [3] Paul Erdős and Carl Pomerance, *On the largest prime factors of  $n$  and  $n + 1$* . Aequationes Mathematicae **17** (1978), 311–321, doi:10.1007/BF01818569.
- [4] Paul Erdős, *On the irrationality of certain series: problems and results*. New Advances in Transcendence Theory, Cambridge University Press, 1988, pp. 102–109, doi:10.1017/CBO9780511897184.009.
- [5] Paul Erdős, *Beweis eines Satzes von Tschebyschef*. Acta Litterarum ac Scientiarum Szeged **5** (1932), 194–198.
- [6] Paul Erdős and Ernst G. Straus, *On the irrationality of certain Ahmes series*. Journal of the Indian Mathematical Society (N.S.) **27** (1964), 129–133. MR0175848.
- [7] Paul Erdős and Ernst G. Straus, *On the irrationality of certain series*. Pacific Journal of Mathematics **55** (1974), no. 1, 85–92, doi:10.2140/pjm.1974.55.85.
- [8] Kevin Ford, Ben Green, Sergei Konyagin, James Maynard and Terence Tao, *Long gaps between primes*. Journal of the American Mathematical Society **31** (2018), 65–105, doi:10.1090/jams/876; arXiv:1412.5029v3.
- [9] John A. Fridy, *Generalized bases for the real numbers*. The Fibonacci Quarterly **4** (1966), no. 3, 193–201.

- [10] A. Ya. Khinchin, *Continued Fractions*. University of Chicago Press, 1964.
- [11] Vjekoslav Kovač and Terence Tao, *On several irrationality problems for Ahmes series*. Acta Mathematica Hungarica **175** (2025), 572–608, doi:[10.1007/s10474-025-01528-0](https://doi.org/10.1007/s10474-025-01528-0); arXiv:[2406.17593](https://arxiv.org/abs/2406.17593)v4. Statement and section numbers refer to arXiv version 4.
- [12] Vivian Kuperberg, *Sums of singular series with large sets and the tail of the distribution of primes*. The Quarterly Journal of Mathematics **74** (2023), no. 4, 1457–1479, doi:[10.1093/qmath/haad030](https://doi.org/10.1093/qmath/haad030); arXiv:[2210.09775](https://arxiv.org/abs/2210.09775)v2. Statement numbers refer to arXiv version 2, 15 June 2023.
- [13] Johan Land, *A conditional proof of the irrationality of  $\sum_{n \geq 1} p_n 2^{-n}$  under a uniform Hardy–Littlewood prime-tuples conjecture*. Research draft, 5 September 2026. Formalisation material accompanies the draft.
- [14] Leonardo de Moura and Sebastian Ullrich, *The Lean 4 theorem prover and programming language*. Automated Deduction – CADE 28, Lecture Notes in Computer Science, Springer, 2021, pp. 625–635, doi:[10.1007/978-3-030-79876-5\\_37](https://doi.org/10.1007/978-3-030-79876-5_37).
- [15] The mathlib Community, *The Lean mathematical library*. Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs, ACM, 2020, pp. 367–381, doi:[10.1145/3372885.3373824](https://doi.org/10.1145/3372885.3373824). Describes a Lean 3-era snapshot; the repository lock identifies the library used by the present Lean 4 sources.
- [16] James Maynard, *Small gaps between primes*. Annals of Mathematics **181** (2015), 383–413, doi:[10.4007/annals.2015.181.1.7](https://doi.org/10.4007/annals.2015.181.1.7).
- [17] Hugh L. Montgomery and Robert C. Vaughan, *Multiplicative Number Theory I: Classical Theory*. Cambridge Studies in Advanced Mathematics, Cambridge University Press, 2007, doi:[10.1017/CBO9780511618314](https://doi.org/10.1017/CBO9780511618314).
- [18] Jan-Christoph Schlage-Puchta, *The irrationality of some number theoretical series*. Acta Arithmetica **126** (2007), no. 4, 295–303, doi:[10.4064/aa126-4-1](https://doi.org/10.4064/aa126-4-1); arXiv:[1105.1451](https://arxiv.org/abs/1105.1451)v1. Published in 2007; arXiv upload is from 2011. Lemma and preprint page locators refer to arXiv version 1.
- [19] Ian Short, *Ford Circles, Continued Fractions, and Rational Approximation*. The American Mathematical Monthly **118** (2011), no. 2, 130–135, doi:[10.4169/amer.math.monthly.118.02.130](https://doi.org/10.4169/amer.math.monthly.118.02.130); arXiv:[0912.1997](https://arxiv.org/abs/0912.1997)v1. Preprint title: Ford circles, continued fractions, and best approximation of the second kind. Preprint theorem locators refer to arXiv version 1.
- [20] Wouter van Doorn and Vjekoslav Kovač, *Lacunary sequences whose reciprocal sums represent all rational numbers in an interval*. Acta Arithmetica **223** (2026), 275–295, doi:[10.4064/aa251001-13-1](https://doi.org/10.4064/aa251001-13-1); arXiv:[2509.24971](https://arxiv.org/abs/2509.24971)v3. Statement numbers refer to arXiv version 3.
- [21] Yitang Zhang, *Bounded gaps between primes*. Annals of Mathematics **179** (2014), 1121–1174, doi:[10.4007/annals.2014.179.3.7](https://doi.org/10.4007/annals.2014.179.3.7).
- [22] Thomas F. Bloom, *Erdős Problem #251*. 2026. Accessed 6 September 2026.
- [23] Erdős Problems contributors, *Erdős Problem #251 discussion thread*. 2026. Accessed 6 September 2026: Tao comment of 7 October 2025 and Land comments of 6 September 2026.
- [24] ChatGPT 5.4 Pro (orchestrated by Vjekoslav Kovač), *On the Erdős problem #251*. Unpublished note, Department of Mathematics, University of Zagreb, 2026. Accessed 6 September 2026.

- [25] The Formal Conjectures Authors, *FormalConjectures.ErdosProblems.251*. 2025. Pinned statement source, not a proof of Problem 251; accessed 28 July 2026.
- [26] Stefan Ringer, *Local gap statistics, telescoping, and normality: a local-pattern approach to Erdős problem 251*. Preprint, 11 September 2026. Mutable main-branch TeX consulted 16 September 2026; the cited conditional statements were rechecked 18 September 2026. Formalisation material accompanies the draft.
- [27] Tonći Crmarić and Vjekoslav Kovač, *On the irrationality of certain super-polynomially decaying series*. Colloquium Mathematicum **179** (2025), 55–68, doi:10.4064/cm9628-5-2025; arXiv:2504.18712v1. Lemma 4 is cited using arXiv version 1.
- [28] Boris Adamczewski, Michael Drmota and Clemens Müllner, *(Logarithmic) densities for automatic sequences along primes and squares*. Transactions of the American Mathematical Society **375** (2022), no. 1, 455–499, doi:10.1090/tran/8476; arXiv:2009.14773v2. Theorems 1.2 and 1.4 refer to arXiv version 2, 13 April 2021; journal publication is 2022.
- [29] Wouter van Doorn, *Partitions with prescribed sum of reciprocals: asymptotic bounds*. 2025; arXiv:2502.02200v2. Version 2, 23 July 2025.
- [30] Szymon Głab and Franciszek Prus-Wiśniowski, *Achievement sets – current results and open problems*. Real Analysis Exchange (2026), doi:10.14321/realanalexch.1766383782; arXiv:2512.17285v1. Advance publication, first available in Project Euclid 8 June 2026. Consulted text remains arXiv:2512.17285v1 (19 December 2025).
- [31] Franciszek Prus-Wiśniowski and Jolanta Ptak, *Achievable Cantorvals almost without reversed Kakeya conditions*. 2024; arXiv:2412.08768v1. Version 1 submitted 11 December 2024. The sparse indices satisfy the overlap inequality, not its strict term-dominating reverse.
- [32] Artur Bartoszewicz, Małgorzata Filipczak and Emilia Szymonik, *Multigeometric sequences and Cantorvals*. Central European Journal of Mathematics **12** (2014), no. 7, 1000–1007, doi:10.2478/s11533-013-0396-4; arXiv:1304.4218v2.
- [33] Vivian Kuperberg, *Sums of singular series along arithmetic progressions and with smooth weights*. International Journal of Number Theory **21** (2025), no. 1, 53–74, doi:10.1142/S1793042125500046; arXiv:2301.06095v1. Preprint uploaded 15 January 2023; journal publication is 2025.
- [34] Abhishek Jha, *The Poisson Tail Conjecture for primes in short intervals*. 2026; arXiv:2605.23014v2. Substantially revised version 2, 12 September 2026, 31 pages. Conditional statements must retain their strong Hardy–Littlewood hypotheses.
- [35] D. H. J. Polymath, *Variants of the Selberg sieve, and bounded intervals containing many primes*. Research in the Mathematical Sciences **1** (2014), article 12, doi:10.1186/s40687-014-0012-7; arXiv:1407.4897v4. Theorem 1.4(i) refers to arXiv version 4 (22 December 2014); the journal numbers it Theorem 4(i). An erratum is recorded at doi:10.1186/s40687-015-0033-x.
- [36] Zbigniew Nitecki, *Cantorvals and Subsum Sets of Null Sequences*. The American Mathematical Monthly **122** (2015), no. 9, 862–870, doi:10.4169/amer.math.monthly.122.9.862; arXiv:1106.3779v2. Consulted preprint: Subsum Sets: Intervals, Cantor Sets, and Cantorvals, version 2 (8 July 2013). Theorem 14 is attributed there to Guthrie–Nymann; its locator is not journal pagination.
- [37] Piotr Miska, Franciszek Prus-Wiśniowski and Jolanta Ptak, *More on Kakeya Conditions for Achievement Sets*. Results in Mathematics **78** (2023), article 113, doi:10.1007/s00025-023-01890-x. Repairs an estimate in the 2021 proof, preserving its uniqueness conclusion, and gives a simpler proof of a weaker theorem without that conclusion.

- [38] Jacek Marchwicki and Piotr Miska, *On Kakeya Conditions for Achievement Sets*. Results in Mathematics **76** (2021), article 181, doi:10.1007/s00025-021-01479-2. Theorem 2.1 is to be read with the proof repair in Miska–Prus-Wiśniowski–Ptak (2023).
- [39] Piotr Nowakowski, *On a new condition implying that an achievement set is a Cantorval and its applications*. 2025; arXiv:2512.17761v1. Version 1, 19 December 2025. Theorem 3.1 requires the Star Procedure of Definition 2 never to break; no application to the present factorial weights is asserted.

## Verification concordance

| Statement       | Formal proof                                      | Statement        | Formal proof                                              |
|-----------------|---------------------------------------------------|------------------|-----------------------------------------------------------|
| Theorem 2.1     | <a href="#">Lean</a> · <a href="#">Comparator</a> | Corollary 2.2    | <a href="#">Lean</a> · <a href="#">Comparator</a>         |
| Proposition 4.1 | <a href="#">Lean</a> · <a href="#">Comparator</a> | Theorem 4.2      | <a href="#">Lean</a> · <a href="#">Comparator</a>         |
| Theorem 4.3     | <a href="#">Lean</a> · <a href="#">Comparator</a> | Corollary 4.4    | <a href="#">Lean</a> · <a href="#">Comparator</a>         |
| Theorem 5.2     | <a href="#">Lean</a> · <a href="#">Comparator</a> | Theorem 5.3      | <a href="#">Lean</a> · <a href="#">Comparator</a>         |
| Lemma 5.4       | <a href="#">Lean</a> · <a href="#">Comparator</a> | Theorem 5.5      | <a href="#">Lean</a> · <a href="#">Comparator</a>         |
| Theorem 5.6     | <a href="#">Lean</a> · <a href="#">Comparator</a> | Theorem 6.1      | <a href="#">Lean</a> · <a href="#">Comparator</a>         |
| Corollary 6.2   | <a href="#">Lean</a> · <a href="#">Comparator</a> | Proposition 6.3  | <a href="#">Lean</a> · <a href="#">Comparator</a>         |
| Theorem 6.4     | <a href="#">Lean</a> · <a href="#">Comparator</a> | Proposition 6.5  | <a href="#">Lean</a> · <a href="#">Comparator</a>         |
| Proposition 6.6 | <a href="#">Lean</a> · <a href="#">Comparator</a> | Proposition 6.7  | <a href="#">Lean</a> · <a href="#">Comparator</a>         |
| Theorem 7.1     | <a href="#">Lean</a> · <a href="#">Comparator</a> | Theorem 7.2      | <a href="#">Lean</a> · <a href="#">Comparator</a> pending |
| Theorem 8.1     | <a href="#">Lean</a> · <a href="#">Comparator</a> | Proposition 8.2  | <a href="#">Lean</a> · <a href="#">Comparator</a>         |
| Theorem 8.3     | <a href="#">Lean</a> · <a href="#">Comparator</a> | Corollary 8.4    | <a href="#">Lean</a> <sup>†</sup>                         |
| Proposition 8.5 | <a href="#">Lean</a> · <a href="#">Comparator</a> | Corollary 8.6    | <a href="#">Lean</a> <sup>†</sup>                         |
| Theorem 8.7     | <a href="#">Lean</a> <sup>†</sup>                 | Theorem 8.8      | <a href="#">Lean</a> · <a href="#">Comparator</a>         |
| Proposition 8.9 | <a href="#">Lean</a> · <a href="#">Comparator</a> | Proposition 8.10 | <a href="#">Lean</a> · <a href="#">Comparator</a>         |
| Proposition D.1 | <a href="#">Lean</a> · <a href="#">Comparator</a> | Proposition D.2  | <a href="#">Lean</a> · <a href="#">Comparator</a>         |
| Proposition D.3 | <a href="#">Lean</a> · <a href="#">Comparator</a> | Proposition D.4  | <a href="#">Lean</a> · <a href="#">Comparator</a>         |
| Proposition D.6 | <a href="#">Lean</a> · <a href="#">Comparator</a> | Theorem E.1      | <a href="#">Lean</a> · <a href="#">Comparator</a>         |