

Irrationality of $F(31/4)$ and the Exact Normalized Hankel Order

Cyclotomic cancellation and a unique least-order moment configuration

WILL COOK*

September 2026

ABSTRACT

For coprime integers $a > b \geq 1$, the Lambert value $F(a/b) = \sum_{n \geq 1} ((a/b)^n - 1)^{-1}$ is irrational whenever $\log b / \log a < \theta^*$, where the explicit constant below satisfies $\theta^* \approx 0.4056830213840605$. In particular, $F((31/4)^r)$ is irrational for every integer $r \geq 1$. The same forms bound its irrationality exponent uniformly in r . Cyclotomic factors are cancelled before the rational-base denominator is cleared; the degree of the resulting polynomial pair determines the cost. The argument does not cover $3/2$. For the normalised Hankel determinant in the second construction we prove

$$\text{ord}_q V_N^* = \frac{N(N-1)(2N-1)}{6}, \quad [q^{\text{ord}_q V_N^*}] V_N^* = \frac{(N!)^2 (N+1)!}{2^N}.$$

The exponent and coefficient come from a unique least-order tuple in a formal moment expansion. A quantitative selector criterion separates local divisibility from the additional estimates needed for a small nonzero real remainder.

1 Introduction

For $t > 1$, expansion of each geometric series gives

$$F(t) = \sum_{n \geq 1} \frac{1}{t^n - 1} = \sum_{n \geq 1} \frac{\tau(n)}{t^n},$$

where $\tau(n)$ is the number of positive divisors of n . Erdős Problem #1049 asks for irrationality at every rational $t > 1$ [1, p. 102]. The theorem below proves an explicit sufficient region and settles the power family in the title.

Cyclotomic cancellation produces positive forms $\Lambda_n = U_n F - V_n$ with integral coefficient polynomials of degree at most W_n , and

$$\log(b^{W_n} \Lambda_n(a/b)) = (C_1 \log b - C_0 \log a) n^2 + o(n^2).$$

A negative exponent yields positive integral linear forms tending to zero.

**Authorship and AI use.* Will Cook built and directed the research infrastructure and reviewed the claims when he could. AI agents did most of the research and drafting. Cook did not independently verify every claim.

The Hankel argument concerns a different family. Its moment expansion makes one increasing index tuple responsible for the first coefficient. The final sections explain what local cancellation can establish at $3/2$ and specify the real estimate that would complete that approach. The complete catalogue of other constructions and their failure witnesses is retained in the long record.

2 A rational base at which F is irrational

Write $\psi_1(u) = \sum_{k \geq 0} (k+u)^{-2}$. Let $\mathcal{I}$ be the thirteen intervals listed in the proof and put

$$C_1 = \frac{1091}{2}, \quad J = \sum_{[u,v) \in \mathcal{I}} (\psi_1(u) - \psi_1(v)), \quad C_0 = 266 - \frac{3}{\pi^2}(225 - J).$$

Thus $\theta^* = C_0/C_1$ and $\mu = C_1/C_0$ are defined exactly.

Theorem 2.1 (rational-base region). *Let $a > b \geq 1$ be coprime integers with*

$$b^\mu < a, \quad \text{equivalently} \quad \frac{\log b}{\log a} < \theta^*,$$

$$\theta^* = \frac{C_0}{C_1} = 0.4056830213840605 \dots,$$

$$\mu = \frac{C_1}{C_0} = 2.4649786835749750 \dots$$

Then $F(a/b)$ is irrational.

The cancellation is performed before homogenisation, so the clearing degree is the degree of the cancelled pair. The value $3/2$ is outside this sufficient region: $J \leq \psi_1(1/14) - \psi_1(1) < 196$ gives $\theta^* < 266/(1091/2) < 1/2 < \log 2 / \log 3$.

Proof of Theorem 2.1. Set

$$a_0 = 14n + 1, \quad a_1 = 12n + 1, \quad a_2 = 14n + 1, \quad \beta = 27n + 2, \quad N = 15n.$$

Use the coefficient pair A_n, B_n of the source identities [3, (8)–(11)], with $H_n = A_n F - B_n$. Put $D_N(X) = \prod_{\ell=1}^N \Phi_\ell(X)$, $M_n = 266n^2 + 34n + 1$, and

$$\Omega_n(X) = \prod_{\ell=2}^N \Phi_\ell(X)^{\nu_\ell}, \quad \nu_\ell = \omega(n/\ell),$$

where the periodic function

$$\omega(x) = \max\{0, \lfloor 14x \rfloor + \lfloor 13x \rfloor - \lfloor 12x \rfloor - \lfloor 15x \rfloor, \\ 2\lfloor 14x \rfloor - \lfloor 13x \rfloor - \lfloor 15x \rfloor\}$$

is zero or one. Its support in $[0, 1)$ consists of

$$\begin{aligned} &[1/14, 1/12), [1/7, 1/6), [3/14, 1/4), [2/7, 1/3), \\ &[5/14, 2/5), [3/7, 7/15), [1/2, 8/15), [4/7, 3/5), \\ &[9/14, 2/3), [5/7, 11/15), [11/14, 4/5), \\ &[6/7, 13/15), [13/14, 14/15). \end{aligned}$$

These are the intervals $\mathcal{I}$ used to define J . The zero-one values and this thirteen-interval support are [Lean source](#).

Integral polynomials and their degrees. The source's polynomial inclusion is Lemma 7, display (23), on p. 161 [3]. Its parameter vector is $n(13, 14, 12, 14, 15, 13)$, its maximum is $15n$, and $\beta - a_1 - a_2 = n > 0$. Its hypotheses therefore hold for every $n \geq 1$. It gives

$$\Lambda_n(X) = X^{-M_n} \frac{D_N(X)}{\Omega_n(X)} H_n(X) = U_n(X)F(X) - V_n(X), \quad U_n, V_n \in \mathbb{Z}[X].$$

This is the polynomial conclusion before the source's subsequent integer-specialisation step in display (24). The coefficientwise interpretation also follows by comparing the source rational coefficients: F is not a rational function, since $F(e^h) = h^{-1} \log(1/h) + O(h^{-1})$ as $h \downarrow 0$. For this estimate split the original sum at $n = 1/h$; the remaining geometric tail is $O(h^{-1})$.

In the source expression for A_n , the degree of the k th summand is

$$d_k = a_0 k + E_k + (a_1 - 1)(k - a_1) + (\beta - k - 1)(k - a_2),$$

where

$$E_k = \frac{a_1(a_1 - 1) - (\beta - a_2)(\beta - a_2 - 1) + (\beta - k)(\beta - k - 1)}{2}.$$

For $a_2 \leq k \leq \beta - 2$, one has $d_{k+1} - d_k = 40n + 1 - k > 0$. There is therefore a unique highest-degree summand, and

$$K_n := \deg A_n = \frac{1091n^2 + 81n + 2}{2}, \quad W_n := \deg U_n = K_n - M_n + \sum_{\ell \leq 15n} (1 - \nu_\ell) \varphi(\ell).$$

For fixed n , the positive representation below gives $H_n(x) = O(1)$ as $x \rightarrow \infty$. Hence $\Lambda_n(x) = O(x^{W_n - K_n})$. Since $F(x) = O(x^{-1})$ and $K_n \geq 1$, the identity $V_n = U_n F - \Lambda_n$ gives $\deg V_n \leq W_n - 1$. Thus W_n clears both coordinates.

The limiting degree cost. The elementary summatory estimate $\sum_{\ell \leq y} \varphi(\ell) = 3y^2/\pi^2 + O(y \log y)$ gives

$$\frac{1}{n^2} \sum_{\ell \leq 15n} \varphi(\ell) \rightarrow \frac{675}{\pi^2}.$$

For $[u, v) \in \mathcal{I}$, the condition $\{n/\ell\} \in [u, v)$ is the disjoint union of intervals $n/(k+v) < \ell \leq n/(k+u)$ for $k \geq 0$. For finitely many k the same summatory estimate applies term by term. The remaining terms have total normalised mass $O(1/K^2)$ after truncation at $k = K$, since they involve only $\ell \leq n/(K+u)$. Letting first n and then K tend to infinity proves

$$\frac{1}{n^2} \sum_{\ell \leq 15n} \nu_\ell \varphi(\ell) \rightarrow \frac{3}{\pi^2} \sum_{[u,v) \in \mathcal{I}} \sum_{k \geq 0} \left(\frac{1}{(k+u)^2} - \frac{1}{(k+v)^2} \right) = \frac{3J}{\pi^2}.$$

Here no contributing index exceeds $14n$, since $u \geq 1/14$. Consequently

$$K_n/n^2 \rightarrow C_1, \quad (K_n - W_n)/n^2 \rightarrow C_0.$$

A positive remainder at every fixed real base. For $x > 1$ and $q = 1/x$, the source identity has the positive representation

$$H_n(x) = \sum_{t \geq 0} q^{a_0 t} \frac{(q^{t+1}; q)_{a_1-1}}{(q; q)_{a_1-1}} \frac{(q; q)_{\beta-a_2-1}}{(q^{a_2+t}; q)_{\beta-a_2}}.$$

The last denominator has length $\beta - a_2$, as required by the source's gamma expression (7) and residues (8). The shorter length in one unnumbered product on printed p. 156 is an indexing discrepancy. Every finite product lies between $P = (q; q)_\infty > 0$ and 1, so

$$P^2 \leq H_n(x) \leq \frac{P^{-2}}{1 - q^{a_0}}.$$

In particular $\log H_n(x) = O_x(1)$ and $\Lambda_n(x) > 0$.

For fixed $x > 1$, the cyclotomic identity

$$\log \Phi_\ell(x) - \varphi(\ell) \log x = \sum_{d|\ell} \mu(d) \log(1 - x^{-\ell/d})$$

has total absolute error $O_x(n)$ over $\ell \leq 15n$. Indeed, it is at most $15nB(x)$, where

$$B(x) = \sum_{d \geq 1} \frac{-\log(1 - x^{-d})}{d} < \infty.$$

Since $0 \leq 1 - \nu_\ell \leq 1$, it follows that

$$\log \Lambda_n(x) = -(K_n - W_n) \log x + O_x(n).$$

Homogenisation. Because U_n, V_n are integral polynomials of degree at most W_n , $b^{W_n} U_n(a/b)$ and $b^{W_n} V_n(a/b)$ are integers. That integrality and the cleared linear-form identity it produces are [Lean source](#). Moreover

$$\begin{aligned} \log(b^{W_n} \Lambda_n(a/b)) &= K_n \log b - (K_n - W_n) \log a + O_{a/b}(n) \\ &= (C_1 \log b - C_0 \log a) n^2 + o(n^2). \end{aligned}$$

The coefficient is negative under the theorem's hypothesis. Thus positive integral linear forms in $F(a/b)$ tend to zero. If $F(a/b) = r/s$ were rational, every such form would have absolute value at least $1/|s|$, a contradiction. The separation bound for a nonzero integral form at a rational target is [Lean source](#). $\square$

Corollary 2.2. $F((31/4)^r)$ is irrational for every integer $r \geq 1$.

Proof. The exact inequalities $31^2 < 4^5$ and $4^{200} < 31^{81}$ give $2/5 < \log 4 / \log 31 < 81/200$. The first term of each trigamma difference yields

$$J \geq J_0 := \sum_{(u,v) \in \mathcal{I}} (u^{-2} - v^{-2}) = \frac{2015640690251}{25971865920}.$$

Using $\pi > 157/50$ gives the rational lower bound

$$\theta^* > \frac{2359630009523263}{5820307922172744} > \frac{81}{200}.$$

Finally, the logarithmic ratio and coprimality are preserved by a common positive integer power. $\square$

Corollary 2.3 (an irrationality measure uniform over powers). *For coprime $a > b \geq 1$ with $\theta = \log b / \log a < \theta^*$ and every integer $r \geq 1$,*

$$\mu_{\text{irr}}(F((a/b)^r)) \leq \frac{1 - \theta}{\theta^* - \theta}.$$

Here $\mu_{\text{irr}}(\xi)$ is the supremum of the exponents ν for which $|\xi - p/q| < q^{-\nu}$ has infinitely many reduced rational solutions. In particular, $\mu_{\text{irr}}(F((31/4)^r)) < 301$ for every $r \geq 1$.

Proof. The raw source coefficient is a sum of $O(n)$ Laurent monomials times two Gaussian polynomials, each of coefficient sum at most 2^{27n+2} . Its coefficient norm is $\exp(O(n))$, and every exponent is at most K_n . The same normalising multiplier used above therefore gives

$$|U_n(x)| \leq x^{W_n} \exp(O_x(n)) \quad (x > 1 \text{ fixed}).$$

Set $\xi = F(a/b)$, $Q_n = b^{W_n} U_n(a/b)$ and $P_n = b^{W_n} V_n(a/b)$. With

$$\alpha = (C_1 - C_0) \log a, \quad \tau = C_0 \log a - C_1 \log b > 0,$$

we have $\log(Q_n \xi - P_n) = -\tau n^2 + o(n^2)$ and $|Q_n| \leq \exp(\alpha n^2 + o(n^2))$.

For integers A, B, p, q with $q > 0$, if $L = A\xi - B$ and $2q|L| \leq 1$, then

$$|L| \leq |A| |\xi - p/q|.$$

When $Ap - Bq = 0$ this is equality. Otherwise the nonzero integer $Ap - Bq$ gives $1/q \leq |L| + |A| |\xi - p/q|$, which proves the inequality. For a fixed $\eta \in (0, \tau)$ choose $n = \lceil \sqrt{\log(2q)/(\tau - \eta)} \rceil$ and apply it to (Q_n, P_n) . The two-sided remainder estimate yields

$$|\xi - p/q| \geq q^{-(\alpha + \tau + 2\eta)/(\tau - \eta) - o(1)}.$$

Let $\eta \downarrow 0$. The resulting bound is $1 + \alpha/\tau = (1 - \theta)/(\theta^* - \theta)$. Taking a common power multiplies α, τ by r , leaving this quotient unchanged; the constants in the approximation inequality may depend on r . For $31/4$, sixteen terms of each trigamma difference and $\pi > 314159/100000$ give $\theta^* > 0.40568$. The logarithm series gives $\log 4 / \log 31 < 0.4036982$. Thus the bound is less than $2981509/9909 < 301$. $\square$

Comparison and scope. Bundschuh and Väänänen's Theorem 2 at $\alpha = -1$ [2, p. 177] gives $\log b / \log a < \theta_{\text{BV}} := 1/2 - 1/\pi^2$. Since $\pi^2 < 10$, one has $\theta_{\text{BV}} < 2/5 < \log 4 / \log 31$. The displayed sufficient regions therefore differ on $[\theta_{\text{BV}}, \theta^*)$. The direction $(14, 12, 14; 27)$, its thirteen intervals and the constant $\mu \approx 2.46497868$ are inherited from [3, p. 162], where μ bounds an integer-base irrationality exponent. The rational-base extension announced in [4, Section 2] has the same shape with an unspecified computable constant. Here the polynomial specialisation identifies μ as admissible for F ; no priority claim is attached.

Negative bases are not treated.

2.1 The base-uniform degree restriction

The preceding construction uses degree and decay estimates valid at every fixed real base greater than one. Such uniformity itself imposes a limit.

Theorem 2.4 (Archimedean cap on base-uniform rank-two families). *Let (U_n, V_n) be pairs in $\mathbb{Z}[X]^2$ satisfying $\Lambda_n(x) = U_n(x)F(x) - V_n(x) \neq 0$, $\deg U_n, \deg V_n \leq \delta n^2(1+o(1))$, $\log \max(H(U_n), H(V_n)) \leq hn^2(1+o(1))$ with H the ℓ^1 coefficient norm, and $\log |\Lambda_n(x)| = -\sigma n^2 \log x (1+o(1))$ for every real $x > 1$, with $\sigma, \delta > 0$ and $h \geq 0$ independent of x . Then with $d_n = \max(\deg U_n, \deg V_n)$, the homogenised forms $b^{d_n} \Lambda_n(a/b)$ tend to zero whenever $\log b / \log a < \sigma / (\sigma + \delta)$, and $\sigma / (\sigma + \delta) \leq 1/2$.*

Proof. Write $H_n = \max(H(U_n), H(V_n))$. Then $|U_n(x)|, |V_n(x)| \leq H_n x^{d_n}$ for real $x > 1$. Suppose $\sigma > \delta$ and choose an integer $p \geq 2$ with $(\sigma - \delta) \log p > h$. Set $a_n = U_n(p)$, $b_n = V_n(p)$ and $L_n = a_n F(p) - b_n$. Hypotheses on height and degree give $|a_n| \leq \exp((h + \delta \log p)n^2 + o(n^2))$, while $|L_n| = \exp(-\sigma \log p n^2 + o(n^2))$. The adjacent integer

$$a_n b_{n+1} - a_{n+1} b_n = a_{n+1} L_n - a_n L_{n+1}$$

is then $o(1)$, hence eventually zero. Also $a_n \neq 0$ for large n : otherwise the nonzero integer $L_n = -b_n$ would have absolute value less than 1. Thus b_n/a_n is eventually a fixed rational r . If $F(p) \neq r$ then $|L_n| \geq |F(p) - r|$; if $F(p) = r$ then $L_n = 0$. Both contradict the hypotheses, so $\sigma \leq \delta$. The integer argument of this paragraph, from the two cross-product limits to the contradiction, is [Lean source](#). The homogenised logarithm satisfies

$$\limsup_{n \rightarrow \infty} n^{-2} \log |b^{d_n} \Lambda_n(a/b)| \leq \delta \log b - \sigma \log(a/b),$$

which is negative on the stated sufficient region. The arithmetic form of that region is [Lean source](#), and the final numerical clause is [Lean source](#). The displayed degree upper bound furnishes a sufficient decay cutoff; the following corollary gives a separate exclusion under the same all-base hypotheses. $\square$

Corollary 2.5 (Undivided forms below the square boundary). *Under the hypotheses of Theorem 2.4, for positive integers a, b with $b < a < b^2$, the undivided forms $b^{d_n} \Lambda_n(a/b)$ do not tend to zero. No limit of d_n/n^2 is assumed.*

Proof. Write $x = a/b$ and suppose $C_n = b^{d_n} \Lambda_n(x) \rightarrow 0$. Eventual nonvanishing gives $\log |C_n| = d_n \log b + \log |\Lambda_n(x)|$ for all sufficiently large n . Also $|C_n| \leq 1$ eventually. The lower side of the remainder asymptotic therefore implies

$$d_n \log b \leq -\log |\Lambda_n(x)| = \sigma \log x n^2 + o(n^2).$$

Set $c = \sigma \log x / \log b$. Since $1 < x < b$, we have $0 < c < \sigma$, and the displayed inequality gives $d_n \leq (c + \varepsilon)n^2$ eventually for each $\varepsilon > 0$. The same family thus satisfies the cap hypotheses with degree upper rate c , its original height bound, and its original nonvanishing and remainder asymptotics at every real base greater than one. The cap yields $\sigma \leq c$, a contradiction. $\square$

The no-decay conclusion is [kernel-checked in Lean](#) under the stated all-base hypotheses.

This strengthens the failure of a sufficient-cutoff test to an exclusion of decay under the stated all-base hypotheses. It does not supply an actual-degree limit or assert divergence. If $d_n/n^2 \rightarrow d$, the separate exact-degree result gives the stronger conclusion $|b^{d_n}\Lambda_n(a/b)| \rightarrow \infty$ in the same strict region. Equality $a = b^2$ remains unclassified. The corollary concerns the undivided forms; it does not exclude base-dependent content division or other irrationality methods outside its hypotheses.

3 Exact normalized-Hankel order in Zudilin's construction

At $x = z = 1$, Zudilin's normalized moments are

$$v_m^* = \sum_{t \geq 0} q^{(m+1)t} \frac{(q; q)_m^3 (q^{t+1}; q)_m}{(q^{m+1+t}; q)_{m+1}}, \quad V_N^* = \det_{0 \leq i, j < N} (v_{i+j}^*).$$

His row transformation proves

$$\text{ord}_q V_N^* \geq \frac{N(N-1)(2N-1)}{6}$$

for every $N \geq 1$ [4, Section 4]. A formal moment expansion identifies the unique least-order term and shows that this estimate is always sharp.

Theorem 3.1 (sharp normalized Hankel order). *For every $N \geq 1$,*

$$\text{ord}_q V_N^* = \frac{N(N-1)(2N-1)}{6},$$

and the coefficient of the first nonzero monomial is

$$[q^{N(N-1)(2N-1)/6}] V_N^* = \frac{(N!)^2 (N+1)!}{2^N}.$$

Proof. Write $P = (q; q)_\infty$ and introduce the coefficientwise formal series

$$G_q(w) = \frac{1}{(w; q)_\infty^3} \sum_{t \geq 0} \frac{w^t}{(q; q)_t} \frac{(q^t w^2; q)_\infty}{(q^t w; q)_\infty^2}, \quad a_k(q) = [w^k] P^4 G_q(w).$$

For fixed w -degree, all operations define power series in q . The product identities $(q; q)_m = P/(q^{m+1}; q)_\infty$ and $(q^a; q)_m = (q^a; q)_\infty / (q^{a+m}; q)_\infty$ give the exact formal identity

$$v_m^* = P^4 G_q(q^{m+1}) = \sum_{k \geq 0} a_k(q) q^{(m+1)k}.$$

At $q = 0$, the term $t = 0$ in G_q is $(1-w^2)/(1-w)^5$ and the terms $t \geq 1$ sum to $w/(1-w)^4$. Consequently

$$G_0(w) = \frac{1+2w}{(1-w)^4}, \quad a_k(0) = \frac{(k+1)^2(k+2)}{2} =: c_k > 0.$$

Apply Cauchy–Binet to a finite truncation of the moment sum and then pass coefficientwise to the limit. This is legitimate because only finitely many increasing index tuples contribute to any fixed q -degree. It gives

$$V_N^* = \sum_{k_0 < \dots < k_{N-1}} \left(\prod_{i=0}^{N-1} a_{k_i}(q) q^{k_i} \right) \prod_{0 \leq i < j < N} (q^{k_i} - q^{k_j})^2.$$

The summand indexed by (k_i) has order

$$\sum_{i=0}^{N-1} (2N - 1 - 2i)k_i.$$

Every weight is positive and $k_i \geq i$. Equality with the least possible order occurs uniquely when $k_i = i$ for every i . That tuple contributes

$$\sum_{i=0}^{N-1} (2N - 1 - 2i)i = \frac{N(N-1)(2N-1)}{6}$$

and leading coefficient

$$\prod_{i=0}^{N-1} c_i = \frac{(N!)^2 (N+1)!}{2^N}.$$

No other tuple can cancel this coefficient. □

Formal order and fixed-base size. The theorem identifies the first formal term. Formal order alone would not control a fixed- q residual: multiplying by $(1 - q)^{N^3}$ preserves that first term and changes the logarithm by a cubic quantity at fixed $0 < q < 1$. The separate positive-measure argument for these same moments proves $V_N^*(q) > 0$ and $\log(V_N^*(q)/(C_N q^{B_N})) = O_q(N)$, where $B_N = N(N-1)(2N-1)/6$ and $C_N = (N!)^2(N+1)!/2^N$. Its complete ordinary proof is in `ZudilinHankelPositiveMeasure.md` and the long record. Neither fact supplies a denominator factor for the 2004 polynomial forms.

4 Local cancellation and the remaining real estimate

Theorem 4.1 (no corridor at base $3/2$). *For all $N \geq 1$ and $K \geq 1$ and all natural Q, D , the tuple $(3, 2, N, K, Q, D)$ is not a coordinatewise corridor.*

Theorem 4.2 (cleared-tail recurrence). *Let $r, s, B, F \in \mathbb{Q}$ with $r \neq 0$, let $c : \mathbb{N} \rightarrow \mathbb{Q}$, and let P_N and U_N be the cleared tail state. Then for every N ,*

$$U_{N+1} = r U_N - B c(N+1) s^{N+1}.$$

Theorem 4.3 (the forcing term). *Let s, B be natural numbers and $c : \mathbb{N} \rightarrow \mathbb{N}$, and put $G_N = B c(N+1) s^{N+1}$.*

1. *If $s \geq 2, B \geq 1$ and $c(N+1) \geq 1$, then $2^{N+1} \leq G_N$.*
2. *If $s = 1$, then $G_N = B c(N+1)$.*

Fix a common width W . For $P \in \mathbb{Z}[X]$ of degree at most W , write

$$H_W(P) = 2^W P(3/2) = \sum_{i=0}^W [P]_i 3^i 2^{W-i}.$$

A specialised row is primitive when its two integer coordinates have gcd one. This normalisation is different from removing the common polynomial coefficient content, and must precede the local count. For depths $R, S \geq 0$, define

$$J_{3,R}(P) = H_W(P) \pmod{3^R}, \quad J_{2,S}(P) = H_W(P) \pmod{2^S}.$$

All four jets of (U, V) vanish precisely when $D = 3^R 2^S$ divides both specialised coordinates. The dependence on the declared width remains fixed when rows are added.

Theorem 4.4 (binary four-jet collision). *Fix a width W and depths R, S , and let $(U_j, V_j)_{j < M}$ be any M pairs of integral polynomials. Call a subset of $\{0, \dots, M-1\}$, equivalently a vector of $\{0, 1\}^M$, a binary selector. If the 2^M binary selectors outnumber the finite four-jet target*

$$(\mathbb{Z}/3^R \mathbb{Z})^2 \times (\mathbb{Z}/2^S \mathbb{Z})^2,$$

then two distinct subsets have the same four-jet sum. Subtracting their indicator vectors gives a nonzero coefficient vector in $\{-1, 0, 1\}^M$ cancelling all four jets. The target has exact cardinality

$$(3^R)^2 (2^S)^2.$$

In particular, if $R > 0$ and $4R + 2S \leq M$, such a collision exists.

Proof. Send each binary selector to the sum of the four-jet signatures it selects. The claimed cardinal inequality and the pigeonhole principle give two distinct selectors in the same fibre. The cardinality formula is the product of the four cyclic-modulus cardinalities. For $R > 0$,

$$(3^R)^2 (2^S)^2 < (4^R)^2 (2^S)^2 = 2^{4R+2S} \leq 2^M,$$

which proves the stated sufficient threshold. The cardinality formula, the collision, the signed $\{-1, 0, 1\}$ vector and the sufficient width are together [Lean source](#). $\square$

The ambient count does not use relations between the two residue coordinates. Vanishing minors can reduce that cost to one coordinate.

Theorem 4.5 (Bézout–Plücker tail collapse). *Let R_0 be a commutative ring and let $w_n = (A_n, B_n) \in R_0^2$. Suppose that every row is unimodular ($u_n A_n + v_n B_n = 1$ for some u_n, v_n) and every adjacent minor vanishes:*

$$A_n B_{n+1} - B_n A_{n+1} = 0 \quad (n \geq 0).$$

Then every pairwise minor $A_i B_j - B_i A_j$ vanishes. In particular, take $R_0 = \mathbb{Z}/(2^S 3^R) \mathbb{Z}$ with $R > 0$. If $S + 2R \leq k$, there are two distinct binary selectors $s, t \in \{0, 1\}^k$ such that

$$\sum_{i < k} s_i w_i = \sum_{i < k} t_i w_i.$$

Thus the sufficient width is $S + 2R$, rather than the ambient two-coordinate width $2S + 4R$.

Proof. A Bézout identity makes each row a unimodular anchor; a unit coordinate is the special case already recorded. Vanishing of the next minor therefore writes the next row as a scalar multiple of the current one; induction places the entire tail on the line through w_0 and proves the pairwise-minor assertion. A determinant-one Bézout shear sends w_0 to $(1, 0)$, so all selector sums have only one free residue coordinate and occupy at most 2^{S3^R} values. Finally

$$2^{S3^R} < 2^{S4^R} = 2^{S+2R} \leq 2^k,$$

and pigeonhole gives the two selectors. The ring-generic minor collapse and the modular selector collision are together [Lean source](#). $\square$

Primitive integer rows are unimodular modulo every modulus. A particular coordinate need not be a unit: $(2, 3)$ modulo 6 is an example. The minor-vanishing hypothesis remains a separate condition on the source rows.

4.1 One minor gcd controls two different costs

Let a rank-two lattice $\Lambda \subset \mathbb{Z}^2$ be generated by primitive rows, and let $g > 0$ be the gcd of their 2×2 minors. Its Smith invariants are $1, g$. Therefore

$$|\text{im}(\Lambda \bmod D)| = \frac{D^2}{\gcd(g, D)}, \quad \left[\mathbb{Z}^2 : \frac{\Lambda \cap D\mathbb{Z}^2}{D} \right] = \frac{g}{\gcd(g, D)}.$$

To verify the formulas, make a unimodular change of coordinates sending Λ to $\mathbb{Z} \oplus g\mathbb{Z}$; that change preserves $D\mathbb{Z}^2$. The first quotient counts the modular signatures. The second is the index remaining after dividing a successful collision by D . Once $D \mid g$, the modular count is D , while the remaining index is g/D .

If two independent divided rows (A_i, B_i) have $|A_i| \leq H$ and $|A_i \zeta - B_i| \leq \varepsilon$, their determinant gives

$$2H\varepsilon \geq \frac{g}{\gcd(g, D)}.$$

This is a restriction on two independent forms in the divided lattice. For one nonzero integral form tending to zero, no additional product $H\varepsilon \rightarrow 0$ is required.

4.2 A collision must have a small nonzero real remainder

A real bin records the analytic requirement alongside the modular signature. The multiplicity to control is the number of equal real values within one modular fibre.

Theorem 4.6 (quantitative bounded-fibre escape). *Let A, B, J be finite sets and let $f : A \rightarrow B$, $g : A \rightarrow \mathbb{R}$ and $\iota : A \rightarrow J$. Suppose that each simultaneous fibre of (f, g) has at most k elements and that, for some $\delta > 0$,*

$$\iota(x) = \iota(y) \implies |g(x) - g(y)| < \delta.$$

If $|B||J|k < |A|$, then some distinct $x, y \in A$ satisfy

$$f(x) = f(y), \quad 0 < |g(x) - g(y)| < \delta.$$

Proof. Partition A by (f, ι) . Some cell has more than k elements, so its g -values cannot all agree. Two unequal values in that cell have the same modular signature and differ by less than δ . $\square$

For primitive integer rows (A_j, B_j) , put $e_j = A_j F(3/2) - B_j$ and apply the theorem to binary selectors, with

$$f(\varepsilon) = \sum_j \varepsilon_j (A_j, B_j) \pmod{D}, \quad g(\varepsilon) = \sum_j \varepsilon_j e_j.$$

Let Q be the number of attained modular signatures and let k bound the simultaneous (f, g) fibres. All selector remainders lie in an interval of length $T = \sum_j |e_j|$. Dividing that interval into half-open bins of width D/n gives the sufficient inequality

$$2^M > Qk \left(\left\lfloor \frac{nT}{D} \right\rfloor + 1 \right). \quad (1)$$

Its conclusion is a signed row sum divisible coordinatewise by D , with a nonzero divided real remainder of absolute value less than $1/n$. More precisely, if modular fibre b has its own real span T_b and exact value multiplicity k_b , it suffices that

$$2^M > \sum_b k_b \left(\left\lfloor \frac{nT_b}{D} \right\rfloor + 1 \right).$$

These are estimates for the primitive real remainders of the declared source family. Bounded unnormalised hypergeometric remainders cannot replace them. A nonzero function need not be nonzero at $3/2$, as the factor $2X - 3$ shows; positive individual remainders need not remain positive after subtraction.

4.3 The source-specific endpoint question

The following sufficient construction keeps source membership, primitive scaling, local cancellation and the real estimate together. Without source and height restrictions, arbitrary polynomial lifts of small rational approximations would simply restate irrationality.

Problem 4.7 (common-width simultaneous endpoint-jet construction). Exhibit an integer constant $C \geq 1$ and, for every sufficiently large positive integer n , positive integers W_n, R_n, S_n, M_n such that

$$n^2 \leq W_n, R_n, S_n \leq Cn^2, \quad 4R_n + 2S_n \leq M_n \leq Cn^2,$$

together with polynomial pairs coming from a named source family with an explicit coefficient-height bound after primitive normalisation, $(U_{n,j}, V_{n,j}) \in \mathbb{Z}[X]^2$ for $0 \leq j < M_n$, each of degree at most the common declared width W_n , whose specialised integer rows are primitive:

$$\gcd(H_{W_n}(U_{n,j}), H_{W_n}(V_{n,j})) = 1.$$

Find a nonzero vector $\lambda^{(n)} \in \{-1, 0, 1\}^{M_n}$ for which, on putting

$$U_n = \sum_{j < M_n} \lambda_j^{(n)} U_{n,j}, \quad V_n = \sum_{j < M_n} \lambda_j^{(n)} V_{n,j},$$

the pair (U_n, V_n) is not $(0, 0)$, all four common-width jets vanish,

$$J_{3,R_n}(U_n) = J_{3,R_n}(V_n) = 0, \quad J_{2,S_n}(U_n) = J_{2,S_n}(V_n) = 0,$$

where every jet in this display is formed using the declared width W_n , and the resulting divided integer linear form

$$A_n = \frac{H_{W_n}(U_n)}{3^{R_n}2^{S_n}}, \quad B_n = \frac{H_{W_n}(V_n)}{3^{R_n}2^{S_n}}, \quad \rho_n = A_n F(3/2) - B_n$$

satisfies the explicit analytic condition

$$0 < |\rho_n| < \frac{1}{n}.$$

The jet equations make A_n, B_n integers. If $F(3/2) = a/b$ were rational, a nonzero ρ_n would have absolute value at least $1/|b|$, contradicting $|\rho_n| < 1/n$ for large n . The required analytic inequality is exactly the displayed scalar condition. For irrationality it suffices to obtain such forms along any unbounded sequence of n ; the all-sufficiently-large- n formulation is a stronger construction requirement. Height estimates become relevant when a specific construction uses them to obtain this scalar inequality, or when an independent-row determinant is invoked.

For the literal two-parameter source deformations, the long record keeps the row-wise primitive normalisation, the two local minor valuations and the remaining real-error minimum together. The narrow regular-scale family has an ordinary determinant obstruction to bounded divided errors; that obstruction does not exclude sparse scales or the wider deformation. Thus the next estimate must concern the real remainders of the chosen family. Increasing the number of modular collisions alone does not establish it.

Functional equations. The classification of Bell and Smertnig implies that $L(z) = \sum_{n \geq 1} \tau(n)z^n$ is not k -Mahler for any $k \geq 2$ [6, Introduction]. The earlier simultaneous-2/3 obstruction is consequently subsumed by this known single-base result. A construction using additional functions or functional relations must specify those functions and its closure conditions; the single-base statement is not an obstruction to every approximation method.

Statements and declarations

Artefact and data availability. The [pinned formal-source revision](#) contains the Lean sources, the fixed toolchain, and the library manifest used in the verification. The ordinary proofs used here are printed with their hypotheses.

Funding and competing interests. This work received no external funding. The author declares no competing interests.

Acknowledgements. The problem numbering and status follow the Erdős Problems catalogue maintained by Thomas Bloom [5].

A Guide to the formal sources

Each linked phrase opens its Lean declaration at the pinned source revision `bb4e24651b37`. The declarations of this note live in seven modules: `RationalBaseLambert`, `QApéryDiagonalNonEquivalence`, `RationalPadéArithmetic`, `ZudilinConeArithmetic`, `ZudilinHeightRegion`, `HermitePadéNoGo`, and `BezoutPluckerJets`. The first contains the corridor, cleared-tail recurrence, and elementary $7/2$ certificate; the second checks the finite $n = 0$ diagonal residual; the remaining four separate the Padé exponent arithmetic, endpoint arithmetic, logarithmic comparisons, rectangular exponent model, and Bézout–Plücker tail collapse. The link coordinates are validated against that pinned revision, so they remain correct as later work moves lines in the working tree.

References

- [1] P. Erdős, *On the irrationality of certain series: problems and results*, in A. Baker (ed.), *New Advances in Transcendence Theory*, Cambridge UP, 1988, pp. 102–109, doi:[10.1017/CBO9780511897184.009](https://doi.org/10.1017/CBO9780511897184.009).
- [2] P. Bundschuh and K. Väänänen, *Arithmetical investigations of a certain infinite product*, *Compositio Math.* **91** (1994), no. 2, 175–199.
- [3] W. Zudilin, *Heine’s basic transform and a permutation group for q -harmonic series*, *Acta Arith.* **111** (2004), no. 2, 153–164, doi:[10.4064/aa111-2-4](https://doi.org/10.4064/aa111-2-4).
- [4] W. Zudilin, *On the irrationality of generalized q -logarithm*, arXiv:1601.02688; *Res. Number Theory* **2** (2016), doi:[10.1007/s40993-016-0042-x](https://doi.org/10.1007/s40993-016-0042-x). The remark that the results extend to non-integer $p = r/s$, $|p| > 1$, under an assumption $\log |r| > c \log |s|$ for a computable $c > 0$, is at the end of Section 2; no value of c is computed there, and the remark is made for the generalized q -logarithm of that paper.
- [5] T. F. Bloom, *Erdős Problem #1049*, erdosproblems.com/1049, accessed 28 July 2026 (page displays “last edited 28 September 2025”). The current record labels the problem open, cites [Er88c, p. 102] and [Er48], and explicitly describes its status as the website owner’s present assessment rather than a literature-completeness guarantee.
- [6] J. Bell and D. Smertnig, *Mahler series with multiplicative coefficient sequences*, arXiv:2603.23456v1, 24 March 2026. The introduction explicitly includes the divisor and totient functions among the examples which are not k -Mahler for any $k \geq 2$.