

Cresko Laboratory Procedures and Protocols

Cresko Laboratory

Friday, December 8, 2023

Table of contents

How to use this book	5
I General Laboratory Protocols	6
1 Contact Information	8
II Molecular Protocols	9
2 cDNA basic	11
2.1 Introduction	11
2.2 Materials:	11
2.3 Solutions:	11
2.4 Procedure:	11
First strand synthesis	11
Reaction can be scaled up to accommodate more starting RNA	12
3 2x Turbo	13
3.1 Introduction	13
3.2 Materials:	13
3.3 Solutions:	13
3.4 Procedure:	14
4 Paraformaldehyde	15
4.1 Introduction	15
4.2 Materials:	15
4.3 Solutions:	15
4.4 Procedure:	15
5 Alizarin Staining	17
5.1 Introduction	17
5.2 Materials:	17
5.3 Solutions:	17
5.4 Procedure:	18

III Microbiology	20
6 Placeholder_Microbiology	22
6.1 xxx	22
xxx	22
IV Vertebrate Husbandry	23
7 Aquarium Cleaning	25
7.1 Introduction	25
7.2 Materials:	25
7.3 Solutions:	25
7.4 Procedure:	26
7.5 Air difuser cleaning:	26
8 Artemia Decapsulation	27
8.1 Introduction	27
8.2 Materials:	27
8.3 Solutions:	28
8.4 Procedure:	28
V Daphnia Husbandry	30
9 Placeholder_Daphnia	32
9.1 xxx	32
xxx	32
VI Bioinformatic	33
10 A field guide to base R	35
10.1 Introduction	35
Prerequisites	35
10.2 Selecting multiple elements with [.	36
Subsetting vectors	36
10.3 Summary	37
11 Summary	38
References	39

Appendices	40
A Sbf1 Barcodes in 96 Well Plate	40
B Gel electrophoresis tips	48
B.1 Introduction	48
B.2 Common Mistakes	49
B.3 Potentially dangerous and/or destructive mistakes:	53
B.4 Conscientiousness	54

How to use this book

This is a Quarto book that contains all of the Procedures and Protocols for the Cresko Laboratory in the Institute of Ecology and Evolution at the University of Oregon.

The book is organized into major sections that contain

- General Laboratory Protocols or the lab
- More detailed Laboratory Protocols
- Husbandry protocols for vertebrate animals primarily stickleback and pipefish, but also zebrafish
- Husbandry protocols for *Daphnia*
- Bioinformatic protocols including how to get on to **Talapas**

You can scroll through the book using the index on the left, but also use the search field to find all relevant protocols.

There are also useful appendices at the end, as well as a section for the references cited throughout the book.

This book was written in Markdown using Quarto. To learn more about Quarto books visit <https://quarto.org/docs/books>.

Part I

General Laboratory Protocols

This section of the book contains general protocols for working in the laboratory.

1 Contact Information

Col1	Col2	Col3
Cresko Laboratory	541-346-5189	Phone
Bill Cresko	541-285-5446	Cell
Mark Currey	541-505-0006	Cell
Susie Bassham	xxxx	Cell

Part II

Molecular Protocols

This section of the book contains common protocols used for molecular biology and genomics in the laboratory. These include standard protocols such as setting up creating reagents, setting up PCRs and running gels, as well as advanced protocols such as creating constructs.

2 cDNA basic

2.1 Introduction

- **Purpose:** This procedure describes how to synthesis cDNA for use with PCR.
- **Procedure Type:** Molecular
- **Species:** N/A

2.2 Materials:

- 2 µl Oligo d(T)23 VN (50 µM, NEB; anchored-dT primer)*
- X µl up to 5 µg total RNA
- 1 µl 10 mM dNTP
- water
- 2 µl 10x RT buffer (Invitrogen)
- 4 µl 25 mM MgCl₂
- 2 µl 0.1 mM DTT – Invitrogen
- 1 µl RNase inhibitor – e.g., RNaseOUT (Invitrogen)
- 1 µl Superscript III reverse transcriptase (200 u/µl – Invitrogen)

2.3 Solutions:

NONE

2.4 Procedure:

First strand synthesis

Combine:

- 2 µl Oligo d(T)23 VN (50 µM, NEB; anchored-dT primer)*
- X µl up to 5 µg total RNA

- 1 μ l 10 mM dNTP mix
- Water (if necessary) to bring total to 10 μ l

Heat to 65°C for 5 min., then ice

Collect contents at bottom of tube by brief centrifugation.

Add:

- 2 μ l 10x RT buffer (Invitrogen)
- 4 μ l 25 mM MgCl₂
- 2 μ l 0.1 mM DTT – Invitrogen
- 1 μ l RNase inhibitor – e.g., RNaseOUT (Invitrogen)
- 1 μ l Superscript III reverse transcriptase (200 u/ μ l – Invitrogen)

Mix by gentle aspiration

- 25°C for 5 min.

Reaction can be scaled up to accommodate more starting RNA

Synthesis: Incubate at 50°C for 50 min.

Inactivation: 85°C for 5 min. Chill on ice, collect contents to bottom by short spin.

Destroy RNA template: 1 μ l RNase H (2 u/ μ l), incubate at 37°C for 20 min.

Proceed to PCR. Depending on expression level, may be able to use a dilution of cDNA as template – try 1:50 dilution in EB, use 2 μ l as template in a 20 μ l reaction. Don't dilute your entire amount of cDNA, as some products may require a higher concentration of template.

3 2x Turbo

3.1 Introduction

- **Purpose:** This procedure describes how to create 2x Turbo PCR mix.
- **Procedure Type:** Molecular
- **Species:** N/A

3.2 Materials:

- 33,000 μ l npH₂O
- 2000 μ l MgSO₄ (100mM)
- 1600 μ l 1M Tris-HCl (pH 8.6)
- 800 μ l 1M KCl
- 800 μ l 1M (NH₄)₂SO₄
- 800 μ l Triton-X 100 (10%)
- 400 μ l DMSO (100 %)
- 120 μ l dATP (100mM)
- 120 μ l dGTP (100mM)
- 120 μ l dTTP (100mM)
- 120 μ l dCTP (100mM)
- 80 μ l 100mg/ml BSA

Total = 40 ml of buffer

3.3 Solutions:

NONE

3.4 Procedure:

- Mix above reagents together
- Place in 1.5 ml ependorph tubes
- Store at -20C

4 Paraformaldehyde

4.1 Introduction

- **Purpose:** This procedure describes how to make 8% paraformaldehyde. This protocol is the one I have used and makes use of pH to get the PFA into solution relatively quickly - then you readjust the pH. It's for 8% - then you can add 1:1 2x PBS.
- **Procedure Type:** Molecular
- **Species:** N/A

4.2 Materials:

- xxx

Total = 40 ml of buffer

4.3 Solutions:

NONE

4.4 Procedure:

HUMAN HEALTH WARNING

Paraformaldehyde can be hazardous to your health - make sure you prepare in the fume hood.

- Add 40 g Paraformaldehyde to 450 ml distilled water (or scale for desired final volume).
- Add 1 ul of 10 N NaOH per ml of water (i.e. 500 ul for 500 ml).
- Apply medium heat while stirring at medium speed to dissolve - approx 15-20 min.
- Solution should not go above 60° C.

- Eventually, granules will fully dissolve and the solution will become translucent.

i DO NOT LET THE SOLUTION STIR BEYOND THIS POINT


It will form a fuzzy precipitate that reduces the solution strength after filtering.

- Once the granules have dissolved and the solution clears, turn off the heat and equilibrate to pH 7.4 with approx 1.5 ml of 20% HCl (or scale, depending on target volume).
- Bring volume to 500 ml (or scaled volume) with distilled water.
- Filter while still warm to 0.45 um (or 0.2 um). Aliquot and store at -20° C.

5 Alizarin Staining

5.1 Introduction

- **Purpose:** Alizarin staining of fixed adult stickleback.
- **Procedure Type:** Molecular
- **Species:**
 - Threespine stickleback, *Gasterosteus aculeatus*
 - xxx
- **Authors**
 - xxx

 Schedule for Cleaning

PLACEHOLDER

5.2 Materials:

- Alizarin red S
- KOH
- H₂O₂
- NaOH
- MESAB
- PFA

5.3 Solutions:

- **0.5% Alizarin red S Stock:**
 - To make 50 mls add 0.25g alizarin red S powder to 50 ml water.
- **0.025% Alizarin Stain**

- To make 100 mls: Add 500µl 0.5% alizarin red S (stock) to 99.5ml 1% KOH
 - 1 Liter: Add 5ml 0.5% alizarin red S (stock) to 995ml (1 liter) 1%KOH
- **3% H2O2/0.5%KOH:** Mix and keep at 4C; Before using, bring to room temperature to hold down introducing bubbles under the skin: 0.5ml 6%H2O2 & 0.5ml 1%KOH.
- **MESAB: Tricaine:** 3-amino benzoic acid ethyl ester from Sigma (Cat # A-5040). Mix in fish safe container with a stir bar:
 - 400 mg tricaine powder
 - 800 mg Na2HPO4 (anhydrous)
 - 100 ml glass distilled water
 - Adjust to ~pH 7 with a drop at a time of 1N NaOH or 1N HCl if needed but it's usually right if you weigh the sodium phosphate carefully and measure the water with a graduated cylinder.
 - For storage: Aliquot into 6 x 25 ml fish safe plastic bottles and store at 4C. Label with date made and use within a couple of weeks.
- **8% PFA:**
 - 8 g Pelleted PFA (Ted Pella, Inc.; cat# 18501)
 - 90 ml dH2O
 - 25 drops 1N NaOH
 - Heat at very low heat and stir until solution clears.
 - Add 25 drops 1N HCl. pH should be 7.0-7.2.
 - Filter and store at 4C not more than 1 week.
 - Use as 4% PFA: dilute 1:1 with 2X PBS, do not store solution more than a few hours.
- **2X PBS**
 - 1.6% NaCl
 - 0.04% KCl
 - 0.04 M PO4 pH 7.0- 7.3

5.4 Procedure:

- **Day One**
 - 2h-8h at R/T depending on size on shaker.
 - Without agitation and with lid open until eyes start to lighten and all skin pigment is gone (usually about an hour or more)
- **Day Two**
 - xxx
 - xxx

— XXX

Part III

Microbiology

This section of the book contains protocols that involve the use of microbes, mostly in the context of host-microbe interactions, but also standard protocols for cloning of nucleic acids using microbes.

6 Placeholder_Microbiology

6.1 xxx

XXXX

xxx

XXXXX

Part IV

Vertebrate Husbandry

This section of the manual contains protocols for the safe and ethical husbandry and use of vertebrate animals, particular the fish models stickleback, zebrafish and syngnathids.

7 Aquarium Cleaning

7.1 Introduction

- **Purpose:** This procedure describes how to clean 20 gallon glass tanks.
- **Procedure Type:** Husbandry
- **Species:**
 - Threespine stickleback, (*Gasterosteus aculeatus*),
 - Gulf pipefish (*Syngnathus scovelli*)
-

Schedule for Cleaning

Tank cleaning is to be done ONLY Monday - Friday

7.2 Materials:

- Scrub pad or sponge
- Cart (you may or may not want to use)
- Old clothes (this can be messy)
- Personal protection equipment (Splash proof glasses or face shield).

7.3 Solutions:

- **Bleach solution:** Make a 10% bleach solution in a 2 gallon bucket. Add 4.5 L of water. Add 0.5 L of bleach and gently stir.
- **Sodium thiosulfate:** Make a 3% solution of sodium thiosulfate in a separate 2 gallon bucket. Add 5 L of water (to line) and 150g (marked on dispenser) of sodium thiosulfate. Mix

Note: When using bleach and/or sodium thiosulfate. Eye protection is required. Please use splash proof glasses or a face shield when using bleach and sodium thiosulfate.

7.4 Procedure:

- Complete bleaching and cleaning of tank. This needs to be done to each tank every 2 months.
- Remove fish from tank and put them into a clean tank. Tanks that are emptied of fish need to be cleaned and sterilized before another batch of fish can be introduced.
- Drain the tank and remove it from the rack. Clean air diffuser as instructed below.
 - Clean the tank and all parts thoroughly with a scrub pad, taking care not to damage the silicon water seals on the inside (algae should be left if very gentle rubbing will not remove it).
 - Squirt about 10 – 20 mls of bleach into the tank. Wash the bleach water thoroughly around the inside of the tank by hand using a pad or sponge exposing all inside portions of the tank to bleach.
 - Rinse the tank thoroughly with hot tap water. Rinse the tank with sodium thiosulfate, and then rinse it again with hot water. Put a few thiosulfate crystals into the tank and leave it.
 - Reassemble the tank and put it back on the rack. Fill with system water and allow water to recirculate for about 30 minutes before adding fish. Watch fish for 15min to look for any signs of distress.
 - Using a dry erase marker record date/time on the front of the tank when system water is turned back on.
- Initial the check list that you have completed the tank cleaning.

7.5 Air difuser cleaning:

- Remove dirty air diffusers from tanks and rinse with tap water to remove excess algae and debris.
- Place in 10% bleach solution for 15-30 minutes.
- Rinse the corner filters with hot water for 5 and then place into 3% sodium thiosulfate for 5 minutes.
- Rinse with hot water for 5 minutes.
- When cleaned air diffusers are placed back into aquaria, observe fish for 15 min for signs of distress.

8 Artemia Decapsulation


8.1 Introduction

- **Purpose:** This procedure describes standard practices for decapsulating brine shrimp. Although brine shrimp can be hatched, collected and then fed to fish, the cysts are often hard to separate from the newly hatched brine shrimp and can be ingested by stickleback and pipefish. To reduce this phenomenon we can decapsulate brine shrimp in advance, and then leave them in a suspended state in the freezer for an extended period of time (~ xxx weeks) before they are hatched.
- **Procedure Type:** Husbandry
- **Species:**
 - Threespine stickleback, (*Gasterosteus aculeatus*),
 - Gulf pipefish (*Syngnathus scovelli*)
- **Authors**
 - Mark C. Currey

8.2 Materials:

- 15 oz can of dried Artemia cysts (approximately 430 g)
- 4.3 L ~6% laundry grade bleach
- Rock Salt (NaCl)
- 125 ml 40% Lye (NaOH) solution
- 30.0 g Sodium thiosulfate ($\text{Na}_2\text{S}_2\text{O}_3$)
- 16 L Hatching Cone with aeration
- 125 μm mesh bag (Aquatic Eco-Systems PMB3, 125 micron x 18")
- Several 3-5 L beakers
- (1-2) Squirt bottles - squeeze type

8.3 Solutions:

 Be ready

Solutions should be prepared in advance.

- Bleach, ~6% laundry grade
 - 25 ppt Salt Solution
 - Combine: 50 g Rock Salt (NaCl) To 2.0 L with tap water
 - Stir to dissolve completely.
- 40% Lye (NaOH) solution
 - Combine: 200 g Lye (NaOH) To 500 mL with tap water
 - Stir to dissolve completely.
 - Store in refrigerator (4°C)
- Buffered Salt Solution
 - Combine: 2L, 25 ppt Salt Solution
 - 125 mL 40% Lye Solution, pre-chilled to 4°C
- 1.0% Sodium Thiosulfate
 - Combine: 30 g sodium thiosulfate To 3.0 L with tap water
 - Stir to dissolve.
- Saturated Brine
 - Combine: ~25g Rock Salt to 4.0 L with tap water
 - Aerate to dissolve.

8.4 Procedure:

1. **Cyst hydration:** Hydrate one full can of dried cyst in 5 L of tap water in a hatching cone with aeration for 1 hour at room temp. Examine the cyst under a dissecting scope with top lighting before proceeding. Dry cysts are dimpled, resembling a deflated basketball, whereas fully hydrated cysts are completely spherical in shape. The cysts must be fully hydrated prior to the de-capsulation step. If cysts are not completely spherical after 1 hour, continue the hydration process (for a maximum of 2 hours), checking the progress of the cysts under a microscope every 15 min.
2. **Filter and rinse cysts:** Collect the hydrated cyst in a 125 um mesh bag and rinse with cool tap water.

3. **Transfer cysts back to the cone:** Add the Buffered Salt Solution to the cone and aerate (save back a filled squirt bottle of salt solution to help transfer cysts to cone). Transfer cysts into cone.
4. **De-capsulation:** Add the bleach (4.3 L) to the cone and continue aeration. Watch the cysts turn from brown to grey to orange, When the cysts are 90% orange, stop the reaction by quickly siphoning the cysts through a 125 um mesh bag and rinsing well with cool tap water.
5. **Neutralization residual chlorine:** To neutralize any residual chlorine transfer the mesh bag to a clean 4 L beaker and pour the 1.0% Sodium Thiosulfate (3L) into the bag. Soak the cysts in the sodium thiosulfate solution for ~1 min, then rinse the cysts with de-ionized tap water. Rinse until discharge turns clear.
6. **Dehydration for long-term storage:** Transfer the cysts back to the cone with 4 L of saturated brine and aerate until salt is dissolved. Transfer dehydrated cyst to (5 or 6) 1 L Nalgene bottles filled with 200 - 300 grams of salt. Add enough salt so that it does not dissolve when de-capsulated brine is added. Fill the bottles with de-capsulated brine. Store in refrigerator. The de-capsulated brine will store for at least 1 month. Hatch brine as you would capsulated brine (see Hatching and Feeding Brine SOP).

Part V

Daphnia Husbandry

This section of the manual contains protocols for the safe and ethical husbandry and use of invertebrate animals, particular the nematode worm *C. remanei* and water fleas of the genus *Daphnia*

9 Placeholder_Daphnia

9.1 xxx

XXXX

xxx

XXXXX

Part VI

Bioinformatic

This section of the book contains protocols for basic bioinformatic skills such as using our laboratory cluster ‘Genome’, as well as our account Nereus on the UO supercomputer Talapas.

Note that there are several appendices that contain greater details and training on things such as the use of command line, R and Python, markdown and literature programming, and documentation using Quarto and Jupyter notebooks.

See Knuth (1984) for additional discussion of literate programming.

10 A field guide to base R

10.1 Introduction

To finish off the programming section, we're going to give you a quick tour of the most important base R functions that we don't otherwise discuss in the book. These tools are particularly useful as you do more programming and will help you read code you'll encounter in the wild.

This is a good place to remind you that the tidyverse is not the only way to solve data science problems. We teach the tidyverse in this book because tidyverse packages share a common design philosophy, increasing the consistency across functions, and making each new function or package a little easier to learn and use. It's not possible to use the tidyverse without using base R, so we've actually already taught you a **lot** of base R functions: from `library()` to load packages, to `sum()` and `mean()` for numeric summaries, to the factor, date, and POSIXct data types, and of course all the basic operators like `+`, `-`, `/`, `*`, `|`, `&`, and `!`. What we haven't focused on so far is base R workflows, so we will highlight a few of those in this chapter.

After you read this book, you'll learn other approaches to the same problems using base R, `data.table`, and other packages. You'll undoubtedly encounter these other approaches when you start reading R code written by others, particularly if you're using StackOverflow. It's 100% okay to write code that uses a mix of approaches, and don't let anyone tell you otherwise!

In this chapter, we'll focus on four big topics: subsetting with `[]`, subsetting with `[[` and `$`, the apply family of functions, and `for` loops. To finish off, we'll briefly discuss two essential plotting functions.

Prerequisites

This package focuses on base R so doesn't have any real prerequisites, but we'll load the tidyverse in order to explain some of the differences.

```
library(tidyverse)
```

10.2 Selecting multiple elements with `[]`

`[]` is used to extract sub-components from vectors and data frames, and is called like `x[i]` or `x[i, j]`. In this section, we'll introduce you to the power of `[]`, first showing you how you can use it with vectors, then how the same principles extend in a straightforward way to two-dimensional (2d) structures like data frames. We'll then help you cement that knowledge by showing how various dplyr verbs are special cases of `[]`.

Subsetting vectors

There are five main types of things that you can subset a vector with, i.e., that can be the `i` in `x[i]`:

1. **A vector of positive integers.** Subsetting with positive integers keeps the elements at those positions:

```
x <- c("one", "two", "three", "four", "five")
x[c(3, 2, 5)]
```

```
[1] "three" "two"   "five"
```

By repeating a position, you can actually make a longer output than input, making the term “subsetting” a bit of a misnomer.

```
x[c(1, 1, 5, 5, 5, 2)]
```

```
[1] "one"  "one"  "five" "five" "five" "two"
```

2. **A vector of negative integers.** Negative values drop the elements at the specified positions:

```
x[c(-1, -3, -5)]
```

```
[1] "two"  "four"
```

3. **A logical vector.** Subsetting with a logical vector keeps all values corresponding to a TRUE value. This is most often useful in conjunction with the comparison functions.

```
x <- c(10, 3, NA, 5, 8, 1, NA)

# All non-missing values of x
x[!is.na(x)]
```

```
[1] 10 3 5 8 1
```

```
# All even (or missing!) values of x
x[x %% 2 == 0]
```

```
[1] 10 NA 8 NA
```

Unlike `filter()`, NA indices will be included in the output as NAs.

4. **A character vector.** If you have a named vector, you can subset it with a character vector:

```
x <- c(abc = 1, def = 2, xyz = 5)
x[c("xyz", "def")]
```

```
xyz def
5 2
```

As with subsetting with positive integers, you can use a character vector to duplicate individual entries.

5. **Nothing.** The final type of subsetting is nothing, `x[]`, which returns the complete `x`. This is not useful for subsetting vectors, but as we'll see shortly, it is useful when subsetting 2d structures like tibbles.

10.3 Summary

In this chapter, we've shown you a selection of base R functions useful for subsetting and iteration. Compared to approaches discussed elsewhere in the book, these functions tend to have more of a “vector” flavor than a “data frame” flavor because base R functions tend to take individual vectors, rather than a data frame and some column specification. This often makes life easier for programming and so becomes more important as you write more functions and begin to write your own packages.

This chapter concludes the programming section of the book. You've made a solid start on your journey to becoming not just a data scientist who uses R, but a data scientist who can *program* in R. We hope these chapters have sparked your interest in programming and that you're looking forward to learning more outside of this book.

11 Summary

In summary, this book has no content whatsoever.

1 + **1**

[1] 2

References

Knuth, Donald E. 1984. “Literate Programming.” *Comput. J.* 27 (2): 97–111. <https://doi.org/10.1093/comjnl/27.2.97>.

A Sbf1 Barcodes in 96 Well Plate

well	Name Barcode(top)	Final top sequence	well	Name (bottom)	Final bottom sequence
A1	AAACSGI- AAACGG- top	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT	A1	SGI- AAACGG- bot	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT
A2	AACGSGI- AACGTT- top	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT	A2	SGI- AACGTT- bot	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT
A3	AACTSGI- AACTGA- top	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT	A3	SGI- AACTGA- bot	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT
A4	AAGASGI- AAGACG- top	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT	A4	SGI- AAGACG- bot	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT
A5	AAGCSGI- AAGCTA- top	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT	A5	SGI- AAGCTA- bot	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT
A6	AATATSGI- AATATC- top	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT	A6	SGI- AATATC- bot	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT
A7	AATGSGI- AATGAG- top	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT	A7	SGI- AATGAG- bot	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT
A8	ACAASGI- ACAAGA- top	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT	A8	SGI- ACAAGA- bot	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT
A9	ACAGSGI- ACAGCG- top	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT	A9	SGI- ACAGCG- bot	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT
A10	ACATASGI- ACATAC- top	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT	A10	SGI- ACATAC- bot	ACACTCTTTCCCTACACGACGCTTTCCTTCG/5PAA/CTCAATCGATCC*AGAAGAGCGT

Name			Name		
well	Barcode	Final top sequence	well	(bottom)	Final bottom sequence
A11	ACCATG-S6H- top	ACACTCTTTCCCTACACGACGAGCTGCTTAAG	A11	ACCATG-S6H- bot	ACACTCTTTCCCTACACGACGAGCTGCTTAAG
A12	ACCCG-S6H- top	ACACTCTTTCCCTACACGACGAGCTGCTTAAG	A12	ACCCG-S6H- bot	ACACTCTTTCCCTACACGACGAGCTGCTTAAG
B1	ACTCTT-S6H- top	ACACTCTTTCCCTACACGACGAGCTGCTTAAG	B1	ACTCTT-S6H- bot	ACACTCTTTCCCTACACGACGAGCTGCTTAAG
B2	ACTGGC-S6H- top	ACACTCTTTCCCTACACGACGAGCTGCTTAAG	B2	ACTGGC-S6H- bot	ACACTCTTTCCCTACACGACGAGCTGCTTAAG
B3	AGCCAT-S6H- top	ACACTCTTTCCCTACACGACGAGCTGCTTAAG	B3	AGCCAT-S6H- bot	ACACTCTTTCCCTACACGACGAGCTGCTTAAG
B4	AGCGCA-S6H- top	ACACTCTTTCCCTACACGACGAGCTGCTTAAG	B4	AGCGCA-S6H- bot	ACACTCTTTCCCTACACGACGAGCTGCTTAAG
B5	AGGGTC-S6H- top	ACACTCTTTCCCTACACGACGAGCTGCTTAAG	B5	AGGGTC-S6H- bot	ACACTCTTTCCCTACACGACGAGCTGCTTAAG
B6	AGGTGT-S6H- top	ACACTCTTTCCCTACACGACGAGCTGCTTAAG	B6	AGGTGT-S6H- bot	ACACTCTTTCCCTACACGACGAGCTGCTTAAG
B7	AGTAGG-S6H- top	ACACTCTTTCCCTACACGACGAGCTGCTTAAG	B7	AGTAGG-S6H- bot	ACACTCTTTCCCTACACGACGAGCTGCTTAAG
B8	AGTTAA-S6H- top	ACACTCTTTCCCTACACGACGAGCTGCTTAAG	B8	AGTTAA-S6H- bot	ACACTCTTTCCCTACACGACGAGCTGCTTAAG
B9	ATAGTA-S6H- top	ACACTCTTTCCCTACACGACGAGCTGCTTAAG	B9	ATAGTA-S6H- bot	ACACTCTTTCCCTACACGACGAGCTGCTTAAG
B10	ATCAAA-S6H- top	ACACTCTTTCCCTACACGACGAGCTGCTTAAG	B10	ATCAAA-S6H- bot	ACACTCTTTCCCTACACGACGAGCTGCTTAAG
B11	ATGCAC-S6H- top	ACACTCTTTCCCTACACGACGAGCTGCTTAAG	B11	ATGCAC-S6H- bot	ACACTCTTTCCCTACACGACGAGCTGCTTAAG

Name			Name		
well	Barcode	Final top sequence	well	(bottom)	Final bottom sequence
B12	ATGTTG-SBH- ATGTTG- top	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT	B12	ATGTTG-SBH- ATGTTG- bot	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT
C1	ATTCCG-SBH- ATTCCG- top	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT	C1	ATTCCG-SBH- ATTCCG- bot	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT
C2	CAAAAA-SBH- CAAAAA- top	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT	C2	CAAAAA-SBH- CAAAAA- bot	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT
C3	CAATCG-SBH- CAATCG- top	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT	C3	CAATCG-SBH- CAATCG- bot	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT
C4	CACCTC-SBH- CACCTC- top	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT	C4	CACCTC-SBH- CACCTC- bot	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT
C5	CAGGCA-SBH- CAGGCA- top	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT	C5	CAGGCA-SBH- CAGGCA- bot	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT
C6	CATACT-SBH- CATACT- top	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT	C6	CATACT-SBH- CATACT- bot	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT
C7	CCATTT-SBH- CCATTT- top	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT	C7	CCATTT-SBH- CCATTT- bot	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT
C8	CCCGGT-SBH- CCCGGT- top	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT	C8	CCCGGT-SBH- CCCGGT- bot	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT
C9	CCCTAA-SBH- CCCTAA- top	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT	C9	CCCTAA-SBH- CCCTAA- bot	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT
C10	CCGAGG-SBH- CCGAGG- top	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT	C10	CCGAGG-SBH- CCGAGG- bot	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT
C11	CCGCAT-SBH- CCGCAT- top	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT	C11	CCGCAT-SBH- CCGCAT- bot	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT
C12	CCTAAC-SBH- CCTAAC- top	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT	C12	CCTAAC-SBH- CCTAAC- bot	ACACTCTTTCCCTACACGACGAGTGGTCTTCCGATGCTCAGGAAGAGCGT

Name			Name		
well	Barcode	Final top sequence	well	(bottom)	Final bottom sequence
D1	CGAGGCG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCT	D1	CGAGGCG- bot	ACACTCTTTCCCTACACGACGCTCTTCCGATCT
D2	CGCAGCG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCT	D2	CGCAGCG- bot	ACACTCTTTCCCTACACGACGCTCTTCCGATCT
D3	CGCGTGG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCT	D3	CGCGTGG- bot	ACACTCTTTCCCTACACGACGCTCTTCCGATCT
D4	CGGTCCG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCT	D4	CGGTCCG- bot	ACACTCTTTCCCTACACGACGCTCTTCCGATCT
D5	CGTCTAG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCT	D5	CGTCTAG- bot	ACACTCTTTCCCTACACGACGCTCTTCCGATCT
D6	CGTGATG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCT	D6	CGTGATG- bot	ACACTCTTTCCCTACACGACGCTCTTCCGATCT
D7	CTACAGG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCT	D7	CTACAGG- bot	ACACTCTTTCCCTACACGACGCTCTTCCGATCT
D8	CTCGCCG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCT	D8	CTCGCCG- bot	ACACTCTTTCCCTACACGACGCTCTTCCGATCT
D9	CTGCGAG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCT	D9	CTGCGAG- bot	ACACTCTTTCCCTACACGACGCTCTTCCGATCT
D10	CTGGTGG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCT	D10	CTGGTGG- bot	ACACTCTTTCCCTACACGACGCTCTTCCGATCT
D11	CTTATGG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCT	D11	CTTATGG- bot	ACACTCTTTCCCTACACGACGCTCTTCCGATCT
D12	CTTTGCG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCT	D12	CTTTGCG- bot	ACACTCTTTCCCTACACGACGCTCTTCCGATCT
E1	GAAATGG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCT	E1	GAAATGG- bot	ACACTCTTTCCCTACACGACGCTCTTCCGATCT

Name			Name		
well	Barcode	Final top sequence	well	(bottom)	Final bottom sequence
E2	GAACSAH- GAACCA- top	ACACTCTTTCCCTACACGACGCTTTCTTCGTAATCTCATGCTG	E2	SAH- GAACCA- bot	GAACGAGCGGTTGTAATCTCATGCTG
E3	GACGSAH- GACGAC- top	ACACTCTTTCCCTACACGACGCTTTCTTCGTAATCTCATGCTG	E3	SAH- GACGAC- bot	GACGAGCGGTTGTAATCTCATGCTG
E4	GACTSAH- GACTCT- top	ACACTCTTTCCCTACACGACGCTTTCTTCGTAATCTCATGCTG	E4	SAH- GACTCT- bot	GACTAGCGGTTGTAATCTCATGCTG
E5	GAGASAH- GAGAGA- top	ACACTCTTTCCCTACACGACGCTTTCTTCGTAATCTCATGCTG	E5	SAH- GAGAGA- bot	GAGAGAGCGGTTGTAATCTCATGCTG
E6	GATCSAH- GATCGT- top	ACACTCTTTCCCTACACGACGCTTTCTTCGTAATCTCATGCTG	E6	SAH- GATCGT- bot	GATCAGCGGTTGTAATCTCATGCTG
E7	GCAGSAH- GCAGAT- top	ACACTCTTTCCCTACACGACGCTTTCTTCGTAATCTCATGCTG	E7	SAH- GCAGAT- bot	GCAGAGCGGTTGTAATCTCATGCTG
E8	GCATSAH- GCATGG- top	ACACTCTTTCCCTACACGACGCTTTCTTCGTAATCTCATGCTG	E8	SAH- GCATGG- bot	GCATAGCGGTTGTAATCTCATGCTG
E9	GCCGSAH- GCCGTA- top	ACACTCTTTCCCTACACGACGCTTTCTTCGTAATCTCATGCTG	E9	SAH- GCCGTA- bot	GCCGAGCGGTTGTAATCTCATGCTG
E10	GCGASAH- GCGACC- top	ACACTCTTTCCCTACACGACGCTTTCTTCGTAATCTCATGCTG	E10	SAH- GCGACC- bot	GCGAGCGGTTGTAATCTCATGCTG
E11	GCGCSAH- GCGCTG- top	ACACTCTTTCCCTACACGACGCTTTCTTCGTAATCTCATGCTG	E11	SAH- GCGCTG- bot	GCGAGCGGTTGTAATCTCATGCTG
E12	GCTCSAH- GCTCAA- top	ACACTCTTTCCCTACACGACGCTTTCTTCGTAATCTCATGCTG	E12	SAH- GCTCAA- bot	GCTAGCGGTTGTAATCTCATGCTG
F1	GGACSAH- GGACTT- top	ACACTCTTTCCCTACACGACGCTTTCTTCGTAATCTCATGCTG	F1	SAH- GGACTT- bot	GGAGAGCGGTTGTAATCTCATGCTG
F2	GGCASAH- GGCAAG- top	ACACTCTTTCCCTACACGACGCTTTCTTCGTAATCTCATGCTG	F2	SAH- GGCAAG- bot	GGAGAGCGGTTGTAATCTCATGCTG

Name			Name		
well	Barcode	Final top sequence	well	(bottom)	Final bottom sequence
F3	GGGCSMI- GGGCGC- top	ACACTCTTTCCCTACACG	F3	SMI-CTTC/5PAA/CTGCCCCAGTTC* GGGCGC- bot	AGGAAGAGCGT
F4	GGGGSMI- GGGGCG- top	ACACTCTTTCCCTACACG	F4	SMI-CTTC/5PAA/CTGCCCCAGTTC* GGGGCG- bot	AGGAAGAGCGT
F5	GGTASMI- GGTACA- top	ACACTCTTTCCCTACACG	F5	SMI-CTTC/5PAA/CTGCAACATATTC* GGTACA- bot	AGGAAGAGCGT
F6	GGTTSMI- GGTTTG- top	ACACTCTTTCCCTACACG	F6	SMI-CTTC/5PAA/CTGCAACATATTC* GGTTTG- bot	AGGAAGAGCGT
F7	GTAASMI- GTAAGT- top	ACACTCTTTCCCTACACG	F7	SMI-CTTC/5PAA/CTGCAACATATTC* GTAAGT- bot	AGGAAGAGCGT
F8	GTATSMI- GTATCC- top	ACACTCTTTCCCTACACG	F8	SMI-CTTC/5PAA/CTGATACCATTC* GTATCC- bot	AGGAAGAGCGT
F9	GTCASMI- GTCATC- top	ACACTCTTTCCCTACACG	F9	SMI-CTTC/5PAA/CTATTCATATTC* GTCATC- bot	AGGAAGAGCGT
F10	GTGCSMI- GTGCCT- top	ACACTCTTTCCCTACACG	F10	SMI-CTTC/5PAA/CTGTCACATATTC* GTGCCT- bot	AGGAAGAGCGT
F11	GTGTASMI- GTGTAA- top	ACACTCTTTCCCTACACG	F11	SMI-CTTC/5PAA/CTTATCATATTC* GTGTAA- bot	AGGAAGAGCGT
F12	GTTGSMI- GTTGGA- top	ACACTCTTTCCCTACACG	F12	SMI-CTTC/5PAA/CTTCAACATATTC* GTTGGA- bot	AGGAAGAGCGT
G1	TAAGSMI- TAAGCT- top	ACACTCTTTCCCTACACG	G1	SMI-CTTC/5PAA/CTGATTCATATTC* TAAGCT- bot	AGGAAGAGCGT
G2	TAATSMI- TAATTC- top	ACACTCTTTCCCTACACG	G2	SMI-CTTC/5PAA/CTAATTAATATTC* TAATTC- bot	AGGAAGAGCGT
G3	TACASMI- TACACA- top	ACACTCTTTCCCTACACG	G3	SMI-CTTC/5PAA/CTTAGACAGTTC* TACACA- bot	AGGAAGAGCGT

Name			Name		
well	Barcode	Final top sequence	well	(bottom)	Final bottom sequence
G4	TACGSGH- TACGGG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCTG	G4	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG
G5	TAGTASBH- TAGTAT- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCTG	G5	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG
G6	TATCASHH- TATCAC- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCTG	G6	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG
G7	TCAASGH- TCAAAG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCTG	G7	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG
G8	TCCTSGH- TCCTGC- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCTG	G8	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG
G9	TCGASBH- TCGATT- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCTG	G9	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG
G10	TCGCSAH- TCGCCA- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCTG	G10	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG
G11	TCGGSBH- TCGGAC- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCTG	G11	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG
G12	TCTCSGH- TCTCGG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCTG	G12	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG
H1	TCTTSBH- TCTTCT- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCTG	H1	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG
H2	TGAASGH- TGAACC- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCTG	H2	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG
H3	TGACASHH- TGACAA- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCTG	H3	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG
H4	TGCCSGH- TGCCCG- top	ACACTCTTTCCCTACACGACGCTCTTCCGATCTG	H4	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG	CTTCTTCGTAAGTACGACGAGGAGAGCGGTTCG

Name			Name		
well	Barcode	Final top sequence	well	(bottom)	Final bottom sequence
H5	TGCTTSAI- TGCTTA- top	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT	H5	TTTGTGTC- TTTGTGTC- bot	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT
H6	TGGGGA- TGGGGA- top	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT	H6	TTTGTGTC- TTTGTGTC- bot	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT
H7	TTATGSAI- TTATGA- top	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT	H7	TTATGSAI- TTATGA- top	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT
H8	TTCCGSAI- TTCCGT- top	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT	H8	TTCCGSAI- TTCCGT- top	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT
H9	TTCTAGSAI- TTCTAG- top	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT	H9	TTCTAGSAI- TTCTAG- top	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT
H10	TTGAGSAI- TTGAGC- top	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT	H10	TTGAGSAI- TTGAGC- top	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT
H11	TTTAASAI- TTTAAT- top	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT	H11	TTTAASAI- TTTAAT- top	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT
H12	TTTGSAI- TTTGTC- top	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT	H12	TTTGSAI- TTTGTC- top	ACACTCTTTCCCTACACGACGAGTTGCTCTTCCGATCTCGAAGAGCGT

B Gel electrophoresis tips

B.1 Introduction

- **Purpose:** To help members of the laboratory run the perfect gels - every time.
- **Procedure Type:** Molecular
- **Species:** N/A
- **Authors**

– Susan Bassham

The effect of electrophoresis is to separate DNA fragments by size in an agarose matrix and buffer using an electrical current. DNA is negatively charged and will migrate toward the positive pole. Many parameters can affect how the DNA moves through the gel: buffer composition, voltage, length of the gel, percentage of agarose in the gel (in other words, the density of the gel matrix), presence of salt in the DNA, protein bound to DNA, and other factors. Some of these conditions also affect which size ranges of DNA will effectively be “resolved” (i.e., separated enough from one another so you can see them as distinct fragment sizes). “Safe-view” is a dye we use in the gel that fluoresces under UV or blue light when bound to DNA, allowing us to see and photograph it when viewed through a special orange filter to cut out the background light. The research goals of using electrophoresis might include:

- measuring the size of fragments in a DNA sample such as the products of PCR or assessing the intactness of the purified DNA from a tissue extraction
- separating fragments from one another so that a particular amplicon or size range of DNA can be purified out of the gel for other downstream applications (such as for cloning a PCR fragment for making probes or for Sanger sequencing, or for size selecting a smear of fragments for making a RAD library) while excluding the other fragment sizes
- checking for the presence or absence of a particular product of PCR as in a screen for orientation of a cloned fragment in a plasmid present in different bacterial colonies, a screen for an insertion in a transgenic animal, a screen to determine the genetic sex or the mitotype of a fish.

B.2 Common Mistakes

Warning

These common mistakes will give you a big headache when trying to run gels

- **Boiling over the agarose when making a gel.**
 - *Result:* failure to monitor the agarose when you are trying to dissolve it in the microwave can easily cause it to boil over because powdered substances create a lot of “nodes of nucleation” for bubbles to form. This is especially easy to accidentally do with higher percentages of agarose. It wastes expensive agarose and creates a mess in the microwave. What remains in the bottle will be of an indeterminate percentage/gel stiffness.
- **Not making sure the tape is adhering well to the mold before casting a gel.**
 - *Result:* liquid agarose will leak out causing a mess and reagent waste. Make sure the mold is dry beforehand, and use the back edge of a comb, e.g., to run across the tape and make sure it has firm contact.
- **Positioning the comb teeth too close to one side or to the bottom of the mold when setting up for pouring a gel.**
 - *Result:* the bottoms (or that side) of wells may be torn when the comb is pulled out, causing the loaded samples to leak out into the buffer during gel loading. Note: even when the comb teeth are not too close to the bottom, low percentage gels are soft and will benefit from having the comb pulled while submerged in the tank - otherwise the suction created by surface tension between the comb plastic and the agarose as you pull up on the comb can tear the bottoms out of the wells, causing loss of samples during loading.
- **Not using the right % of agarose for your DNA size range of interest.**
 - *Result:* bands you care about won't resolve optimally (i.e., you might not be able to accurately measure the sizes of your bands and might not be able to tell if something is one band or multiple bands). Pour a higher percentage gel to resolve large fragments (e.g. less than 1%) or a lower percentage gel (i.e. between 1 and 4%) to resolve small to very small ones, depending on your needs/expectations for what size bands you will see, how many of them, or how important it is to accurately estimate their sizes.
- **Not homogeneously mixing the Safeview into the agarose before pouring.**
 - *Result:* DNA might not be visible in all parts of the gel.

- **Over-cooling the agarose before pouring.**
 - *Result:* Safeview will not be evenly distributed in the gel, and/or the gel have lumps and not be of uniform thickness because some of the agarose will have already started polymerizing.
- **Running a gel with a lot of bubbles in it.**
 - *Result:* DNA migration can be impeded/distorted by bubbles in the gel. Usually as a gel is cooling, bubbles will migrate to the top where (usually) they will be out of the path of the DNA unless the DNA volume fills the wells to the top. But bubbles can sometimes occur deeper in the gel – particularly in a very high % agarose gel. Try to nudge bubbles out of the way before the gel solidifies, either by raking them with a comb that is not on a holder or nudging individual bubbles with a pipette tip before the gel congeals. If the gel is already solid, avoid loading in lanes that will run across a bubble.
- **Running a gel that wasn't mixed homogeneously before pouring in the mold.**
 - *Result:* there will be lumps in the gel of more dense agarose that will cause distortion of how the DNA migrates.
- **Accidentally contacting the agarose with a pipette tip.**
 - *Result:* the bottom or side of the well can become perforated, causing the sample to leak out the bottom or into the next well. If the tip is pressed against the agarose, the sample can be forcefully and suddenly expelled and blast out of the well.
- **Overloading the wells with too much sample volume.**
 - *Result:* samples can become cross-contaminated by DNA from adjacent wells.
- **Overloading the lane with too much DNA.**
 - *Result:* DNA can become retarded during running and the apparent size will not be accurately gauged by the ladder.
- **Forgetting to load a DNA ladder.**
 - *Result:* you won't know if your DNA bands are the right size when you look at and photograph your gel.
- **Not using gel loading mix in the DNA or using ladder that isn't premixed with gel loading mix.**

- *Result:* The DNA or ladder (which is also DNA) will be lost mostly during loading without the gel loading mix that makes it sink to the bottom of the wells in the gel and stay there while you are loading. Both your samples and the ladder are DNA in an aqueous solution that is about the same density as the buffer (usually) – therefore they both should have gel loading mix in them in order to be denser than the tank buffer. The gel loading mix also includes convenient, charged dyes that migrate at different rates to help you see that your gel is running and how far it has progressed. The gel loading mix types only really matter if one type happens to have a dye that migrates exactly like your band of interest - in that case, you may want to switch to a different mix so that the dye doesn't block the fluorescence of your band during visualization.
- **Not thoroughly mixing gel loading mix into your DNA before loading.**
 - *Result:* you could lose part of your DNA to floating out of the well during loading. (This can also happen if there are other reasons your DNA is not very dense, such as if there is residual ethanol in it from preceding processing. In that case, you will see your DNA rapidly floating up to the surface as soon as you start expelling it from the tip).
- **Adding premixed DNA ladder to your samples instead of gel loading mix.**
 - *Result:* ladder will appear in every lane, obscuring your bands and wasting expensive ladder.
- **Forgetting to start the current on your loaded gel.**
 - *Result:* DNA will diffuse both out of the wells and into the gel in all directions causing loss of some of the DNA and blurriness of the remaining DNA once the gel is run. Diffusion can also happen if it takes a long time to load the gel. If there are a great number of samples to load – such as in a two- or three-comb gel, it is often best to load one tier and run the gel for 5 minutes so the DNA enters the gel and diffuses more slowly before moving on to load the next tier. Diffusion can happen more quickly if the buffer is warm from a previous run. If the buffer is warm, replace it before trying to load another gel in the same box.
- **Not keeping track of the order of sample loading.**
 - *Result:* you won't know which lane corresponds to which sample. This can happen, for example, when loading samples from a strip of PCR tubes, where their order can accidentally be rotated 180 degrees, or when loading from a PCR plate, where the plate orientation can be rotated 180 or 90 degrees (e.g., wells are loaded with respect to rows versus columns).

- **Running your gel backwards by accidentally reversing the positive and negative electrodes.**
 - *Result*, your DNA will migrate out of the end of the gel nearest the wells (the “top” of the gel) and be lost into the tank buffer. Always check that your electrodes are hooked up to the correct leads relative to the terminals at the power source, that your gel is oriented so that the DNA will run toward the positive pole. If you have already loaded your gel in the wrong orientation, do not lift the gel out of the tank, but just reverse the leads (red to black) to correct the current. Remember that DNA is negatively charged and will run to the positive (red) pole. When you start the current, the negative electrode (anode) at the well end of the gel should be making noticeably more bubbles than the positive electrode (cathode) at the bottom end of the gel. That’s because electrolysis of water (H₂O) will produce twice as much free hydrogen at the anode than the cathode makes free oxygen. Making a habit of checking that tells you two things: that current is really flowing and that you do have the right orientation of poles.
- **Running a gel at too low a voltage.**
 - *Result*: low molecular weight bands may look fuzzy and faint because they will be diffusing in random directions as they migrate.
- **Running DNA that is too salty - e.g., DNA in Phusion buffer or in NEB restriction buffer 3 (or 3.1) etc.**
 - *Result*: a “salt front” will form where DNA in those salty buffers will be slowed (“retarded”) relative to the ladder, making the ladder a useless measure of the actual size of the DNA. Bands will be compressed at a “front” and will be a weird shape like a smile or a frown rather than a straight band.
 - If you know your DNA is in a salty buffer, you can remove this problem by cleaning the DNA first (i.e., via a cleanup column or with paramagnetic beads), or you can mitigate the problem by diluting only a few microliters (e.g. 5 µl) of your DNA into water and gel loading mix before loading (assuming you have a high enough concentration of DNA that it can still be seen if you load only a small fraction of it).
- **Running a gel made up in a different buffer (or different concentration of buffer) than the buffer in the gel tank; this includes using old buffer that has been evaporating in the tank through multiple runs or over time.**
 - *Result*: bands will not migrate as expected. “Fronts” may form where the leading edge of the migrating DNA is compressed.
- **Losing your gel of the end of the gel mold when you are taking it out of the tank.**

- *Result:* the gel could shatter and be unsalvageable. *Solution:* transport gels in a dish and be especially careful when moving flabby, low percentage agarose gels (under 1%).

B.3 Potentially dangerous and/or destructive mistakes:

Warning

Running gels is so standard in a laboratory that we can take it for granted, but these mistakes can be costly in terms of ruining a gel at best, or posing human safety risks at worst

- **Heating a bottle with a lid on it.**
 - *Result:* possible explosion. Just leave the lid off when heating agarose.
- **Adding Safeview to molten agarose that is too hot.**
 - *Result:* much of the Safeview will be degraded by heat, causing your DNA to be hard to see or image. **The plastic gel mold could be permanently warped if the agarose is too hot.** The tape may fail, causing a mess in the fume hood and reagent wastage. A swirled bottle should be just comfortable (70 degrees C or a bit less).
- **Running a gel at too high a voltage.**
 - *Result:* **buffer could overheat during running and permanently warp the plastic of the gel box.** Bands might look smear and not be well resolved.
- **Dissolving agarose in water instead of electrophoresis buffer or putting water in the electrophoresis tank instead of buffer.**
 - *Result -* total failure for DNA to migrate into the gel, loss of samples. **If there is only water and not buffer in the tank, it could overheat and permanently warp the plastic of the gel box.**
- **Over-running your gel.**
 - *Result:* the DNA might run off the bottom of the gel and be lost. If you had more than one tier of wells in the gel, the DNA in the top tier will run into the zone of the next tier down, causing distortion and blurriness of the bands as they cross the wells of the lower tier and creating confusion in interpretation because very small fragments from the top tier will be overlapping very large fragments of the lower

tier. If the gel runs for a long time, the **buffer could overheat and permanently warp the plastic of the gel box.**

B.4 Conscientiousness

Warning

We all work in our laboratory together. Please think of your labmates when you are done running your gel.

Please really clean the gel molds and combs after use. The next person shouldn't have to fish another lab mate's combs and molds out of the sink and clean and dry them before they can pour their gel. After pulling combs from a polymerized gel, make sure to really rub them under a flow of water; a skin of polymerized agarose left on the combs (higher percentages of agarose are especially prone to this) will mean the next person has to clean this off before they can pour a gel. Likewise, make sure the gel melting bottle is rinsed immediately after you pour your gel. If residual agarose is allowed to dry in the bottle, it can create an almost insoluble blob in the gel of the next person to use the bottle. If you see that some of your agarose has polymerized in the bottle, add water to the bottle and microwave it for a few minutes to dissolve and discard the residue.